
Kan Kajimoto

List of Publications by Year in descending order

Source: https://exaly.com/author-pdf/9086476/publications.pdf Version: 2024-02-01



ΚλΝ ΚΛΙΙΜΟΤΟ

#	Article	IF	CITATIONS
1	Long-term patency of small-diameter vascular graft made from fibroin, a silk-based biodegradable material. Journal of Vascular Surgery, 2010, 51, 155-164.	1.1	197
2	Effectiveness of Statin-Eluting Stent on Early Inflammatory Response and Neointimal Thickness in a Porcine Coronary Model. Circulation Journal, 2008, 72, 832-838.	1.6	53
3	Short-term 20-mg atorvastatin therapy reduces key inflammatory factors including c-Jun N-terminal kinase and dendritic cells and matrix metalloproteinase expression in human abdominal aortic aneurysmal wall. Atherosclerosis, 2009, 206, 505-511.	0.8	52
4	Metabolic Syndrome Predicts 10-Year Mortality in Non-Diabetic Patients Following Coronary Artery Bypass Surgery. Circulation Journal, 2008, 72, 1481-1486.	1.6	43
5	Coronary Artery Bypass Revascularization Using Bilateral Internal Thoracic Arteries in Diabetic Patients: A Systematic Review and Meta-Analysis. Annals of Thoracic Surgery, 2015, 99, 1097-1104.	1.3	42
6	Impact of Metabolic Syndrome on 10-Year Clinical Outcomes Among Patients With Acute Coronary Syndrome. Circulation Journal, 2009, 73, 1454-1458.	1.6	41
7	Prognostic Value of the Metabolic Syndrome for Long-Term Outcomes in Patients Undergoing Percutaneous Coronary Intervention. Circulation Journal, 2006, 70, 1531-1537.	1.6	40
8	Mortality risk of triglyceride levels in patients with coronary artery disease. Heart, 2013, 99, 22-29.	2.9	40
9	Probucol therapy improves long-term (>10-year) survival after complete revascularization: A propensity analysis. Atherosclerosis, 2012, 220, 463-469.	0.8	35
10	Pioglitazone attenuates neointimal thickening via suppression of the early inflammatory response in a porcine coronary after stenting. Atherosclerosis, 2008, 197, 612-619.	0.8	28
11	Impact of Metabolic Syndrome among Patients with and without Diabetes Mellitus on Long-Term Outcomes after Percutaneous Coronary Intervention. Hypertension Research, 2008, 31, 235-241.	2.7	26
12	Metabolic syndrome is an independent risk factor for stroke and acute renal failure after coronary artery bypass grafting. Journal of Thoracic and Cardiovascular Surgery, 2009, 137, 658-663.	0.8	26
13	Influence of diabetes on >10-year outcomes after percutaneous coronary intervention. Heart and Vessels, 2008, 23, 149-154.	1.2	24
14	Prognostic impact of chronic kidney disease on 10-year clinical outcomes among patients with acute coronary syndrome. Journal of Cardiology, 2012, 60, 438-442.	1.9	23
15	Therapy with statins and aspirin enhances long-term outcome of percutaneous coronary intervention. Heart and Vessels, 2008, 23, 35-39.	1.2	18
16	Meta-analysis of Randomized Controlled Trials on the Treatment of Unprotected Left Main Coronary Artery Disease: One-Year Outcomes with Coronary Artery Bypass Grafting Versus Percutaneous Coronary Artery Intervention with Drug-Eluting Stent. Journal of Cardiac Surgery, 2012, 27, 152-157.	0.7	15
17	Comparing outcomes after off-pump coronary artery bypass versus drug-eluting stent in diabetic patients. Journal of Cardiology, 2012, 59, 195-201.	1.9	15
18	Relationship between the metabolic syndrome and the incidence of stroke after complete coronary revascularization over a 10-year follow-up period. Atherosclerosis, 2009, 207, 195-199.	0.8	13

Καν Καιιμότο

#	Article	IF	CITATIONS
19	The adverse prognostic significance of the metabolic syndrome with and without hypertension in patients who underwent complete coronary revascularization. Journal of Hypertension, 2009, 27, 1017-1024.	0.5	12
20	The Impact of Pravastatin Therapy on Long-Term Outcome in Patients With Metabolic Syndrome Undergoing Complete Coronary Revascularization. Circulation Journal, 2009, 73, 2104-2109.	1.6	12
21	Prognostic significance of glomerular filtration rate estimated by the Japanese equation among patients who underwent complete coronary revascularization. Hypertension Research, 2011, 34, 378-383.	2.7	11
22	Non-high-density lipoprotein cholesterol is a practical predictor of long-term cardiac death after coronary artery bypass grafting. Atherosclerosis, 2012, 221, 206-211.	0.8	11
23	Push-Up Technique and Anatomical Deployment With the Endurant Stent-Graft System for Severely Angulated Aneurysm Necks. Journal of Endovascular Therapy, 2017, 24, 435-439.	1.5	9
24	Synergistic Effects of Calcium-Channel and Angiotensin-Receptor Blockers on Endothelial Function and Inflammatory Responses in a Porcine Drug-Eluting Stent Model. Circulation Journal, 2010, 74, 1704-1710.	1.6	8
25	Mid-term Results for the Maze Procedure in Patients with Non-mitral Valvular Atrial Fibrillation. Annals of Thoracic and Cardiovascular Surgery, 2011, 17, 356-362.	0.8	6
26	Association of Low Glomerular Filtration Rate With the Incidence of Stroke in Patients Following Complete Coronary Revascularization. Circulation Journal, 2011, 75, 2372-2378.	1.6	6
27	String-Sign in Left Internal Thoracic Artery Is Associated With Regression in Left Main Trunk Stenosis After Coronary Artery Bypass. International Heart Journal, 2011, 52, 84-87.	1.0	6
28	Long-term outcomes of women with coronary artery disease following complete coronary revascularization. Journal of Cardiology, 2011, 58, 158-164.	1.9	5
29	Is Off-Pump CABG Really Inferior to On-Pump Strategies for Long-Term Survival?. Journal of the American College of Cardiology, 2014, 64, 1181.	2.8	5
30	Dipyridamole Therapy Improves Long-Term Survival After Complete Revascularization in Patients With Impaired Cardiac Function A Propensity Analysis. Circulation Journal, 2008, 72, 1588-1593.	1.6	4
31	Propensity Score Analysis of 10-Year Long-term Outcome After Bypass Surgery or Plain Old Balloon Angioplasty in Patients With Metabolic Syndrome. International Heart Journal, 2011, 52, 372-376.	1.0	4
32	Skeletonization with an ultrasonic scalpel is as safe as a non-skeletonized dissection in preserving the endothelial function of the human gastroepiploic artery. Interactive Cardiovascular and Thoracic Surgery, 2008, 8, 216-220.	1.1	3
33	Skeletonized Gastroepiploic Artery for Off-Pump Coronary Artery Bypass Grafting. Heart Surgery Forum, 2004, 7, E164-E169.	0.5	3
34	Free triiodothyronine (fT3) and B-type natriuretic peptide (BNP) predict in-hospital mortality after valve surgery. General Thoracic and Cardiovascular Surgery, 2020, 68, 585-595.	0.9	2
35	Propensity analysis of 12 years outcome after bypass graft or balloon angioplasty in patients with multivessel coronary artery disease. Journal of Cardiology, 2008, 52, 186-194.	1.9	1
36	Current State-of-the-art of Coronary Artery Bypass Surgery. Journal of the Japanese Coronary Association, 2014, 20, 295-303.	0.0	1

Καν Καιιμότο

#	Article	IF	CITATIONS
37	Independent Effect of Low Flow on Outcomes in Patients Undergoing Aortic Valve Replacement for Severe Aortic Stenosis. Circulation Journal, 2018, 82, 2199-2205.	1.6	1
38	Neointima on the scar site after the left atrial appendage amputation. Journal of Cardiac Surgery, 2019, 34, 855-855.	0.7	1
39	Coronary Artery Bypass Grafting in Patients with Chronic Kidney Disease: Chronic Kidney Disease Has an Independent Adverse Effect on the Long-Term Outcome of Coronary Artery Bypass Grafting. BioMed Research International, 2022, 2022, 1-14.	1.9	1
40	A new technique that prevents paravalvular leakage after aortic valve replacement using a rapidâ€deployment valve system. Journal of Cardiac Surgery, 2021, 36, 2225-2232.	0.7	0