

# William Yeo

## List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/9165279/publications.pdf>

Version: 2024-02-01

43  
papers

1,368  
citations

687363

13  
h-index

345221

36  
g-index

48  
all docs

48  
docs citations

48  
times ranked

1399  
citing authors

#	ARTICLE	IF	CITATIONS
1	Clinical and Radiological Outcomes of Minimally Invasive Versus Open Transforaminal Lumbar Interbody Fusion. <i>Spine</i> , 2009, 34, 1385-1389.	2.0	383
2	Clinical and radiological outcomes of open versus minimally invasive transforaminal lumbar interbody fusion. <i>European Spine Journal</i> , 2012, 21, 2265-2270.	2.2	237
3	Five-Year Outcomes of Minimally Invasive Versus Open Transforaminal Lumbar Interbody Fusion. <i>Spine</i> , 2013, 38, 2049-2055.	2.0	204
4	A proposed set of metrics for standardized outcome reporting in the management of low back pain. <i>Monthly Notices of the Royal Astronomical Society: Letters</i> , 2015, 86, 523-533.	3.3	120
5	Clinical and Radiological Outcomes Comparing Percutaneous Chevron-Akin Osteotomies vs Open Scarf-Akin Osteotomies for Hallux Valgus. <i>Foot and Ankle International</i> , 2018, 39, 311-317.	2.3	69
6	Short-term outcomes of arthroscopic TightRope® fixation are better than hook plate fixation in acute unstable acromioclavicular joint dislocations. <i>European Journal of Orthopaedic Surgery and Traumatology</i> , 2018, 28, 869-875.	1.4	35
7	Intraoperative reduction does not result in better outcomes in low-grade lumbar spondylolisthesis with neurogenic symptoms after minimally invasive transforaminal lumbar interbody fusion—a 5-year follow-up study. <i>Spine Journal</i> , 2016, 16, 182-190.	1.3	31
8	The Association Between Psychological Resilience and Physical Function Among Older Adults With Hip Fracture Surgery. <i>Journal of the American Medical Directors Association</i> , 2020, 21, 260-266.e2.	2.5	30
9	Surgical site infection after primary total knee arthroplasty is associated with a longer duration of surgery. <i>Journal of Orthopaedic Surgery</i> , 2018, 26, 230949901878564.	1.0	27
10	Single-Level Minimally Invasive Transforaminal Lumbar Interbody Fusion Provides Sustained Improvements in Clinical and Radiological Outcomes up to 5 Years Postoperatively in Patients with Neurogenic Symptoms Secondary to Spondylolisthesis. <i>Asian Spine Journal</i> , 2017, 11, 204-212.	2.0	19
11	Elderly Patients Achieving Clinical and Radiological Outcomes Comparable with Those of Younger Patients Following Minimally Invasive Transforaminal Lumbar Interbody Fusion. <i>Asian Spine Journal</i> , 2017, 11, 230-242.	2.0	19
12	Measurement properties of Pain Catastrophizing Scale in patients with knee osteoarthritis. <i>Clinical Rheumatology</i> , 2021, 40, 295-301.	2.2	18
13	Poor Baseline Mental Health Does Not Influence Improvement in Patient-reported Outcomes, Satisfaction, and Return to Work Two Years After Single-level Anterior Cervical Discectomy and Fusion. <i>Spine</i> , 2019, 44, 839-847.	2.0	17
14	Preoperative Leg Pain Score Predicts Patient Satisfaction After Transforaminal Lumbar Interbody Fusion Surgery. <i>Global Spine Journal</i> , 2018, 8, 354-358.	2.3	15
15	The Influence of Body Mass Index on Functional Outcomes, Satisfaction, and Return to Work After Single-level Minimally-invasive Transforaminal Lumbar Interbody Fusion. <i>Spine</i> , 2019, 44, 809-817.	2.0	12
16	Bone Morphogenic Protein Is a Viable Adjunct for Fusion in Minimally Invasive Transforaminal Lumbar Interbody Fusion. <i>Asian Spine Journal</i> , 2016, 10, 1091.	2.0	12
17	The Impact of Diabetes on Patient Outcomes After Total Knee Arthroplasty in an Asian Population. <i>Journal of Arthroplasty</i> , 2018, 33, 3186-3189.	3.1	11
18	Predictors for Rehabilitation Outcome in Asian Geriatric Hip Fracture Patients. <i>Journal of Orthopaedic Surgery</i> , 2016, 24, 153-157.	1.0	10

#	ARTICLE	IF	CITATIONS
19	Association between preoperative anaemia and blood transfusion with long-term functional and quality of life outcomes amongst patients undergoing primary total knee arthroplasty in Singapore: a single-centre retrospective study. <i>Quality of Life Research</i> , 2019, 28, 85-98.	3.1	10
20	Patients With Poor Baseline Mental Health May Experience Significant Improvements in Pain and Disability After Minimally Invasive Transforaminal Lumbar Interbody Fusion. <i>Clinical Spine Surgery</i> , 2020, 33, 205-214.	1.3	9
21	Pre-operative pressure pain thresholds do not meaningfully explain satisfaction or improvement in pain after knee replacement: a cohort study. <i>Osteoarthritis and Cartilage</i> , 2019, 27, 49-58.	1.3	9
22	Time Taken to Return to Work Does Not Influence Outcomes of Minimally Invasive Transforaminal Lumbar Interbody Fusion. <i>Spine</i> , 2019, 44, 503-509.	2.0	8
23	A comparison of contiguous two-level anterior cervical discectomy and fusion using a structural allograft versus a Polyetheretherketone (PEEK) cage: the results of a three-year follow-up. <i>BMC Musculoskeletal Disorders</i> , 2020, 21, 331.	1.9	8
24	Outcomes of arthroscopic capsular release for the diabetic frozen shoulder in Asian patients. <i>Journal of Orthopaedic Surgery</i> , 2018, 26, 230949901875757.	1.0	7
25	Subtalar arthroscopy and fluoroscopy in percutaneous fixation of intra-articular calcaneal fractures. <i>Journal of Orthopaedic Surgery</i> , 2017, 25, 230949901668499.	1.0	6
26	The Effect of Body Mass Index on Long-Term Patient-Reported Outcome Scores after Anterior Cervical Discectomy and Fusion in an Asian Population: A 2-Year Study. <i>Asian Spine Journal</i> , 2021, 15, 512-522.	2.0	6
27	Predictors of Outcomes After Single-level Anterior Cervical Discectomy and Fusion for Cervical Spondylotic Myelopathy. <i>Clinical Spine Surgery</i> , 2020, 33, E525-E532.	1.3	6
28	Associated lumbar scoliosis does not affect outcomes in patients undergoing focal minimally invasive surgery-transforaminal lumbar interbody fusion (MISTLIF) for neurogenic symptoms—a minimum 2-year follow-up study. <i>Spine Journal</i> , 2017, 17, 34-43.	1.3	5
29	Perioperative Factors Influencing Postoperative Satisfaction After Lateral Access Surgery for Degenerative Lumbar Spondylolisthesis. <i>International Journal of Spine Surgery</i> , 2019, 13, 415-422.	1.5	5
30	The Role of Prefracture Health Status in Physical and Mental Function After Hip Fracture Surgery. <i>Journal of the American Medical Directors Association</i> , 2018, 19, 989-994.e2.	2.5	4
31	The Diagnostic Value of Magnetic Resonance Imaging in Identifying Unsuspected Malignancy in Patients Undergoing Percutaneous Vertebral Augmentation for Vertebral Compression Fractures. <i>International Journal of Spine Surgery</i> , 2019, 13, 464-469.	1.5	4
32	Arthroscopic undersurface rotator cuff repair versus conventional arthroscopic double-row rotator cuff repair — Comparable results at 2-year follow-up. <i>Journal of Orthopaedic Surgery</i> , 2018, 26, 230949901875757.	1.0	3
33	Validity and Reliability of the American Orthopaedic Foot and Ankle Society Score for the English-Literate Singapore Population With Hallux Valgus. <i>Foot and Ankle Specialist</i> , 2022, , 193864002210794.	1.0	3
34	Reduction in foraminal height after lateral access surgery does not affect quality of life: A 2-year outcome study on lateral lumbar interbody fusion. <i>Journal of Orthopaedic Surgery</i> , 2019, 27, 230949901982933.	1.0	2
35	A Comparison of Clinical Outcomes of Two Methods of Femoral Hamstring Graft Pin Fixation in Anterior Cruciate Ligament Reconstruction. <i>Proceedings of Singapore Healthcare</i> , 2013, 22, 248-252.	0.6	1
36	AB1324—MEASUREMENT PROPERTIES OF THE 10-ITEM CONNOR-DAVIDSON RESILIENCE SCALE AMONG PATIENTS WITH TOTAL KNEE REPLACEMENT BASED ON ITEM RESPONSE THEORY. , 2019, , .		1

#	ARTICLE	IF	CITATIONS
37	Clinical and radiographic outcomes of computer-navigated total knee arthroplasty are not adversely affected by body mass index. <i>Journal of Orthopaedics</i> , 2020, 19, 54-58.	1.3	1
38	No Difference in Functional Outcome but Higher Revision Rate Among Smokers Undergoing Cervical Artificial Disc Replacement: Analysis of a Spine Registry. <i>International Journal of Spine Surgery</i> , 2020, 14, 916-923.	1.5	1
39	The incidence of infection in routine knee arthroscopy without the use of prophylactic antibiotics is low. <i>Proceedings of Singapore Healthcare</i> , 2019, 28, 110-114.	0.6	0
40	THU0445â€¦ASSOCIATION BETWEEN PATIENTâ€™S EXPECTATION AND SATISFACTION FOLLOWING TOTAL KNEE REPLACEMENT FOR OSTEOARTHRITIS. , 2019, , .		0
41	THU0439â€¦MEASUREMENT PROPERTIES OF PAIN CATASTROPHIZING SCALE IN PATIENTS WITH KNEE OSTEOARTHRITIS. , 2019, , .		0
42	Women Do Not Have Poorer Outcomes After Minimally Invasive Lumbar Fusion Surgery: A Five-Year Follow-Up Study. <i>International Journal of Spine Surgery</i> , 2020, 14, 756-761.	1.5	0
43	Radiofrequency Plantar Fascia Coblation With and Without Gastrocnemius Recession in the Management of Recalcitrant Plantar Fasciitis. <i>Foot and Ankle International</i> , 0, , 107110072210918.	2.3	0