

Shylie Fordyce Mackintosh

List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/9192344/publications.pdf>

Version: 2024-02-01

74
papers

2,489
citations

304743

22
h-index

206112

48
g-index

74
all docs

74
docs citations

74
times ranked

3224
citing authors

#	ARTICLE	IF	CITATIONS
1	Oral Testosterone Supplementation Increases Muscle and Decreases Fat Mass in Healthy Elderly Males With Low-Normal Gonadal Status. <i>Journals of Gerontology - Series A Biological Sciences and Medical Sciences</i> , 2003, 58, M618-M625.	3.6	294
2	Falls after Stroke. <i>International Journal of Stroke</i> , 2012, 7, 482-490.	5.9	244
3	Does the 'Otago exercise programme' reduce mortality and falls in older adults?: a systematic review and meta-analysis. <i>Age and Ageing</i> , 2010, 39, 681-687.	1.6	241
4	Balance Score and a History of Falls in Hospital Predict Recurrent Falls in the 6 Months Following Stroke Rehabilitation. <i>Archives of Physical Medicine and Rehabilitation</i> , 2006, 87, 1583-1589.	0.9	205
5	Falls and injury prevention should be part of every stroke rehabilitation plan. <i>Clinical Rehabilitation</i> , 2005, 19, 441-451.	2.2	166
6	What Works in Falls Prevention After Stroke?. <i>Stroke</i> , 2010, 41, 1715-1722.	2.0	137
7	Aerobic Exercise to Improve Cognitive Function in Adults With Neurological Disorders: A Systematic Review. <i>Archives of Physical Medicine and Rehabilitation</i> , 2011, 92, 1044-1052.	0.9	111
8	Foot orthoses for adults with flexible pes planus: a systematic review. <i>Journal of Foot and Ankle Research</i> , 2014, 7, 23.	1.9	80
9	Paediatric flexible flat foot: how are we measuring it and are we getting it right? A systematic review. <i>Journal of Foot and Ankle Research</i> , 2018, 11, 21.	1.9	78
10	Effects of a Multifactorial Falls Prevention Program for People With Stroke Returning Home After Rehabilitation: A Randomized Controlled Trial. <i>Archives of Physical Medicine and Rehabilitation</i> , 2012, 93, 1648-1655.	0.9	70
11	Differences in foot kinematics between young and older adults during walking. <i>Gait and Posture</i> , 2014, 39, 689-694.	1.4	70
12	Risk Factors for Falls in People With a Lower Limb Amputation: A Systematic Review. <i>PM and R</i> , 2017, 9, 170.	1.6	67
13	Functional Independence Measure. <i>Australian Journal of Physiotherapy</i> , 2009, 55, 65.	0.9	51
14	Consensus-based recommendations of Australian podiatrists for the prescription of foot orthoses for symptomatic flexible pes planus in adults. <i>Journal of Foot and Ankle Research</i> , 2014, 7, 49.	1.9	36
15	Experiences of health care for older people who need support to live at home: A systematic review of the qualitative literature. <i>Geriatric Nursing</i> , 2017, 38, 315-324.	1.9	35
16	TECHNOLOGY IN EDUCATION: WHAT IS APPROPRIATE FOR RURAL AND REMOTE ALLIED HEALTH PROFESSIONALS?. <i>Australian Journal of Rural Health</i> , 1998, 6, 189-193.	1.5	34
17	Fall Recovery Subactivity Recognition With RGB-D Cameras. <i>IEEE Transactions on Industrial Informatics</i> , 2016, 12, 2312-2320.	11.3	34
18	Use of the Theoretical Domains Framework to Develop an Intervention to Improve Physical Therapist Management of the Risk of Falls After Discharge. <i>Physical Therapy</i> , 2014, 94, 1660-1675.	2.4	33

#	ARTICLE	IF	CITATIONS
19	Developing physiotherapy student safety skills in readiness for clinical placement using standardised patients compared with peer-role play: a pilot non-randomised controlled trial. <i>BMC Medical Education</i> , 2017, 17, 133.	2.4	28
20	Access to rehabilitation for patients with stroke in Australia. <i>Medical Journal of Australia</i> , 2019, 210, 21-26.	1.7	28
21	â€œI understand all the major thingsâ€™: how older people with limited English proficiency decide their need for a professional interpreter during health care after stroke. <i>Ethnicity and Health</i> , 2013, 18, 610-625.	2.5	23
22	Exercise class participation among residents in low-level residential aged care could be enhanced: a qualitative study. <i>Australian Journal of Physiotherapy</i> , 2008, 54, 111-117.	0.9	22
23	Qualitative studies using inâ€œdepth interviews with older people from multiple language groups: methodological systematic review. <i>Journal of Advanced Nursing</i> , 2012, 68, 22-35.	3.3	22
24	Walking aid use after discharge following hip fracture is rarely reviewed and often inappropriate: an observational study. <i>Journal of Physiotherapy</i> , 2010, 56, 267-272.	1.7	21
25	Repeatability of stance phase kinematics from a multi-segment foot model in people aged 50 years and older. <i>Gait and Posture</i> , 2013, 38, 349-351.	1.4	21
26	Individual nutrition therapy and exercise regime: A controlled trial of injured, vulnerable elderly (INTERACTIVE trial). <i>BMC Geriatrics</i> , 2008, 8, 4.	2.7	19
27	Altered dynamic foot kinematics in people with medial knee osteoarthritis during walking: A cross-sectional study. <i>Knee</i> , 2014, 21, 1101-1106.	1.6	19
28	Injuries in Young Female Elite Gymnasts. <i>Physiotherapy</i> , 1992, 78, 804-808.	0.4	16
29	Reconsidering inherent requirements: a contribution to the debate from the clinical placement experience of a physiotherapy student with vision impairment. <i>BMC Medical Education</i> , 2016, 16, 74.	2.4	16
30	Determining Current Physical Therapist Management of Hip Fracture in an Acute Care Hospital and Physical Therapists' Rationale for This Management. <i>Physical Therapy</i> , 2011, 91, 1490-1502.	2.4	15
31	The FLASSH study: protocol for a randomised controlled trial evaluating falls prevention after stroke and two sub-studies. <i>BMC Neurology</i> , 2009, 9, 14.	1.8	14
32	Adults with flexible pes planus and the approach to the prescription of customised foot orthoses in clinical practice: A clinical records audit. <i>Foot</i> , 2015, 25, 101-109.	1.1	14
33	Improvements in knee biomechanics during walking are associated with increased physical activity after total knee arthroplasty. <i>Journal of Orthopaedic Research</i> , 2015, 33, 1818-1825.	2.3	13
34	Cost-effectiveness of individualized nutrition and exercise therapy for rehabilitation following hip fracture. <i>Journal of Rehabilitation Medicine</i> , 2016, 48, 378-385.	1.1	13
35	Harnessing group composition-related effects in pain management programs: a review and recommendations. <i>Pain Management</i> , 2016, 6, 161-173.	1.5	12
36	Patient Self-report Measures of Chronic Pain Consultation Measures: A Systematic Review. <i>Clinical Journal of Pain</i> , 2010, 26, 235-243.	1.9	11

#	ARTICLE	IF	CITATIONS
37	Sociodemographic Factors Associated With Self-Reported Exercise and Physical Activity Behaviors and Attitudes of South Australians. <i>Journal of Aging and Health</i> , 2012, 24, 287-306.	1.7	11
38	A balance screening tool for older people: Reliability and validity. <i>International Journal of Therapy and Rehabilitation</i> , 2006, 13, 558-561.	0.3	10
39	Regular group exercise is associated with improved mood but not quality of life following stroke. <i>PeerJ</i> , 2014, 2, e331.	2.0	10
40	Improvement of Physical Therapist Assessment of Risk of Falls in the Hospital and Discharge Handover Through an Intervention to Modify Clinical Behavior. <i>Physical Therapy</i> , 2016, 96, 764-773.	2.4	9
41	Validity study of a novel test protocol for the identification of submaximal muscular effort. <i>Isokinetics and Exercise Science</i> , 1996, 6, 139-144.	0.4	8
42	Balance, fear of falling, pain and gait aid use by low care older people: pilot study. <i>Australasian Journal on Ageing</i> , 2004, 23, 77-81.	0.9	8
43	An isokinetic muscle strengthening program for adults with an acquired brain injury leads to meaningful improvements in physical function. <i>Brain Injury</i> , 2010, 24, 970-977.	1.2	8
44	The effect of limited English proficiency on falls risk and falls prevention after stroke. <i>Age and Ageing</i> , 2012, 41, 104-107.	1.6	8
45	Physical activity habits and preferences in the month prior to a first-ever stroke. <i>PeerJ</i> , 2014, 2, e489.	2.0	8
46	The Experience of Acupuncture Care from the Perspective of People with Chronic Low Back Pain: A Grounded Theory Study. <i>Acupuncture in Medicine</i> , 2014, 32, 333-339.	1.0	7
47	Are group size and composition associated with treatment outcomes in group cognitive behavioural therapy for chronic pain?. <i>Pain</i> , 2018, 159, 783-792.	4.2	7
48	Relationships between adiposity and postural control in girls during balance tasks of varying difficulty. <i>Obesity Research and Clinical Practice</i> , 2019, 13, 358-364.	1.8	7
49	Acupuncturists's perspectives on outcome measures to evaluate acupuncture care for chronic low back pain. <i>Complementary Therapies in Medicine</i> , 2010, 18, 28-41.	2.7	6
50	Isokinetic strength training of lower limb muscles following acquired brain injury. <i>Brain Injury</i> , 2010, 24, 1399-1407.	1.2	6
51	Falls efficacy and fear of falling in stroke: issues with measurement and interpretation. <i>Disability and Rehabilitation</i> , 2012, 34, 704-704.	1.8	6
52	The journey to total hip or knee replacement. <i>Australian Health Review</i> , 2012, 36, 130.	1.1	6
53	Improving mood through physical activity for carers and care recipients (IMPACCT): protocol for a randomised trial. <i>Journal of Physiotherapy</i> , 2013, 59, 125.	1.7	6
54	Visibility and meanings of partnership in health care for older people who need support to live at home. <i>Scandinavian Journal of Caring Sciences</i> , 2018, 32, 1027-1037.	2.1	6

#	ARTICLE	IF	CITATIONS
55	Veterinary Surgeons' Perceptions of Animal Physiotherapy. <i>Physiotherapy</i> , 1993, 79, 312-316.	0.4	5
56	Snakes and ladders: the barriers and facilitators of elective hip- and knee-replacement surgery in Australian public hospitals. <i>Australian Health Review</i> , 2013, 37, 166.	1.1	5
57	Telephone and face-to-face interviews generate similar falls circumstances information from community-dwelling adults with stroke. <i>Australian and New Zealand Journal of Public Health</i> , 2009, 33, 295-296.	1.8	4
58	Effect of Individual Nutrition Therapy and Exercise Regime on Gait Speed, Physical Function, Strength and Balance, Body Composition, Energy and Protein, in Injured, Vulnerable Elderly: A Multisite Randomized Controlled Trial (INTERACTIVE). <i>Nutrients</i> , 2021, 13, 3182.	4.1	4
59	A Survey of Animal Physiotherapy Practice in Britain. <i>Physiotherapy</i> , 1994, 80, 285-289.	0.4	3
60	A comparison of electronic and paper-based clinical skills assessment: Systematic review. <i>Medical Teacher</i> , 2019, 41, 1151-1159.	1.8	3
61	Scope of occupational therapy practice for adults with both Down syndrome and dementia: A cross-sectional survey. <i>Australian Occupational Therapy Journal</i> , 2020, 67, 218-228.	1.1	3
62	Can student-peers teach using simulated-based learning as well as faculty: A non-equivalent posttest-only study. <i>Nurse Education Today</i> , 2020, 91, 104470.	3.3	2
63	Is the psychological composition of the therapeutic group associated with individual outcomes in group cognitive behavioural therapy for chronic pain?. <i>British Journal of Pain</i> , 2021, 15, 69-81.	1.5	2
64	Occupational therapists and physiotherapists weighing up the dignity of risk for people living with a brain injury: grounded theory. <i>Disability and Rehabilitation</i> , 2022, 44, 7145-7151.	1.8	2
65	Hydrotherapy and Tai Chi each provide clinical improvements for older people with osteoarthritis. <i>Australian Journal of Physiotherapy</i> , 2008, 54, 143.	0.9	1
66	Is there a relationship between postural alignment and mobility for adults after acquired brain injury? A systematic review. <i>Brain Injury</i> , 2017, 31, 571-580.	1.2	1
67	Development of autonomy on placement: perceptions of physiotherapy students and educators in Australia and the United Kingdom. <i>Physiotherapy Theory and Practice</i> , 2021, , 1-11.	1.3	1
68	Improving physical mobility is critical for wellbeing in people with severe impairment after an acquired brain injury: a qualitative study. <i>Brain Injury</i> , 2022, 36, 232-238.	1.2	1
69	Improvement in postural alignment is associated with recovery of mobility after complex acquired brain injury: An observational study. <i>Physiotherapy Theory and Practice</i> , 2022, , 1-13.	1.3	1
70	Are group identity and sense of belonging relevant for group pain management programmes? An exploratory pilot study. <i>British Journal of Pain</i> , 2022, 16, 528-537.	1.5	1
71	A multidisciplinary evaluation and management unit reduces mortality in sick, frail older patients. <i>Australian Journal of Physiotherapy</i> , 2002, 48, 321.	0.9	0
72	Expectations Confounded: Patients Are Satisfied with THR and TKR Delivery in Australian Public Hospitals—A Multisite Case Study. , 2012, 2012, 1-6.		0

#	ARTICLE	IF	CITATIONS
73	Changes in the six-minute walk test and foot health status using customised foot orthoses vs sham inserts in an adult flat foot population. A pilot randomised control trial.. The Allied Health Scholar, 2020, 1, 87-115.	0.1	0
74	Development and feasibility testing of an <scp>evidenceâ€based</scp> occupational therapy program for adults with both Down syndrome and dementia. Journal of Policy and Practice in Intellectual Disabilities, 0, , .	2.7	0