

Wei Xu

List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/9196392/publications.pdf>

Version: 2024-02-01

29
papers

384
citations

933447

10
h-index

839539

18
g-index

30
all docs

30
docs citations

30
times ranked

672
citing authors

#	ARTICLE	IF	CITATIONS
1	Prognostic significance of serum immunoglobulin paraprotein in mucosa-associated lymphoid tissue (MALT) lymphoma. <i>British Journal of Haematology</i> , 2022, 196, 1353-1361.	2.5	4
2	Prognostic value of 25-hydroxy vitamin D in extranodal NK/T cell lymphoma. <i>Annals of Hematology</i> , 2021, 100, 445-453.	1.8	9
3	Circulating low CD4+/CD8+ ratio is associated with poor prognosis in Waldenstrom macroglobulinemia patients. <i>Annals of Hematology</i> , 2021, 100, 995-1002.	1.8	5
4	Low serum cholesterol levels predict inferior prognosis and improve prognostic index scoring for peripheral T-cell lymphoma, unspecified. <i>Leukemia Research</i> , 2021, 103, 106534.	0.8	4
5	25-Hydroxy vitamin D deficiency predicts inferior prognosis in Hodgkin lymphoma. <i>Leukemia Research</i> , 2021, 105, 106580.	0.8	6
6	Pretreatment whole blood Epstein-Barr virus DNA predicts prognosis in Hodgkin lymphoma. <i>Leukemia Research</i> , 2021, 107, 106607.	0.8	5
7	Prognostic nutritional index, a novel biomarker which predicts worse prognosis in diffuse large B cell lymphoma. <i>Leukemia Research</i> , 2021, 110, 106664.	0.8	5
8	Circ-RPL15: a plasma circular RNA as novel oncogenic driver to promote progression of chronic lymphocytic leukemia. <i>Leukemia</i> , 2020, 34, 919-923.	7.2	41
9	25-Hydroxy vitamin D deficiency predicts inferior prognosis in mantle cell lymphoma. <i>Journal of Cancer Research and Clinical Oncology</i> , 2020, 146, 1003-1009.	2.5	13
10	98% IGHV gene identity is the optimal cutoff to dichotomize the prognosis of Chinese patients with chronic lymphocytic leukemia. <i>Cancer Medicine</i> , 2020, 9, 999-1007.	2.8	8
11	The prognostic value of 25-hydroxy vitamin D deficiency and its interaction with c-Myc expression in diffuse large B cell lymphoma. <i>Annals of Hematology</i> , 2020, 99, 2377-2384.	1.8	5
12	Interim response in diffuse large B cell lymphoma on CT: what is the optimal size reduction (\hat{I}^{SPD}) for predicting outcome?. <i>European Radiology</i> , 2020, 30, 3094-3100.	4.5	0
13	The impacts of zanubrutinib on immune cells in patients with chronic lymphocytic leukemia/small lymphocytic lymphoma. <i>Hematological Oncology</i> , 2019, 37, 392-400.	1.7	39
14	<p>Diabetes mellitus predicts inferior survival in diffuse large B-cell lymphoma: a propensity score-matched analysis</p>. <i>Cancer Management and Research</i> , 2019, Volume 11, 2849-2870.	1.9	11
15	Association of red blood cell distribution width and outcomes in patients with mantle cell lymphoma. <i>Cancer Medicine</i> , 2019, 8, 2751-2758.	2.8	9
16	A new SH2D1A mutation in a female adult XLP disease with hemophagocytic lymphohistiocytosis and NK-cell leukemia. <i>Annals of Hematology</i> , 2019, 98, 2829-2831.	1.8	5
17	The percentage of cells with 17p deletion and the size of 17p deletion subclones show prognostic significance in chronic lymphocytic leukemia. <i>Genes Chromosomes and Cancer</i> , 2019, 58, 43-51.	2.8	5
18	Circulating tumor DNA: clinical roles in diffuse large B cell lymphoma. <i>Annals of Hematology</i> , 2019, 98, 255-269.	1.8	17

#	ARTICLE	IF	CITATIONS
19	DA-EPOCH-R improves the outcome over that of R-CHOP regimen for DLBCL patients below 60 years, GCB phenotype, and those with high-risk IPI, but not for double expressor lymphoma. <i>Journal of Cancer Research and Clinical Oncology</i> , 2019, 145, 117-127.	2.5	17
20	Low T3 syndrome as a predictor of poor prognosis in chronic lymphocytic leukemia. <i>International Journal of Cancer</i> , 2018, 143, 466-477.	5.1	30
21	NOTCH1 mutation and its prognostic significance in Chinese chronic lymphocytic leukemia: a retrospective study of 317 cases. <i>Cancer Medicine</i> , 2018, 7, 1689-1696.	2.8	7
22	Elevated absolute NK cell counts in peripheral blood predict good prognosis in chronic lymphocytic leukemia. <i>Journal of Cancer Research and Clinical Oncology</i> , 2018, 144, 449-457.	2.5	31
23	Prognostic significance of serum immunoglobulin paraprotein in patients with diffuse large B cell lymphoma. <i>British Journal of Haematology</i> , 2018, 182, 131-134.	2.5	10
24	NDRG2 mRNA levels and miR-28-5p and miR-650 activity in chronic lymphocytic leukemia. <i>BMC Cancer</i> , 2018, 18, 1009.	2.6	15
25	The prognostic role of HBV infection in chronic lymphocytic leukemia. <i>Journal of Cancer Research and Clinical Oncology</i> , 2018, 144, 1309-1315.	2.5	4
26	Low serum cholesterol levels predict inferior prognosis and improve NCCN-IPI scoring in diffuse large B cell lymphoma. <i>International Journal of Cancer</i> , 2018, 143, 1884-1895.	5.1	27
27	Hyperfibrinogenemia is a poor prognostic factor in diffuse large B cell lymphoma. <i>Annals of Hematology</i> , 2018, 97, 1841-1849.	1.8	7
28	Definition of disease progression risk stratification in untreated chronic lymphocytic leukemia using combined clinical, molecular and virological variables. <i>Hematological Oncology</i> , 2018, 36, 656-662.	1.7	2
29	Low T3 syndrome is a strong prognostic predictor in diffuse large B cell lymphoma. <i>British Journal of Haematology</i> , 2017, 177, 95-105.	2.5	43