

Frédéric Gaymard

List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/9229866/publications.pdf>

Version: 2024-02-01

25
papers

3,170
citations

394421

19
h-index

580821

25
g-index

25
all docs

25
docs citations

25
times ranked

2935
citing authors

#	ARTICLE	IF	CITATIONS
1	Non-autonomous stomatal control by pavement cell turgor via the K ⁺ channel subunit <i>AtKC1</i> . <i>Plant Cell</i> , 2022, 34, 2019-2037.	6.6	18
2	B3 Transcription Factors Determine Iron Distribution and FERRITIN Gene Expression in Embryo but Do Not Control Total Seed Iron Content. <i>Frontiers in Plant Science</i> , 2022, 13, .	3.6	5
3	Coumarin accumulation and trafficking in <i>Arabidopsis thaliana</i> : a complex and dynamic process. <i>New Phytologist</i> , 2021, 229, 2062-2079.	7.3	54
4	2000 years of agriculture in the Atacama desert lead to changes in the distribution and concentration of iron in maize. <i>Scientific Reports</i> , 2021, 11, 17322.	3.3	6
5	The Transcription Factor bHLH121 Interacts with bHLH105 (ILR3) and Its Closest Homologs to Regulate Iron Homeostasis in Arabidopsis. <i>Plant Cell</i> , 2020, 32, 508-524.	6.6	111
6	Sulphur availability modulates Arabidopsis thaliana responses to iron deficiency. <i>PLoS ONE</i> , 2020, 15, e0237998.	2.5	16
7	A Global Proteomic Approach Sheds New Light on Potential Iron-Sulfur Client Proteins of the Chloroplastic Maturation Factor NFU3. <i>International Journal of Molecular Sciences</i> , 2020, 21, 8121.	4.1	5
8	Transcriptional Regulation of Iron Distribution in Seeds: A Perspective. <i>Frontiers in Plant Science</i> , 2020, 11, 725.	3.6	6
9	Identification of client iron-sulfur proteins of the chloroplastic NFU2 transfer protein in Arabidopsis thaliana. <i>Journal of Experimental Botany</i> , 2020, 71, 4171-4187.	4.8	25
10	The Transcriptional Control of Iron Homeostasis in Plants: A Tale of bHLH Transcription Factors?. <i>Frontiers in Plant Science</i> , 2019, 10, 6.	3.6	146
11	Iron-sulfur protein NFU2 is required for branched-chain amino acid synthesis in Arabidopsis roots. <i>Journal of Experimental Botany</i> , 2019, 70, 1875-1889.	4.8	25
12	Transcriptional integration of the responses to iron availability in Arabidopsis by the bHLH factor ILR3. <i>New Phytologist</i> , 2019, 223, 1433-1446.	7.3	92
13	Accumulation and Secretion of Coumarinolignans and other Coumarins in Arabidopsis thaliana Roots in Response to Iron Deficiency at High pH. <i>Frontiers in Plant Science</i> , 2016, 7, 1711.	3.6	105
14	Facilitated Fe Nutrition by Phenolic Compounds Excreted by the Arabidopsis ABCG37/PDR9 Transporter Requires the IRT1/FRO2 High-Affinity Root Fe ²⁺ Transport System. <i>Molecular Plant</i> , 2016, 9, 485-488.	8.3	105
15	Iron- and Ferritin-Dependent Reactive Oxygen Species Distribution: Impact on Arabidopsis Root System Architecture. <i>Molecular Plant</i> , 2015, 8, 439-453.	8.3	106
16	Iron nutrition, biomass production, and plant product quality. <i>Trends in Plant Science</i> , 2015, 20, 33-40.	8.8	435
17	Involvement of the ABCG37 transporter in secretion of scopoletin and derivatives by <i>Arabidopsis</i> roots in response to iron deficiency. <i>New Phytologist</i> , 2014, 201, 155-167.	7.3	322
18	Monothiol Glutaredoxin-BolA Interactions: Redox Control of Arabidopsis thaliana BolA2 and SufE1. <i>Molecular Plant</i> , 2014, 7, 187-205.	8.3	70

#	ARTICLE	IF	CITATIONS
19	Regulated expression of Arabidopsis shaker K ⁺ channel genes involved in K ⁺ uptake and distribution in the plant. <i>Plant Molecular Biology</i> , 2003, 51, 773-787.	3.9	221
20	The Arabidopsis outward K ⁺ channel GORK is involved in regulation of stomatal movements and plant transpiration. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 2003, 100, 5549-5554.	7.1	388
21	Guard Cell Inward K ⁺ Channel Activity in Arabidopsis Involves Expression of the Twin Channel Subunits KAT1 and KAT2. <i>Journal of Biological Chemistry</i> , 2001, 276, 3215-3221.	3.4	217
22	Biochemical characterization of the Arabidopsis K ⁺ channels KAT1 and AKT1 expressed or co-expressed in insect cells. <i>Plant Journal</i> , 2000, 23, 527-538.	5.7	39
23	A Shaker-like K ⁺ Channel with Weak Rectification Is Expressed in Both Source and Sink Phloem Tissues of Arabidopsis. <i>Plant Cell</i> , 2000, 12, 837-851.	6.6	196
24	A Shaker-Like K ⁺ Channel with Weak Rectification Is Expressed in Both Source and Sink Phloem Tissues of Arabidopsis. <i>Plant Cell</i> , 2000, 12, 837.	6.6	120
25	Tissue-specific expression of Arabidopsis AKT1 gene is consistent with a role in K ⁺ nutrition. <i>Plant Journal</i> , 1996, 9, 195-203.	5.7	337