

Akiyoshi Uezumi

List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/9237438/publications.pdf>

Version: 2024-02-01

62
papers

5,091
citations

172457

29
h-index

138484

58
g-index

63
all docs

63
docs citations

63
times ranked

5879
citing authors

| # | ARTICLE | IF | CITATIONS |
|----|---|------|-----------|
| 1 | Relayed signaling between mesenchymal progenitors and muscle stem cells ensures adaptive stem cell response to increased mechanical load. <i>Cell Stem Cell</i> , 2022, 29, 265-280.e6. | 11.1 | 36 |
| 2 | Galectin-3 promotes the adipogenic differentiation of PDGFR β ⁺ cells and ectopic fat formation in regenerating muscle. <i>Development (Cambridge)</i> , 2022, 149, . | 2.5 | 5 |
| 3 | Increased MFC β at neuromuscular junctions is an exacerbating factor for sarcopenia-associated denervation. <i>Aging Cell</i> , 2022, 21, e13536. | 6.7 | 7 |
| 4 | Detection of muscle stem cell-derived myonuclei in murine overloaded muscles. <i>STAR Protocols</i> , 2022, 3, 101307. | 1.2 | 1 |
| 5 | Desloratadine inhibits heterotopic ossification by suppression of BMP2 β mad1/5/8 signaling. <i>Journal of Orthopaedic Research</i> , 2021, 39, 1297-1304. | 2.3 | 9 |
| 6 | Retinoic Acid Receptor Agonists Suppress Muscle Fatty Infiltration in Mice. <i>American Journal of Sports Medicine</i> , 2021, 49, 332-339. | 4.2 | 8 |
| 7 | Collagen-VI supplementation by cell transplantation improves muscle regeneration in Ullrich congenital muscular dystrophy model mice. <i>Stem Cell Research and Therapy</i> , 2021, 12, 446. | 5.5 | 11 |
| 8 | Transgenic Expression of Bmp3b in Mesenchymal Progenitors Mitigates Age-Related Muscle Mass Loss and Neuromuscular Junction Degeneration. <i>International Journal of Molecular Sciences</i> , 2021, 22, 10246. | 4.1 | 6 |
| 9 | Liver fibrosis-induced muscle atrophy is mediated by elevated levels of circulating TNF β . <i>Cell Death and Disease</i> , 2021, 12, 11. | 6.3 | 14 |
| 10 | Mesenchymal Bmp3b expression maintains skeletal muscle integrity and decreases in age-related sarcopenia. <i>Journal of Clinical Investigation</i> , 2021, 131, . | 8.2 | 63 |
| 11 | Dlk1 regulates quiescence in calcitonin receptor-mutant muscle stem cells. <i>Stem Cells</i> , 2021, 39, 306-317. | 3.2 | 5 |
| 12 | Measurement of Lateral Transmission of Force in the Extensor Digitorum Longus Muscle of Young and Old Mice. <i>International Journal of Molecular Sciences</i> , 2021, 22, 12356. | 4.1 | 3 |
| 13 | Methods for Accurate Assessment of Myofiber Maturity During Skeletal Muscle Regeneration. <i>Frontiers in Cell and Developmental Biology</i> , 2020, 8, 267. | 3.7 | 42 |
| 14 | Implication of basal lamina dependency in survival of Nrf2-null muscle stem cells via an antioxidative-independent mechanism. <i>Journal of Cellular Physiology</i> , 2019, 234, 1689-1698. | 4.1 | 10 |
| 15 | Tbx1 regulates inherited metabolic and myogenic abilities of progenitor cells derived from slow- and fast-type muscle. <i>Cell Death and Differentiation</i> , 2019, 26, 1024-1036. | 11.2 | 23 |
| 16 | Expression and Functional Analyses of Dlk1 in Muscle Stem Cells and Mesenchymal Progenitors during Muscle Regeneration. <i>International Journal of Molecular Sciences</i> , 2019, 20, 3269. | 4.1 | 11 |
| 17 | Reduced expression of calcitonin receptor is closely associated with age-related loss of the muscle stem cell pool. <i>JCSM Rapid Communications</i> , 2019, 2, 1-13. | 1.6 | 4 |
| 18 | Cell-autonomous and redundant roles of Hey1 and HeyL in muscle stem cells: HeyL requires Hes1 to bind diverse DNA sites. <i>Development (Cambridge)</i> , 2019, 146, . | 2.5 | 34 |

| # | ARTICLE | IF | CITATIONS |
|----|---|------|-----------|
| 19 | The CalcR-PKA-Yap1 Axis Is Critical for Maintaining Quiescence in Muscle Stem Cells. <i>Cell Reports</i> , 2019, 29, 2154-2163.e5. | 6.4 | 38 |
| 20 | <i>Myogenin</i> promoter-associated IncRNA <i>Myoparr</i> is essential for myogenic differentiation. <i>EMBO Reports</i> , 2019, 20, . | 4.5 | 46 |
| 21 | Sustained expression of HeyL is critical for the proliferation of muscle stem cells in overloaded muscle. <i>ELife</i> , 2019, 8, . | 6.0 | 40 |
| 22 | UBL3 modification influences protein sorting to small extracellular vesicles. <i>Nature Communications</i> , 2018, 9, 3936. | 12.8 | 53 |
| 23 | Gm7325 Transcription Is Regulated by MyoD in Activated Muscle Satellite Cells. <i>Biophysical Journal</i> , 2018, 114, 628a. | 0.5 | 0 |
| 24 | Promethazine Hydrochloride Inhibits Ectopic Fat Cell Formation in Skeletal Muscle. <i>American Journal of Pathology</i> , 2017, 187, 2627-2634. | 3.8 | 12 |
| 25 | Disuse Atrophy Accompanied by Intramuscular Ectopic Adipogenesis in Vastus Medialis Muscle of Advanced Osteoarthritis Patients. <i>American Journal of Pathology</i> , 2017, 187, 2674-2685. | 3.8 | 12 |
| 26 | <i>Gm7325</i> is MyoD-dependently expressed in activated muscle satellite cells. <i>Biomedical Research</i> , 2017, 38, 215-219. | 0.9 | 8 |
| 27 | Notch ligands regulate the muscle stem-like state ex vivo but are not sufficient for retaining regenerative capacity. <i>PLoS ONE</i> , 2017, 12, e0177516. | 2.5 | 30 |
| 28 | Cell-Surface Protein Profiling Identifies Distinctive Markers of Progenitor Cells in Human Skeletal Muscle. <i>Stem Cell Reports</i> , 2016, 7, 263-278. | 4.8 | 95 |
| 29 | Identification, Isolation, and Characterization of Mesenchymal Progenitors in Mouse and Human Skeletal Muscle. <i>Methods in Molecular Biology</i> , 2016, 1460, 241-253. | 0.9 | 20 |
| 30 | Toward Regenerative Medicine for Muscular Dystrophies. , 2016, , 103-122. | | 0 |
| 31 | Calcitonin Receptor Signaling Inhibits Muscle Stem Cells from Escaping the Quiescent State and the Niche. <i>Cell Reports</i> , 2015, 13, 302-314. | 6.4 | 88 |
| 32 | Evidence of Notch-Hesr-Nrf2 Axis in Muscle Stem Cells, but Absence of Nrf2 Has No Effect on Their Quiescent and Undifferentiated State. <i>PLoS ONE</i> , 2015, 10, e0138517. | 2.5 | 11 |
| 33 | Doublecortin marks a new population of transiently amplifying muscle progenitor cells and is required for myofiber maturation during skeletal muscle regeneration. <i>Development (Cambridge)</i> , 2015, 142, 51-61. | 2.5 | 29 |
| 34 | Pro-Insulin-Like Growth Factor-II Ameliorates Age-Related Inefficient Regenerative Response by Orchestrating Self-Reinforcement Mechanism of Muscle Regeneration. <i>Stem Cells</i> , 2015, 33, 2456-2468. | 3.2 | 22 |
| 35 | Identification and characterization of PDGFR α ⁺ mesenchymal progenitors in human skeletal muscle. <i>Cell Death and Disease</i> , 2014, 5, e1186-e1186. | 6.3 | 241 |
| 36 | Roles of nonmyogenic mesenchymal progenitors in pathogenesis and regeneration of skeletal muscle. <i>Frontiers in Physiology</i> , 2014, 5, 68. | 2.8 | 114 |

| # | ARTICLE | IF | CITATIONS |
|----|--|------|-----------|
| 37 | Adult stem cell and mesenchymal progenitor theories of aging. <i>Frontiers in Cell and Developmental Biology</i> , 2014, 2, 10. | 3.7 | 37 |
| 38 | Human skeletal muscle-derived PDGFR ^{hi} cells formed heterotopic ossification. <i>Osteoarthritis and Cartilage</i> , 2014, 22, S130-S131. | 1.3 | 1 |
| 39 | Imatinib attenuates severe mouse dystrophy and inhibits proliferation and fibrosis-marker expression in muscle mesenchymal progenitors. <i>Neuromuscular Disorders</i> , 2013, 23, 349-356. | 0.6 | 55 |
| 40 | Osteogenic Differentiation Capacity of Human Skeletal Muscle-Derived Progenitor Cells. <i>PLoS ONE</i> , 2013, 8, e56641. | 2.5 | 75 |
| 41 | Calcitonin receptor and Odz4 are differently expressed in Pax7-positive cells during skeletal muscle regeneration. <i>Journal of Molecular Histology</i> , 2012, 43, 581-587. | 2.2 | 20 |
| 42 | Chondrogenic differentiation potential of CD56+ satellite cell and PDGFR ^{hi} mesenchymal stem cell derived from human skeletal muscle. <i>Osteoarthritis and Cartilage</i> , 2012, 20, S270-S271. | 1.3 | 0 |
| 43 | Fibrosis and adipogenesis originate from a common mesenchymal progenitor in skeletal muscle. <i>Journal of Cell Science</i> , 2011, 124, 3654-3664. | 2.0 | 517 |
| 44 | Hesr1 and Hesr3 are essential to generate undifferentiated quiescent satellite cells and to maintain satellite cell numbers. <i>Development (Cambridge)</i> , 2011, 138, 4609-4619. | 2.5 | 125 |
| 45 | Mesenchymal progenitors distinct from satellite cells contribute to ectopic fat cell formation in skeletal muscle. <i>Nature Cell Biology</i> , 2010, 12, 143-152. | 10.3 | 1,013 |
| 46 | Activin signaling as an emerging target for therapeutic interventions. <i>Cell Communication and Signaling</i> , 2009, 7, 15. | 6.5 | 153 |
| 47 | CD90-positive cells, an additional cell population, produce laminin $\beta 2$ upon transplantation to <i>mdx</i> mice. <i>Experimental Cell Research</i> , 2008, 314, 193-203. | 2.6 | 23 |
| 48 | Suppression of macrophage functions impairs skeletal muscle regeneration with severe fibrosis. <i>Experimental Cell Research</i> , 2008, 314, 3232-3244. | 2.6 | 183 |
| 49 | Transgenic expression of a myostatin inhibitor derived from follistatin increases skeletal muscle mass and ameliorates dystrophic pathology in <i>mdx</i> mice. <i>FASEB Journal</i> , 2008, 22, 477-487. | 0.5 | 171 |
| 50 | Muscle CD31 ^{hi} CD45 ^{hi} Side Population Cells Promote Muscle Regeneration by Stimulating Proliferation and Migration of Myoblasts. <i>American Journal of Pathology</i> , 2008, 173, 781-791. | 3.8 | 75 |
| 51 | Signal Transduction Pathway through Activin Receptors as a Therapeutic Target of Musculoskeletal Diseases and Cancer. <i>Endocrine Journal</i> , 2008, 55, 11-21. | 1.6 | 147 |
| 52 | Autologous Transplantation of SM/C-2.6+ Satellite Cells Transduced with Micro-dystrophin CS1 cDNA by Lentiviral Vector into <i>mdx</i> Mice. <i>Molecular Therapy</i> , 2007, 15, 2178-2185. | 8.2 | 82 |
| 53 | Cardiac side population cells have a potential to migrate and differentiate into cardiomyocytes <i>in vitro</i> and <i>in vivo</i> . <i>Journal of Cell Biology</i> , 2007, 176, 329-341. | 5.2 | 308 |
| 54 | Molecular Signature of Quiescent Satellite Cells in Adult Skeletal Muscle. <i>Stem Cells</i> , 2007, 25, 2448-2459. | 3.2 | 402 |

| # | ARTICLE | IF | CITATIONS |
|----|---|-----|-----------|
| 55 | NO production results in suspension-induced muscle atrophy through dislocation of neuronal NOS. <i>Journal of Clinical Investigation</i> , 2007, 117, 2468-2476. | 8.2 | 157 |
| 56 | Functional heterogeneity of side population cells in skeletal muscle. <i>Biochemical and Biophysical Research Communications</i> , 2006, 341, 864-873. | 2.1 | 110 |
| 57 | Inhibitors of the TGF- β Superfamily and their Clinical Applications. <i>Mini-Reviews in Medicinal Chemistry</i> , 2006, 6, 1255-1261. | 2.4 | 31 |
| 58 | 36. Transplantation of SM/C-2.6+ Satellite Cells Transduced with Micro-Dystrophin CS1 cDNA by Lentiviral Vector into mdx Mice. <i>Molecular Therapy</i> , 2006, 13, S15. | 8.2 | 0 |
| 59 | Participation of Bone Marrow-Derived Cells in Fibrotic Changes in Denervated Skeletal Muscle. <i>American Journal of Pathology</i> , 2005, 166, 1721-1732. | 3.8 | 20 |
| 60 | Mac-1low early myeloid cells in the bone marrow-derived SP fraction migrate into injured skeletal muscle and participate in muscle regeneration. <i>Biochemical and Biophysical Research Communications</i> , 2004, 321, 1050-1061. | 2.1 | 50 |
| 61 | Purification and cell-surface marker characterization of quiescent satellite cells from murine skeletal muscle by a novel monoclonal antibody. <i>Experimental Cell Research</i> , 2004, 296, 245-255. | 2.6 | 179 |
| 62 | Collagen-VI Supplementation by Cell Transplantation Improves Muscle Regeneration in Ullrich Congenital Muscular Dystrophy Model Mice. <i>SSRN Electronic Journal</i> , 0, , . | 0.4 | 0 |