

# Marzia Malcangio

## List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/9238667/publications.pdf>

Version: 2024-02-01

111  
papers

9,979  
citations

26630

56  
h-index

34986

98  
g-index

113  
all docs

113  
docs citations

113  
times ranked

9483  
citing authors

#	ARTICLE	IF	CITATIONS
1	Response to Mylius et al.. Pain, 2022, 163, e495-e495.	4.2	0
2	Pain-resolving microglia. Science, 2022, 376, 33-34.	12.6	9
3	MicroRNAâ€21â€5p functions via RECK/MMP9 as a proalgesic regulator of the blood nerve barrier in nerve injury. Annals of the New York Academy of Sciences, 2022, 1515, 184-195.	3.8	6
4	Changes in bloodâ€spinal cord barrier permeability and neuroimmune interactions in the underlying mechanisms of chronic pain. Pain Reports, 2021, 6, e879.	2.7	20
5	REPRINTED WITH PERMISSION OF IASP â€ PAIN 162 (2021) 999â€1006: Pain in the neurodegenerating brain: insights into pharmacotherapy for Alzheimer disease and Parkinson disease. BÂ³l, 2021, 22, 46-55.	0.1	0
6	Fractalkine/CX3CR1 Pathway in Neuropathic Pain: An Update. Frontiers in Pain Research, 2021, 2, 684684.	2.0	17
7	Microglial heterogeneity in chronic pain. Brain, Behavior, and Immunity, 2021, 96, 279-289.	4.1	24
8	Pain in the neurodegenerating brain: insights into pharmacotherapy for Alzheimer disease and Parkinson disease. Pain, 2021, 162, 999-1006.	4.2	23
9	Changes in vascular permeability in the spinal cord contribute to chemotherapy-induced neuropathic pain. Brain, Behavior, and Immunity, 2020, 83, 248-259.	4.1	26
10	Translational value of preclinical models for rheumatoid arthritis pain. Pain, 2020, 161, 1399-1400.	4.2	3
11	Imbalance of proresolving lipid mediators in persistent allodynia dissociated from signs of clinical arthritis. Pain, 2020, 161, 2155-2166.	4.2	28
12	The Role of Spinal Cord CX3CL1/CX3CR1 Signalling in Chronic Pain. Current Tissue Microenvironment Reports, 2020, 1, 23-29.	3.2	4
13	Cathepsin S as a potential therapeutic target for chronic pain. Medicine in Drug Discovery, 2020, 7, 100047.	4.5	9
14	The role of microRNAs in neurons and neuroimmune communication in the dorsal root ganglia in chronic pain. Neuroscience Letters, 2020, 735, 135230.	2.1	4
15	Cathepsin S acts via protease-activated receptor 2 to activate sensory neurons and induce itch-like behaviour. Neurobiology of Pain (Cambridge, Mass ), 2019, 6, 100032.	2.5	23
16	Role of the immune system in neuropathic pain. Scandinavian Journal of Pain, 2019, 20, 33-37.	1.3	131
17	Pain in Parkinson's disease: new concepts in pathogenesis and treatment. Current Opinion in Neurology, 2019, 32, 579-588.	3.6	61
18	A refined rat primary neonatal microglial culture method that reduces time, cost and animal use. Journal of Neuroscience Methods, 2018, 304, 92-102.	2.5	8

#	ARTICLE	IF	CITATIONS
19	GABAB receptors and pain. <i>Neuropharmacology</i> , 2018, 136, 102-105.	4.1	52
20	Role of TrkA signalling and mast cells in the initiation of osteoarthritis pain in the monoiodoacetate model. <i>Osteoarthritis and Cartilage</i> , 2018, 26, 84-94.	1.3	45
21	A novel interaction between CX3CR1 and CCR2 signalling in monocytes constitutes an underlying mechanism for persistent vincristine-induced pain. <i>Journal of Neuroinflammation</i> , 2018, 15, 101.	7.2	41
22	The therapeutic potential of targeting chemokine signalling in the treatment of chronic pain. <i>Journal of Neurochemistry</i> , 2017, 141, 520-531.	3.9	36
23	Inflammatory pain control by blocking oxidized phospholipid-mediated TRP channel activation. <i>Scientific Reports</i> , 2017, 7, 5447.	3.3	53
24	Spinal mechanisms of neuropathic pain: Is there a P2X4-BDNF controversy?. <i>Neurobiology of Pain (Cambridge, Mass )</i> , 2017, 1, 1-5.	2.5	20
25	Exosomal cargo including microRNA regulates sensory neuron to macrophage communication after nerve trauma. <i>Nature Communications</i> , 2017, 8, 1778.	12.8	224
26	The Therapeutic Potential of Monocyte/Macrophage Manipulation in the Treatment of Chemotherapy-Induced Painful Neuropathy. <i>Frontiers in Molecular Neuroscience</i> , 2017, 10, 397.	2.9	35
27	Microglia and chronic pain. <i>Pain</i> , 2016, 157, 1002-1003.	4.2	14
28	Role of extracellular calcitonin gene-related peptide in spinal cord mechanisms of cancer-induced bone pain. <i>Pain</i> , 2016, 157, 666-676.	4.2	27
29	Selective Cathepsin S Inhibition with MIV-247 Attenuates Mechanical Allodynia and Enhances the Antiallodynic Effects of Gabapentin and Pregabalin in a Mouse Model of Neuropathic Pain. <i>Journal of Pharmacology and Experimental Therapeutics</i> , 2016, 358, 387-396.	2.5	33
30	The Monoiodoacetate Model of Osteoarthritis Pain in the Mouse. <i>Journal of Visualized Experiments</i> , 2016, , .	0.3	81
31	Reduced thermal sensitivity and increased opioidergic tone in the TASTPM mouse model of Alzheimer's disease. <i>Pain</i> , 2016, 157, 2285-2296.	4.2	22
32	Neuron-immune mechanisms contribute to pain in early stages of arthritis. <i>Journal of Neuroinflammation</i> , 2016, 13, 96.	7.2	81
33	Environmental cold exposure increases blood flow and affects pain sensitivity in the knee joints of CFA-induced arthritic mice in a TRPA1-dependent manner. <i>Arthritis Research and Therapy</i> , 2016, 18, 7.	3.5	39
34	Chronic Pain: New Insights in Molecular and Cellular Mechanisms. <i>BioMed Research International</i> , 2015, 2015, 1-2.	1.9	8
35	Calcitonin Gene-Related Peptide-Expressing Sensory Neurons and Spinal Microglial Reactivity Contribute to Pain States in Collagen-Induced Arthritis. <i>Arthritis and Rheumatology</i> , 2015, 67, 1668-1677.	5.6	51
36	The Role of G-Protein Receptor 84 in Experimental Neuropathic Pain. <i>Journal of Neuroscience</i> , 2015, 35, 8959-8969.	3.6	48

#	ARTICLE	IF	CITATIONS
37	Development of monosodium acetate-induced osteoarthritis and inflammatory pain in ageing mice. <i>Age</i> , 2015, 37, 9792.	3.0	22
38	The Role of Glia in the Spinal Cord in Neuropathic and Inflammatory Pain. <i>Handbook of Experimental Pharmacology</i> , 2015, 227, 145-170.	1.8	199
39	Selective Activation of Microglia Facilitates Synaptic Strength. <i>Journal of Neuroscience</i> , 2015, 35, 4552-4570.	3.6	142
40	Fractalkine/CX3CR1 signaling during neuropathic pain. <i>Frontiers in Cellular Neuroscience</i> , 2014, 8, 121.	3.7	122
41	Monocytes expressing CX3CR1 orchestrate the development of vincristine-induced pain. <i>Journal of Clinical Investigation</i> , 2014, 124, 2023-2036.	8.2	140
42	Astrocytesâ€”Multitaskers in chronic pain. <i>European Journal of Pharmacology</i> , 2013, 716, 120-128.	3.5	50
43	Painâ€”like behaviour and spinal changes in the monosodium iodoacetate model of osteoarthritis in <scp>C57Bl</scp>/6 mice. <i>European Journal of Pain</i> , 2013, 17, 514-526.	2.8	77
44	microRNAs in nociceptive circuits as predictors of future clinical applications. <i>Frontiers in Molecular Neuroscience</i> , 2013, 6, 33.	2.9	70
45	Neuropathic pain and cytokines: current perspectives. <i>Journal of Pain Research</i> , 2013, 6, 803.	2.0	244
46	Distinct Nav1.7-dependent pain sensations require different sets of sensory and sympathetic neurons. <i>Nature Communications</i> , 2012, 3, 791.	12.8	228
47	Assessment and treatment of pain in people with dementia. <i>Nature Reviews Neurology</i> , 2012, 8, 264-274.	10.1	270
48	Chemokine mediated neuronâ€”glia communication and aberrant signalling in neuropathic pain states. <i>Current Opinion in Pharmacology</i> , 2012, 12, 67-73.	3.5	93
49	Spinal cathepsin S and fractalkine contribute to chronic pain in the collagenâ€”induced arthritis model. <i>Arthritis and Rheumatism</i> , 2012, 64, 2038-2047.	6.7	74
50	Microglial signalling mechanisms: Cathepsin S and Fractalkine. <i>Experimental Neurology</i> , 2012, 234, 283-292.	4.1	118
51	Fractalkine/CX3CR1 Signalling in Chronic Pain and Inflammation. <i>Current Pharmaceutical Biotechnology</i> , 2011, 12, 1707-1714.	1.6	72
52	Glatiramer acetate attenuates neuropathic allodynia through modulation of adaptive immune cells. <i>Journal of Neuroimmunology</i> , 2011, 234, 19-26.	2.3	40
53	A distinct role for transient receptor potential ankyrin 1, in addition to transient receptor potential vanilloid 1, in tumor necrosis factor $\alpha$ -induced inflammatory hyperalgesia and Freund's complete adjuvant-induced monoarthritis. <i>Arthritis and Rheumatism</i> , 2011, 63, 819-829.	6.7	151
54	Cathepsin S release from primary cultured microglia is regulated by the P2X7 receptor. <i>Glia</i> , 2010, 58, 1710-1726.	4.9	122

#	ARTICLE	IF	CITATIONS
55	Reduced inflammatory and neuropathic pain and decreased spinal microglial response in fractalkine receptor (CX3CR1) knockout mice. <i>Journal of Neurochemistry</i> , 2010, 114, 1143-1157.	3.9	124
56	P2X7-Dependent Release of Interleukin-1 $\beta$ and Nociception in the Spinal Cord following Lipopolysaccharide. <i>Journal of Neuroscience</i> , 2010, 30, 573-582.	3.6	261
57	Systemic blockade of P2X3 and P2X2/3 receptors attenuates bone cancer pain behaviour in rats. <i>Brain</i> , 2010, 133, 2549-2564.	7.6	110
58	Cathepsin S Inhibition Attenuates Neuropathic Pain and Microglial Response Associated with Spinal Cord Injury. <i>Open Pain Journal</i> , 2010, 3, 117-122.	0.4	7
59	The Liberation of Fractalkine in the Dorsal Horn Requires Microglial Cathepsin S. <i>Journal of Neuroscience</i> , 2009, 29, 6945-6954.	3.6	188
60	Chemokines and pain mechanisms. <i>Brain Research Reviews</i> , 2009, 60, 125-134.	9.0	241
61	MAP kinase and pain. <i>Brain Research Reviews</i> , 2009, 60, 135-148.	9.0	872
62	Rapid isolation and culture of primary microglia from adult mouse spinal cord. <i>Journal of Neuroscience Methods</i> , 2009, 183, 223-237.	2.5	36
63	Gabapentin reverses microglial activation in the spinal cord of streptozotocin-induced diabetic rats. <i>European Journal of Pain</i> , 2009, 13, 807-811.	2.8	127
64	Current Challenges in Glia-Pain Biology. <i>Neuron</i> , 2009, 64, 46-54.	8.1	295
65	Hydrogen peroxide is a novel mediator of inflammatory hyperalgesia, acting via transient receptor potential vanilloid 1-dependent and independent mechanisms. <i>Pain</i> , 2009, 141, 135-142.	4.2	93
66	The Cathepsin S/Fractalkine Pair: New Players in Spinal Cord Neuropathic Pain Mechanisms. , 2009, , 455-471.		1
67	Overcoming hERG issues for brain-penetrating cathepsin S inhibitors: 2-Cyanopyrimidines. Part 2. <i>Bioorganic and Medicinal Chemistry Letters</i> , 2008, 18, 5280-5284.	2.2	25
68	Spinal changes associated with mechanical hypersensitivity in a model of Guillain-Barré syndrome. <i>Neuroscience Letters</i> , 2008, 437, 98-102.	2.1	34
69	Discovery of Orally Bioavailable Cathepsin S Inhibitors for the Reversal of Neuropathic Pain. <i>Journal of Medicinal Chemistry</i> , 2008, 51, 5502-5505.	6.4	36
70	Phosphatidylinositol 3-Kinase Is a Key Mediator of Central Sensitization in Painful Inflammatory Conditions. <i>Journal of Neuroscience</i> , 2008, 28, 4261-4270.	3.6	131
71	Specific Antinociceptive Activity of Cholest-4-en-3-one, Oxime (TRO19622) in Experimental Models of Painful Diabetic and Chemotherapy-Induced Neuropathy. <i>Journal of Pharmacology and Experimental Therapeutics</i> , 2008, 326, 623-632.	2.5	65
72	Inhibition of spinal microglial cathepsin S for the reversal of neuropathic pain. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 2007, 104, 10655-10660.	7.1	410

#	ARTICLE	IF	CITATIONS
73	Role of the cysteine protease cathepsin S in neuropathic hyperalgesia. <i>Pain</i> , 2007, 130, 225-234.	4.2	119
74	Role of spinal microglia in rat models of peripheral nerve injury and inflammation. <i>European Journal of Pain</i> , 2007, 11, 223-230.	2.8	213
75	Rapid co-release of interleukin 1 $\beta$ and caspase 1 in spinal cord inflammation. <i>Journal of Neurochemistry</i> , 2006, 99, 868-880.	3.9	97
76	Artemin has potent neurotrophic actions on injured C-fibres. <i>Journal of the Peripheral Nervous System</i> , 2006, 11, 330-345.	3.1	42
77	Brain-derived neurotrophic factor induces NMDA receptor subunit one phosphorylation via ERK and PKC in the rat spinal cord. <i>European Journal of Neuroscience</i> , 2004, 20, 1769-1778.	2.6	138
78	Brain-derived neurotrophic factor as a drug target for CNS disorders. <i>Expert Opinion on Therapeutic Targets</i> , 2004, 8, 391-399.	3.4	114
79	Pain related behaviour in two models of osteoarthritis in the rat knee. <i>Pain</i> , 2004, 112, 83-93.	4.2	356
80	Release of BDNF and GABA in the dorsal horn of neuropathic rats. <i>European Journal of Neuroscience</i> , 2003, 18, 1169-1174.	2.6	132
81	Basal and activity-induced release of substance P from primary afferent fibres in NK1 receptor knockout mice: evidence for negative feedback. <i>Neuropharmacology</i> , 2003, 45, 1101-1110.	4.1	22
82	A common thread for pain and memory synapses? Brain-derived neurotrophic factor and trkB receptors. <i>Trends in Pharmacological Sciences</i> , 2003, 24, 116-121.	8.7	141
83	The signaling components of sensory fiber transmission involved in the activation of ERK MAP kinase in the mouse dorsal horn. <i>Molecular and Cellular Neurosciences</i> , 2003, 24, 259-270.	2.2	74
84	GDNF and somatostatin in sensory neurones. <i>Current Opinion in Pharmacology</i> , 2003, 3, 41-45.	3.5	31
85	Mechanism by which Brain-Derived Neurotrophic Factor Increases Dopamine Release from the Rabbit Retina. , 2003, 44, 791.		26
86	A novel control mechanism based on GDNF modulation of somatostatin release from sensory neurones. <i>FASEB Journal</i> , 2002, 16, 730-732.	0.5	20
87	BDNF Modulates Sensory Neuron Synaptic Activity by a Facilitation of GABA Transmission in the Dorsal Horn. <i>Molecular and Cellular Neurosciences</i> , 2002, 21, 51-62.	2.2	92
88	Noxious Stimulation Induces Trk Receptor and Downstream ERK Phosphorylation in Spinal Dorsal Horn. <i>Molecular and Cellular Neurosciences</i> , 2002, 21, 684-695.	2.2	121
89	BDNF: a neuromodulator in nociceptive pathways?. <i>Brain Research Reviews</i> , 2002, 40, 240-249.	9.0	189
90	Effect of brain-derived neurotrophic factor on the release of substance P from rat spinal cord. <i>NeuroReport</i> , 2001, 12, 21-24.	1.2	16

#	ARTICLE	IF	CITATIONS
91	Brain-Derived Neurotrophic Factor Is Released in the Dorsal Horn by Distinctive Patterns of Afferent Fiber Stimulation. <i>Journal of Neuroscience</i> , 2001, 21, 4469-4477.	3.6	272
92	Intrathecally delivered glial cell line-derived neurotrophic factor produces electrically evoked release of somatostatin in the dorsal horn of the spinal cord. <i>Journal of Neurochemistry</i> , 2001, 78, 221-229.	3.9	22
93	Intrathecally injected neurotrophins and the release of substance P from the rat isolated spinal cord. <i>European Journal of Neuroscience</i> , 2000, 12, 139-144.	2.6	60
94	Abnormal substance P release from the spinal cord following injury to primary sensory neurons. <i>European Journal of Neuroscience</i> , 2000, 12, 397-399.	2.6	95
95	Peptide autoreceptors: does an autoreceptor for substance P exist?. <i>Trends in Pharmacological Sciences</i> , 1999, 20, 405-407.	8.7	59
96	NMDA receptor activation modulates evoked release of substance P from rat spinal cord. <i>British Journal of Pharmacology</i> , 1998, 125, 1625-1626.	5.4	62
97	A pharmacologic analysis of mechanical hyperalgesia in streptozotocin/diabetic rats. <i>Pain</i> , 1998, 76, 151-157.	4.2	207
98	±-Lipoic acid corrects neuropeptide deficits in diabetic rats via induction of trophic support. <i>Neuroscience Letters</i> , 1997, 222, 191-194.	2.1	72
99	Nerve Growth Factor- and Neurotrophin-3-Induced Changes in Nociceptive Threshold and the Release of Substance P from the Rat Isolated Spinal Cord. <i>Journal of Neuroscience</i> , 1997, 17, 8459-8467.	3.6	101
100	Nerve Growth Factor Treatment Increases Stimulus-evoked Release of Sensory Neuropeptides in the Rat Spinal Cord. <i>European Journal of Neuroscience</i> , 1997, 9, 1101-1104.	2.6	62
101	Neurotrophic factorsâ€™ regulation of neuronal phenotype. <i>Neuroscience Research Communications</i> , 1997, 21, 57-66.	0.2	3
102	GABA, glutamate and substance Pâ€™like immunoreactivity release: effects of novel GABA <sub>B</sub> antagonists. <i>British Journal of Pharmacology</i> , 1996, 118, 1153-1160.	5.4	61
103	Calcitonin gene-related peptide content, basal outflow and electrically-evoked release from monoarthritic rat spinal cord in vitro. <i>Pain</i> , 1996, 66, 351-358.	4.2	36
104	Effect of interleukin-1 <sup>2</sup> on the release of substance P from rat isolated spinal cord. <i>European Journal of Pharmacology</i> , 1996, 299, 113-118.	3.5	74
105	Evidence for release of glutamic acid, aspartic acid and substance P but not <sup>3</sup> -aminobutyric acid from primary afferent fibres in rat spinal cord. <i>European Journal of Pharmacology</i> , 1996, 302, 27-36.	3.5	28
106	Possible Therapeutic Application of GABAB Receptor Agonists and Antagonists. <i>Clinical Neuropharmacology</i> , 1995, 18, 285-305.	0.7	40
107	Chronic (-)baclofen or CGP 36742 alters GABAB receptor sensitivity in rat brain and spinal cord. <i>NeuroReport</i> , 1995, 6, 399.	1.2	39
108	Effect of the tachykinin NK <sub>1</sub> receptor antagonists, RP 67580 and SR 140333, on electricallyâ€™evoked substance P release from rat spinal cord. <i>British Journal of Pharmacology</i> , 1994, 113, 635-641.	5.4	33

#	ARTICLE	IF	CITATIONS
109	Spinal cord SP release and hyperalgesia in monoarthritic rats: involvement of the GABA <sub>B</sub> receptor system. <i>British Journal of Pharmacology</i> , 1994, 113, 1561-1566.	5.4	58
110	Plasticity of GABAB receptor in rat spinal cord detected by autoradiography. <i>European Journal of Pharmacology</i> , 1993, 250, 153-156.	3.5	38
111	GABAB receptor-mediated inhibition of forskolin-stimulated cyclic AMP accumulation in rat spinal cord. <i>Neuroscience Letters</i> , 1993, 158, 189-192.	2.1	17