## Sallie E Lamb

List of Publications by Year in descending order

Source: https://exaly.com/author-pdf/9264147/publications.pdf Version: 2024-02-01



| #  | Article   | IF  | CITATIONS |
|----|---|-----|-----------|
| 1  | Components, design and effectiveness of digital physical rehabilitation interventions for older people: A systematic review. Journal of Telemedicine and Telecare, 2022, 28, 162-176.   | 2.7 | 8         |
| 2  | The Effect of Therapeutic Exercise Interventions on Physical and Psychosocial Outcomes in Adults<br>Aged 80 Years and Older: A Systematic Review and Meta-Analysis. Journal of Aging and Physical Activity,<br>2022, 30, 517-534.   | 1.0 | 2         |
| 3  | Surgical fixation with K-wires versus casting in adults with fracture of distal radius: DRAFFT2 multicentre randomised clinical trial. BMJ, The, 2022, 376, e068041.  | 6.0 | 8         |
| 4  | Effects of a Home-Based Physical Training and Activity Promotion Program in Community-Dwelling<br>Older Persons with Cognitive Impairment after Discharge from Rehabilitation: A Randomized<br>Controlled Trial. Journals of Gerontology - Series A Biological Sciences and Medical Sciences, 2022,<br>77, 2435-2444. | 3.6 | 5         |
| 5  | Effect of an osteoporotic fracture prevention program on fracture incidence in routine care: a cluster-randomized trial. BMC Medicine, 2022, 20, 49.  | 5.5 | 4         |
| 6  | Moulded cast compared with K-wire fixation after manipulation of an acute dorsally displaced distal radius fracture: the DRAFFT 2 RCT. Health Technology Assessment, 2022, 26, 1-80.  | 2.8 | 2         |
| 7  | Flexibility and resistance exercises versus usual care for improving pain and function after distal radius fracture in adults aged 50 years or over: protocol for the WISE randomised multicentre feasibility trial. Pilot and Feasibility Studies, 2022, 8, 55.  | 1.2 | 2         |
| 8  | SWAT 110: Printing the primary outcomE on Pink PapER versus standard paper to increase participant engagement to postal questionnaires (PEPPER). Research Methods in Medicine & Health Sciences, 2022, 3, 49-54.  | 1.2 | 0         |
| 9  | The Clinical Effectiveness of a Physiotherapy Delivered Physical and Psychological Group Intervention for Older Adults With Neurogenic Claudication: The BOOST Randomized Controlled Trial. Journals of Gerontology - Series A Biological Sciences and Medical Sciences, 2022, 77, 1654-1664.                         | 3.6 | 9         |
| 10 | Efficacy of Liposomal Bupivacaine and Bupivacaine Hydrochloride vs Bupivacaine Hydrochloride Alone<br>as a Periarticular Anesthetic for Patients Undergoing Knee Replacement. JAMA Surgery, 2022, 157, 481.   | 4.3 | 12        |
| 11 | Exercise to prevent shoulder problems after breast cancer surgery: the PROSPER RCT. Health Technology Assessment, 2022, 26, 1-124.  | 2.8 | 9         |
| 12 | P135 Lower limb muscle strength and balance in older adults with distal radius fracture: a systematic review. Rheumatology, 2022, 61, .   | 1.9 | 0         |
| 13 | Anti-tumour necrosis factor therapy for early-stage Dupuytren's disease (RIDD): a phase 2b,<br>randomised, double-blind, placebo-controlled trial. Lancet Rheumatology, The, 2022, 4, e407-e416.  | 3.9 | 10        |
| 14 | Understanding prescribed dose in hand strengthening exercise for rheumatoid arthritis: A secondary<br>analysis of the SARAH trial. Musculoskeletal Care, 2022, 20, 899-907.   | 1.4 | 2         |
| 15 | The effectiveness of rehabilitation interventions including outdoor mobility on older adults'<br>physical activity, endurance, outdoor mobility and falls-related self-efficacy: systematic review and<br>meta-analysis. Age and Ageing, 2022, 51, .  | 1.6 | 10        |
| 16 | Nonlinear Dynamic Measures of Walking in Healthy Older Adults: A Systematic Scoping Review.<br>Sensors, 2022, 22, 4408.   | 3.8 | 3         |
| 17 | Interventions for preventing falls in older people living in the community. The Cochrane Library, 2021, 2021, CD007146.   | 2.8 | 1,914     |
| 18 | Exercise Effects on Falls, Fractures, Hospitalizations, and Mortality in Older Adults With Dementia:<br>An Individual-Level Patient Data Meta-analysis. Journals of Gerontology - Series A Biological Sciences<br>and Medical Sciences, 2021, 76, e203-e212.  | 3.6 | 8         |

| #  | Article  | IF  | CITATIONS |
|----|--|-----|-----------|
| 19 | ltems for consideration in a reporting guideline for mediation analyses: a Delphi study. BMJ<br>Evidence-Based Medicine, 2021, 26, 106-106.  | 3.5 | 8         |
| 20 | Characteristics of L-PRP preparations for treating Achilles tendon rupture within the PATH-2 study.<br>Platelets, 2021, 32, 273-279.   | 2.3 | 8         |
| 21 | The findings of a surgical hip fracture trial were generalizable to the UK national hip fracture database. Journal of Clinical Epidemiology, 2021, 131, 141-151.   | 5.0 | 3         |
| 22 | The evidence for cognitive behavioural therapy in any condition, population or context: a meta-review of systematic reviews and panoramic meta-analysis. Psychological Medicine, 2021, 51, 21-29.  | 4.5 | 46        |
| 23 | Identification of subgroup effect with an individual participant data meta-analysis of randomised controlled trials of three different types of therapist-delivered care in low back pain. BMC Musculoskeletal Disorders, 2021, 22, 191.                           | 1.9 | 4         |
| 24 | Cognitive–behavioural therapy for a variety of conditions: an overview of systematic reviews and panoramic meta-analysis. Health Technology Assessment, 2021, 25, 1-378.   | 2.8 | 22        |
| 25 | The Severe Asthma Questionnaire: sensitivity to change and minimal clinically important difference.<br>European Respiratory Journal, 2021, 57, 2100300.  | 6.7 | 5         |
| 26 | Factors associated with persistent opioid use after an upper extremity fracture. Bone & Joint Open, 2021, 2, 119-124.  | 2.6 | 8         |
| 27 | Monitoring, implementation and reporting of interventions in a selection of trials assessing exercise therapy for the management of shoulder subacromial pain: a cross-sectional investigation. BMJ Open, 2021, 11, e044462.                                       | 1.9 | 3         |
| 28 | Disentangling contextual effects from musculoskeletal treatments. Osteoarthritis and Cartilage, 2021, 29, 297-299.   | 1.3 | 23        |
| 29 | A hand exercise mobile app for people with rheumatoid arthritis in Turkey: design, development and usability study. Rheumatology International, 2021, 41, 1151-1160.   | 3.0 | 5         |
| 30 | Adrenaline to improve survival in out-of-hospital cardiac arrest: the PARAMEDIC2 RCT. Health Technology Assessment, 2021, 25, 1-166.   | 2.8 | 6         |
| 31 | Psychometric Properties of the Proxy-Reported Life-Space Assessment in Institutionalized Settings<br>(LSA-IS-Proxy) for Older Persons with and without Cognitive Impairment. International Journal of<br>Environmental Research and Public Health, 2021, 18, 3872. | 2.6 | 5         |
| 32 | Fall prevention interventions in primary care to reduce fractures and falls in people aged 70 years and over: the PreFIT three-arm cluster RCT. Health Technology Assessment, 2021, 25, 1-114.   | 2.8 | 12        |
| 33 | Hand Osteoarthritis: investigating Pain Effects of estrogen-containing therapy (HOPE-e): a protocol for a feasibility randomised placebo-controlled trial. Pilot and Feasibility Studies, 2021, 7, 133.  | 1.2 | 1         |
| 34 | Effectiveness of community-based rehabilitation interventions incorporating outdoor mobility on ambulatory ability and falls-related self-efficacy after hip fracture: a systematic review and meta-analysis. Archives of Osteoporosis, 2021, 16, 99.              | 2.4 | 3         |
| 35 | Getting Recovery Right After Neck Dissection (GRRAND-F): mixed-methods feasibility study to design a pragmatic randomised controlled trial protocol. BMJ Open, 2021, 11, e045741.  | 1.9 | 2         |
| 36 | Rehabilitation to enable recovery from COVID-19: a rapid systematic review. Physiotherapy, 2021, 111, 4-22.  | 0.4 | 48        |

| #  | Article  | IF   | CITATIONS |
|----|--|------|-----------|
| 37 | Experiences and perceptions of trial participants and healthcare professionals in the UK Frozen Shoulder Trial (UK FROST): a nested qualitative study. BMJ Open, 2021, 11, e040829.  | 1.9  | О         |
| 38 | Statistical analysis plan for a pragmatic phase III randomised controlled trial examining behaviour change physiotherapy intervention to increase physical activity following hip and knee replacements: the PEP-TALK trial. Trials, 2021, 22, 467.                                  | 1.6  | 1         |
| 39 | Progressive exercise compared with best practice advice, with or without corticosteroid injection,<br>for the treatment of patients with rotator cuff disorders (CRASP): a multicentre, pragmatic, 2 × 2<br>factorial, randomised controlled trial. Lancet, The, 2021, 398, 416-428. | 13.7 | 53        |
| 40 | Home environmental assessments and modification delivered by occupational therapists to reduce falls in people aged 65 years and over: the OTIS RCT. Health Technology Assessment, 2021, 25, 1-118.  | 2.8  | 8         |
| 41 | Measuring the success of blinding in placebo-controlled trials: Should we be so quick to dismiss it?.<br>Journal of Clinical Epidemiology, 2021, 135, 176-181.   | 5.0  | 12        |
| 42 | Progressive exercise compared with best-practice advice, with or without corticosteroid injection, for rotator cuff disorders: the GRASP factorial RCT. Health Technology Assessment, 2021, 25, 1-158.   | 2.8  | 10        |
| 43 | Home-based rehabilitation programme compared with traditional physiotherapy for patients at risk of poor outcome after knee arthroplasty: the CORKA randomised controlled trial. BMJ Open, 2021, 11, e052598.  | 1.9  | 11        |
| 44 | A Guideline for Reporting Mediation Analyses of Randomized Trials and Observational Studies. JAMA -<br>Journal of the American Medical Association, 2021, 326, 1045.   | 7.4  | 169       |
| 45 | Research Note: Transporting causal effects from randomised trials to target populations to improve external validity. Journal of Physiotherapy, 2021, 67, 315-318.   | 1.7  | Ο         |
| 46 | Effectiveness of interventions for the management of primary frozen shoulder. Bone & Joint Open, 2021, 2, 773-784.   | 2.6  | 12        |
| 47 | Risk Factors for Mobility Decline in Community-Dwelling Older Adults: A Systematic Literature<br>Review. Journal of Aging and Physical Activity, 2021, 29, 1053-1066.  | 1.0  | 7         |
| 48 | Multi-Factorial Fall Prevention Strategies: Where to Next?. , 2021, , 386-395.   |      | 0         |
| 49 | The Prevalence of Back and Leg Pain and the Cross-sectional Association With Adverse Health<br>Outcomes in Community Dwelling Older Adults in England. Spine, 2021, 46, 54-61.   | 2.0  | 7         |
| 50 | Unicompartmental compared with total knee replacement for patients with multimorbidities: a cohort study using propensity score stratification and inverse probability weighting. Health Technology Assessment, 2021, 25, 1-126.   | 2.8  | 6         |
| 51 | Exercise versus usual care after non-reconstructive breast cancer surgery (UK PROSPER): multicentre randomised controlled trial and economic evaluation. BMJ, The, 2021, 375, e066542.   | 6.0  | 18        |
| 52 | A feasibility study to assess the design of a multicentre randomized controlled trial of the clinical and cost-effectiveness of a caregiving intervention for people following hip fracture surgery. Bone & Joint Open, 2021, 2, 909-920.  | 2.6  | 5         |
| 53 | Evaluation of Clinical Practice Guidelines on Fall Prevention and Management for Older Adults. JAMA<br>Network Open, 2021, 4, e2138911.  | 5.9  | 121       |
| 54 | Multifactorial interventions for preventing falls in older people living in the community: a systematic<br>review and meta-analysis of 41 trials and almost 20 000 participants. British Journal of Sports<br>Medicine, 2020, 54, 1340-1350.   | 6.7  | 77        |

| #  | Article   | IF   | CITATIONS |
|----|---|------|-----------|
| 55 | Development and implementation of the physiotherapy-led exercise interventions for the treatment of rotator cuff disorders for the †Getting it Right: Addressing Shoulder Pain' (GRASP) trial. Physiotherapy, 2020, 107, 252-266.   | 0.4  | 8         |
| 56 | Physiotherapy for primary frozen shoulder in secondary care: Developing and implementing stand-alone and post operative protocols for UK FROST and inferences for wider practice. Physiotherapy, 2020, 107, 150-160.  | 0.4  | 17        |
| 57 | Physiotherapy rehabilitation for osteoporotic vertebral fracture—a randomised controlled trial and economic evaluation (PROVE trial). Osteoporosis International, 2020, 31, 277-289.  | 3.1  | 19        |
| 58 | Exercise for preventing falls in older people living in the community: an abridged Cochrane systematic review. British Journal of Sports Medicine, 2020, 54, 885-891.   | 6.7  | 155       |
| 59 | Management of adults with primary frozen shoulder in secondary care (UK FROST): a multicentre, pragmatic, three-arm, superiority randomised clinical trial. Lancet, The, 2020, 396, 977-989.  | 13.7 | 97        |
| 60 | Translating the Strengthening and Stretching for Rheumatoid Arthritis of the Hand Programme from clinical trial to clinical practice: An effectiveness–implementation study. Hand Therapy, 2020, 25, 87-97.   | 1.4  | 2         |
| 61 | Behaviour change physiotherapy intervention to increase physical activity following hip and knee<br>replacement (PEP-TALK): study protocol for a pragmatic randomised controlled trial. BMJ Open, 2020,<br>10, e035014.   | 1.9  | 6         |
| 62 | Better Outcomes for Older people with Spinal Trouble (BOOST) trial: statistical analysis plan for a randomised controlled trial of a combined physical and psychological intervention for older adults with neurogenic claudication. Trials, 2020, 21, 667.   | 1.6  | 2         |
| 63 | Evaluation of the implementation of the Back Skills Training (BeST) programme using online training: a cohort implementation study. Physiotherapy, 2020, 109, 4-12.   | 0.4  | 2         |
| 64 | Cost-effectiveness of the PDSAFE personalised physiotherapy intervention for fall prevention in<br>Parkinson's: an economic evaluation alongside a randomised controlled trial. BMC Neurology, 2020,<br>20, 295.  | 1.8  | 6         |
| 65 | Understanding the mechanisms of a combined physical and psychological intervention for people with neurogenic claudication: protocol for a causal mediation analysis of the BOOST trial. BMJ Open, 2020, 10, e037121.   | 1.9  | 1         |
| 66 | Cost-effectiveness of adrenaline for out-of-hospital cardiac arrest. Critical Care, 2020, 24, 579.  | 5.8  | 15        |
| 67 | Cohort profile: Oxford Pain, Activity and Lifestyle (OPAL) Study, a prospective cohort study of older<br>adults in England. BMJ Open, 2020, 10, e037516.  | 1.9  | 8         |
| 68 | Physical Activity and Exercise in Mild Cognitive Impairment and Dementia: An Umbrella Review of<br>Intervention and Observational Studies. Journal of the American Medical Directors Association, 2020,<br>21, 1415-1422.e6.  | 2.5  | 97        |
| 69 | TIDieR-Placebo: A guide and checklist for reporting placebo and sham controls. PLoS Medicine, 2020, 17, e1003294.   | 8.4  | 52        |
| 70 | Early Psychological and Social Factors Explain the Recovery Trajectory After Distal Radial Fracture.<br>Journal of Bone and Joint Surgery - Series A, 2020, 102, 788-795.   | 3.0  | 48        |
| 71 | Progressive exercise compared with best practice advice, with or without corticosteroid injection, for the treatment of rotator cuff disorders: statistical analysis plan for the Getting it Right:<br>Addressing Shoulder Pain (GRASP) 2 × 2 factorial multicentre randomised controlled trial. Trials, 2020, 21, 767. | 1.6  | 2         |
| 72 | Synergistic effects of hip/knee osteoarthritis and comorbidities on mobility and self-care limitations among older adults: Cross-sectional analysis of the Oxford pain, Activity and Lifestyle study. Journal of Comorbidity, 2020, 10, 2235042X2097452.  | 3.9  | 1         |

Sallie E Lamb

| #  | Article   | IF               | CITATIONS         |
|----|---|------------------|-------------------|
| 73 | Will We Do If We Can? Habitual Qualitative and Quantitative Physical Activity in Multi-Morbid, Older<br>Persons with Cognitive Impairment. Sensors, 2020, 20, 7208.   | 3.8              | 1                 |
| 74 | Musculoskeletal pain and loneliness, social support and social engagement among older adults:<br>Analysis of the Oxford Pain, Activity and Lifestyle cohort. Musculoskeletal Care, 2020, 19, 269-277.   | 1.4              | 12                |
| 75 | The online version of an evidence-based hand exercise program for people with rheumatoid arthritis:<br>A mixed-method, proof-of-concept study. Journal of Hand Therapy, 2020, , .   | 1.5              | 7                 |
| 76 | Screening and Intervention to Prevent Falls and Fractures in Older People. New England Journal of Medicine, 2020, 383, 1848-1859.   | 27.0             | 74                |
| 77 | Comparison of the clinical and cost effectiveness of two management strategies (rehabilitation) Tj ETQq1 1 0.784 protocol for the ACL SNNAP randomised controlled trial. Trials, 2020, 21, 405.   | ·314 rgBT<br>1.6 | /Overlock 1<br>20 |
| 78 | The impact of a patient's engagement in their health on the magnitude of limitations and experience following upper limb fractures. Bone and Joint Journal, 2020, 102-B, 42-47.   | 4.4              | 21                |
| 79 | Participant expectations and experiences of a tailored physiotherapy intervention for people with Parkinson's and a history of falls. Disability and Rehabilitation, 2020, , 1-9.   | 1.8              | 6                 |
| 80 | Exercise for people with a fragility fracture of the pelvis or lower limb: a systematic review of interventions evaluated in clinical trials and reporting quality. BMC Musculoskeletal Disorders, 2020, 21, 435.                                       | 1.9              | 6                 |
| 81 | Plaster cast versus functional brace for non-surgical treatment of Achilles tendon rupture (UKSTAR):<br>a multicentre randomised controlled trial and economic evaluation. Lancet, The, 2020, 395, 441-448.   | 13.7             | 41                |
| 82 | Enhanced rehabilitation and care models for adults with dementia following hip fracture surgery.<br>The Cochrane Library, 2020, 2020, CD010569.   | 2.8              | 28                |
| 83 | Development of A Guideline for Reporting Mediation Analyses (AGReMA). BMC Medical Research<br>Methodology, 2020, 20, 19.  | 3.1              | 11                |
| 84 | Is Protocolised Weaning that Includes Early Extubation Onto Non-Invasive Ventilation More Cost<br>Effective Than Protocolised Weaning Without Non-Invasive Ventilation? Findings from the Breathe<br>Study. PharmacoEconomics - Open, 2020, 4, 697-710. | 1.8              | 3                 |
| 85 | The importance of different frailty domains in a population based sample in England. BMC Geriatrics, 2020, 20, 16.  | 2.7              | 12                |
| 86 | Bayesian adaptive designs for multi-arm trials: an orthopaedic case study. Trials, 2020, 21, 83.  | 1.6              | 17                |
| 87 | Plaster cast versus functional bracing for Achilles tendon rupture: the UKSTAR RCT. Health<br>Technology Assessment, 2020, 24, 1-86.  | 2.8              | 10                |
| 88 | Surgical treatments compared with early structured physiotherapy in secondary care for adults with primary frozen shoulder: the UK FROST three-arm RCT. Health Technology Assessment, 2020, 24, 1-162.  | 2.8              | 11                |
| 89 | Moderators of Cognitive Outcomes from an Exercise Program in People with Mild to Moderate<br>Dementia. Journal of the American Geriatrics Society, 2020, 68, 2095-2100.   | 2.6              | 1                 |
| 90 | Outpatient physiotherapy versus home-based rehabilitation for patients at risk of poor outcomes<br>after knee arthroplasty: CORKA RCT. Health Technology Assessment, 2020, 24, 1-116.   | 2.8              | 62                |

| #   | Article  | IF  | CITATIONS |
|-----|--|-----|-----------|
| 91  | Evaluating the impact of a champion on implementation of the Back Skills Training (BeST) programme<br>in Canada: a mixed methods feasibility study protocol. BMJ Open, 2020, 10, e040834.                                | 1.9 | 0         |
| 92  | The different influence of high levels of physical activity on the incidence of knee OA in overweight<br>and obese men and women-a gender specific analysis. Osteoarthritis and Cartilage, 2019, 27, 1430-1436.          | 1.3 | 9         |
| 93  | Psychosocial factors affecting variation in patient-reported outcomes after elbow fractures. Journal of Shoulder and Elbow Surgery, 2019, 28, 1431-1440.   | 2.6 | 24        |
| 94  | Caregiver Interventions for Adults Discharged from the Hospital: Systematic Review and Metaâ€Analysis.<br>Journal of the American Geriatrics Society, 2019, 67, 1960-1969.   | 2.6 | 19        |
| 95  | Inadequate description of placebo and sham controls in a systematicÂreview of recent trials. European<br>Journal of Clinical Investigation, 2019, 49, e13169.  | 3.4 | 11        |
| 96  | Investigating causal mechanisms in randomised controlled trials. Trials, 2019, 20, 524.  | 1.6 | 25        |
| 97  | Comparison of alternative falls data collection methods in the Prevention of Falls Injury Trial<br>(PreFIT). Journal of Clinical Epidemiology, 2019, 106, 32-40.   | 5.0 | 21        |
| 98  | Using Bayesian adaptive designs to improve phase III trials: a respiratory care example. BMC Medical<br>Research Methodology, 2019, 19, 99.  | 3.1 | 27        |
| 99  | Surgical fixation with K-wires versus plaster casting in the treatment of dorsally displaced distal radius fractures: protocol for Distal Radius Acute Fracture Fixation Trial 2 (DRAFFT 2). BMJ Open, 2019, 9, e028474. | 1.9 | 10        |
| 100 | Caregiver-Delivered Stroke Rehabilitation in Rural China. Stroke, 2019, 50, 1825-1830.   | 2.0 | 51        |
| 101 | An overview of systematic reviews found suboptimal reporting and methodological limitations of mediation studies investigating causal mechanisms. Journal of Clinical Epidemiology, 2019, 111, 60-68.e1.                 | 5.0 | 23        |
| 102 | Effectiveness of scoliosis-specific exercises for adolescent idiopathic scoliosis compared with other non-surgical interventions: a systematic review and meta-analysis. Physiotherapy, 2019, 105, 214-234.              | 0.4 | 28        |
| 103 | Development and delivery of the BOOST (Better Outcomes for Older adults with Spinal Trouble) intervention for older adults with neurogenic claudication. Physiotherapy, 2019, 105, 262-274.                              | 0.4 | 8         |
| 104 | Duration of Treatment Effect Should Be Considered in the Design and Interpretation of Clinical<br>Trials: Results of a Discrete Choice Experiment. Medical Decision Making, 2019, 39, 461-473.                           | 2.4 | 9         |
| 105 | Multicentre, randomised controlled trial of PDSAFE, a physiotherapist-delivered fall prevention<br>programme for people with Parkinson's. Journal of Neurology, Neurosurgery and Psychiatry, 2019, 90,<br>774-782.       | 1.9 | 56        |
| 106 | 7â€Platelet rich plasma for acute achilles tendon rupture: a double-blind, multicentre, randomised, placebo-controlled trial. , 2019, , .  |     | 0         |
| 107 | Platelet rich plasma injection for acute Achilles tendon rupture: PATH-2 randomised, placebo controlled, superiority trial. BMJ: British Medical Journal, 2019, 367, l6132.  | 2.3 | 51        |
| 108 | Exercise for preventing falls in older people living in the community. The Cochrane Library, 2019, 2019, CD012424.   | 2.8 | 529       |

| #   | Article   | IF  | CITATIONS |
|-----|---|-----|-----------|
| 109 | Short and long-term clinical effectiveness and cost-effectiveness of a late-phase community-based<br>balance and gait exercise program following hip fracture. The EVA-Hip Randomised Controlled Trial.<br>PLoS ONE, 2019, 14, e0224971.  | 2.5 | 25        |
| 110 | Progressive functional exercise versus best practice advice for adults aged 50 years or over after ankle fracture: protocol for a pilot randomised controlled trial in the UK - the Ankle Fracture Treatment: Enhancing Rehabilitation (AFTER) study. BMJ Open, 2019, 9, e030877. | 1.9 | 8         |
| 111 | Construct Validity and Precision of Different Patient-reported Outcome Measures During Recovery<br>After Upper Extremity Fractures. Clinical Orthopaedics and Related Research, 2019, 477, 2521-2530.   | 1.5 | 20        |
| 112 | Study of Peri-Articular Anaesthetic for Replacement of the Knee (SPAARK): study protocol for a patient-blinded, randomised controlled superiority trial of liposomal bupivacaine. Trials, 2019, 20, 732.  | 1.6 | 2         |
| 113 | Relationship Between Magnitude of Limitations and Patient Experience During Recovery from<br>Upper-Extremity Fracture. JBJS Open Access, 2019, 4, e0002.  | 1.5 | 9         |
| 114 | Does Structured Exercise Improve Cognitive Impairment in People with Mild to Moderate Dementia? A<br>Cost-Effectiveness Analysis from a Confirmatory Randomised Controlled Trial: The Dementia and<br>Physical Activity (DAPA) Trial. PharmacoEconomics - Open, 2019, 3, 215-227. | 1.8 | 11        |
| 115 | On Being (Not) Old: Agency, Selfâ€care, and Lifeâ€course Aspirations in the United States. Medical<br>Anthropology Quarterly, 2019, 33, 263-281.  | 1.4 | 15        |
| 116 | Process evaluation of complex interventions tested in randomised controlled trials in musculoskeletal disorders: a systematic review protocol. BMJ Open, 2019, 9, e028160.  | 1.9 | 5         |
| 117 | Predicting patient-reported and objectively measured functional outcome 6 months after ankle<br>fracture in people aged 60 years or over in the UK: prognostic model development and internal<br>validation. BMJ Open, 2019, 9, e029813.  | 1.9 | 6         |
| 118 | Platelet-rich plasma injection for adults with acute Achilles tendon rupture: the PATH-2 RCT. Efficacy and Mechanism Evaluation, 2019, 6, 1-98.   | 0.7 | 8         |
| 119 | Exercise- and strategy-based physiotherapy-delivered intervention for preventing repeat falls in people with Parkinson's: the PDSAFE RCT. Health Technology Assessment, 2019, 23, 1-150.  | 2.8 | 44        |
| 120 | Exercise or manual physiotherapy compared with a single session of physiotherapy for osteoporotic vertebral fracture: three-arm PROVE RCT. Health Technology Assessment, 2019, 23, 1-318.   | 2.8 | 13        |
| 121 | Protocolised non-invasive compared with invasive weaning from mechanical ventilation for adults in intensive care: the Breathe RCT. Health Technology Assessment, 2019, 23, 1-114.  | 2.8 | 4         |
| 122 | Treatment of first-time traumatic anterior shoulder dislocation: the UK TASH-D cohort study. Health<br>Technology Assessment, 2019, 23, 1-104.  | 2.8 | 12        |
| 123 | Current UK Practices on Health Economics Analysis Plans (HEAPs): Are We Using Heaps of Them?.<br>Pharmacoeconomics, 2018, 36, 253-257.  | 3.3 | 11        |
| 124 | Reproducible and replicable pain research: a critical review. Pain, 2018, 159, 1683-1689.   | 4.2 | 80        |
| 125 | Three-Year Follow-up of a Trial of Close Contact Casting vs Surgery for Initial Treatment of Unstable<br>Ankle Fractures in Older Adults. JAMA - Journal of the American Medical Association, 2018, 319, 1274.  | 7.4 | 19        |
| 126 | Randomised controlled trial of exercise to prevent shoulder problems in women undergoing breast<br>cancer treatment: study protocol for the prevention of shoulder problems trial (UK PROSPER). BMJ<br>Open, 2018, 8, e019078.  | 1.9 | 22        |

| #   | Article   | IF  | CITATIONS |
|-----|---|-----|-----------|
| 127 | Physiotherapist-delivered cognitive-behavioural interventions are effective for low back pain, but can<br>they be replicated in clinical practice? A systematic review. Disability and Rehabilitation, 2018, 40, 1-9.   | 1.8 | 105       |
| 128 | Exploring physiotherapists' experiences of implementing a cognitive behavioural approach for<br>managing low back pain and identifying barriers to long-term implementation. Physiotherapy, 2018, 104,<br>107-115.  | 0.4 | 26        |
| 129 | Development and delivery of an exercise programme for falls prevention: the Prevention of Falls<br>Injury Trial (PreFIT). Physiotherapy, 2018, 104, 72-79.  | 0.4 | 7         |
| 130 | Cost-Effectiveness of the Pdsafe Home-Based Personalized Physiotherapy Intervention to Prevent Falls<br>Among People with Parkinson's: an Economic Evaluation Alongside a Randomised Controlled Trial.<br>Value in Health, 2018, 21, S93-S94.                             | 0.3 | 0         |
| 131 | Development and prospective external validation of a tool to predict poor recovery at 9 months after<br>acute ankle sprain in UK emergency departments: the SPRAINED prognostic model. BMJ Open, 2018, 8,<br>e022802.   | 1.9 | 5         |
| 132 | Effectiveness of cognitive–behavioural therapy: a protocol for an overview of systematic reviews and meta-analyses. BMJ Open, 2018, 8, e025761.   | 1.9 | 13        |
| 133 | Sample size calculations are poorly conducted and reported in manyÂrandomized trials of hip and knee<br>osteoarthritis: results of aÂsystematic review. Journal of Clinical Epidemiology, 2018, 104, 52-61.   | 5.0 | 22        |
| 134 | Better Outcomes for Older people with Spinal Trouble (BOOST) Trial: a randomised controlled trial of a combined physical and psychological intervention for older adults with neurogenic claudication, a protocol. BMJ Open, 2018, 8, e022205.                            | 1.9 | 16        |
| 135 | What Factors Are Associated With Disability After Upper Extremity Injuries? A Systematic Review.<br>Clinical Orthopaedics and Related Research, 2018, 476, 2190-2215.   | 1.5 | 92        |
| 136 | Effect of Protocolized Weaning With Early Extubation to Noninvasive Ventilation vs Invasive Weaning<br>on Time to Liberation From Mechanical Ventilation Among Patients With Respiratory Failure. JAMA -<br>Journal of the American Medical Association, 2018, 320, 1881. | 7.4 | 68        |
| 137 | Platelet-rich plasma in Achilles tendon healing 2 (PATH-2) trial: statistical analysis plan for a<br>multicentre, double-blinded, parallel-group, placebo-controlled randomised clinical trial. Trials, 2018,<br>19, 464.   | 1.6 | 4         |
| 138 | Can occupational therapist-led home environmental assessment prevent falls in older people? A modified cohort randomised controlled trial protocol. BMJ Open, 2018, 8, e022488.   | 1.9 | 18        |
| 139 | Predictors of falls and fractures leading to hospitalization in people with schizophrenia spectrum disorder: A large representative cohort study. Schizophrenia Research, 2018, 201, 70-78.   | 2.0 | 27        |
| 140 | Dementia And Physical Activity (DAPA) trial of moderate to high intensity exercise training for people with dementia: randomised controlled trial. BMJ: British Medical Journal, 2018, 361, k1675.  | 2.3 | 258       |
| 141 | Cost-Effectiveness of a Multifaceted Podiatry Intervention for the Prevention of Falls in Older<br>People: The REducing Falls with Orthoses and a Multifaceted Podiatry Intervention Trial Findings.<br>Gerontology, 2018, 64, 503-512.                                   | 2.8 | 10        |
| 142 | Out-of-hours discharge from intensive care, in-hospital mortality and intensive care readmission rates: a systematic review and meta-analysis. Intensive Care Medicine, 2018, 44, 1115-1129.  | 8.2 | 47        |
| 143 | The differential influence of high levels of physical activity on knee OA in overweight and obese men<br>and women. Osteoarthritis and Cartilage, 2018, 26, S11-S12.  | 1.3 | 1         |
| 144 | Participation in a trial in the emergency situation: a qualitative study of patient experience in the UK<br>WOLLF trial. Trials, 2018, 19, 328.   | 1.6 | 10        |

| #   | Article   | IF  | CITATIONS |
|-----|---|-----|-----------|
| 145 | Multifactorial and multiple component interventions for preventing falls in older people living in the community. The Cochrane Library, 2018, 2018, CD012221.   | 2.8 | 234       |
| 146 | Development of an exercise intervention for the prevention of musculoskeletal shoulder problems<br>after breast cancer treatment: the prevention of shoulder problems trial (UK PROSPER). BMC Health<br>Services Research, 2018, 18, 463. | 2.2 | 22        |
| 147 | Exercise for rheumatoid arthritis of the hand. The Cochrane Library, 2018, 2018, CD003832.  | 2.8 | 24        |
| 148 | Anti-Tumour Necrosis Factor Therapy for Dupuytren's Disease: A Randomised Dose Response Proof of<br>Concept Phase 2a Clinical Trial. EBioMedicine, 2018, 33, 282-288.   | 6.1 | 39        |
| 149 | Predictors of Falls and Fractures Leading to Hospitalization in People With Dementia: A Representative<br>Cohort Study. Journal of the American Medical Directors Association, 2018, 19, 607-612.   | 2.5 | 46        |
| 150 | O32 Impact evaluation of an online training on the Strengthening And Stretching for Rheumatoid<br>Arthritis of the Hand (SARAH) programme recommended in the NICE guidelines for rheumatoid<br>arthritis. Rheumatology, 2018, 57, .       | 1.9 | 0         |
| 151 | Effect of Negative Pressure Wound Therapy vs Standard Wound Management on 12-Month Disability<br>Among Adults With Severe Open Fracture of the Lower Limb. JAMA - Journal of the American Medical<br>Association, 2018, 319, 2280.        | 7.4 | 137       |
| 152 | A study update newsletter or Post-it® note did not increase postal questionnaire response rates in a falls prevention trial: an embedded randomised factorial trial. F1000Research, 2018, 7, 1083.  | 1.6 | 3         |
| 153 | An Online Hand Exercise Intervention for Adults With Rheumatoid Arthritis (mySARAH): Design,<br>Development, and Usability Testing. Journal of Medical Internet Research, 2018, 20, e10457.   | 4.3 | 20        |
| 154 | Intramedullary nail fixation versus locking plate fixation for adults with a fracture of the distal tibia: the UK FixDT RCT. Health Technology Assessment, 2018, 22, 1-148.   | 2.8 | 16        |
| 155 | Aerobic and strength training exercise programme for cognitive impairment in people with mild to moderate dementia: the DAPA RCT. Health Technology Assessment, 2018, 22, 1-202.  | 2.8 | 37        |
| 156 | Negative-pressure wound therapy versus standard dressings for adults with an open lower limb fracture: the WOLLF RCT. Health Technology Assessment, 2018, 22, 1-162.  | 2.8 | 42        |
| 157 | A nested randomised controlled trial of a newsletter and Post-it® note did not increase postal questionnaire response rates in a falls prevention trial. F1000Research, 2018, 7, 1083.  | 1.6 | 3         |
| 158 | Prognostic models for identifying risk of poor outcome in people with acute ankle sprains: the SPRAINED development and external validation study. Health Technology Assessment, 2018, 22, 1-112.   | 2.8 | 124       |
| 159 | Is Vitamin D Supplementation Effective for Low Back Pain? A Systematic Review and Meta-Analysis. Pain<br>Physician, 2018, 21, 121-145.  | 0.4 | 13        |
| 160 | A qualitative study of older people's experience of living with neurogenic claudication to inform the development of a physiotherapy intervention. Disability and Rehabilitation, 2017, 39, 1009-1017.                                    | 1.8 | 13        |
| 161 | Relationship Between Low Bone Mineral Density and Fractures With Incident Cardiovascular Disease:<br>A Systematic Review and Meta-Analysis. Journal of Bone and Mineral Research, 2017, 32, 1126-1135.                                    | 2.8 | 109       |
| 162 | Severity of Hypoxemia and Effect of High-Frequency Oscillatory Ventilation in Acute Respiratory<br>Distress Syndrome. American Journal of Respiratory and Critical Care Medicine, 2017, 196, 727-733.                                     | 5.6 | 82        |

| #   | Article   | IF  | CITATIONS |
|-----|---|-----|-----------|
| 163 | Hand exercises for patients with rheumatoid arthritis: an extended follow-up of the SARAH randomised controlled trial. BMJ Open, 2017, 7, e013121.  | 1.9 | 19        |
| 164 | The design and development of a complex multifactorial falls assessment intervention for falls prevention: The Prevention of Falls Injury Trial (PreFIT). BMC Geriatrics, 2017, 17, 116.  | 2.7 | 16        |
| 165 | Explaining How Cognitive Behavioral Approaches Work for Low Back Pain. Spine, 2017, 42, E1031-E1039.  | 2.0 | 37        |
| 166 | Can Mapping Algorithms Based on Raw Scores Overestimate QALYs Gained by Treatment? A Comparison of Mappings Between the Roland–Morris Disability Questionnaire and the EQ-5D-3L Based on Raw and Differenced Score Data. Pharmacoeconomics, 2017, 35, 549-559.  | 3.3 | 5         |
| 167 | Clinical and cost-effectiveness of progressive exercise compared with best practice advice, with or without corticosteroid injection, for the treatment of rotator cuff disorders: protocol for a 2x2 factorial randomised controlled trial (the GRASP trial). BMJ Open, 2017, 7, e018004.                      | 1.9 | 20        |
| 168 | Intrarater reliability and agreement of linear encoder derived heel-rise endurance test outcome measures in healthy adults. Journal of Electromyography and Kinesiology, 2017, 36, 34-39.   | 1.7 | 17        |
| 169 | Measuring physical capacity and performance in older people. Best Practice and Research in Clinical Rheumatology, 2017, 31, 243-254.  | 3.3 | 22        |
| 170 | Incidence of shoulder dislocations in the UK, 1995–2015: a population-based cohort study. BMJ Open, 2017, 7, e016112.   | 1.9 | 70        |
| 171 | Platelet rich Plasma in Achilles Tendon Healing 2 (PATH-2) trial: protocol for a multicentre,<br>participant and assessor-blinded, parallel-group randomised clinical trial comparing platelet-rich<br>plasma (PRP) injection versus placebo injection for Achilles tendon rupture. BMJ Open, 2017, 7, e018135. | 1.9 | 16        |
| 172 | Effect of Locking Plate Fixation vs Intramedullary Nail Fixation on 6-Month Disability Among Adults<br>With Displaced Fracture of the Distal Tibia. JAMA - Journal of the American Medical Association, 2017,<br>318, 1767.   | 7.4 | 55        |
| 173 | Cast versus functional brace in the rehabilitation of patients treated non-operatively for a rupture of the Achilles tendon: protocol for the UK study of tendo achilles rehabilitation (UK STAR) multi-centre randomised trial. BMJ Open, 2017, 7, e019628.  | 1.9 | 6         |
| 174 | An empirical evaluation of the SF-12, SF-6D, EQ-5D and Michigan Hand Outcome Questionnaire in patients with rheumatoid arthritis of the hand. Health and Quality of Life Outcomes, 2017, 15, 20.  | 2.4 | 50        |
| 175 | An optimised patient information sheet did not significantly increase recruitment or retention in a falls prevention study: an embedded randomised recruitment trial. Trials, 2017, 18, 144.  | 1.6 | 30        |
| 176 | A longitudinal, qualitative study exploring sustained adherence to a hand exercise programme for rheumatoid arthritis evaluated in the SARAH trial. Disability and Rehabilitation, 2017, 39, 1856-1863.   | 1.8 | 12        |
| 177 | Mediating Effect of Changes in Hand Impairments on Hand Function in Patients With Rheumatoid<br>Arthritis: Exploring the Mechanisms of an Effective Exercise Program. Arthritis Care and Research,<br>2017, 69, 982-988.  | 3.4 | 8         |
| 178 | Advancing Physical Therapist Interventions by Investigating Causal Mechanisms. Physical Therapy, 2017, 97, 1119-1121.   | 2.4 | 13        |
| 179 | Cohort Randomised Controlled Trial of a Multifaceted Podiatry Intervention for the Prevention of Falls in Older People (The REFORM Trial). PLoS ONE, 2017, 12, e0168712.  | 2.5 | 37        |
| 180 | Core outcome measures for interventions to prevent or slow the progress of dementia for people<br>living with mild to moderate dementia: Systematic review and consensus recommendations. PLoS ONE,<br>2017, 12, e0179521.  | 2.5 | 48        |

| #   | Article   | IF                | CITATIONS      |
|-----|---|-------------------|----------------|
| 181 | Prognostic factors for recovery following acute lateral ankle ligament sprain: a systematic review.<br>BMC Musculoskeletal Disorders, 2017, 18, 421.  | 1.9               | 34             |
| 182 | A systematic review and meta-analysis of online versus alternative methods for training licensed health care professionals to deliver clinical interventions. BMC Medical Education, 2017, 17, 227.   | 2.4               | 74             |
| 183 | Current practice in methodology and reporting of the sample size calculation in randomised trials of hip and knee osteoarthritis: a protocol for a systematic review. Trials, 2017, 18, 466.  | 1.6               | 4              |
| 184 | United Kingdom Frozen Shoulder Trial (UK FROST), multi-centre, randomised, 12 month, parallel group,<br>superiority study to compare the clinical and cost-effectiveness of Early Structured Physiotherapy<br>versus manipulation under anaesthesia versus arthroscopic capsular release for patients referred to<br>secondary care with a primary frozen shoulder: study protocol for a randomised controlled trial.<br>Trials, 2017, 18, 614. | 1.6               | 25             |
| 185 | A Systematic Review of Outcome Measures Assessing Disability Following Upper Extremity Trauma.<br>Journal of the American Academy of Orthopaedic Surgeons Global Research and Reviews, 2017, 1, e021.   | 0.7               | 15             |
| 186 | Study protocol: A multi-centre, double blind, randomised, placebo-controlled, parallel group, phase II<br>trial (RIDD) to determine the efficacy of intra-nodular injection of anti-TNF to control disease<br>progression in early Dupuytren's disease, with an embedded dose response study Wellcome Open<br>Research, 2017, 2, 37.  | 1.8               | 13             |
| 187 | Study protocol: A multi-centre, double blind, randomised, placebo-controlled, parallel group, phase II<br>trial (RIDD) to determine the efficacy of intra-nodular injection of anti-TNF to control disease<br>progression in early Dupuytren's disease, with an embedded dose response study Wellcome Open<br>Research, 2017, 2, 37.  | 1.8               | 15             |
| 188 | A Web-Based Training Resource for Therapists to Deliver an Evidence-Based Exercise Program for<br>Rheumatoid Arthritis of the Hand (iSARAH): Design, Development, and Usability Testing. Journal of<br>Medical Internet Research, 2017, 19, e411.   | 4.3               | 13             |
| 189 | Prehospital randomised assessment of a mechanical compression device in out-of-hospital cardiac arrest (PARAMEDIC): a pragmatic, cluster randomised trial and economic evaluation. Health Technology Assessment, 2017, 21, 1-176.   | 2.8               | 28             |
| 190 | Clinical effectiveness and cost-effectiveness of a multifaceted podiatry intervention for falls prevention in older people: a multicentre cohort randomised controlled trial (the REducing Falls) Tj ETQq0 0 0 rg   | BT /Overlo<br>2.8 | ock 10 Tf 50 3 |
| 191 | 21, 1-198.<br>Development of a core outcome set for disease modification trials in mild to moderate dementia: a systematic review, patient and public consultation and consensus recommendations. Health Technology Assessment, 2017, 21, 1-192.  | 2.8               | 37             |
| 192 | Mapping the Association between Vitamin D and Low Back Pain: A Systematic Review and Meta-Analysis<br>of Observational Studies. Pain Physician, 2017, 20, 611-640.  | 0.4               | 20             |
| 193 | Economic and Health-Related Quality of Life Outcomes of Whiplash Associated Disorders. Spine, 2016, 41, 1378-1386.  | 2.0               | 21             |
| 194 | COmmunity-based Rehabilitation after Knee Arthroplasty (CORKA): study protocol for a randomised controlled trial. Trials, 2016, 17, 501.  | 1.6               | 7              |
| 195 | Using mixed methods evaluation to assess the feasibility of online clinical training in evidence based interventions: a case study of cognitive behavioural treatment for low back pain. BMC Medical Education, 2016, 16, 163.  | 2.4               | 20             |
| 196 | A cluster randomised controlled trial of advice, exercise or multifactorial assessment to prevent falls and fractures in community-dwelling older adults: protocol for the prevention of falls injury trial (PreFIT). BMJ Open, 2016, 6, e009362.   | 1.9               | 24             |
| 197 | One-year resource utilisation, costs and quality of life in patients with acute respiratory distress<br>syndrome (ARDS): secondary analysis of a randomised controlled trial. Journal of Intensive Care, 2016,<br>4, 56.  | 2.9               | 40             |
| 198 | Close Contact Casting vs Surgery for Initial Treatment of Unstable Ankle Fractures in Older Adults.<br>JAMA - Journal of the American Medical Association, 2016, 316, 1455.   | 7.4               | 94             |

| #   | Article   | IF   | CITATIONS |
|-----|---|------|-----------|
| 199 | A randomized controlled trial on rehabilitation through caregiver-delivered nurse-organized service<br>programs for disabled stroke patients in rural china (the RECOVER trial): design and rationale.<br>International Journal of Stroke, 2016, 11, 823-830.   | 5.9  | 17        |
| 200 | What Value Can Qualitative Research Add to Quantitative Research Design? An Example From an<br>Adolescent Idiopathic Scoliosis Trial Feasibility Study. Qualitative Health Research, 2016, 26, 1838-1850.   | 2.1  | 20        |
| 201 | Pre-hospital Assessment of the Role of Adrenaline: Measuring the Effectiveness of Drug<br>administration In Cardiac arrest (PARAMEDIC-2): Trial protocol. Resuscitation, 2016, 108, 75-81.  | 3.0  | 43        |
| 202 | Dementia and Physical Activity (DAPA) - an exercise intervention to improve cognition in people with mild to moderate dementia: study protocol for a randomized controlled trial. Trials, 2016, 17, 165.  | 1.6  | 19        |
| 203 | Appraising the uptake and use of recommendations for a common outcome data set for clinical trials:<br>a case study in fall injury prevention. Trials, 2016, 17, 131.   | 1.6  | 23        |
| 204 | The application of multilevel modelling to account for the influence of walking speed in gait analysis.<br>Gait and Posture, 2016, 43, 216-219.   | 1.4  | 12        |
| 205 | Ageing, Muscle Power and Physical Function: A Systematic Review and Implications for Pragmatic Training Interventions. Sports Medicine, 2016, 46, 1311-1332.  | 6.5  | 157       |
| 206 | The Immediate Effects of Different Types of Ankle Support Introduced 6 Weeks After Surgical Internal<br>Fixation for Ankle Fracture on Gait and Pain: A Randomized Crossover Trial. Journal of Orthopaedic<br>and Sports Physical Therapy, 2016, 46, 157-167.   | 3.5  | 2         |
| 207 | The Ankle Injury Management (AIM) trial: a pragmatic, multicentre, equivalence randomised controlled trial and economic evaluation comparing close contact casting with open surgical reduction and internal fixation in the treatment of unstable ankle fractures in patients aged over 60 years. Health Technology Assessment. 2016. 20. 1-158. | 2.8  | 73        |
| 208 | Identifying back pain subgroups: developing and applying approaches using individual patient data collected within clinical trials. Programme Grants for Applied Research, 2016, 4, 1-278.  | 1.0  | 9         |
| 209 | Protocol for a randomised controlled trial of standard wound management versus negative pressure wound therapy in the treatment of adult patients with an open fracture of the lower limb: UK Wound management of Open Lower Limb Fractures (UK WOLFF). BMJ Open, 2015, 5, e009087.   | 1.9  | 23        |
| 210 | A multi-centre, randomised controlled trial of the effectiveness of PDSAFE to prevent falls among people with Parkinson's: study protocol. BMC Neurology, 2015, 15, 81.   | 1.8  | 10        |
| 211 | A randomised controlled trial and cost-effectiveness analysis of high-frequency oscillatory ventilation against conventional artificial ventilation for adults with acute respiratory distress syndrome. The OSCAR (OSCillation in ARDS) study. Health Technology Assessment, 2015, 19, 1-178.  | 2.8  | 34        |
| 212 | Quantitative biomechanical comparison of ankle fracture casting methods. Biomedizinische Technik, 2015, 60, 263-7.  | 0.8  | 2         |
| 213 | Exercises to improve function of the rheumatoid hand (SARAH): a randomised controlled trial. Lancet,<br>The, 2015, 385, 421-429.  | 13.7 | 107       |
| 214 | Comprehensive geriatric care for patients with hip fractures: a prospective, randomised, controlled trial. Lancet, The, 2015, 385, 1623-1633.   | 13.7 | 449       |
| 215 | Patterns of interaction between factors that enhance or inhibit recovery from chronic low back pain. Disability and Rehabilitation, 2015, 37, 194-206.  | 1.8  | 3         |
| 216 | Mechanical versus manual chest compression for out-of-hospital cardiac arrest (PARAMEDIC): a pragmatic, cluster randomised controlled trial. Lancet, The, 2015, 385, 947-955.   | 13.7 | 362       |

| #   | Article   | IF   | CITATIONS |
|-----|---|------|-----------|
| 217 | Predictors of attendance to group exercise: a cohort study of older adults in long-term care facilities. BMC Geriatrics, 2015, 15, 37.  | 2.7  | 19        |
| 218 | "lf I can get over that, I can get over anythingâ€â€"understanding how individuals with acute whiplash<br>disorders form beliefs about pain and recovery: a qualitative study. Physiotherapy, 2015, 101, 178-186.   | 0.4  | 19        |
| 219 | The case for stepped-wedge studies: a trial of falls prevention. Lancet, The, 2015, 385, 2556-2557.   | 13.7 | 6         |
| 220 | UK Fixation of Distal Tibia Fractures (UK FixDT): protocol for a randomised controlled trial of<br>â€ <sup>−</sup> locking' plate fixation versus intramedullary nail fixation in the treatment of adult patients with a<br>displaced fracture of the distal tibia. BMJ Open, 2015, 5, e009162. | 1.9  | 11        |
| 221 | Development of an exercise intervention to improve cognition in people with mild to moderate<br>dementia: Dementia And Physical Activity (DAPA) Trial, registration ISRCTN32612072. Physiotherapy, 2015,<br>101, 126-134.   | 0.4  | 22        |
| 222 | Better balance, fewer falls. BMJ, The, 2015, 351, h3930.  | 6.0  | 1         |
| 223 | The effects of ankle supports on gait in adults: A randomized cross-over study. Journal of Electromyography and Kinesiology, 2015, 25, 973-981.   | 1.7  | 2         |
| 224 | Risk factors for chronic disability in a cohort of patients with acute whiplash associated disorders seeking physiotherapy treatment for persisting symptoms. Physiotherapy, 2015, 101, 34-43.  | 0.4  | 33        |
| 225 | Recommendations for a Core Outcome Set for Measuring Standing Balance in Adult Populations: A<br>Consensus-Based Approach. PLoS ONE, 2015, 10, e0120568.  | 2.5  | 66        |
| 226 | The Effectiveness of Cognitive Behavioural Treatment for Non-Specific Low Back Pain: A Systematic Review and Meta-Analysis. PLoS ONE, 2015, 10, e0134192.   | 2.5  | 149       |
| 227 | UK DRAFFT: a randomised controlled trial of percutaneous fixation with Kirschner wires versus volar locking-plate fixation in the treatment of adult patients with a dorsally displaced fracture of the distal radius. Health Technology Assessment, 2015, 19, 1-124.                           | 2.8  | 72        |
| 228 | Strengthening And stretching for Rheumatoid Arthritis of the Hand (SARAH). A randomised controlled trial and economic evaluation. Health Technology Assessment, 2015, 19, 1-222.  | 2.8  | 43        |
| 229 | Active Treatment for Idiopathic Adolescent Scoliosis (ACTIvATeS): a feasibility study. Health Technology Assessment, 2015, 19, 1-242.   | 2.8  | 16        |
| 230 | The Beta Agonist Lung Injury Trial Prevention. A Randomized Controlled Trial. American Journal of<br>Respiratory and Critical Care Medicine, 2014, 189, 674-683.  | 5.6  | 64        |
| 231 | The REFORM study protocol: a cohort randomised controlled trial of a multifaceted podiatry intervention for the prevention of falls in older people. BMJ Open, 2014, 4, e006977.  | 1.9  | 21        |
| 232 | Percutaneous fixation with Kirschner wires versus volar locking plate fixation in adults with<br>dorsally displaced fracture of distal radius: randomised controlled trial. BMJ, The, 2014, 349,<br>g4807-g4807.  | 6.0  | 167       |
| 233 | The administration of intermittent parathyroid hormone affects functional recovery from pertrochanteric fractured neck of femur: a protocol for a prospective mixed method pilot study with randomisation of treatment allocation and blinded assessment (FRACTT). BMJ Open, 2014, 4, e004389.  | 1.9  | 5         |
| 234 | Better reporting of interventions: template for intervention description and replication (TIDieR) checklist and guide. BMJ, The, 2014, 348, g1687-g1687.  | 6.0  | 5,661     |

| #   | Article  | IF   | CITATIONS |
|-----|--|------|-----------|
| 235 | Case typologies, chronic illness and primary health care. Journal of Evaluation in Clinical Practice, 2014, 20, 513-521.   | 1.8  | 3         |
| 236 | Ankle Injury Management (AIM): design of a pragmatic multi-centre equivalence randomised controlled trial comparing Close Contact Casting (CCC) to Open surgical Reduction and Internal Fixation (ORIF) in the treatment of unstable ankle fractures in patients over 60Âyears. BMC Musculoskeletal Disorders, 2014, 15, 79. | 1.9  | 14        |
| 237 | High Prevalence of Falls, Fear of Falling, and Impaired Balance in Older Adults with Pain in the United<br>States: Findings from the 2011 National Health and Aging Trends Study. Journal of the American<br>Geriatrics Society, 2014, 62, 1844-1852.  | 2.6  | 114       |
| 238 | Properties of patient-reported outcome measures in individuals following acute whiplash injury.<br>Health and Quality of Life Outcomes, 2014, 12, 38.  | 2.4  | 16        |
| 239 | Treatment compliance and effectiveness of a cognitive behavioural intervention for low back pain: a complier average causal effect approach to the BeST data set. BMC Musculoskeletal Disorders, 2014, 15, 17.   | 1.9  | 24        |
| 240 | Physiotherapy Rehabilitation for Osteoporotic Vertebral Fracture (PROVE): study protocol for a randomised controlled trial. Trials, 2014, 15, 22.  | 1.6  | 18        |
| 241 | Matching patients to an intervention for back pain: classifying patients using a latent class approach.<br>Journal of Evaluation in Clinical Practice, 2014, 20, 544-550.  | 1.8  | 10        |
| 242 | Operative versus non-operative treatment for closed, displaced, intra-articular fractures of the calcaneus: randomised controlled trial. BMJ, The, 2014, 349, g4483-g4483.   | 6.0  | 226       |
| 243 | Mapping between the Roland Morris Questionnaire and Generic Preference-Based Measures. Value in Health, 2014, 17, 686-695.   | 0.3  | 20        |
| 244 | Early Ankle Movement Versus Immobilization in the Postoperative Management of Ankle Fracture in<br>Adults: A Systematic Review and Meta-analysis. Journal of Orthopaedic and Sports Physical Therapy,<br>2014, 44, 690-C7.   | 3.5  | 41        |
| 245 | Permanent personhood or meaningful decline? Toward a critical anthropology of successful aging.<br>Journal of Aging Studies, 2014, 29, 41-52.  | 1.4  | 215       |
| 246 | Emergency department treatments and physiotherapy for acute whiplash: a pragmatic, two-step, randomised controlled trial. Lancet, The, 2013, 381, 546-556.   | 13.7 | 102       |
| 247 | Factors associated with mobility outcomes in older people post-ankle fracture: An observational cohort study focussing on peripheral vessel function. Injury, 2013, 44, 987-993.   | 1.7  | 6         |
| 248 | Exercise for depression in elderly people $\hat{a} \in $ "Authors' reply. Lancet, The, 2013, 382, 1325.  | 13.7 | 0         |
| 249 | High-Frequency Oscillation for Acute Respiratory Distress Syndrome. New England Journal of Medicine, 2013, 368, 806-813.   | 27.0 | 1,024     |
| 250 | Exercise for depression in elderly residents of care homes: a cluster-randomised controlled trial.<br>Lancet, The, 2013, 382, 41-49.   | 13.7 | 108       |
| 251 | Change in Vision, Visual Disability, and Health After Cataract Surgery. Optometry and Vision Science, 2013, 90, 392-399.   | 1.2  | 15        |
| 252 | Exercise for depression in care home residents: a randomised controlled trial with cost-effectiveness analysis (OPERA). Health Technology Assessment, 2013, 17, 1-281.   | 2.8  | 41        |

| #   | Article  | IF   | CITATIONS |
|-----|--|------|-----------|
| 253 | Beta-Agonist Lung injury TrIal-2 (BALTI-2): a multicentre, randomised, double-blind, placebo-controlled<br>trial and economic evaluation of intravenous infusion of salbutamol versus placebo in patients with<br>acute respiratory distress syndrome. Health Technology Assessment, 2013, 17, v-vi, 1-87. | 2.8  | 39        |
| 254 | Managing Injuries of the Neck Trial (MINT): a randomised controlled trial of treatments for whiplash<br>injuries. Health Technology Assessment, 2012, 16, iii-iv, 1-141.   | 2.8  | 28        |
| 255 | Effect of intravenous Î <sup>2</sup> -2 agonist treatment on clinical outcomes in acute respiratory distress syndrome (BALTI-2): a multicentre, randomised controlled trial. Lancet, The, 2012, 379, 229-235.  | 13.7 | 307       |
| 256 | Intravenous salbutamol in ARDS and increased mortality – Authors' reply. Lancet, The, 2012, 379, 1876.   | 13.7 | 0         |
| 257 | Effect of exercise on depression severity in older people: systematic review and meta-analysis of randomised controlled trials. British Journal of Psychiatry, 2012, 201, 180-185.   | 2.8  | 324       |
| 258 | A systematic review of patient-reported outcome measures used to assess Achilles tendon rupture management: What's being used and should we be using it?. British Journal of Sports Medicine, 2012, 46, 1102-1109.   | 6.7  | 60        |
| 259 | Telephone interviews can be used to collect follow-up data subsequent to no response to postal questionnaires in clinical trials. Journal of Clinical Epidemiology, 2012, 65, 90-99.   | 5.0  | 14        |
| 260 | Strengthening and stretching for rheumatoid arthritis of the hand (SARAH): design of a randomised controlled trial of a hand and upper limb exercise intervention - ISRCTN89936343. BMC Musculoskeletal Disorders, 2012, 13, 230.  | 1.9  | 12        |
| 261 | Development and delivery of an exercise intervention for rheumatoid arthritis: Strengthening and stretching for rheumatoid arthritis of the hand (SARAH) trial. Physiotherapy, 2012, 98, 121-130.  | 0.4  | 45        |
| 262 | Group cognitive behavioural interventions for low back pain in primary care: Extended follow-up of the Back Skills Training Trial (ISRCTN54717854). Pain, 2012, 153, 494-501.  | 4.2  | 43        |
| 263 | The Achilles tendon total rupture score: a study of responsiveness, internal consistency and convergent validity on patients with acute Achilles tendon ruptures. Health and Quality of Life Outcomes, 2012, 10, 24.   | 2.4  | 57        |
| 264 | Assessment of Physical Activity in Older People With and Without Cognitive Impairment. Journal of Aging and Physical Activity, 2011, 19, 347-372.  | 1.0  | 35        |
| 265 | Exploring the use of the â€~Adapted Miracle Question' in the assessment of low back pain. Physiotherapy, 2011, 97, 203-208.  | 0.4  | 7         |
| 266 | Effect of in-hospital comprehensive geriatric assessment (CGA) in older people with hip fracture. The protocol of the Trondheim Hip Fracture Trial. BMC Geriatrics, 2011, 11, 18.  | 2.7  | 47        |
| 267 | UK DRAFFT - A randomised controlled trial of percutaneous fixation with kirschner wires versus volar locking-plate fixation in the treatment of adult patients with a dorsally displaced fracture of the distal radius. BMC Musculoskeletal Disorders, 2011, 12, 201.                                      | 1.9  | 22        |
| 268 | Beta Agonist Lung Injury TrIal-2 (BALTI-2) trial protocol: A randomised, double-blind,<br>placebo-controlled of intravenous infusion of salbutamol in the acute respiratory distress<br>syndrome. Trials, 2011, 12, 113.   | 1.6  | 17        |
| 269 | Reporting of complex interventions in clinical trials: development of a taxonomy to classify and describe fall-prevention interventions. Trials, 2011, 12, 125.  | 1.6  | 151       |
| 270 | The OPERA trial: protocol for a randomised trial of an exercise intervention for older people in residential and nursing accommodation. Trials, 2011, 12, 27.  | 1.6  | 29        |

| #   | Article   | IF                 | CITATIONS            |
|-----|---|--------------------|----------------------|
| 271 | The Beta Agonist Lung Injury TrIal (BALTI) - prevention trial protocol. Trials, 2011, 12, 79.   | 1.6                | 13                   |
| 272 | Predicting response to a cognitive–behavioral approach to treating low back pain: Secondary analysis<br>of the BeST data set. Arthritis Care and Research, 2011, 63, 1271-1279.   | 3.4                | 20                   |
| 273 | In-Shoe Plantar Pressures Within Ankle-Foot Orthoses. American Journal of Sports Medicine, 2011, 39, 2679-2685.   | 4.2                | 13                   |
| 274 | Sarcopenia and some simple approaches to modifying the consequence of ageing. Clinical Medicine, 2011, 11, 170-172.   | 1.9                | 0                    |
| 275 | Rethinking the Generation Gap: Age and Agency in Middle-Class Kolkata. Journal of Aging, Humanities, and the Arts, 2010, 4, 83-97.  | 0.0                | 5                    |
| 276 | Effect of a Statewide Fall Prevention Program on Incidence of Femoral Fractures in Residents of<br>Longâ€Term Care Facilities. Journal of the American Geriatrics Society, 2010, 58, 70-75.   | 2.6                | 22                   |
| 277 | A cognitive-behavioural programme for the management of low back pain in primary care: a description and justification of the intervention used in the Back Skills Training Trial (BeST; ISRCTN) Tj ETQq1 1 C                         | .7 <b>8</b> 4814 r | g <b>B1</b> 6/Overlo |
| 278 | Prehospital Randomised Assessment of a Mechanical Compression Device In Cardiac Arrest<br>(PaRAMeDIC) Trial Protocol. Scandinavian Journal of Trauma, Resuscitation and Emergency Medicine,<br>2010, 18, 58.                          | 2.6                | 45                   |
| 279 | A multicentred randomised controlled trial of a primary care-based cognitive behavioural programme<br>for low back pain. The Back Skills Training (BeST) trial. Health Technology Assessment, 2010, 14, 1-253,<br>iii-iv.             | 2.8                | 122                  |
| 280 | Kognitive Gruppen-Verhaltenstherapie in der primÄ <b>r</b> en Gesundheitsversorgung bei LWS-Syndrom: eine<br>randomisierte, kontrollierte Studie und Analyse der Kostenwirksamkeit. Osteopathische Medizin, 2010,<br>11, 24-26.       | 0.2                | 1                    |
| 281 | Recovery of Muscle Strength and Power After Limb-Lengthening Surgery. Archives of Physical Medicine and Rehabilitation, 2010, 91, 384-388.  | 0.9                | 20                   |
| 282 | Group cognitive behavioural treatment for low-back pain in primary care: a randomised controlled trial and cost-effectiveness analysis. Lancet, The, 2010, 375, 916-923.  | 13.7               | 347                  |
| 283 | Developing Evidence for How to Tailor Medical Interventions for the Individual Patient. Qualitative<br>Health Research, 2010, 20, 1629-1641.  | 2.1                | 39                   |
| 284 | Development and delivery of a physiotherapy intervention for the early management of whiplash<br>injuries: The Managing Injuries of Neck Trial (MINT) Intervention. Physiotherapy, 2009, 95, 15-23.                                   | 0.4                | 13                   |
| 285 | Group versus individual sessions delivered by a physiotherapist for female urinary incontinence: an interview study with women attending group sessions nested within a randomised controlled trial. BMC Women's Health, 2009, 9, 25. | 2.0                | 5                    |
| 286 | Group treatments for sensitive health care problems: a randomised controlled trial of group versus<br>individual physiotherapy sessions for female urinary incontinence. BMC Women's Health, 2009, 9, 26.                             | 2.0                | 30                   |
| 287 | Does a monetary incentive improve the response to a postal questionnaire in a randomised controlled trial? The MINT incentive study. Trials, 2009, 10, 44.  | 1.6                | 30                   |
| 288 | Interventions for preventing falls in older people living in the community. , 2009, , CD007146.   |                    | 656                  |

| #   | Article  | IF   | CITATIONS |
|-----|--|------|-----------|
| 289 | Multidisciplinary assessment of elderly people with a history of multiple falls reduces the risk of<br>further falls. Australian Journal of Physiotherapy, 2009, 55, 139.  | 0.9  | 4         |
| 290 | Mechanical supports for acute, severe ankle sprain: a pragmatic, multicentre, randomised controlled<br>trial. Lancet, The, 2009, 373, 575-581.   | 13.7 | 191       |
| 291 | 10-day below-knee cast for management of severe ankle sprains – Authors' reply. Lancet, The, 2009, 373,<br>1602-1603.  | 13.7 | 1         |
| 292 | The Prognostic Value of Symptom Responses in the Conservative Management of Spinal Pain. Spine, 2009, 34, 2686-2699.   | 2.0  | 16        |
| 293 | EuroQol EQ-5D and condition-specific measures of health outcome in women with urinary incontinence: reliability, validity and responsiveness. Quality of Life Research, 2008, 17, 475-483.   | 3.1  | 50        |
| 294 | Hip Fractures in Institutionalized Elderly People: Incidence Rates and Excess Mortality. Journal of<br>Bone and Mineral Research, 2008, 23, 1825-1831.   | 2.8  | 131       |
| 295 | Prevention of Falls in Nursing Homes: Subgroup Analyses of a Randomized Fall Prevention Trial.<br>Journal of the American Geriatrics Society, 2008, 56, 1092-1097.   | 2.6  | 48        |
| 296 | Response and nonâ€response to postal questionnaire followâ€up in a clinical trial – a qualitative study<br>of the patient's perspective. Journal of Evaluation in Clinical Practice, 2008, 14, 226-235.  | 1.8  | 32        |
| 297 | A national survey of services for the prevention and management of falls in the UK. BMC Health<br>Services Research, 2008, 8, 233.   | 2.2  | 23        |
| 298 | A systematic literature review of psychological factors and the development of late whiplash syndrome. Pain, 2008, 135, 20-30.   | 4.2  | 107       |
| 299 | Chronic disease management programme in people with severe knee osteoarthritis: efficacy and moderators of response. Clinical Rehabilitation, 2008, 22, 169-178.   | 2.2  | 17        |
| 300 | The Optimal Sequence and Selection of Screening Test Items to Predict Fall Risk in Older Disabled<br>Women: The Women's Health and Aging Study. Journals of Gerontology - Series A Biological Sciences<br>and Medical Sciences, 2008, 63, 1082-1088. | 3.6  | 54        |
| 301 | Responsiveness of a Patient Specific Outcome Measure Compared With the Oswestry Disability Index v2.1 and Roland and Morris Disability Questionnaire for Patients With Subacute and Chronic Low Back Pain. Spine, 2008, 33, 2450-2457.               | 2.0  | 79        |
| 302 | Leg Extension Power Deficit and Mobility Limitation in Women Recovering from Hip Fracture.<br>American Journal of Physical Medicine and Rehabilitation, 2008, 87, 363-370.   | 1.4  | 34        |
| 303 | Title is missing!. Journal of Rehabilitation Research and Development, 2008, 45, 1105.   | 1.6  | 161       |
| 304 | A Systematic Literature Review of Physical Prognostic Factors for the Development of Late Whiplash<br>Syndrome. Spine, 2007, 32, E764-E780.  | 2.0  | 76        |
| 305 | Design considerations in a clinical trial of a cognitive behavioural intervention for the management<br>of low back pain in primary care: Back Skills Training Trial. BMC Musculoskeletal Disorders, 2007, 8, 14.                                    | 1.9  | 15        |
| 306 | Managing Injuries of the Neck Trial (MINT): design of a randomised controlled trial of treatments for<br>whiplash associated disorders. BMC Musculoskeletal Disorders, 2007, 8, 7.   | 1.9  | 28        |

| #   | Article   | IF  | CITATIONS |
|-----|---|-----|-----------|
| 307 | A Validation Study of the New Zealand Score for Hip and Knee Surgery. Clinical Orthopaedics and Related Research, 2007, 464, 190-195.   | 1.5 | 11        |
| 308 | Systematic review of definitions and methods of measuring falls in randomised controlled fall prevention trials. Age and Ageing, 2006, 35, 5-10.  | 1.6 | 377       |
| 309 | Cost-Utility Analysis of Physiotherapy Treatment Compared With Physiotherapy Advice in Low Back<br>Pain. Spine, 2006, 31, 1381-1387.  | 2.0 | 59        |
| 310 | A RESPONSE LETTER TO DRS. HAUSDORFF AND YOGEV. Journal of the American Geriatrics Society, 2006, 54, 865-866.   | 2.6 | 12        |
| 311 | ASYMMETRICAL LOWER EXTREMITY POWER DEFICIT AS A RISK FACTOR FOR INJURIOUS FALLS IN HEALTHY OLDER WOMEN. Journal of the American Geriatrics Society, 2006, 54, 551-553.  | 2.6 | 33        |
| 312 | Personal meanings in the construction of need for total knee replacement surgery. Social Science and Medicine, 2006, 63, 43-53.   | 3.8 | 55        |
| 313 | Maximising response to postal questionnaires – A systematic review of randomised trials in health research. BMC Medical Research Methodology, 2006, 6, 5.   | 3.1 | 274       |
| 314 | Re: 'Falls Definition Validation'. Age and Ageing, 2006, 36, 111-112.   | 1.6 | 6         |
| 315 | Suitability of Physical Activity Questionnaires for Older Adults in Fall-Prevention Trials: A Systematic Review. Journal of Aging and Physical Activity, 2005, 13, 461-481.                                       | 1.0 | 55        |
| 316 | Measuring the Psychological Outcomes of Falling: A Systematic Review. Journal of the American<br>Geriatrics Society, 2005, 53, 501-510.   | 2.6 | 384       |
| 317 | Development of a Common Outcome Data Set for Fall Injury Prevention Trials: The Prevention of Falls<br>Network Europe Consensus. Journal of the American Geriatrics Society, 2005, 53, 1618-1622.                 | 2.6 | 1,615     |
| 318 | A critical appraisal and defence of the Oxfordshire low back pain trial. Physiotherapy, 2005, 91, 1-2.  | 0.4 | 3         |
| 319 | Clinical and cost effectiveness of mechanical support for severe ankle sprains: design of a randomised controlled trial in the emergency department [ISRCTN 37807450]. BMC Musculoskeletal Disorders, 2005, 6, 1. | 1.9 | 46        |
| 320 | Effective Measurement of the Functional Progress of Stroke Clients. British Journal of Occupational Therapy, 2005, 68, 253-259.   | 0.9 | 11        |
| 321 | Randomised controlled trial of physiotherapy compared with advice for low back pain. BMJ: British<br>Medical Journal, 2004, 329, 708.   | 2.3 | 163       |
| 322 | Association between radiographic joint space narrowing, function, pain and muscle power in severe osteoarthritis of the knee. Clinical Rehabilitation, 2004, 18, 793-800.   | 2.2 | 109       |
| 323 | Multi-item outcome measures for lateral ligament injury of the ankle: a structured review. Journal of Evaluation in Clinical Practice, 2004, 10, 339-352.   | 1.8 | 26        |
| 324 | Cryotherapy after total knee replacement: a survey of current practice. Physiotherapy Research<br>International, 2003, 8, 111-120.  | 1.5 | 33        |

| #   | Article   | IF  | CITATIONS |
|-----|---|-----|-----------|
| 325 | Recovery of mobility after knee arthroplasty. Journal of Arthroplasty, 2003, 18, 575-582.   | 3.1 | 40        |
| 326 | Risk Factors for Falling in Home-Dwelling Older Women With Stroke. Stroke, 2003, 34, 494-501.   | 2.0 | 242       |
| 327 | Changes in muscle torque following anterior cruciate ligament reconstruction: A comparison<br>between hamstrings and patella tendon graft procedures on 45 patients. Acta Orthopaedica, 2002, 73,<br>546-552. | 1.4 | 54        |
| 328 | Neuromuscular stimulation of the quadriceps muscle after hip fracture: A randomized controlled trial. Archives of Physical Medicine and Rehabilitation, 2002, 83, 1087-1092.                                  | 0.9 | 50        |
| 329 | Being a Widow and Other Life Stories: The Interplay between Lives and Words. Anthropology and Humanism, 2001, 26, 16-34.  | 0.7 | 50        |
| 330 | Loss of Knee Range of Motion in Leg Lengthening. Journal of Orthopaedic and Sports Physical Therapy, 2001, 31, 238-246.   | 3.5 | 30        |
| 331 | HP0035â€Association between tess functional questionnaire and timed tests of function. , 2001, , .  |     | 0         |
| 332 | HP0012â€The discriminatory validity of the new zealand score for hip and knee surgery in a uk population. , 2001, , .   |     | 0         |
| 333 | Effects of Neuromuscular Stimulation on the Contractile Efficiency of the First Dorsal Interosseous<br>Muscle of the Hand. Physiotherapy, 2000, 86, 380.  | 0.4 | 1         |
| 334 | How Much Function Do Functional Restoration Programmes Restore?. Physiotherapy, 2000, 86, 496.  | 0.4 | 0         |
| 335 | A Functional Restoration Programme for Chronic Low Back Pain. Physiotherapy, 2000, 86, 285-293.   | 0.4 | 32        |
| 336 | Research Assessment Exercise 2001: what is the big deal?. International Journal of Therapy and Rehabilitation, 2000, 7, 373-373.  | 0.1 | 1         |
| 337 | Interventions for preventing falls in elderly people. , 2000, , CD000340.   |     | 695       |
| 338 | Physiotherapy Outcome Measures for Rehabilitation of Elderly People. Physiotherapy, 1998, 84, 216-221.  | 0.4 | 22        |
| 339 | Does intereferential therapy affect blood flow?. Clinical Rehabilitation, 1994, 8, 213-218.   | 2.2 | 8         |
| 340 | Exercise $\hat{a} \in$ "The Other Root of our Profession. Physiotherapy, 1993, 79, 772.   | 0.4 | 13        |
| 341 | Immobilisation versus early ankle movement for treating acute lateral ankle ligament injuries in adults. The Cochrane Library, 0, , .   | 2.8 | 0         |
| 342 | Multifactorial and multiple component interventions for preventing falls in older people living in the community. The Cochrane Library, 0, , .  | 2.8 | 8         |

| #   | Article  | lF  | CITATIONS |
|-----|--|-----|-----------|
| 343 | Exercise for preventing falls in older people living in the community. The Cochrane Library, 0, , .  | 2.8 | 9         |
| 344 | Environmental interventions for preventing falls in older people living in the community. The Cochrane Library, 0, , .   | 2.8 | 23        |
| 345 | Home hazard assessment and environmental modification to prevent falls in older people: the OTIS trial. F1000Research, 0, 10, 500.   | 1.6 | 2         |
| 346 | Better reporting of interventions: template for intervention description and replication (TIDieR) checklist and guide. , 0, .  |     | 1         |
| 347 | Can behavioural change interventions improve self-efficacy and exercise adherence among people with Parkinson's? A systematic review protocol. HRB Open Research, 0, 5, 15.  | 0.6 | Ο         |
| 348 | Anti-TNF (adalimumab) injection for the treatment of adults with frozen shoulder during the pain<br>predominant stage protocol for a multi-centre, randomised, double blind, parallel group, feasibility<br>trial. NIHR Open Research, 0, 2, 28. | 0.0 | 1         |