

# Haoqiang Zhang

## List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/927042/publications.pdf>

Version: 2024-02-01

16  
papers

119  
citations

1478505

6  
h-index

1372567

10  
g-index

16  
all docs

16  
docs citations

16  
times ranked

129  
citing authors

#	ARTICLE	IF	CITATIONS
1	Free Triiodothyronine Levels are Related to Executive Function and Scene Memory in Type 2 Diabetes Mellitus Patients Without Diagnosed Thyroid Diseases. <i>Diabetes, Metabolic Syndrome and Obesity: Targets and Therapy</i> , 2022, Volume 15, 1041-1050.	2.4	6
2	Emodin Attenuated the Kidney Damage of High-Fat-Diet Mice via the Upregulation of Glucagon-Like Peptide-1 Receptor. <i>BioMed Research International</i> , 2021, 2021, 1-9.	1.9	2
3	Increased Plasma Level of 24S-Hydroxycholesterol and Polymorphism of CYP46A1 SNP (rs754203) Are Associated With Mild Cognitive Impairment in Patients With Type 2 Diabetes. <i>Frontiers in Aging Neuroscience</i> , 2021, 13, 619916.	3.4	9
4	Glucagon-like peptide-1 attenuated carboxymethyl lysine induced neuronal apoptosis via peroxisome proliferation activated receptor-1 $\beta$ . <i>Aging</i> , 2021, 13, 19013-19027.	3.1	6
5	Elevated Plasma Free Fatty Acid Susceptible to Early Cognitive Impairment in Type 2 Diabetes Mellitus. <i>Journal of Alzheimer's Disease</i> , 2021, 82, 1345-1356.	2.6	3
6	Inverted U-shaped correlation between serum low-density lipoprotein cholesterol levels and cognitive functions of patients with type 2 diabetes mellitus. <i>Lipids in Health and Disease</i> , 2021, 20, 103.	3.0	5
7	Decreased Plasma Level of Lipoprotein Lipase Predicted Verbal Disfluency in Chinese Type 2 Diabetes Mellitus Patients with Early Cognitive Deficits. <i>Current Alzheimer Research</i> , 2021, 18, 656-666.	1.4	1
8	Liraglutide improved the cognitive function of diabetic mice via the receptor of advanced glycation end products down-regulation. <i>Aging</i> , 2021, 13, 525-536.	3.1	12
9	Association of Low-Density Lipoprotein Receptor-Related Protein 1 and Its rs1799986 Polymorphism With Mild Cognitive Impairment in Chinese Patients With Type 2 Diabetes. <i>Frontiers in Neuroscience</i> , 2020, 14, 743.	2.8	6
10	Ginsenoside Rb1 Alleviated High-Fat-Diet-Induced Hepatocytic Apoptosis via Peroxisome Proliferator-Activated Receptor $\alpha$ . <i>BioMed Research International</i> , 2020, 2020, 1-9.	1.9	12
11	The CC Genotype of Insulin-Induced Gene 2 rs7566605 Is a Protective Factor of Hypercholesterolemia Susceptible to Mild Cognitive Impairment, Especially to the Executive Function of Patients with Type 2 Diabetes Mellitus. <i>BioMed Research International</i> , 2020, 2020, 1-7.	1.9	4
12	Chronic hyperglycemia induces tau hyperphosphorylation by downregulating OGT-involved O-GlcNAcylation in vivo and in vitro. <i>Brain Research Bulletin</i> , 2020, 156, 76-85.	3.0	18
13	Higher Plasma Level of Nampt Presaging Memory Dysfunction in Chinese Type 2 Diabetes Patients with Mild Cognitive Impairment. <i>Journal of Alzheimer's Disease</i> , 2019, 70, 303-314.	2.6	8
14	Interleukin 29 activates expression of tissue inhibitor of metalloproteinase 1 in macrophages via toll-like receptor 2. <i>Molecular Medicine Reports</i> , 2018, 17, 8363-8368.	2.4	9
15	Toll-like receptor 2 mediates deposition of collagen I in adipose tissue of high fat diet-induced obese mice. <i>Molecular Medicine Reports</i> , 2018, 17, 5958-5963.	2.4	8
16	Ginsenoside Rb1 increases insulin sensitivity through suppressing 11 $\beta$ -hydroxysteroid dehydrogenase type I. <i>American Journal of Translational Research (discontinued)</i> , 2017, 9, 1049-1057.	0.0	10