

Maria Pia Abruzzese

List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/9304314/publications.pdf>

Version: 2024-02-01

8
papers

405
citations

1163117
8
h-index

1588992
8
g-index

8
all docs

8
docs citations

8
times ranked

814
citing authors

#	ARTICLE	IF	CITATIONS
1	Genotoxic stress modulates the release of exosomes from multiple myeloma cells capable of activating NK cell cytokine production: Role of HSP70/TLR2/NF- κ B axis. <i>Oncolimmunology</i> , 2017, 6, e1279372.	4.6	100
2	The IMiDs targets IKZF-1/3 and IRF4 as novel negative regulators of NK cell-activating ligands expression in multiple myeloma. <i>Oncotarget</i> , 2015, 6, 23609-23630.	1.8	78
3	Inhibition of bromodomain and extra-terminal (BET) proteins increases NKG2D ligand MICA expression and sensitivity to NK cell-mediated cytotoxicity in multiple myeloma cells: role of cMYC-IRF4-miR-125b interplay. <i>Journal of Hematology and Oncology</i> , 2016, 9, 134.	17.0	72
4	Nitric oxide donors increase PVR/CD155 DNAM-1 ligand expression in multiple myeloma cells: role of DNA damage response activation. <i>BMC Cancer</i> , 2015, 15, 17.	2.6	54
5	Immunoregulatory and Effector Activities of Nitric Oxide and Reactive Nitrogen Species in Cancer. <i>Current Medicinal Chemistry</i> , 2016, 23, 2618-2636.	2.4	42
6	p38 MAPK differentially controls NK activating ligands at transcriptional and post-transcriptional level on multiple myeloma cells. <i>Oncolimmunology</i> , 2017, 6, e1264564.	4.6	29
7	Activation of liver X receptor up-regulates the expression of the NKG2D ligands MICA and MICB in multiple myeloma through different molecular mechanisms. <i>FASEB Journal</i> , 2019, 33, 9489-9504.	0.5	19
8	The homeobox transcription factor MEIS2 is a regulator of cancer cell survival and IMiDs activity in Multiple Myeloma: modulation by Bromodomain and Extra-Terminal (BET) protein inhibitors. <i>Cell Death and Disease</i> , 2019, 10, 324.	6.3	11