

# L BÃ¼nger

## List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/9315834/publications.pdf>

Version: 2024-02-01

134  
papers

3,233  
citations

159585

30  
h-index

197818

49  
g-index

141  
all docs

141  
docs citations

141  
times ranked

3358  
citing authors

#	ARTICLE	IF	CITATIONS
1	Prediction of intramuscular fat in lamb by visible and near-infrared spectroscopy in an abattoir environment. <i>Meat Science</i> , 2021, 171, 108286.	5.5	12
2	Predicting the shear value and intramuscular fat in meat from Nellore cattle using Vis-NIR spectroscopy. <i>Meat Science</i> , 2020, 163, 108077.	5.5	11
3	Prediction of intramuscular fat content and shear force in Texel lamb loins using combinations of different X-ray computed tomography (CT) scanning techniques. <i>Meat Science</i> , 2018, 140, 78-85.	5.5	8
4	Analysis of single nucleotide polymorphisms variation associated with important economic and computed tomography measured traits in Texel sheep. <i>Animal</i> , 2018, 12, 915-922.	3.3	12
5	Effects of feed allowance and indispensable amino acid reduction on feed intake, growth performance and carcass characteristics of growing pigs. <i>PLoS ONE</i> , 2018, 13, e0195645.	2.5	38
6	Absolute Radiation Thermometry in the NIR. <i>International Journal of Thermophysics</i> , 2017, 38, 1.	2.1	6
7	Prediction of intramuscular fat content using CT scanning of packaged lamb cuts and relationships with meat eating quality. <i>Meat Science</i> , 2017, 123, 112-119.	5.5	33
8	Baseline Muscle Mass Is a Poor Predictor of Functional Overload-Induced Gain in the Mouse Model. <i>Frontiers in Physiology</i> , 2016, 7, 534.	2.8	6
9	Myostatin dysfunction is associated with reduction in overload induced hypertrophy of soleus muscle in mice. <i>Scandinavian Journal of Medicine and Science in Sports</i> , 2016, 26, 894-901.	2.9	9
10	Oxidative costs of reproduction in mouse strains selected for different levels of food intake and which differ in reproductive performance. <i>Scientific Reports</i> , 2016, 6, 36353.	3.3	16
11	Heterogeneous variances and genetics by environment interactions in genetic evaluation of crossbred lambs. <i>Animal</i> , 2015, 9, 380-387.	3.3	4
12	The effects of different farm environments on the performance of Texel sheep. <i>Animal</i> , 2015, 9, 1624-1634.	3.3	5
13	Non-invasive methods for the determination of body and carcass composition in livestock: dual-energy X-ray absorptiometry, computed tomography, magnetic resonance imaging and ultrasound: invited review. <i>Animal</i> , 2015, 9, 1250-1264.	3.3	135
14	Genome-wide association study of footrot in Texel sheep. <i>Genetics Selection Evolution</i> , 2015, 47, 35.	3.0	20
15	Effects of low protein diets on performance of pigs with a lean genotype between 40 and 115 kg liveweight. <i>Animal Production Science</i> , 2015, 55, 461.	1.3	5
16	Myostatin dysfunction impairs force generation in extensor digitorum longus muscle and increases exercise-induced protein efflux from extensor digitorum longus and soleus muscles. <i>Applied Physiology, Nutrition and Metabolism</i> , 2015, 40, 817-821.	1.9	4
17	Traceability of a CCD-Camera System for High-Temperature Measurements. <i>International Journal of Thermophysics</i> , 2015, 36, 1784-1802.	2.1	10
18	Characterisation of terminal sire sheep farm systems, based on a range of environmental factors: a case study in the context of genotype by environment interactions using Charollais lambs. <i>Animal</i> , 2014, 8, 867-876.	3.3	5

#	ARTICLE	IF	CITATIONS
19	Effect and mode of action of the Texel muscling QTL (TM-QTL) on carcass traits in purebred Texel lambs. <i>Animal</i> , 2014, 8, 1053-1061.	3.3	4
20	Limits to sustained energy intake. XXII. Reproductive performance of two selected mouse lines with different thermal conductance. <i>Journal of Experimental Biology</i> , 2014, 217, 3718-32.	1.7	6
21	Effect of the Texel muscling QTL (TM-QTL) on spine characteristics in purebred Texel lambs. <i>Small Ruminant Research</i> , 2014, 117, 34-40.	1.2	8
22	Prediction of intramuscular fat levels in Texel lamb loins using X-ray computed tomography scanning. <i>Meat Science</i> , 2014, 98, 263-271.	5.5	20
23	Genetic improvement of hill sheep â€œ Impacts on profitability and greenhouse gas emissions. <i>Small Ruminant Research</i> , 2014, 120, 27-34.	1.2	13
24	Evaluating invasive and non-invasive methods to determine fat content in the laboratory mouse. <i>Open Life Sciences</i> , 2014, 10, .	1.4	2
25	Effects of low protein diets on pigs with a lean genotype. 1. Carcass composition measured by dissection and muscle fatty acid composition. <i>Meat Science</i> , 2013, 95, 123-128.	5.5	60
26	The relationship between video image analysis (VIA), visual classification, and saleable meat yield of sirloin and fillet cuts of beef carcasses differing in breed and gender. <i>Livestock Science</i> , 2013, 158, 169-178.	1.6	12
27	Effects of low protein diets on pigs with a lean genotype 2. Compositional traits measured with computed tomography (CT). <i>Meat Science</i> , 2013, 95, 129-136.	5.5	21
28	Comparison of the Richardsonâ€™Lucy method and a classical approach for spectrometer bandpass correction. <i>Metrologia</i> , 2013, 50, 107-118.	1.2	40
29	Analyses of muscle spindles in the soleus of six inbred mouse strains. <i>Journal of Anatomy</i> , 2013, 223, 289-296.	1.5	12
30	Genetic and genomic analyses of musculoskeletal differences between BEH and BEL strains. <i>Physiological Genomics</i> , 2013, 45, 940-947.	2.3	14
31	Between- and within-breed variations of spine characteristics in sheep1. <i>Journal of Animal Science</i> , 2013, 91, 995-1004.	0.5	25
32	Genetic evaluation of days to harvest in crossbred lambs1. <i>Journal of Animal Science</i> , 2013, 91, 5153-5160.	0.5	1
33	Index selection in terminal sires improves lamb performance at finishing1. <i>Journal of Animal Science</i> , 2013, 91, 38-43.	0.5	9
34	Meta-analysis of effects of gender in combination with carcass weight and breed on pork quality1. <i>Journal of Animal Science</i> , 2013, 91, 1480-1492.	0.5	41
35	Index selection in terminal sires improves early lamb growth1. <i>Journal of Animal Science</i> , 2012, 90, 142-151.	0.5	7
36	Investigation into the presence of genotype by environment (GÃ—E) interactions in Scottish Blackface lamb weaning traits. , 2012, , 23-31.		0

#	ARTICLE	IF	CITATIONS
37	The effect of sex on some carcass and meat quality traits in Texel ewe and ram lambs. <i>Animal Production Science</i> , 2012, 52, 601.	1.3	16
38	Genetic Parameters for Fitness and Neonatal Behavior Traits in Sheep. <i>Behavior Genetics</i> , 2012, 42, 899-911.	2.1	27
39	Parallel Selection Mapping Using Artificially Selected Mice Reveals Body Weight Control Loci. <i>Current Biology</i> , 2012, 22, 794-800.	3.9	82
40	Bayesian meta-analysis of the effect of fasting, transport and lairage times on four attributes of pork meat quality. <i>Meat Science</i> , 2012, 90, 584-598.	5.5	22
41	A review of the development and use of video image analysis (VIA) for beef carcass evaluation as an alternative to the current EUROP system and other subjective systems. <i>Meat Science</i> , 2012, 92, 307-318.	5.5	62
42	Investigation into the presence of genotype by environment interactions (GÄ—E) in Scottish Blackface lamb traits. <i>Small Ruminant Research</i> , 2012, 105, 46-52.	1.2	12
43	The effect of the Texel Muscling QTL on live and carcass weight in Texel lambs. <i>Small Ruminant Research</i> , 2012, 105, 117-121.	1.2	2
44	Factors affecting dystocia and offspring vigour in different sheep genotypes. <i>Preventive Veterinary Medicine</i> , 2012, 103, 257-264.	1.9	37
45	Meta-analysis of the effects of dietary vitamin E supplementation on Î±-tocopherol concentration and lipid oxidation in pork. <i>Meat Science</i> , 2011, 87, 305-314.	5.5	23
46	Evaluating the effects of a single copy of a mutation in the myostatin gene (c.*1232G>A) on carcass traits in crossbred lambs. <i>Meat Science</i> , 2011, 87, 412-418.	5.5	22
47	Genotypic effects of the Texel Muscling QTL (TM-QTL) on meat quality in purebred Texel lambs. <i>Meat Science</i> , 2011, 89, 125-132.	5.5	12
48	A Stratified Transcriptomics Analysis of Polygenic Fat and Lean Mouse Adipose Tissues Identifies Novel Candidate Obesity Genes. <i>PLoS ONE</i> , 2011, 6, e23944.	2.5	48
49	Development and validation of on-farm behavioural scoring systems to assess birth assistance and lamb vigour. <i>Animal</i> , 2011, 5, 776-783.	3.3	24
50	Interactive effects of protein nutrition, genetic growth potential and <i>Heligmosomoides bakeri</i> infection pressure on resilience and resistance in mice. <i>Parasitology</i> , 2011, 138, 1305-1315.	1.5	9
51	Evaluating the effects of the c.*1232G > A mutation and TM-QTL in TexelÄ—Welsh Mountain lambs using ultrasound and video image analyses. <i>Small Ruminant Research</i> , 2011, 99, 99-109.	1.2	11
52	The effect of gestational undernutrition on maternal weight change and fetal weight in lines of mice selected for different growth characteristics. <i>British Journal of Nutrition</i> , 2011, 105, 539-548.	2.3	1
53	The effects of a loin muscling quantitative trait locus (LoinMAXâ„¢) on carcass and VIA-based traits in crossbred lambs. <i>Animal</i> , 2010, 4, 407-416.	3.3	14
54	Characterisation of white line degeneration in sheep and evidence for genetic influences on its occurrence. <i>Veterinary Research Communications</i> , 2010, 34, 481-489.	1.6	16

#	ARTICLE	IF	CITATIONS
55	Comparison of repeatability and multiple trait threshold models for litter size in sheep using observed and simulated data in Bayesian analyses. <i>Journal of Animal Breeding and Genetics</i> , 2010, 127, 261-271.	2.0	5
56	Genetic parameters for carcass dimensional measurements from Video Image Analysis and their association with conformation and fat class scores. <i>Livestock Science</i> , 2010, 128, 92-100.	1.6	14
57	The prediction of carcass composition and tissue distribution in beef cattle using ultrasound scanning at the start and/or end of the finishing period. <i>Livestock Science</i> , 2010, 131, 193-202.	1.6	17
58	The effect of the Texel muscling QTL (TM-QTL) on meat quality traits in crossbred lambs. <i>Meat Science</i> , 2010, 85, 684-690.	5.5	11
59	The effect of conditioning period on loin muscle tenderness in crossbred lambs with or without the Texel muscling QTL (TM-QTL). <i>Meat Science</i> , 2010, 85, 715-720.	5.5	7
60	Meta-analysis of effects of dietary vitamin E and post slaughter storage conditions on changes of redness (a*) of pork. <i>Archives Animal Breeding</i> , 2010, 53, 564-577.	1.4	3
61	The effects of three muscling Quantitative Trait Loci on growth patterns of crossbred lambs. <i>Proceedings of the British Society of Animal Science</i> , 2009, 2009, 40-40.	0.0	0
62	Genetic relationship between longevity and objectively or subjectively assessed performance traits in sheep using linear censored models1. <i>Journal of Animal Science</i> , 2009, 87, 3482-3489.	0.5	17
63	Use of meat quality information in breeding programmes. , 2009, , 264-291.		3
64	Effects of the Texel muscling quantitative trait locus on carcass traits in crossbred lambs. <i>Animal</i> , 2009, 3, 189-199.	3.3	17
65	Muscle fibre characteristics of two contrasting sheep breeds: Scottish Blackface and Texel. <i>Meat Science</i> , 2009, 81, 372-381.	5.5	16
66	Genetic parameters for carcass composition and performance data in crossbred lambs measured by Video Image Analysis. <i>Meat Science</i> , 2009, 81, 619-625.	5.5	7
67	Prediction of lamb carcass composition and meat quality using combinations of post-mortem measurements. <i>Meat Science</i> , 2009, 81, 711-719.	5.5	31
68	Evaluation of Video Image Analysis (VIA) technology to predict meat yield of sheep carcasses on-line under UK abattoir conditions. <i>Meat Science</i> , 2009, 82, 94-100.	5.5	45
69	Prediction of lamb meat eating quality in two divergent breeds using various live animal and carcass measurements. <i>Meat Science</i> , 2009, 83, 366-375.	5.5	23
70	Immune responses to macroparasites are sensitive to the interaction between genetic growth potential and protein nutrition in mice. <i>Proceedings of the Nutrition Society</i> , 2009, 68, .	1.0	0
71	Genetic growth potential interacts with nutrition on the ability of mice to cope with <i>Heligmosomoides bakeri</i> infection. <i>Parasitology</i> , 2009, 136, 1043-1055.	1.5	15
72	Effects of a quantitative trait locus for increased muscularity on carcass traits measured by subjective conformation and fat class scores and video image analysis in crossbred lambs. <i>Animal</i> , 2009, 3, 1532-1543.	3.3	9

#	ARTICLE	IF	CITATIONS
73	Breeding for resistance to footrot – the use of hoof lesion scoring to quantify footrot in sheep. <i>Veterinary Research Communications</i> , 2008, 32, 583-589.	1.6	29
74	Divergent Physical Activity and Novel Alternative Responses to High Fat Feeding in Polygenic Fat and Lean Mice. <i>Behavior Genetics</i> , 2008, 38, 292-300.	2.1	23
75	Muscularity and eating quality of lambs: Effects of breed, sex and selection of sires using muscularity measurements by computed tomography. <i>Meat Science</i> , 2008, 79, 105-112.	5.5	32
76	The use of various live animal measurements to predict carcass and meat quality in two divergent lamb breeds. <i>Meat Science</i> , 2008, 80, 1138-1149.	5.5	45
77	Breeding for resistance to mastitis in United Kingdom sheep, a review and economic appraisal. <i>Veterinary Record</i> , 2008, 162, 369-376.	0.3	39
78	Relationships between lamb carcass quality traits measured by X-ray computed tomography and current UK hill sheep breeding goals. <i>Animal</i> , 2008, 2, 36-43.	3.3	12
79	Genetic and phenotypic aspects of foot lesion scores in sheep of different breeds and ages. <i>Animal</i> , 2008, 2, 1289-1296.	3.3	30
80	The effects of selection indices for sustainable hill sheep production on carcass composition and muscularity of lambs, measured using X-ray computed tomography. <i>Animal</i> , 2008, 2, 27-35.	3.3	14
81	Lack of myostatin results in excessive muscle growth but impaired force generation. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 2007, 104, 1835-1840.	7.1	341
82	Associations of polymorphisms of the ovine prion protein gene with growth, carcass, and computerized tomography traits in Scottish Blackface lambs <sup>1</sup> . <i>Journal of Animal Science</i> , 2007, 85, 632-640.	0.5	16
83	Changes in carcass traits during growth in lambs of two contrasting breeds, measured using computer tomography. <i>Livestock Science</i> , 2007, 107, 37-52.	1.6	20
84	Accuracy of in vivo muscularity indices measured by computed tomography and their association with carcass quality in lambs. <i>Meat Science</i> , 2007, 75, 533-542.	5.5	31
85	Prediction of sensory characteristics of lamb meat samples by near infrared reflectance spectroscopy. <i>Meat Science</i> , 2007, 76, 509-516.	5.5	124
86	Fine Mapping of Mouse QTLs for Fatness Using SNP Data. <i>OMICS A Journal of Integrative Biology</i> , 2007, 11, 341-350.	2.0	3
87	Evaluation of Video Image Analysis (VIA) technology to predict meat yield of sheep carcasses online under abattoir conditions. <i>Proceedings of the British Society of Animal Science</i> , 2007, 2007, 108-108.	0.0	2
88	Interactive effects of selection for growth and protein supply on the consequences of gastrointestinal parasitism on growth performance in mice. <i>Proceedings of the British Society of Animal Science</i> , 2007, 2007, 92-92.	0.0	3
89	Quantitative Trait Loci for Regional Adiposity in Mouse Lines Divergently Selected for Food Intake. <i>Obesity</i> , 2007, 15, 2994-3004.	3.0	7
90	A genetic investigation of various growth models to describe growth of lambs of two contrasting breeds <sup>1</sup> . <i>Journal of Animal Science</i> , 2006, 84, 2642-2654.	0.5	58

#	ARTICLE	IF	CITATIONS
91	In vivo prediction of internal fat weight in Scottish Blackface lambs, using computer tomography. <i>Journal of Animal Breeding and Genetics</i> , 2006, 123, 105-113.	2.0	18
92	Phenotypic characterisation of extreme growth-selected mouse lines: An important prerequisite for future QTL analysis. <i>Open Life Sciences</i> , 2006, 1, 345-375.	1.4	0
93	Testing selection indices for sustainable hill sheep production – lamb growth and carcass traits. <i>Animal Science</i> , 2006, 82, 445-453.	1.3	25
94	In vivo measurements of muscle volume by automatic image analysis of spiral computed tomography scans. <i>Animal Science</i> , 2006, 82, 545-553.	1.3	24
95	Microarray gene expression analysis of the Fob3b obesity QTL identifies positional candidate gene Sqle and perturbed cholesterol and glycolysis pathways. <i>Physiological Genomics</i> , 2005, 20, 224-232.	2.3	39
96	Mice with Low Metabolic Rates Are Not Susceptible to Weight Gain When Fed a High-Fat Diet. <i>Obesity</i> , 2005, 13, 556-566.	4.0	21
97	A paternally imprinted QTL for mature body mass on mouse Chromosome 8. <i>Mammalian Genome</i> , 2005, 16, 567-577.	2.2	17
98	Effects of the Compact Mutant Myostatin Allele Mstn Cmpt-dl1Abc Introgressed into a High Growth Mouse Line on Skeletal Muscle Cellularity. <i>Journal of Muscle Research and Cell Motility</i> , 2005, 26, 103-112.	2.0	42
99	Relationships between quantitative and reproductive fitness traits in animals. <i>Philosophical Transactions of the Royal Society B: Biological Sciences</i> , 2005, 360, 1489-1502.	4.0	32
100	A Polygenic Model of the Metabolic Syndrome With Reduced Circulating and Intra-Adipose Glucocorticoid Action. <i>Diabetes</i> , 2005, 54, 3371-3378.	0.6	62
101	GENETICS OF BODY COMPOSITION AND METABOLIC RATE. , 2005, , 131-160.		4
102	Marker-assisted introgression of the Compact mutant myostatin allele MstnCmpt-dl1Abc into a mouse line with extreme growth effects on body composition and muscularity. <i>Genetical Research</i> , 2004, 84, 161-173.	0.9	28
103	Growth selection in mice reveals conserved and redundant expression patterns of the insulin-like growth factor system. <i>General and Comparative Endocrinology</i> , 2004, 136, 248-259.	1.8	16
104	Identification and reciprocal introgression of a QTL affecting body mass in mice. <i>Genetics Selection Evolution</i> , 2004, 36, 577-91.	3.0	2
105	Genetic complexity of an obesity QTL (Fob3) revealed by detailed genetic mapping. <i>Mammalian Genome</i> , 2004, 15, 472-481.	2.2	37
106	Identification and reciprocal introgression of a QTL affecting body mass in mice. <i>Genetics Selection Evolution</i> , 2004, 36, 577-591.	3.0	1
107	Genetic-statistical analysis of growth in selected and unselected mouse lines. <i>Journal of Experimental Animal Science</i> , 2003, 42, 218-232.	0.5	22
108	Long-term divergent selection on body fatness in mice indicates a regulation system that is independent of leptin production and reception. <i>FASEB Journal</i> , 2003, 17, 85-87.	0.5	18

#	ARTICLE	IF	CITATIONS
109	Intrinsic properties of muscle satellite cells are changed in response to long-term selection of mice for different growth traits. <i>Cell and Tissue Research</i> , 2002, 310, 339-348.	2.9	15
110	Inbred lines of mice derived from long-term growth selected lines: unique resources for mapping growth genes. <i>Mammalian Genome</i> , 2001, 12, 678-686.	2.2	64
111	Thermoregulatory responses of two mouse <i>Mus musculus</i> strains selectively bred for high and low food intake. <i>Journal of Comparative Physiology B: Biochemical, Systemic, and Environmental Physiology</i> , 2001, 171, 661-668.	1.5	32
112	Characterization of a Major X-Linked Quantitative Trait Locus Influencing Body Weight of Mice. , 2001, 92, 355-357.		10
113	Inbred lines of mice derived from long-term growth selected lines: unique resources for mapping growth genes. <i>Mammalian Genome</i> , 2001, 012, 0678-0686.	2.2	12
114	Resting metabolic rate and morphology in mice ( <i>Mus musculus</i> ) selected for high and low food intake. <i>Journal of Experimental Biology</i> , 2001, 204, 777-784.	1.7	123
115	Resting metabolic rate and morphology in mice ( <i>Mus musculus</i> ) selected for high and low food intake. <i>Journal of Experimental Biology</i> , 2001, 204, 777-84.	1.7	85
116	Analysis of response to 20 generations of selection for body composition in mice: fit to infinitesimal model assumptions. <i>Genetics Selection Evolution</i> , 2000, 32, 3-21.	3.0	36
117	Mapping of obesity QTLs in a cross between mouse lines divergently selected on fat content. <i>Mammalian Genome</i> , 2000, 11, 2-7.	2.2	74
118	Polymerase chain reaction-restriction fragment length polymorphism (PCR-RFLP) assay for the mouse leptin receptor ( <i>Leprdb</i> ) mutation. <i>Laboratory Animals</i> , 1999, 33, 380-384.	1.0	24
119	Neuropeptide Y gene expression in lines of mice subjected to long-term divergent selection on fat content. <i>Journal of Molecular Endocrinology</i> , 1999, 23, 77-83.	2.5	6
120	Inbred lines of mice derived from long-term divergent selection on fat content and body weight. <i>Mammalian Genome</i> , 1999, 10, 645-648.	2.2	42
121	Leptin levels in lines of mice developed by long-term divergent selection on fat content. <i>Genetical Research</i> , 1999, 73, 37-44.	0.9	13
122	Role of growth hormone in the genetic change of mice divergently selected for body weight and fatness. <i>Genetical Research</i> , 1999, 74, 351-360.	0.9	14
123	Effects of thyroid hormone deficiency on mice selected for increased and decreased body weight and fatness. <i>Genetical Research</i> , 1998, 72, 39-53.	0.9	9
124	Long-term selection for protein amount over 70 generations in mice. <i>Genetical Research</i> , 1998, 72, 93-109.	0.9	28
125	Effects of thyroid hormone deficiency on mice selected for increased and decreased body weight and fatness. <i>Genetical Research</i> , 1998, 72, 59-72.	0.9	0
126	Effects of leptin administration on long-term selected fat mice. <i>Genetical Research</i> , 1997, 69, 215-225.	0.9	17



#	ARTICLE	IF	CITATIONS
127	Effects of leptin administration on lines of mice selected long-term for fatness. Genetical Research, 1997, 70, 79-89.	0.9	0
128	Effects of selection on food intake in the adult mouse. Journal of Animal Breeding and Genetics, 1997, 114, 419-434.	2.0	26
129	Prenatal growth in lines of mice selected for body weight. Journal of Animal Breeding and Genetics, 1996, 113, 535-543.	2.0	1
130	Analysis of a long-term selection experiment with an exponential model. Journal of Animal Breeding and Genetics, 1994, 111, 1-13.	2.0	17
131	Selection for litter weight on the 21st day after long-term selection for first litter performance in laboratory mice. Journal of Animal Breeding and Genetics, 1990, 107, 161-168.	2.0	2
132	Kreuzungswirkungen bei Merkmalen der Fruchtbarkeit und des Wachstums nach Langzeitselektion auf Erstwurfleistung bei der Labormaus. Journal of Animal Breeding and Genetics, 1990, 107, 241-248.	2.0	1
133	Zur Selektionswürdigkeit von Merkmalen der Muskelstruktur – Modellversuch mit Labormäusen. Journal of Animal Breeding and Genetics, 1989, 106, 208-216.	2.0	3
134	Use of X-Ray Computed Tomography (CT) in UK Sheep Production and Breeding. , 0, , .		10