

Suk Kyoon An

List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/9372451/publications.pdf>

Version: 2024-02-01

108
papers

3,139
citations

147801

31
h-index

182427

51
g-index

112
all docs

112
docs citations

112
times ranked

4637
citing authors

#	ARTICLE	IF	CITATIONS
1	Impact of data extraction errors in meta-analyses on the association between depression and peripheral inflammatory biomarkers: an umbrella review. <i>Psychological Medicine</i> , 2023, 53, 2017-2030.	4.5	4
2	The stress-vulnerability model on the path to schizophrenia: Interaction between BDNF methylation and schizotypy on the resting-state brain network. <i>NPJ Schizophrenia</i> , 2022, 8, .	3.6	0
3	Associations of Self-Consciousness with Insomnia Symptoms. <i>Chronobiology in Medicine</i> , 2021, 3, 25-30.	0.4	0
4	“Reading the Mind in the Eyes Test” Translated and Korean Versions. <i>Psychiatry Investigation</i> , 2021, 18, 295-303.	1.6	7
5	Empathic Tendency and Theory of Mind Skills in Young Individuals with Schizophrenia: Its Associations with Self-Reported Schizotypy and Executive Function. <i>Korean Journal of Schizophrenia Research</i> , 2021, 24, 26-35.	0.3	0
6	Increased resting-state cerebellar-cortical connectivity in breast cancer survivors with cognitive complaints after chemotherapy. <i>Scientific Reports</i> , 2021, 11, 12105.	3.3	6
7	Shame and guilt in youth at ultra-high risk for psychosis. <i>Comprehensive Psychiatry</i> , 2021, 108, 152241.	3.1	5
8	Sex-specific association of hair cortisol concentration with stress-related psychological factors in healthy young adults. <i>Biology of Sex Differences</i> , 2021, 12, 56.	4.1	4
9	Empathy and Theory of Mind in Ultra-High Risk for Psychosis: Relations With Schizotypy and Executive Function. <i>Psychiatry Investigation</i> , 2021, 18, 1109-1116.	1.6	2
10	Effect of Childhood Trauma on the Association Between Stress-Related Psychological Factors and Hair Cortisol Level in Young Adults. <i>Psychiatry Investigation</i> , 2021, 18, 1131-1136.	1.6	1
11	T57. IMPAIRED FACIAL EMOTION RECOGNITION IN INDIVIDUALS AT ULTRA-HIGH RISK FOR PSYCHOSIS: CORRELATIONS WITH SCHIZOTYPY AND PARANOID LEVEL. <i>Schizophrenia Bulletin</i> , 2020, 46, S253-S253.	4.3	0
12	Deactivation of anterior cingulate cortex during virtual social interaction in obsessive-compulsive disorder. <i>Psychiatry Research - Neuroimaging</i> , 2020, 304, 111154.	1.8	3
13	Impaired Facial Emotion Recognition in Individuals at Ultra-High Risk for Psychosis and Associations With Schizotypy and Paranoia Level. <i>Frontiers in Psychiatry</i> , 2020, 11, 577.	2.6	16
14	S73. RELATIONSHIP OF COGNITIVE ABILITY AND PERSONALITY TRAITS WITH HOSTILE ATTRIBUTION BIAS IN NONCLINICAL SUBJECTS: THEORY OF MIND AS A MEDIATOR. <i>Schizophrenia Bulletin</i> , 2020, 46, S62-S62.	4.3	0
15	Changes in resting-state brain connectivity following computerized cognitive behavioral therapy for insomnia in dialysis patients: A pilot study. <i>General Hospital Psychiatry</i> , 2020, 66, 24-29.	2.4	10
16	Reduced activation of the ventromedial prefrontal cortex during self-referential processing in individuals at ultra-high risk for psychosis. <i>Australian and New Zealand Journal of Psychiatry</i> , 2020, 54, 528-538.	2.3	6
17	Reading the Mind in the Eyes Test: Relationship with Neurocognition and Facial Emotion Recognition in Non-Clinical Youths. <i>Psychiatry Investigation</i> , 2020, 17, 835-839.	1.6	7
18	Sex differences in the association between social relationships and insomnia symptoms. <i>Journal of Clinical Sleep Medicine</i> , 2020, 16, 1871-1881.	2.6	0

#	ARTICLE	IF	CITATIONS
19	Factors Associated With Psychosocial Functioning and Outcome of Individuals With Recent-Onset Schizophrenia and at Ultra-High Risk for Psychosis. <i>Frontiers in Psychiatry</i> , 2019, 10, 459.	2.6	14
20	Associations of systemic inflammation with frontotemporal functional network connectivity and out-degree social-network size in community-dwelling older adults. <i>Brain, Behavior, and Immunity</i> , 2019, 79, 309-313.	4.1	8
21	Individualized Prediction of Transition to Psychosis in 1,676 Individuals at Clinical High Risk: Development and Validation of a Multivariable Prediction Model Based on Individual Patient Data Meta-Analysis. <i>Frontiers in Psychiatry</i> , 2019, 10, 345.	2.6	29
22	Reduced DNA Methylation of the Oxytocin Receptor Gene Is Associated With Anhedonia-Asociality in Women With Recent-Onset Schizophrenia and Ultra-high Risk for Psychosis. <i>Schizophrenia Bulletin</i> , 2019, 45, 1279-1290.	4.3	27
23	Population-based dementia prediction model using Korean public health examination data: A cohort study. <i>PLoS ONE</i> , 2019, 14, e0211957.	2.5	9
24	Oxytocin receptor gene variants are associated with emotion recognition and resilience, but not with false-belief reasoning performance in healthy young Korean volunteers. <i>CNS Neuroscience and Therapeutics</i> , 2019, 25, 519-526.	3.9	10
25	Psychotic conversion of individuals at ultra-high risk for psychosis: The potential roles of schizotypy and basic symptoms. <i>Microbial Biotechnology</i> , 2019, 13, 546-554.	1.7	22
26	Suicidal ideation in individuals at ultra-high risk for psychosis and its association with suspiciousness independent of depression. <i>Microbial Biotechnology</i> , 2019, 13, 539-545.	1.7	13
27	Cox Proportional Hazard Regression Versus a Deep Learning Algorithm in the Prediction of Dementia: An Analysis Based on Periodic Health Examination. <i>JMIR Medical Informatics</i> , 2019, 7, e13139.	2.6	16
28	The Associations between Anger Expression and Insomnia among Community-Dwelling Older Adults in Korea. <i>Chronobiology in Medicine</i> , 2019, 1, 86-93.	0.4	0
29	Coping styles in individuals at ultra-high risk for psychosis: Associations with cognitive appraisals. <i>Psychiatry Research</i> , 2018, 264, 162-168.	3.3	11
30	Genetic variation in cytokine genes and risk for transition to psychosis among individuals at ultra-high risk. <i>Schizophrenia Research</i> , 2018, 195, 589-590.	2.0	6
31	Ultra-High Risk for Psychosis : Clinical Characteristics and Diagnosis. <i>Journal of Korean Neuropsychiatric Association</i> , 2018, 57, 210.	0.5	2
32	Fragile Self and Malevolent Others: Biased Attribution Styles in Individuals at Ultra-High Risk for Psychosis. <i>Psychiatry Investigation</i> , 2018, 15, 796-804.	1.6	7
33	Aberrant cerebro-cerebellar functional connectivity and minimal self-disturbance in individuals at ultra-high risk for psychosis and with first-episode schizophrenia. <i>Schizophrenia Research</i> , 2018, 202, 138-140.	2.0	19
34	Cognitive Behavioral Therapy for Insomnia Reduces Hypnotic Prescriptions. <i>Psychiatry Investigation</i> , 2018, 15, 499-504.	1.6	11
35	Aberrant Tendency of Noncurrent Emotional Experiences in Individuals at Ultra-High Risk for Psychosis. <i>Psychiatry Investigation</i> , 2018, 15, 876-883.	1.6	8
36	Predicting Working Memory Capacity in Older Subjects Using Quantitative Electroencephalography. <i>Psychiatry Investigation</i> , 2018, 15, 790-795.	1.6	0

#	ARTICLE	IF	CITATIONS
37	Prevalence of Metabolic Syndrome in Patients with Schizophrenia in Korea: A Multicenter Nationwide Cross-Sectional Study. <i>Psychiatry Investigation</i> , 2017, 14, 44.	1.6	27
38	Impaired Social and Role Function in Ultra-High Risk for Psychosis and First-Episode Schizophrenia: Its Relations with Negative Symptoms. <i>Psychiatry Investigation</i> , 2017, 14, 186.	1.6	25
39	Impaired Social and Role Function in Ultra-High Risk for Psychosis and First-Episode Schizophrenia: Its Relations with Negative Symptoms. <i>Psychiatry Investigation</i> , 2017, 14, 539.	1.6	31
40	Violent behavior in individuals with schizophrenia. <i>Journal of the Korean Medical Association</i> , 2016, 59, 947.	0.3	0
41	PM456. Aberrant cortico-cerebellar connectivity of the default mode network in individuals at ultra-high risk for psychosis: a resting-state fMRI study. <i>International Journal of Neuropsychopharmacology</i> , 2016, 19, 65-66.	2.1	0
42	PM462. Impaired self-referential processing in patients with first-episode schizophrenia: an event-related potential study. <i>International Journal of Neuropsychopharmacology</i> , 2016, 19, 68-68.	2.1	0
43	Associations between actigraphy-assessed sleep, inflammatory markers, and insulin resistance in the Midlife Development in the United States (MIDUS) study. <i>Sleep Medicine</i> , 2016, 27-28, 72-79.	1.6	32
44	Experiential pleasure deficits in the prodrome: A study of emotional experiences in individuals at ultra-high risk for psychosis and recent-onset schizophrenia. <i>Comprehensive Psychiatry</i> , 2016, 68, 209-216.	3.1	20
45	Common variants of HTR3 genes are associated with obsessive-compulsive disorder and its phenotypic expression. <i>Scientific Reports</i> , 2016, 6, 32564.	3.3	10
46	Heterogeneity of Psychosis Risk Within Individuals at Clinical High Risk. <i>JAMA Psychiatry</i> , 2016, 73, 113.	11.0	354
47	Frontostriatal Connectivity Changes in Major Depressive Disorder After Repetitive Transcranial Magnetic Stimulation. <i>Journal of Clinical Psychiatry</i> , 2016, 77, e1137-e1143.	2.2	54
48	Relationships between chronotypes and affective temperaments in healthy young adults. <i>Journal of Affective Disorders</i> , 2015, 175, 256-259.	4.1	65
49	Neurocognitive impairments in individuals at ultra-high risk for psychosis: Who will really convert?. <i>Australian and New Zealand Journal of Psychiatry</i> , 2015, 49, 462-470.	2.3	54
50	Impaired facial emotion recognition in individuals at ultra-high risk for psychosis and with first-episode schizophrenia, and their associations with neurocognitive deficits and self-reported schizotypy. <i>Schizophrenia Research</i> , 2015, 165, 60-65.	2.0	45
51	Emotional Dysregulation, Attributional Bias, Neurocognitive Impairment in Individuals at Ultra-High Risk for Psychosis and with Schizophrenia : Its Association with Paranoia. <i>Korean Journal of Schizophrenia Research</i> , 2014, 17, 63.	0.3	2
52	Antipsychotic polypharmacy and high-dose prescription in schizophrenia: a 5-year comparison. <i>Australian and New Zealand Journal of Psychiatry</i> , 2014, 48, 52-60.	2.3	51
53	Reduced Binding Potential of GABA-A/Benzodiazepine Receptors in Individuals at Ultra-high Risk for Psychosis: An [18F]-Fluoroflumazenil Positron Emission Tomography Study. <i>Schizophrenia Bulletin</i> , 2014, 40, 548-557.	4.3	33
54	Distinct functional connectivity of limbic network in the washing type obsessive-compulsive disorder. <i>Progress in Neuro-Psychopharmacology and Biological Psychiatry</i> , 2014, 53, 149-155.	4.8	19

#	ARTICLE	IF	CITATIONS
55	Impact of Delirium on Clinical Outcomes in Intensive Care Unit Patients: An Observational Study in a Korean General Hospital. <i>Journal of Korean Neuropsychiatric Association</i> , 2014, 53, 418.	0.5	7
56	Theory of Mind as a Mediator of Reasoning and Facial Emotion Recognition: Findings from 200 Healthy People. <i>Psychiatry Investigation</i> , 2014, 11, 105.	1.6	15
57	Theta- γ coupling during a working memory task as compared to a simple vigilance task. <i>Neuroscience Letters</i> , 2013, 532, 39-43.	2.1	41
58	Small-world networks in individuals at ultra-high risk for psychosis and first-episode schizophrenia during a working memory task. <i>Neuroscience Letters</i> , 2013, 535, 35-39.	2.1	25
59	Temperament and character in individuals at ultra-high risk for psychosis and with first-episode schizophrenia: Associations with psychopathology, psychosocial functioning, and aspects of psychological health. <i>Comprehensive Psychiatry</i> , 2013, 54, 1161-1168.	3.1	36
60	Genetic Influence of COMT and BDNF Gene Polymorphisms on Resilience in Healthy College Students. <i>Neuropsychobiology</i> , 2013, 68, 174-180.	1.9	29
61	APOLIPOPROTEIN E GENE POLYMORPHISM, ALCOHOL USE, AND THEIR INTERACTIONS IN COMBAT-RELATED POSTTRAUMATIC STRESS DISORDER. <i>Depression and Anxiety</i> , 2013, 30, 1194-1201.	4.1	35
62	Neural Evidence for Emotional Involvement in Pathological Alcohol Craving. <i>Alcohol and Alcoholism</i> , 2013, 48, 288-294.	1.6	18
63	The relationship between psychosocial functioning and resilience and negative symptoms in individuals at ultra-high risk for psychosis. <i>Australian and New Zealand Journal of Psychiatry</i> , 2013, 47, 762-771.	2.3	74
64	Attributional Style in Healthy Persons: Its Association with 'Theory of Mind' Skills. <i>Psychiatry Investigation</i> , 2013, 10, 34.	1.6	17
65	Increased Intra-Individual Variability of Cognitive Processing in Subjects at Risk Mental State and Schizophrenia Patients. <i>PLoS ONE</i> , 2013, 8, e78354.	2.5	34
66	Aberrant Auditory Processing in Schizophrenia and in Subjects at Ultra-High-Risk for Psychosis. <i>Schizophrenia Bulletin</i> , 2012, 38, 1258-1267.	4.3	42
67	Gamma oscillatory activity in relation to memory ability in older adults. <i>International Journal of Psychophysiology</i> , 2012, 86, 58-65.	1.0	24
68	FKBP5 polymorphisms as vulnerability to anxiety and depression in patients with advanced gastric cancer: A controlled and prospective study. <i>Psychoneuroendocrinology</i> , 2012, 37, 1569-1576.	2.7	40
69	Differential alteration of automatic semantic processing in treated patients affected by bipolar mania and schizophrenia: An N400 study. <i>Progress in Neuro-Psychopharmacology and Biological Psychiatry</i> , 2012, 38, 194-200.	4.8	25
70	Psychometric analysis of the Korean version of the Disgust Scale- γ Revised. <i>Comprehensive Psychiatry</i> , 2012, 53, 648-655.	3.1	11
71	Early intervention in psychosis: Insights from Korea. <i>Asian Journal of Psychiatry</i> , 2012, 5, 98-105.	2.0	46
72	Regional cortical thinning in subjects with high genetic loading for schizophrenia. <i>Schizophrenia Research</i> , 2012, 141, 197-203.	2.0	42

#	ARTICLE	IF	CITATIONS
73	Regional Brain Atrophy and Functional Disconnection in Broca's Area in Individuals at Ultra-High Risk for Psychosis and Schizophrenia. PLoS ONE, 2012, 7, e51975.	2.5	44
74	Working Memory Deficits in Ultra-High Risk for Psychosis and Schizophrenia. Korean Journal of Schizophrenia Research, 2012, 15, 66.	0.3	3
75	Neurocognitive performance in subjects at ultrahigh risk for schizophrenia: a comparison with first-episode schizophrenia. Comprehensive Psychiatry, 2011, 52, 33-40.	3.1	31
76	Emotional Priming With Facial Exposures in Euthymic Patients With Bipolar Disorder. Journal of Nervous and Mental Disease, 2011, 199, 971-977.	1.0	5
77	Coping Strategies and Their Relationship to Psychopathologies in People at Ultra High-Risk for Psychosis and With Schizophrenia. Journal of Nervous and Mental Disease, 2011, 199, 106-110.	1.0	32
78	Biased to contempt and deficits of the negative facial emotion recognition in patients with schizophrenia. International Clinical Psychopharmacology, 2011, 26, e99.	1.7	0
79	Clinical efficacy of individual cognitive therapy in reducing psychiatric symptoms in people at ultra-high risk for psychosis. Microbial Biotechnology, 2011, 5, 174-178.	1.7	21
80	Differential priming effect for subliminal fear and disgust facial expressions. Attention, Perception, and Psychophysics, 2011, 73, 473-481.	1.3	7
81	Longitudinal Patterns of Social Functioning and Conversion to Psychosis in Subjects at Ultra-High Risk. Australian and New Zealand Journal of Psychiatry, 2011, 45, 763-770.	2.3	33
82	Associated Factors of Quality of Life in First-Episode Schizophrenia Patients. Psychiatry Investigation, 2011, 8, 201.	1.6	25
83	Genetic fuzzy classifier for sleep stage identification. Computers in Biology and Medicine, 2010, 40, 629-634.	7.0	76
84	What factors are related to delayed treatment in individuals at high risk for psychosis?. Microbial Biotechnology, 2010, 4, 124-131.	1.7	19
85	A descriptive study of pathways to care of high risk for psychosis adolescents in Korea. Microbial Biotechnology, 2010, 4, 119-123.	1.7	18
86	Structural Brain Alterations in Individuals at Ultra-high Risk for Psychosis: A Review of Magnetic Resonance Imaging Studies and Future Directions. Journal of Korean Medical Science, 2010, 25, 1700.	2.5	48
87	Psychotic Features as the First Manifestation of 22q11.2 Deletion Syndrome. Psychiatry Investigation, 2010, 7, 72.	1.6	8
88	Attribution bias in ultra-high risk for psychosis and first-episode schizophrenia. Schizophrenia Research, 2010, 118, 54-61.	2.0	115
89	Association of DRD4 and COMT Polymorphisms with Disgust Sensitivity in Healthy Volunteers. Neuropsychobiology, 2010, 61, 105-112.	1.9	20
90	Decreased P3 amplitudes elicited by negative facial emotion in manic patients: Selective deficits in emotional processing. Neuroscience Letters, 2010, 481, 92-96.	2.1	26

#	ARTICLE	IF	CITATIONS
91	Reduced visual P300 amplitudes in individuals at ultra-high risk for psychosis and first-episode schizophrenia. <i>Neuroscience Letters</i> , 2010, 486, 156-160.	2.1	37
92	Perception bias of disgust in ambiguous facial expressions in obsessive-compulsive disorder. <i>Psychiatry Research</i> , 2010, 178, 126-131.	3.3	22
93	The Reliability and Validity of the Korean Version of the Structured Interview for Prodromal Syndrome. <i>Psychiatry Investigation</i> , 2010, 7, 257.	1.6	57
94	To discard or not to discard: the neural basis of hoarding symptoms in obsessive-compulsive disorder. <i>Molecular Psychiatry</i> , 2009, 14, 318-331.	7.9	137
95	Individual differences in disgust sensitivity modulate neural responses to aversive/disgusting stimuli. <i>European Journal of Neuroscience</i> , 2008, 27, 3050-3058.	2.6	85
96	Is a neutral face really evaluated as being emotionally neutral?. <i>Psychiatry Research</i> , 2008, 157, 77-85.	3.3	146
97	Alteration of brain metabolites in young alcoholics without structural changes. <i>NeuroReport</i> , 2007, 18, 1511-1514.	1.2	59
98	Neural Responses to Facial Expressions of Disgust but not Fear are Modulated by Washing Symptoms in OCD. <i>Biological Psychiatry</i> , 2007, 61, 1072-1080.	1.3	82
99	Sex Differences in Neural Responses to Disgusting Visual Stimuli: Implications for Disgust-Related Psychiatric Disorders. <i>Biological Psychiatry</i> , 2007, 62, 464-471.	1.3	53
100	Shape deformation of the insula in schizophrenia. <i>NeuroImage</i> , 2006, 32, 220-227.	4.2	25
101	Neural substrates associated with evaluative processing during co-activation of positivity and negativity: A PET investigation. <i>Biological Psychology</i> , 2006, 73, 253-261.	2.2	16
102	Behavioral evidence of blunted and inappropriate affective responses in schizophrenia: Lack of a "negativity bias". <i>Psychiatry Research</i> , 2006, 142, 53-66.	3.3	17
103	Aberrantly flattened responsivity to emotional pictures in paranoid schizophrenia. <i>Psychiatry Research</i> , 2006, 143, 135-145.	3.3	17
104	Differences of Photographs Inducing Craving Between Alcoholics and Non-alcoholics. <i>Yonsei Medical Journal</i> , 2006, 47, 491.	2.2	20
105	Greater Impairment in Negative Emotion Evaluation Ability in Patients with Paranoid Schizophrenia. <i>Yonsei Medical Journal</i> , 2006, 47, 343.	2.2	12
106	Increased P3 Amplitudes Induced by Alcohol-Related Pictures in Patients With Alcohol Dependence. <i>Alcoholism: Clinical and Experimental Research</i> , 2004, 28, 1317-1323.	2.4	72
107	Reduced P3 amplitudes by negative facial emotional photographs in schizophrenia. <i>Schizophrenia Research</i> , 2003, 64, 125-135.	2.0	45
108	Positive and negative symptoms and regional cerebral perfusion in antipsychotic-naive schizophrenic patients: a high-resolution SPECT study. <i>Psychiatry Research - Neuroimaging</i> , 1999, 90, 159-168.	1.8	35