Ulises De la Cruz-Mosso

List of Publications by Year in descending order

Source: https://exaly.com/author-pdf/9373742/publications.pdf Version: 2024-02-01



#	Article	IF	CITATIONS
1	Association of Vitamin D Metabolism Gene Polymorphisms with Autoimmunity: Evidence in Population Genetic Studies. International Journal of Molecular Sciences, 2020, 21, 9626.	4.1	49
2	Macrophage migration inhibitory factor: Association of â^'794 CATT5–8 and â^'173 G>C polymorphisms with TNF-α in systemic lupus erythematosus. Human Immunology, 2014, 75, 433-439.	2.4	39
3	MIF promotes a differential Th1/Th2/Th17 inflammatory response in human primary cell cultures: Predominance of Th17 cytokine profile in PBMC from healthy subjects and increase of IL-6 and TNF-α in PBMC from active SLE patients. Cellular Immunology, 2018, 324, 42-49.	3.0	37
4	Circulating CD36 and oxLDL levels are associated with cardiovascular risk factors in young subjects. BMC Cardiovascular Disorders, 2014, 14, 54.	1.7	34
5	MIF functional polymorphisms (-794 CATT5-8 and -173 G>C) are associated with MIF serum levels, severity and progression in male multiple sclerosis from western Mexican population. Journal of Neuroimmunology, 2018, 320, 117-124.	2.3	26
6	Relationship of Excess Weight with Clinical Activity and Dietary Intake Deficiencies in Systemic Lupus Erythematosus Patients. Nutrients, 2019, 11, 2683.	4.1	25
7	Functional effects of vitamin D: From nutrient to immunomodulator. Critical Reviews in Food Science and Nutrition, 2022, 62, 3042-3062.	10.3	22
8	Association of cardiometabolic risk status with clinical activity and damage in systemic lupus erythematosus patients: A cross-sectional study. Clinical Immunology, 2021, 222, 108637.	3.2	15
9	Association between the â^'794 (CATT) _{5–8} <i>MIF</i> Gene Polymorphism and Susceptil to Acute Coronary Syndrome in a Western Mexican Population. Journal of Immunology Research, 2014, 2014, 1-5.	oility 2.2	14
10	PAI-1 mRNA expression and plasma level in rheumatoid arthritis: relationship with 4G/5G PAI-1 polymorphism. Rheumatology International, 2012, 32, 3951-3956.	3.0	13
11	Expression of MIF and TNFA in psoriatic arthritis: relationship with Th1/Th2/Th17 cytokine profiles and clinical variables. Clinical and Experimental Medicine, 2018, 18, 229-235.	3.6	13
12	Relationship of metabolic syndrome and its components with -844 G/A and HindIII C/G PAI-1 gene polymorphisms in Mexican children. BMC Pediatrics, 2012, 12, 41.	1.7	12
13	High expression of interleukine-1 receptor antagonist in rheumatoid arthritis: Association with IL1RN*2/2 genotype. Autoimmunity, 2017, 50, 468-475.	2.6	11
14	Functional MIF promoter haplotypes modulate Th17-related cytokine expression in peripheral blood mononuclear cells from control subjects and rheumatoid arthritis patients. Cytokine, 2019, 115, 89-96.	3.2	11
15	CRP Serum Levels Are Associated with High Cardiometabolic Risk and Clinical Disease Activity in Systemic Lupus Erythematosus Patients. Journal of Clinical Medicine, 2022, 11, 1849.	2.4	11
16	A potential inflammatory role of IL-31 in psoriatic arthritis: A correlation with Th17 cytokine profile. International Journal of Immunopathology and Pharmacology, 2020, 34, 205873842090718.	2.1	10
17	MIF and TNFαserum levels in rheumatoid arthritis patients treated with disease-modifying antirheumatic drugs: a cross-sectional study. Immunopharmacology and Immunotoxicology, 2015, 37, 207-213.	2.4	9
18	Body adiposity but not insulin resistance is associated with -675 4G/5G polymorphism in the PAI-1 gene in a sample of Mexican children. Jornal De Pediatria, 2013, 89, 492-498.	2.0	7

#	Article	IF	CITATIONS
19	Th1/Th17 Cytokine Profile is Induced by Macrophage Migration Inhibitory Factor in Peripheral Blood Mononuclear Cells from Rheumatoid Arthritis Patients. Current Molecular Medicine, 2019, 18, 679-688.	1.3	7
20	Association of extrapituitary prolactin promoter polymorphism with disease susceptibility and anti-RNP antibodies in Mexican patients with systemic lupus erythematosus. Archives of Medical Science, 2018, 14, 1025-1032.	0.9	5
21	The â^675 4G/5G <i>PAI-1</i> polymorphism confers genetic susceptibility to systemic lupus erythematosus, its clinical manifestations, and comorbidities in Mexican-Mestizo population. Autoimmunity, 2020, 53, 71-77.	2.6	5
22	Presence of Adenovirus-36 DNA in Adipose Tissue of Women: Relationship with Adipocyte Morphology and the Expression of C/EBPβ and HIF-1α. Diabetes, Metabolic Syndrome and Obesity: Targets and Therapy, 2021, Volume 14, 477-486.	2.4	4
23	Influence of Diet and Levels of Zonulin, Lipopolysaccharide and C-Reactive Protein on Cardiometabolic Risk Factors in Young Subjects. Nutrients, 2021, 13, 4472.	4.1	4
24	Association of High Calcitriol Serum Levels and Its Hydroxylation Efficiency Ratio with Disease Risk in SLE Patients with Vitamin D Deficiency. Journal of Immunology Research, 2021, 2021, 1-16.	2.2	4
25	Association of 86Åbp variable number of tandem repeat (VNTR) polymorphism of interleukin-1 receptor antagonist (IL1RN) with susceptibility and clinical activity in rheumatoid arthritis. Clinical Rheumatology, 2017, 36, 1247-1252.	2.2	2
26	Macrophage migration inhibitory factor: A promising oncogenic serological biomarker for oral squamous cell carcinoma. International Journal of Immunopathology and Pharmacology, 2021, 35, 205873842110384.	2.1	2
27	Body adiposity but not insulin resistance is associated with -675 4G/5G polymorphism in the PAI-1 gene in a sample of Mexican children. Jornal De Pediatria (Versão Em Português), 2013, 89, 492-498.	0.2	0