

Nicholas R Medjeral-Thomas

List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/9380791/publications.pdf>

Version: 2024-02-01

19
papers

1,367
citations

623734

14
h-index

839539

18
g-index

20
all docs

20
docs citations

20
times ranked

1487
citing authors

#	ARTICLE	IF	CITATIONS
1	Can Urinary Complement Proteins Stratify Patients to Therapeutic Complement Inhibitors?. <i>Kidney International Reports</i> , 2022, 7, 939-941.	0.8	1
2	Complement and kidney disease, new insights. <i>Current Opinion in Nephrology and Hypertension</i> , 2021, 30, 310-316.	2.0	5
3	Longitudinal proteomic profiling of dialysis patients with COVID-19 reveals markers of severity and predictors of death. <i>ELife</i> , 2021, 10, .	6.0	58
4	O- and N-Glycosylation of Serum Immunoglobulin A is Associated with IgA Nephropathy and Glomerular Function. <i>Journal of the American Society of Nephrology: JASN</i> , 2021, 32, 2455-2465.	6.1	33
5	Protease inhibitor plasma concentrations associate with COVID-19 infection. <i>Oxford Open Immunology</i> , 2021, 2, iqab014.	2.8	7
6	New Insights into Epidemiology and Outcome of Bacterial Infection-Related Glomerulonephritis. <i>Clinical Journal of the American Society of Nephrology: CJASN</i> , 2021, 16, 1149-1151.	4.5	0
7	Complement activation in IgA nephropathy. <i>Seminars in Immunopathology</i> , 2021, 43, 679-690.	6.1	42
8	Complement in IgA Nephropathy: The Role of Complement in the Pathogenesis, Diagnosis, and Future Management of IgA Nephropathy. <i>Advances in Chronic Kidney Disease</i> , 2020, 27, 111-119.	1.4	13
9	The role of complement in IgA nephropathy. <i>Molecular Immunology</i> , 2019, 114, 123-132.	2.2	76
10	Glomerular Complement Factor H-Related Protein 5 (FHR5) Is Highly Prevalent in C3 Glomerulopathy and Associated With Renal Impairment. <i>Kidney International Reports</i> , 2019, 4, 1387-1400.	0.8	17
11	Glomerular membrane attack complex is not a reliable marker of ongoing C5 activation in lupus nephritis. <i>Kidney International</i> , 2019, 95, 655-665.	5.2	33
12	Progressive IgA Nephropathy Is Associated With Low Circulating Mannan-Binding Lectin-Associated Serine Protease-3 (MASP-3) and Increased Glomerular Factor H-Related Protein-5 (FHR5) Deposition. <i>Kidney International Reports</i> , 2018, 3, 426-438.	0.8	57
13	Circulating complement factor H-related protein 5 levels contribute to development and progression of IgA nephropathy. <i>Kidney International</i> , 2018, 94, 150-158.	5.2	57
14	Circulating complement factor H-related proteins 1 and 5 correlate with disease activity in IgA nephropathy. <i>Kidney International</i> , 2017, 92, 942-952.	5.2	99
15	The complement factor H-related proteins. <i>Immunological Reviews</i> , 2016, 274, 191-201.	6.0	54
16	Long-term Outcomes of Rituximab Therapy in Ocular Granulomatosis with Polyangiitis. <i>Ophthalmology</i> , 2015, 122, 1262-1268.	5.2	53
17	C3 Glomerulopathy. <i>Clinical Journal of the American Society of Nephrology: CJASN</i> , 2014, 9, 46-53.	4.5	192
18	A novel CFHR5 fusion protein causes C3 glomerulopathy in a family without Cypriot ancestry. <i>Kidney International</i> , 2014, 85, 933-937.	5.2	61

#	ARTICLE	IF	CITATIONS
19	C3 glomerulopathy: consensus report. <i>Kidney International</i> , 2013, 84, 1079-1089.	5.2	505