
## Donald J Macintyre

List of Publications by Year in descending order

Source: https://exaly.com/author-pdf/939371/publications.pdf Version: 2024-02-01



#	Article	IF	CITATIONS
1	Genome-wide association analyses identify 44 risk variants and refine the genetic architecture of major depression. Nature Genetics, 2018, 50, 668-681.	21.4	2,224
2	Genetic relationship between five psychiatric disorders estimated from genome-wide SNPs. Nature Genetics, 2013, 45, 984-994.	21.4	2,067
3	Genome-wide association study identifies 30 loci associated with bipolar disorder. Nature Genetics, 2019, 51, 793-803.	21.4	1,191
4	Collaborative genome-wide association analysis supports a role for ANK3 and CACNA1C in bipolar disorder. Nature Genetics, 2008, 40, 1056-1058.	21.4	1,102
5	Analysis of shared heritability in common disorders of the brain. Science, 2018, 360, .	12.6	1,085
6	Whole-genome association study of bipolar disorder. Molecular Psychiatry, 2008, 13, 558-569.	7.9	642
7	Genome-wide association study of more than 40,000 bipolar disorder cases provides new insights into the underlying biology. Nature Genetics, 2021, 53, 817-829.	21.4	629
8	Genome-wide association study of major depressive disorder: new results, meta-analysis, and lessons learned. Molecular Psychiatry, 2012, 17, 36-48.	7.9	405
9	Genome-wide association for major depressive disorder: a possible role for the presynaptic protein piccolo. Molecular Psychiatry, 2009, 14, 359-375.	7.9	354
10	Cohort Profile: Generation Scotland: Scottish Family Health Study (GS:SFHS). The study, its participants and their potential for genetic research on health and illness. International Journal of Epidemiology, 2013, 42, 689-700.	1.9	353
11	Wake-up call for British psychiatry. British Journal of Psychiatry, 2008, 193, 6-9.	2.8	183
12	Genome-wide Association for Major Depression Through Age at Onset Stratification: Major Depressive Disorder Working Group of the Psychiatric Genomics Consortium. Biological Psychiatry, 2017, 81, 325-335.	1.3	175
13	Common polygenic risk for autism spectrum disorder (ASD) is associated with cognitive ability in the general population. Molecular Psychiatry, 2016, 21, 419-425.	7.9	145
14	The Genetics of the Mood Disorder Spectrum: Genome-wide Association Analyses of More Than 185,000 Cases and 439,000 Controls. Biological Psychiatry, 2020, 88, 169-184.	1.3	137
15	Chromosomal abnormalities and mental illness. Molecular Psychiatry, 2003, 8, 275-287.	7.9	111
16	A Cytogenetic Abnormality and Rare Coding Variants Identify ABCA13 as a Candidate Gene in Schizophrenia, Bipolar Disorder, and Depression. American Journal of Human Genetics, 2009, 85, 833-846.	6.2	102
17	Epidemiology and Heritability of Major Depressive Disorder, Stratified by Age of Onset, Sex, and Illness Course in Generation Scotland: Scottish Family Health Study (CS:SFHS). PLoS ONE, 2015, 10, e0142197.	2.5	101
18	The Genetic Architecture of Depression in Individuals of East Asian Ancestry. JAMA Psychiatry, 2021, 78, 1258.	11.0	88

DONALD J MACINTYRE

#	Article	IF	CITATIONS
19	Does Childhood Trauma Moderate Polygenic Risk for Depression? A Meta-analysis of 5765 Subjects From the Psychiatric Genomics Consortium. Biological Psychiatry, 2018, 84, 138-147.	1.3	87
20	Genome-wide by environment interaction studies of depressive symptoms and psychosocial stress in UK Biobank and Generation Scotland. Translational Psychiatry, 2019, 9, 14.	4.8	87
21	An Analysis of Two Genome-wide Association Meta-analyses Identifies a New Locus for Broad Depression Phenotype. Biological Psychiatry, 2017, 82, 322-329.	1.3	84
22	Self-reported medication use validated through record linkage to national prescribing data. Journal of Clinical Epidemiology, 2018, 94, 132-142.	5.0	75
23	708 Common and 2010 rare DISC1 locus variants identified in 1542 subjects: analysis for association with psychiatric disorder and cognitive traits. Molecular Psychiatry, 2014, 19, 668-675.	7.9	59
24	The association between lower educational attainment and depression owing to shared genetic effects? Results in ~25 000 subjects. Molecular Psychiatry, 2015, 20, 735-743.	7.9	59
25	Association of polygenic score for major depression with response to lithium in patients with bipolar disorder. Molecular Psychiatry, 2021, 26, 2457-2470.	7.9	44
26	A loss of mature microglial markers without immune activation in schizophrenia. Glia, 2021, 69, 1251-1267.	4.9	43
27	A validation of the diathesis-stress model for depression in Generation Scotland. Translational Psychiatry, 2019, 9, 25.	4.8	40
28	Shared Genetics and Couple-Associated Environment Are Major Contributors to the Risk of Both Clinical and Self-Declared Depression. EBioMedicine, 2016, 14, 161-167.	6.1	32
29	A Combined Pathway and Regional Heritability Analysis Indicates NETRIN1 Pathway Is Associated With Major Depressive Disorder. Biological Psychiatry, 2017, 81, 336-346.	1.3	32
30	Polygenic risk for alcohol dependence associates with alcohol consumption, cognitive function and social deprivation in a populationâ€based cohort. Addiction Biology, 2016, 21, 469-480.	2.6	27
31	Classical Human Leukocyte Antigen Alleles and C4 Haplotypes Are Not Significantly Associated With Depression. Biological Psychiatry, 2020, 87, 419-430.	1.3	27
32	Cohort profile for the STratifying Resilience and Depression Longitudinally (STRADL) study: A depression-focused investigation of Generation Scotland, using detailed clinical, cognitive, and neuroimaging assessments. Wellcome Open Research, 2019, 4, 185.	1.8	27
33	Genome-wide Regional Heritability Mapping Identifies a Locus Within the TOX2 Gene Associated With Major Depressive Disorder. Biological Psychiatry, 2017, 82, 312-321.	1.3	26
34	Major depressive disorder and current psychological distress moderate the effect of polygenic risk for obesity on body mass index. Translational Psychiatry, 2015, 5, e592-e592.	4.8	24
35	Facial emotion recognition in Scottish prisoners. International Journal of Law and Psychiatry, 2012, 35, 57-61.	0.9	21
36	Association of GPR50, an X-linked orphan G protein-coupled receptor, and affective disorder in an independent sample of the Scottish population. Neuroscience Letters, 2010, 475, 169-173.	2.1	18

DONALD J MACINTYRE

#	Article	IF	CITATIONS
37	Alzheimer's disease risk factor complement receptor 1 is associated with depression. Neuroscience Letters, 2012, 510, 6-9.	2.1	18
38	Description of arts therapies practice with adults suffering from depression in the UK: Quantitative results from the nationwide survey. Arts in Psychotherapy, 2013, 40, 458-464.	1.2	17
39	Association of Whole-Genome and NETRIN1 Signaling Pathway–Derived Polygenic Risk Scores for Major Depressive Disorder and White Matter Microstructure in the UK Biobank. Biological Psychiatry: Cognitive Neuroscience and Neuroimaging, 2019, 4, 91-100.	1.5	16
40	Digital Support Platform: a qualitative research study investigating the feasibility of an internet-based, postdiagnostic support platform for families living with dementia. BMJ Open, 2018, 8, e020281.	1.9	15
41	Association of DISC1 variants with age of onset in a population-based sample of recurrent major depression. Molecular Psychiatry, 2013, 18, 745-747.	7.9	14
42	Genetic variation in Hyperpolarization-activated cyclic nucleotide-gated channels and its relationship with neuroticism, cognition and risk of depression. Frontiers in Genetics, 2012, 3, 116.	2.3	12
43	Cohort profile for the STratifying Resilience and Depression Longitudinally (STRADL) study: A depression-focused investigation of Generation Scotland, using detailed clinical, cognitive, and neuroimaging assessments. Wellcome Open Research, 0, 4, 185.	1.8	12
44	Art psychotherapy practice with adults who suffer from depression in the UK: Qualitative findings from a depression-specific questionnaire. Arts in Psychotherapy, 2014, 41, 563-569.	1.2	11
45	Current Versus Lifetime Depression, APOE Variation, and Their Interaction on Cognitive Performance in Younger and Older Adults. Psychosomatic Medicine, 2015, 77, 480-492.	2.0	11
46	Pharmaco-epidemiology of antidepressant exposure in a UK cohort record-linkage study. Journal of Psychopharmacology, 2019, 33, 482-493.	4.0	11
47	Language function following preterm birth: prediction using machine learning. Pediatric Research, 2022, 92, 480-489.	2.3	11
48	Altered metabolic parameters in association with antipsychotic medication use in diabetes: A population based case-control study. Psychoneuroendocrinology, 2016, 66, 214-220.	2.7	10
49	Increased number of T-lymphocytes in post-mortem brain tissue of patients with schizophrenia Schizophrenia Research, 2020, 216, 526-528.	2.0	10
50	Attitudes of patients and family members towards implantable psychiatric medication. Schizophrenia Research, 2008, 105, 279-286.	2.0	9
51	Description of arts therapies practice with adults suffering from depression in the UK: Qualitative findings from the nationwide survey. Arts in Psychotherapy, 2014, 41, 535-544.	1.2	8
52	Evaluation of a brief art psychotherapy group for adults suffering from mild to moderate depression: Pilot pre, post and follow-up study. International Journal of Art Therapy: Inscape, 2017, 22, 106-117.	1.6	7
53	Digital Access in Working-Age and Older Adults and Their Caregivers Attending Psychiatry Outpatient Clinics: Quantitative Survey. JMIR Aging, 2018, 1, e4.	3.0	2
54	Treatment resistant depression in the UK: sub-analysis of a European real-world evidence study. BJPsych Open, 2021, 7, S55-S55.	0.7	0

#	Article	IF	CITATIONS
55	A GP guide to schizophrenia. Practitioner, 2003, 247, 692-4, 696, 698 passim.	0.3	0