

# Gang Li

## List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/9403697/publications.pdf>

Version: 2024-02-01

16  
papers

313  
citations

933447

10  
h-index

996975

15  
g-index

16  
all docs

16  
docs citations

16  
times ranked

259  
citing authors

#	ARTICLE	IF	CITATIONS
1	Population-specific age estimation in Black Americans and Chinese people based on pulp chamber volume of first molars from cone beam computed tomography. <i>International Journal of Legal Medicine</i> , 2022, 136, 811-819.	2.2	5
2	Diagnostic efficacy of CBCT, MRI, and CBCT-MRI fused images in distinguishing articular disc calcification from loose body of temporomandibular joint. <i>Clinical Oral Investigations</i> , 2021, 25, 1907-1914.	3.0	14
3	Age estimation based on 3D pulp chamber segmentation of first molars from cone-beam computed tomography by integrated deep learning and level set. <i>International Journal of Legal Medicine</i> , 2021, 135, 365-373.	2.2	17
4	Specific oral and maxillofacial identifiers in panoramic radiographs used for human identification. <i>Journal of Forensic Sciences</i> , 2021, 66, 910-918.	1.6	8
5	Quantitative assessment of condyle positional changes before and after orthognathic surgery based on fused 3D images from cone beam computed tomography. <i>Clinical Oral Investigations</i> , 2020, 24, 2663-2672.	3.0	15
6	Potential biological damage of human peripheral blood lymphocytes induced by computed tomography examination of the oromaxillofacial region. <i>Oral Surgery, Oral Medicine, Oral Pathology and Oral Radiology</i> , 2020, 130, 708-716.	0.4	0
7	DNA double-strand breaks of human peripheral blood lymphocyte induced by CT examination of oral and maxillofacial region. <i>Clinical Oral Investigations</i> , 2020, 24, 4617-4624.	3.0	3
8	Congenital infiltrating lipomatosis of the face: A subtype of hemifacial hyperplasia. <i>International Journal of Pediatric Otorhinolaryngology</i> , 2019, 125, 107-112.	1.0	15
9	A three-dimensional study of hemimandibular hyperplasia, hemimandibular elongation, solitary condylar hyperplasia, simple mandibular asymmetry and condylar osteoma or osteochondroma. <i>Journal of Cranio-Maxillo-Facial Surgery</i> , 2019, 47, 1665-1675.	1.7	16
10	Effect of fracture orientation on detection accuracy of vertical root fractures in non-endodontically treated teeth using cone beam computed tomography. <i>Clinical Oral Investigations</i> , 2019, 23, 4433-4439.	3.0	6
11	Buccal mucosa cell damage in individuals following dental X-ray examinations. <i>Scientific Reports</i> , 2018, 8, 2509.	3.3	11
12	Age estimation based on pulp cavity/chamber volume of 13 types of tooth from cone beam computed tomography images. <i>International Journal of Legal Medicine</i> , 2016, 130, 1159-1167.	2.2	71
13	Age estimation based on pulp chamber volume of first molars from cone-beam computed tomography images. <i>Forensic Science International</i> , 2015, 253, 133.e1-133.e7.	2.2	78
14	Detection accuracy of proximal caries by phosphor plate and cone-beam computerized tomography images scanned with different resolutions. <i>Clinical Oral Investigations</i> , 2012, 16, 1015-1021.	3.0	23
15	Diagnostic accuracy of proximal caries by digital radiographs: an in vivo and in vitro comparative study. <i>Oral Surgery Oral Medicine Oral Pathology Oral Radiology and Endodontics</i> , 2010, 109, 463-467.	1.4	12
16	Measurement accuracy of marginal bone level in digital radiographs with and without color coding. <i>Acta Odontologica Scandinavica</i> , 2007, 65, 254-258.	1.6	19