

# Jason Jungsik Song

## List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/941485/publications.pdf>

Version: 2024-02-01

156  
papers

1,128  
citations

516710

16  
h-index

642732

23  
g-index

167  
all docs

167  
docs citations

167  
times ranked

1545  
citing authors

#	ARTICLE	IF	CITATIONS
1	Birmingham Vasculitis Activity Score und der Short Form 36-Item Health Survey als Prädiktoren aktueller depressiver Störungen bei Patienten mit antineutrophile-zytoplasmatische-Antikörper-assoziiierter Vaskulitis während der SARS-CoV-2-Pandemie. <i>Zeitschrift Fur Rheumatologie</i> , 2024, 83, 222-229.	1.0	1
2	Reclassification of previously diagnosed GPA patients using the 2022 ACR/EULAR classification criteria. <i>Rheumatology</i> , 2023, 62, 1179-1186.	1.9	8
3	2019 American College of Rheumatology/European League Against Rheumatism classification criteria for IgG4-related disease by Wallace et al. <i>Annals of the Rheumatic Diseases</i> , 2022, 81, e179-e179.	0.9	10
4	Nutrition Risk Index Score at Diagnosis Can Effectively Predict Poor Prognosis in Patients With Antineutrophil Cytoplasmic Antibody-Associated Vasculitis. , 2022, 32, 423-431.		4
5	Drug Survival of Biologic Therapy in Elderly Patients With Rheumatoid Arthritis Compared With Nonelderly Patients. <i>Journal of Clinical Rheumatology</i> , 2022, 28, e81-e88.	0.9	4
6	A retrospective analysis of antineutrophil cytoplasmic antibody-associated vasculitis aiming for an equation prediction end-stage renal disease. <i>Clinical Rheumatology</i> , 2022, 41, 773-781.	2.2	5
7	Association Between Idiopathic Cutaneous Leukocytoclastic Angiitis and ANCA-negative Microscopic Polyangiitis. <i>Journal of Rheumatic Diseases</i> , 2022, 29, 40-45.	1.1	0
8	Clinical application of low erythrocyte sedimentation rate/high C-reactive protein to antineutrophil cytoplasmic antibody-associated vasculitis. <i>Journal of Clinical Laboratory Analysis</i> , 2022, 36, e24237.	2.1	2
9	Anti-Citrullinated Peptide Antibody Expression and Its Association with Clinical Features and Outcomes in Patients with Antineutrophil Cytoplasmic Antibody-Associated Vasculitis. <i>Medicina (Lithuania)</i> , 2022, 58, 558.	2.0	2
10	Effect of numbers of metabolic syndrome components on mortality in patients with antineutrophil cytoplasmic antibody-associated vasculitis with metabolic syndrome.. <i>Clinical and Experimental Rheumatology</i> , 2022, , .	0.8	0
11	IM156, a new AMPK activator, protects against polymicrobial sepsis. <i>Journal of Cellular and Molecular Medicine</i> , 2022, 26, 3378-3386.	3.6	1
12	Incidence and Patterns of Interstitial Lung Disease and Their Clinical Impact on Mortality in Patients with Antineutrophil Cytoplasmic Antibody-Associated Vasculitis: Korean Single-Centre Observational Study. <i>Journal of Immunology Research</i> , 2022, 2022, 1-7.	2.2	3
13	Modified Body Mass Index at Diagnosis is a Useful Predictor of Mortality in Patients With Antineutrophil Cytoplasmic Antibody-associated Vasculitis. <i>Journal of Rheumatic Diseases</i> , 2022, 29, 154-161.	1.1	0
14	Serum albumin, prealbumin, and ischemia-modified albumin levels in patients with ANCA-associated vasculitis: A prospective cohort study. <i>PLoS ONE</i> , 2022, 17, e0271055.	2.5	5
15	Clinical and imaging findings suggestive of histopathological immunoglobulin G4-related disease: a single-center retrospective study. <i>Clinical Rheumatology</i> , 2021, 40, 1423-1430.	2.2	2
16	Systemic inflammation response index predicts all-cause mortality in patients with antineutrophil cytoplasmic antibody-associated vasculitis. <i>International Urology and Nephrology</i> , 2021, 53, 1631-1638.	1.4	4
17	Association Between Serum Alarmin Levels and Disease-specific Indices in Patients With Anti-neutrophil Cytoplasmic Antibody-associated Vasculitis. <i>In Vivo</i> , 2021, 35, 1761-1768.	1.3	1
18	Fibrosis-5 predicts end-stage renal disease in patients with microscopic polyangiitis and granulomatosis with polyangiitis without substantial liver diseases. <i>Clinical and Experimental Medicine</i> , 2021, 21, 399-406.	3.6	4

#	ARTICLE	IF	CITATIONS
19	Fibrinogen to albumin ratio reflects the activity of antineutrophil cytoplasmic antibody-associated vasculitis. <i>Journal of Clinical Laboratory Analysis</i> , 2021, 35, e23731.	2.1	7
20	Correlation between serum cysteine-rich protein 61 and disease activity of antineutrophil cytoplasmic antibody-associated vasculitis. <i>Clinical Rheumatology</i> , 2021, 40, 3703-3710.	2.2	1
21	Reclassification of Korean patients with polymyositis and dermatomyositis based on the Bohan and Peter criteria by the 2017 European League Against Rheumatism/American College of Rheumatology classification criteria for adult and juvenile idiopathic inflammatory myopathies. <i>Korean Journal of Internal Medicine</i> , 2021, 36, 441-446.	1.7	3
22	Association between follistatin-related protein 1 and the functional status of patients with anti-neutrophil cytoplasmic antibody-associated vasculitis. <i>Chinese Medical Journal</i> , 2021, 134, 1168-1174.	2.3	0
23	Total Haemolytic Complement Activity at Diagnosis as an Indicator of the Baseline Activity of Antineutrophil Cytoplasmic Antibody-associated Vasculitis. <i>Journal of Rheumatic Diseases</i> , 2021, 28, 85-93.	1.1	1
24	Detection of intracellular monosodium urate crystals in gout synovial fluid using optical diffraction tomography. <i>Scientific Reports</i> , 2021, 11, 10019.	3.3	9
25	Clinical features of Korean elderly patients with antineutrophil cytoplasmic antibody-associated vasculitis. <i>Korean Journal of Internal Medicine</i> , 2021, 36, 731-741.	1.7	3
26	Novel mortality-predicting index at diagnosis can effectively predict all-cause mortality in patients with antineutrophil cytoplasmic antibody-associated vasculitis. <i>Journal of Clinical Laboratory Analysis</i> , 2021, 35, e23885.	2.1	4
27	Clinical significance of large unstained cell count in estimating the current activity of antineutrophil cytoplasmic antibody-associated vasculitis. <i>International Journal of Clinical Practice</i> , 2021, 75, e14512.	1.7	3
28	Predictive Ability of Serum IL-27 Level for Assessing Activity of Antineutrophil Cytoplasmic Antibody-Associated Vasculitis. <i>Mediators of Inflammation</i> , 2021, 2021, 1-8.	3.0	2
29	Serum adipokine profiles in patients with microscopic polyangiitis and granulomatosis with polyangiitis: An exploratory analysis. <i>PLoS ONE</i> , 2021, 16, e0254226.	2.5	1
30	Efficacy of tacrolimus as maintenance therapy after cyclophosphamide for treating antineutrophil cytoplasmic antibody-associated vasculitis. <i>Medicine (United States)</i> , 2021, 100, e26956.	1.0	0
31	Evaluation of body composition using computed tomography in patients with anti-neutrophil cytoplasmic antibody-associated vasculitis. <i>Korean Journal of Internal Medicine</i> , 2021, 36, 1221-1232.	1.7	1
32	The novel fibrosis index at diagnosis may predict all-cause mortality in patients with antineutrophil cytoplasmic antibody-associated vasculitis without substantial liver diseases. <i>Clinics</i> , 2021, 76, e2501.	1.5	2
33	Antineutrophil Cytoplasmic Antibody Positivity Is Associated with Vascular Involvement in Behçet's Disease. <i>Yonsei Medical Journal</i> , 2021, 62, 149.	2.2	6
34	Male Sex Is a Significant Predictor of All-cause Mortality in Patients with Antineutrophil Cytoplasmic Antibody-associated Vasculitis. <i>Journal of Korean Medical Science</i> , 2021, 36, e120.	2.5	8
35	The significance of cytoplasmic antinuclear antibody patterns in autoimmune liver disease. <i>PLoS ONE</i> , 2021, 16, e0244950.	2.5	5
36	The Efficacy of Mycophenolate Mofetil in Remission Maintenance Therapy for Microscopic Polyangiitis and Granulomatosis with Polyangiitis. <i>Yonsei Medical Journal</i> , 2021, 62, 494.	2.2	2

#	ARTICLE	IF	CITATIONS
37	Efficacy of the fibrosis index for predicting end-stage renal disease in patients with antineutrophil cytoplasmic antibody-associated vasculitis. <i>International Journal of Clinical Practice</i> , 2021, 75, e13929.	1.7	4
38	Serum Clusterin Level Could Reflect the Current Activity of Antineutrophil Cytoplasmic Antibody-Associated Vasculitis. <i>Yonsei Medical Journal</i> , 2021, 62, 1016.	2.2	3
39	Serum progranulin as a predictive marker for high activity of antineutrophil cytoplasmic antibody-associated vasculitis. <i>Journal of Clinical Laboratory Analysis</i> , 2021, 35, e24048.	2.1	1
40	B-cell metabolism regulator IM156 contributes to the mitigation of systemic lupus erythematosus. <i>Korean Journal of Transplantation</i> , 2021, 35, S31-S31.	0.1	0
41	Prevalence of Osteopenia in Drug-Naive Patients With Antineutrophil Cytoplasmic Antibody-Associated Vasculitis. <i>Journal of Clinical Rheumatology</i> , 2021, 27, e330-e335.	0.9	2
42	Serum granzyme B is associated with otorhinolaryngological, pulmonary, and renal involvement of antineutrophil cytoplasmic antibody-associated vasculitis. <i>Journal of Investigative Medicine</i> , 2021, 69, 91-95.	1.6	0
43	Pan-immune-inflammation value at diagnosis independently predicts all-cause mortality in patients with antineutrophil cytoplasmic antibody-associated vasculitis. <i>Clinical and Experimental Rheumatology</i> , 2021, 39 Suppl 129, 88-93.	0.8	1
44	D-dimer predicts poor hospitalisation outcomes in patients with antineutrophil cytoplasmic autoantibody-associated vasculitis. <i>Clinical and Experimental Rheumatology</i> , 2021, 39 Suppl 129, 94-100.	0.8	0
45	Serum galectin-9 could be a potential biomarker in assessing the disease activity of antineutrophil cytoplasmic antibody-associated vasculitis. <i>Clinical and Experimental Rheumatology</i> , 2021, .	0.8	0
46	Significance of antineutrophil cytoplasmic antibody positivity in patients with systemic sclerosis: a single-centre pilot study in Korea. <i>Clinical and Experimental Rheumatology</i> , 2021, 39 Suppl 131, 111-118.	0.8	0
47	D-dimer predicts poor hospitalisation outcomes in patients with antineutrophil cytoplasmic autoantibody-associated vasculitis. <i>Clinical and Experimental Rheumatology</i> , 2021, 39, 94-100.	0.8	3
48	Pan-immune-inflammation value at diagnosis independently predicts all-cause mortality in patients with antineutrophil cytoplasmic antibody-associated vasculitis. <i>Clinical and Experimental Rheumatology</i> , 2021, 39, 88-93.	0.8	14
49	Significance of antineutrophil cytoplasmic antibody positivity in patients with systemic sclerosis: a single-centre pilot study in Korea. <i>Clinical and Experimental Rheumatology</i> , 2021, 39, 111-118.	0.8	0
50	Clinical impact of proteinase 3-antineutrophil cytoplasmic antibody positivity in eosinophilic granulomatosis with polyangiitis. <i>Korean Journal of Internal Medicine</i> , 2021, .	1.7	1
51	Metabolic Syndrome Severity Score, Comparable to Serum Creatinine, Could Predict the Occurrence of End-Stage Kidney Disease in Patients with Antineutrophil Cytoplasmic Antibody-Associated Vasculitis. <i>Journal of Clinical Medicine</i> , 2021, 10, 5744.	2.4	5
52	Multivariable index for assessing the activity and predicting all-cause mortality in antineutrophil cytoplasmic antibody-associated vasculitis. <i>Journal of Clinical Laboratory Analysis</i> , 2020, 34, e23022.	2.1	13
53	Can antineutrophil cytoplasmic antibody positivity at diagnosis predict the poor outcomes of Sjögren's syndrome?. <i>Rheumatology International</i> , 2020, 40, 1063-1070.	3.0	4
54	Clinical implication of chronic paranasal sinusitis for the classification of microscopic polyangiitis. <i>International Journal of Clinical Practice</i> , 2020, 74, e13431.	1.7	2

#	ARTICLE	IF	CITATIONS
55	Double positivity for antineutrophil cytoplasmic antibody (ANCA) and anti-glomerular basement membrane antibody could predict end-stage renal disease in ANCA-associated vasculitis: a monocentric pilot study. <i>Clinical Rheumatology</i> , 2020, 39, 831-840.	2.2	2
56	Atherogenic index of plasma predicts cerebrovascular accident occurrence in antineutrophil cytoplasmic antibody-associated vasculitis. <i>Lipids in Health and Disease</i> , 2020, 19, 184.	3.0	7
57	Serum Amyloid A Is a Biomarker of Disease Activity and Health-Related Quality-of-Life in Patients with Antineutrophil Cytoplasmic Antibody-Associated Vasculitis. <i>Disease Markers</i> , 2020, 2020, 1-9.	1.3	4
58	Definite IgG4-related disease had no overlap with eosinophilic granulomatosis with polyangiitis in Korean patients: a pilot study in one centre. <i>Clinical Rheumatology</i> , 2020, 39, 3009-3015.	2.2	3
59	Association between the antineutrophil cytoplasmic antibody and late coronary arterial occlusive disease in patients with Takayasu arteritis. <i>Journal of Cardiology</i> , 2020, 76, 407-412.	1.9	3
60	Clinical implication of plasma exchange on life-threatening antineutrophil cytoplasmic antibody-associated vasculitis. <i>BMC Pulmonary Medicine</i> , 2020, 20, 147.	2.0	2
61	Clinical characteristics and long-term outcomes of Libmanâ€Sacks endocarditis in patients with systemic lupus erythematosus. <i>Lupus</i> , 2020, 29, 1115-1120.	1.6	11
62	Non-histologic factors discriminating proliferative lupus nephritis from membranous lupus nephritis. <i>Arthritis Research and Therapy</i> , 2020, 22, 138.	3.5	5
63	Comparison of clinical features and outcomes between patients with early and delayed lupus nephritis. <i>BMC Nephrology</i> , 2020, 21, 258.	1.8	5
64	Hyperuricemia is associated with decreased renal function and occurrence of end-stage renal disease in patients with microscopic polyangiitis and granulomatosis with polyangiitis: a retrospective study. <i>Rheumatology International</i> , 2020, 40, 1089-1099.	3.0	7
65	Clinical characteristics associated with drug-free sustained remission in patients with rheumatoid arthritis: Data from Korean Intensive Management of Early Rheumatoid Arthritis (KIMERA). <i>Seminars in Arthritis and Rheumatism</i> , 2020, 50, 1414-1420.	3.4	3
66	Serum interleukin-16 significantly correlates with the Vasculitis Damage Index in antineutrophil cytoplasmic antibody-associated vasculitis. <i>Arthritis Research and Therapy</i> , 2020, 22, 73.	3.5	6
67	Serum Mannose-Binding Lectin Levels Are Correlated with the Disease Activity of Antineutrophil Cytoplasmic Antibody-Associated Vasculitis: A Single-Center Study. <i>Tohoku Journal of Experimental Medicine</i> , 2020, 251, 117-123.	1.2	7
68	Worse Renal Presentation and Prognosis in Initial-Onset Lupus Nephritis than Early-Onset Lupus Nephritis. <i>Yonsei Medical Journal</i> , 2020, 61, 951.	2.2	4
69	Rituximab Biosimilar Prevents Poor Outcomes of Microscopic Polyangiitis and Granulomatosis with Polyangiitis as Effectively as Rituximab Originator. <i>Yonsei Medical Journal</i> , 2020, 61, 712.	2.2	5
70	Soluble Lectin-Like Oxidized Low-Density Lipoprotein Receptor 1 Is Inversely Correlated with the Activity of ANCA-Associated Vasculitis. <i>Yonsei Medical Journal</i> , 2020, 61, 720.	2.2	2
71	Lipid Profiles in Anti-neutrophil Cytoplasmic Antibody-associated Vasculitis: A Cross-sectional Analysis. <i>Journal of Rheumatic Diseases</i> , 2020, 27, 261-269.	1.1	4
72	Pregnancy Morbidities in Korean Patients with Takayasu Arteritis: A Monocentric Pilot Study. <i>Yonsei Medical Journal</i> , 2020, 61, 970.	2.2	3

#	ARTICLE	IF	CITATIONS
73	MiR-451 suppresses inflammatory responses in ankylosing spondylitis by targeting macrophage migration inhibitory factor. <i>Clinical and Experimental Rheumatology</i> , 2020, 38, 275-281.	0.8	4
74	Will the HALP score help to assess the activity and predict the prognosis of antineutrophil cytoplasmic antibody-associated vasculitis?. <i>Clinical and Experimental Rheumatology</i> , 2020, 38 Suppl 124, 236-237.	0.8	2
75	Application of the 2019 classification criteria for systemic lupus erythematosus to patients with established ANCA-associated vasculitis. <i>Clinical and Experimental Rheumatology</i> , 2020, 38 Suppl 124, 243-244.	0.8	0
76	Predictor of depressive disorders in patients with antineutrophil cytoplasmic antibody-associated vasculitis. <i>Clinical Rheumatology</i> , 2019, 38, 3485-3491.	2.2	7
77	Evaluation of macrophage activation syndrome in hospitalised patients with Kikuchi-Fujimoto disease based on the 2016 EULAR/ACR/PRINTO classification criteria. <i>PLoS ONE</i> , 2019, 14, e0219970.	2.5	8
78	Could hypereosinophilia at diagnosis estimate the current activity or predict relapse in systemic immunosuppressive drug-naïve patients with eosinophilic granulomatosis with polyangiitis?. <i>Rheumatology International</i> , 2019, 39, 1899-1905.	3.0	3
79	Serum Aminoacyl-tRNA Synthetase-Interacting Multifunctional Protein-1 Can Predict Severe Antineutrophil Cytoplasmic Antibody-Associated Vasculitis: A Pilot Monocentric Study. <i>BioMed Research International</i> , 2019, 2019, 1-6.	1.9	7
80	Comparison of the Clinical Implications among Five Different Nutritional Indices in Patients with Lupus Nephritis. <i>Nutrients</i> , 2019, 11, 1456.	4.1	14
81	Risk of Cancers in Antineutrophil Cytoplasmic Antibody-Associated Vasculitis: Results from the Korea National Health Insurance Claims Database 2010–2018. <i>Journal of Clinical Medicine</i> , 2019, 8, 1871.	2.4	10
82	Clinical characteristics of patients with systemic lupus erythematosus showing a false-positive result of syphilis screening. <i>Rheumatology International</i> , 2019, 39, 1859-1866.	3.0	4
83	Anti-Smith antibody is associated with disease activity in patients with new-onset systemic lupus erythematosus. <i>Rheumatology International</i> , 2019, 39, 1937-1944.	3.0	26
84	Fibrosis-4 index at diagnosis is associated with all-cause mortality in patients with microscopic polyangiitis and granulomatosis with polyangiitis. <i>BMC Gastroenterology</i> , 2019, 19, 90.	2.0	13
85	Anti-phospholipid antibody syndrome occurrence in patients with persistent anti-phospholipid antibodies. <i>Rheumatology International</i> , 2019, 39, 1359-1367.	3.0	4
86	Comparison of Radiological and Histological Findings of Lung Parenchyma in Patients with Antineutrophil Cytoplasmic Antibody-Associated Vasculitis. <i>Yonsei Medical Journal</i> , 2019, 60, 454.	2.2	10
87	Should nasal biopsy inevitably be performed for classifying granulomatosis with polyangiitis in patients with rhinosinusitis? A retrospective chart review study. <i>Rheumatology International</i> , 2019, 39, 885-892.	3.0	3
88	Ex Vivo Interferon Gamma Production by Peripheral Immune Cells Predicts Survival in Lung Adenocarcinoma. <i>Clinical Lung Cancer</i> , 2019, 20, e299-e308.	2.6	2
89	Uterine Artery Embolization in Patients With Autoimmune Disease: A Matched Case-Control Study. <i>American Journal of Roentgenology</i> , 2019, 212, 1148-1153.	2.2	1
90	Serum interleukin-21 positivity could indicate the current activity of antineutrophil cytoplasmic antibody-associated vasculitis: a monocentric prospective study. <i>Clinical Rheumatology</i> , 2019, 38, 1685-1690.	2.2	5



#	ARTICLE	IF	CITATIONS
91	Prognostic nutritional index is associated with disease severity and relapse in ANCA-associated vasculitis. <i>International Journal of Rheumatic Diseases</i> , 2019, 22, 797-804.	1.9	16
92	ANCA positivity at the time of renal biopsy is associated with chronicity index of lupus nephritis. <i>Rheumatology International</i> , 2019, 39, 879-884.	3.0	9
93	Persistent antiphospholipid antibodies are associated with thrombotic events in ANCA-associated vasculitis: A retrospective monocentric study. <i>Nefrologia</i> , 2019, 39, 395-401.	0.4	6
94	FRI0651...THE CLINICAL IMPLICATION OF NASAL BIOPSY FOR CLASSIFYING GRANULOMATOSIS WITH POLYANGIITIS IN PATIENTS WITH RHINOSINUSITIS: A SINGLE CENTRE RETROSPECTIVE STUDY. , 2019, , .		0
95	AB0525...ANTI-SMITH ANTIBODY IS ASSOCIATED WITH DISEASE ACTIVITY IN PATIENTS WITH NEW-ONSET SYSTEMIC LUPUS ERYTHEMATOSUS. , 2019, , .		0
96	THU0576...PROGNOSTIC FACTORS PREDICTING THE SURVIVAL OF PATIENTS WITH MACROPHAGE ACTIVATION SYNDROME. , 2019, , .		0
97	125...Treatment outcome in lupus nephritis patients treated with mycophenolate mofetil: from a real-world clinical practice. , 2019, , .		0
98	AB0347...CLINICAL RELEVANCE OF ANTI-CARBAMYLATED PROTEIN ANTIBODY: IS IT SAME AS CAUCASIANS IN ASIANS?. , 2019, , .		0
99	Clinical role of albumin to globulin ratio in microscopic polyangiitis: a retrospective monocentric study. <i>Clinical Rheumatology</i> , 2019, 38, 487-494.	2.2	11
100	Low serum complement 3 level is associated with severe ANCA-associated vasculitis at diagnosis. <i>Clinical and Experimental Nephrology</i> , 2019, 23, 223-230.	1.6	20
101	Systemic immune-inflammatory index could estimate the cross-sectional high activity and the poor outcomes in immunosuppressive drug-naïve patients with antineutrophil cytoplasmic antibody-associated vasculitis. <i>Nephrology</i> , 2019, 24, 711-717.	1.6	42
102	No overlap between IgG4-related disease and microscopic polyangiitis and granulomatosis with polyangiitis despite elevated serum IgG4 at diagnosis: a retrospective monocentric study. <i>Clinical Rheumatology</i> , 2019, 38, 1147-1154.	2.2	21
103	Controlling Nutritional Status Score is Associated with All-Cause Mortality in Patients with Antineutrophil Cytoplasmic Antibody-Associated Vasculitis. <i>Yonsei Medical Journal</i> , 2019, 60, 1164.	2.2	8
104	Metabolic Reprogramming by the Excessive AMPK Activation Exacerbates Antigen-Specific Memory CD8 <sup>+</sup> T Cell Differentiation after Acute Lymphocytic Choriomeningitis Virus Infection. <i>Immune Network</i> , 2019, 19, e11.	3.6	7
105	Serum soluble programmed cell death protein 1 could predict the current activity and severity of antineutrophil cytoplasmic antibody-associated vasculitis: a monocentric prospective study. <i>Clinical and Experimental Rheumatology</i> , 2019, 37 Suppl 117, 116-121.	0.8	2
106	Risk of Primary Spontaneous Pneumothorax According to Chest Configuration. <i>Thoracic and Cardiovascular Surgeon</i> , 2018, 66, 583-588.	1.0	7
107	Birmingham vasculitis activity and chest manifestation at diagnosis can predict hospitalised infection in ANCA-associated vasculitis. <i>Clinical Rheumatology</i> , 2018, 37, 2133-2141.	2.2	12
108	Mean platelet volume can estimate the current vasculitis activity of microscopic polyangiitis. <i>Rheumatology International</i> , 2018, 38, 1095-1101.	3.0	9

#	ARTICLE	IF	CITATIONS
109	Decreased muscle mass is independently associated with knee pain in female patients with radiographically mild osteoarthritis: a nationwide cross-sectional study (KNHANES 2010-2011). <i>Clinical Rheumatology</i> , 2018, 37, 1333-1340.	2.2	15
110	Hemoglobin A1c, Not Glycated Albumin, Can Independently Reflect the Ankylosing Spondylitis Disease Activity Score. <i>Journal of Rheumatic Diseases</i> , 2018, 25, 131.	1.1	2
111	Treat-to-Target Strategy for Asian Patients with Early Rheumatoid Arthritis: Result of a Multicenter Trial in Korea. <i>Journal of Korean Medical Science</i> , 2018, 33, e346.	2.5	5
112	A novel antimicrobial peptide acting via formyl peptide receptor 2 shows therapeutic effects against rheumatoid arthritis. <i>Scientific Reports</i> , 2018, 8, 14664.	3.3	20
113	Risk factors associated with inadequate control of disease activity in elderly patients with rheumatoid arthritis: Results from a nationwide Korean College of Rheumatology BIOlogics (KOBIO) registry. <i>PLoS ONE</i> , 2018, 13, e0205651.	2.5	13
114	Red Blood Cell Distribution Width Can Predict Vasculitis Activity and Poor Prognosis in Granulomatosis with Polyangiitis. <i>Yonsei Medical Journal</i> , 2018, 59, 294.	2.2	6
115	Neutrophil to lymphocyte ratio at diagnosis can estimate vasculitis activity and poor prognosis in patients with ANCA-associated vasculitis: a retrospective study. <i>BMC Nephrology</i> , 2018, 19, 187.	1.8	32
116	Safety of Tocilizumab in Rheumatoid Arthritis Patients with Resolved Hepatitis B Virus Infection: Data from Real-World Experience. <i>Yonsei Medical Journal</i> , 2018, 59, 452.	2.2	27
117	Delta Neutrophil Index Is Associated with Vasculitis Activity and Risk of Relapse in ANCA-Associated Vasculitis. <i>Yonsei Medical Journal</i> , 2018, 59, 397.	2.2	16
118	Platelet to lymphocyte ratio is associated with the current activity of ANCA-associated vasculitis at diagnosis: a retrospective monocentric study. <i>Rheumatology International</i> , 2018, 38, 1865-1871.	3.0	28
119	Serum leucine-rich $\alpha_2$ -glycoprotein is elevated in patients with systemic lupus erythematosus and correlates with disease activity. <i>Clinica Chimica Acta</i> , 2018, 486, 253-258.	1.1	10
120	Reclassification of polyarteritis nodosa based on the 1990 ACR criteria using the 2007 EMA algorithm modified by the 2012 CHCC definitions. <i>Clinical and Experimental Rheumatology</i> , 2018, 36 Suppl 111, 165-166.	0.8	2
121	The utility of the ACR/EULAR 2017 provisional classification criteria for granulomatosis with polyangiitis in Korean patients with antineutrophil cytoplasmic antibody-associated vasculitis. <i>Clinical and Experimental Rheumatology</i> , 2018, 36 Suppl 111, 85-87.	0.8	14
122	Renal outcome of kidney-transplantation in Korean recipients with ANCA-associated vasculitis. <i>Clinical and Experimental Rheumatology</i> , 2018, 36 Suppl 111, 115-120.	0.8	2
123	Serum aminoacyl-tRNA synthetase-interacting multifunctional protein-1 (AIMP1), a novel disease activity predictive biomarker of systemic lupus erythematosus. <i>Clinical and Experimental Rheumatology</i> , 2018, 36, 533-539.	0.8	2
124	Cancer development in Korean patients with ANCA-associated vasculitis: a single centre study. <i>Clinical and Experimental Rheumatology</i> , 2018, 36 Suppl 111, 73-77.	0.8	3
125	The initial predictors of death in 153 patients with ANCA-associated vasculitis in a single Korean centre. <i>Clinical and Experimental Rheumatology</i> , 2018, 36 Suppl 111, 65-72.	0.8	9
126	Serum anti-lysozyme is associated with disease activity of Behçet's disease. <i>International Journal of Rheumatic Diseases</i> , 2017, 20, 261-268.	1.9	8



#	ARTICLE	IF	CITATIONS
127	Five factor score of more than 1 is associated with relapse during the first 2 year follow up in patients with eosinophilic granulomatosis with polyangiitis. <i>International Journal of Rheumatic Diseases</i> , 2017, 20, 1261-1268.	1.9	12
128	In-hospital mortality in febrile lupus patients based on 2016 EULAR/ACR/PRINTO classification criteria for macrophage activation syndrome. <i>Seminars in Arthritis and Rheumatism</i> , 2017, 47, 216-221.	3.4	28
129	Evaluation of Spleen Glucose Metabolism Using <sup>18</sup> F-FDG PET/CT in Patients with Febrile Autoimmune Disease. <i>Journal of Nuclear Medicine</i> , 2017, 58, 507-513.	5.0	33
130	Application of the 2016 EULAR/ACR/PRINTO Classification Criteria for Macrophage Activation Syndrome in Patients with Adult-onset Still Disease. <i>Journal of Rheumatology</i> , 2017, 44, 996-1003.	2.0	43
131	Delta neutrophil index contributes to the differential diagnosis between acute gout attack and cellulitis within 24 hours after hospitalization. <i>Rheumatology</i> , 2017, 56, kew471.	1.9	11
132	Malignancies in Korean patients with immunoglobulin G4-related disease. <i>International Journal of Rheumatic Diseases</i> , 2017, 20, 1028-1035.	1.9	35
133	Birmingham vasculitis activity score at diagnosis is a significant predictor of relapse of polyarteritis nodosa. <i>Rheumatology International</i> , 2017, 37, 685-694.	3.0	11
134	Echocardiographic features in patients with ANCA-associated vasculitis within 3 months before and after diagnosis. <i>Clinical Rheumatology</i> , 2017, 36, 2751-2759.	2.2	9
135	The clinical utility of splenic fluorodeoxyglucose uptake for diagnosis and prognosis in patients with macrophage activation syndrome. <i>Medicine (United States)</i> , 2017, 96, e7901.	1.0	6
136	Birmingham vasculitis activity score of more than 9.5 at diagnosis is an independent predictor of refractory disease in granulomatosis with polyangiitis. <i>International Journal of Rheumatic Diseases</i> , 2017, 20, 1593-1605.	1.9	17
137	Clinical significance of delta neutrophil index in the differential diagnosis between septic arthritis and acute gout attack within 24 hours after hospitalization. <i>Medicine (United States)</i> , 2017, 96, e7431.	1.0	7
138	Defective autophagy activity and its association with spinal damage in patients with ankylosing spondylitis. <i>Joint Bone Spine</i> , 2017, 84, 583-587.	1.6	12
139	Fever as an initial manifestation of spondyloarthritis: A retrospective study. <i>PLoS ONE</i> , 2017, 12, e0184323.	2.5	5
140	Decreased ex vivo production of interferon-gamma is associated with severity and poor prognosis in patients with lupus. <i>Arthritis Research and Therapy</i> , 2017, 19, 193.	3.5	14
141	Platelet Distribution Width and Mean Platelet Volume Are Not Correlated with the Disease Activity Indices of Ankylosing Spondylitis. <i>Journal of Rheumatic Diseases</i> , 2017, 24, 143.	1.1	0
142	Subsequent Thrombotic Outcomes in Patients with Ischemic Stroke with Antiphospholipid Antibody Positivity. <i>Yonsei Medical Journal</i> , 2017, 58, 1128.	2.2	6
143	Predictors of mortality in autoimmune disease patients with concurrent cytomegalovirus infections detected by quantitative real-time PCR. <i>PLoS ONE</i> , 2017, 12, e0181590.	2.5	9
144	Chest and renal involvements, Birmingham vascular activity score more than 13.5 and five factor score (1996) more than 1 at diagnosis are significant predictors of relapse of microscopic polyangiitis. <i>Clinical and Experimental Rheumatology</i> , 2017, 35 Suppl 103, 47-54.	0.8	11

#	ARTICLE	IF	CITATIONS
145	Differential expressions of NOD-like receptors and their associations with inflammatory responses in rheumatoid arthritis. <i>Clinical and Experimental Rheumatology</i> , 2017, 35, 630-637.	0.8	18
146	Clinical and prognostic features of Korean patients with MPO-ANCA, PR3-ANCA and ANCA-negative vasculitis. <i>Clinical and Experimental Rheumatology</i> , 2017, 35 Suppl 103, 111-118.	0.8	16
147	Anti- $\epsilon$ m is associated with the early poor outcome of lupus nephritis. <i>International Journal of Rheumatic Diseases</i> , 2016, 19, 897-902.	1.9	22
148	Brain meningioma in a patient with systemic lupus erythematosus. <i>Yeungnam University Journal of Medicine</i> , 2016, 33, 159.	1.4	1
149	Application of the 2013 ACR/EULAR classification criteria for systemic sclerosis to patients with Raynaud's phenomenon. <i>Arthritis Research and Therapy</i> , 2015, 17, 77.	3.5	20
150	Disease duration and Medsger's severity score are associated with significant liver fibrosis in patients with systemic sclerosis. <i>Clinical and Experimental Rheumatology</i> , 2015, 33, S68-74.	0.8	5
151	Active Tuberculosis Risk With Tumor Necrosis Factor Inhibitors After Treating Latent Tuberculosis. <i>Journal of Clinical Rheumatology</i> , 2014, 20, 68-73.	0.9	12
152	Serum galectin-9 could be a potential biomarker in assessing the disease activity of antineutrophil cytoplasmic antibody-associated vasculitis. <i>Clinical and Experimental Rheumatology</i> , 0, , .	0.8	1
153	Effect of numbers of metabolic syndrome components on mortality in patients with antineutrophil cytoplasmic antibody-associated vasculitis with metabolic syndrome. <i>Clinical and Experimental Rheumatology</i> , 0, , .	0.8	1
154	Application of the 2022 ACR/EULAR criteria for microscopic polyangiitis to patients with previously diagnosed microscopic polyangiitis. <i>Clinical and Experimental Rheumatology</i> , 0, , .	0.8	2
155	Fatty Liver Index Independently Predicts All-Cause Mortality in Patients With Antineutrophil Cytoplasmic Antibody-Associated Vasculitis but No Substantial Liver Disease. <i>Frontiers in Cardiovascular Medicine</i> , 0, 9, .	2.4	3
156	Predicting the depressive status using empirical dietary inflammatory index in patients with antineutrophil cytoplasmic antibody-associated vasculitis. <i>Journal of Clinical Laboratory Analysis</i> , 0, , .	2.1	1