

Sabine Fillinger

List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/9423225/publications.pdf>

Version: 2024-02-01

34
papers

4,662
citations

236925
25
h-index

395702
33
g-index

40
all docs

40
docs citations

40
times ranked

5156
citing authors

| # | ARTICLE | IF | CITATIONS |
|----|--|-----|-----------|
| 1 | Directed evolution predicts cytochrome <i>b_L</i> G37V target site modification as probable adaptive mechanism towards the <i>Qil</i> fungicide fenpicoxamid in <i>Zymoseptoria tritici</i> . Environmental Microbiology, 2022, 24, 1117-1132. | 3.8 | 13 |
| 2 | Comparative quantitative proteomics of osmotic signal transduction mutants in <i>Botrytis cinerea</i> explain mutant phenotypes and highlight interaction with cAMP and Ca ²⁺ signalling pathways. Journal of Proteomics, 2020, 212, 103580. | 2.4 | 5 |
| 3 | Fungicide Sensitivity Shifting of <i>Zymoseptoria tritici</i> in the Finnish-Baltic Region and a Novel Insertion in the MFS1 Promoter. Frontiers in Plant Science, 2020, 11, 385. | 3.6 | 21 |
| 4 | Plasticity of the <i>MFS1</i> Promoter Leads to Multidrug Resistance in the Wheat Pathogen <i>Zymoseptoria tritici</i> . MSphere, 2017, 2, . | 2.9 | 75 |
| 5 | Phenylpyrroles: 30 Years, Two Molecules and (Nearly) No Resistance. Frontiers in Microbiology, 2016, 7, 2014. | 3.5 | 82 |
| 6 | Proposal for a unified nomenclature for target-site mutations associated with resistance to fungicides. Pest Management Science, 2016, 72, 1449-1459. | 3.4 | 76 |
| 7 | <i>Botrytis</i> , the Good, the Bad and the Ugly. , 2016, , 1-15. | | 49 |
| 8 | Chemical Control and Resistance Management of <i>Botrytis</i> Diseases. , 2016, , 189-216. | | 42 |
| 9 | Fungicide efflux and the <i>MgMFS</i> 1 transporter contribute to the multidrug resistance phenotype in <i>Zymoseptoria tritici</i> field isolates. Environmental Microbiology, 2015, 17, 2805-2823. | 3.8 | 140 |
| 10 | Phosphoproteome profiles of the phytopathogenic fungi <i>Alternaria brassicicola</i> and <i>Botrytis cinerea</i> during exponential growth in axenic cultures. Proteomics, 2014, 14, 1639-1645. | 2.2 | 13 |
| 11 | Site-directed mutagenesis of the <i>P225</i> , <i>N230</i> and <i>H272</i> residues of succinate dehydrogenase subunit <i>B</i> from <i>Botrytis cinerea</i> highlights different roles in enzyme activity and inhibitor binding. Environmental Microbiology, 2014, 16, 2253-2266. | 3.8 | 90 |
| 12 | Role of sterol 3-oxoreductase sensitivity in susceptibility to the fungicide fenhexamid in <i>Botrytis cinerea</i> and other phytopathogenic fungi. Pest Management Science, 2013, 69, 642-651. | 3.4 | 20 |
| 13 | A Functional Bikaverin Biosynthesis Gene Cluster in Rare Strains of <i>Botrytis cinerea</i> Is Positively Controlled by VELVET. PLoS ONE, 2013, 8, e53729. | 2.5 | 69 |
| 14 | Functional and Structural Comparison of Pyrrolnitrin- and Iprodione-Induced Modifications in the Class III Histidine-Kinase Bos1 of <i>Botrytis cinerea</i> . PLoS ONE, 2012, 7, e42520. | 2.5 | 62 |
| 15 | Strong resistance to the fungicide fenhexamid entails a fitness cost in <i>Botrytis cinerea</i> , as shown by comparisons of isogenic strains. Pest Management Science, 2012, 68, 684-691. | 3.4 | 49 |
| 16 | Genomic Analysis of the Necrotrophic Fungal Pathogens <i>Sclerotinia sclerotiorum</i> and <i>Botrytis cinerea</i> . PLoS Genetics, 2011, 7, e1002230. | 3.5 | 902 |
| 17 | The osmosensing signal transduction pathway from <i>Botrytis cinerea</i> regulates cell wall integrity and MAP kinase pathways control melanin biosynthesis with influence of light. Fungal Genetics and Biology, 2011, 48, 377-387. | 2.1 | 66 |
| 18 | Fungicide-Driven Evolution and Molecular Basis of Multidrug Resistance in Field Populations of the Grey Mould Fungus <i>Botrytis cinerea</i> . PLoS Pathogens, 2009, 5, e1000696. | 4.7 | 329 |

| # | ARTICLE | IF | CITATIONS |
|----|--|-----|-----------|
| 19 | The HOG1-like MAP kinase Sak1 of <i>Botrytis cinerea</i> is negatively regulated by the upstream histidine kinase Bos1 and is not involved in dicarboximide- and phenylpyrrole-resistance. <i>Fungal Genetics and Biology</i> , 2008, 45, 1062-1074. | 2.1 | 100 |
| 20 | Genetic Analysis of Fenhexamid-Resistant Field Isolates of the Phytopathogenic Fungus <i>Botrytis cinerea</i> . <i>Antimicrobial Agents and Chemotherapy</i> , 2008, 52, 3933-3940. | 3.2 | 97 |
| 21 | A Class III Histidine Kinase Acts as a Novel Virulence Factor in <i>Botrytis cinerea</i> . <i>Molecular Plant-Microbe Interactions</i> , 2006, 19, 1042-1050. | 2.6 | 149 |
| 22 | Expressed sequence tags from the phytopathogenic fungus <i>Botrytis cinerea</i> . <i>European Journal of Plant Pathology</i> , 2005, 111, 139-146. | 1.7 | 20 |
| 23 | Glycerol dehydrogenase, encoded by <i>gldB</i> is essential for osmotolerance in <i>Aspergillus nidulans</i> . <i>Molecular Microbiology</i> , 2003, 49, 131-141. | 2.5 | 62 |
| 24 | Essential <i>Bacillus subtilis</i> genes. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 2003, 100, 4678-4683. | 7.1 | 1,261 |
| 25 | The <i>bddDC</i> Operon of <i>Bacillus subtilis</i> Encodes Thiol-disulfide Oxidoreductases Required for Competence Development. <i>Journal of Biological Chemistry</i> , 2002, 277, 6994-7001. | 3.4 | 85 |
| 26 | cAMP and ras signalling independently control spore germination in the filamentous fungus <i>Aspergillus nidulans</i> . <i>Molecular Microbiology</i> , 2002, 44, 1001-1016. | 2.5 | 170 |
| 27 | Molecular and physiological characterization of the NAD-dependent glycerol 3-phosphate dehydrogenase in the filamentous fungus <i>Aspergillus nidulans</i> . <i>Molecular Microbiology</i> , 2001, 39, 145-157. | 2.5 | 58 |
| 28 | Trehalose is required for the acquisition of tolerance to a variety of stresses in the filamentous fungus <i>Aspergillus nidulans</i> The GenBank accession number for the sequence reported in this paper is AF043230. <i>Microbiology (United Kingdom)</i> , 2001, 147, 1851-1862. | 1.8 | 187 |
| 29 | In vivo studies of upstream regulatory cis-acting elements of the <i>alcR</i> gene encoding the transactivator of the ethanol regulon in <i>Aspergillus nidulans</i> . <i>Molecular Microbiology</i> , 2000, 36, 123-131. | 2.5 | 40 |
| 30 | Two Glyceraldehyde-3-phosphate Dehydrogenases with Opposite Physiological Roles in a Nonphotosynthetic Bacterium. <i>Journal of Biological Chemistry</i> , 2000, 275, 14031-14037. | 3.4 | 173 |
| 31 | Histidinol Phosphate Phosphatase, Catalyzing the Penultimate Step of the Histidine Biosynthesis Pathway, Is Encoded by <i>ytvP</i> (<i>hisJ</i>) in <i>Bacillus subtilis</i> . <i>Journal of Bacteriology</i> , 1999, 181, 3277-3280. | 2.2 | 22 |
| 32 | The Zinc Binuclear Cluster Activator AlcR Is Able to Bind to Single Sites but Requires Multiple Repeated Sites for Synergistic Activation of the <i>alcA</i> Gene in <i>Aspergillus nidulans</i> . <i>Journal of Biological Chemistry</i> , 1997, 272, 22859-22865. | 3.4 | 43 |
| 33 | A newly identified gene cluster in <i>Aspergillus nidulans</i> comprises five novel genes localized in the <i>alc</i> region that are controlled both by the specific transactivator AlcR and the general carbon catabolite repressor CreA. <i>Molecular Microbiology</i> , 1996, 20, 475-488. | 2.5 | 52 |
| 34 | The basal level of transcription of the <i>alc</i> genes in the ethanol regulon in <i>Aspergillus nidulans</i> is controlled both by the specific transactivator AlcR and the general carbon catabolite repressor CreA. <i>FEBS Letters</i> , 1995, 368, 547-550. | 2.8 | 35 |