Christian Meyer

List of Publications by Year in descending order

Source: https://exaly.com/author-pdf/9442314/publications.pdf

Version: 2024-02-01

169 papers 6,773 citations

43 h-index 71 g-index

214 all docs

214 docs citations

times ranked

214

7339 citing authors

#	Article	IF	CITATIONS
1	Severity of alcohol dependence and mortality after 20Âyears in an adult general population sample. International Journal of Methods in Psychiatric Research, 2022, 31, e1915.	2.1	6
2	The Role of Tobacco Smoking in the Efficacy of Brief Alcohol Intervention: Results from a Randomized Controlled Trial. International Journal of Environmental Research and Public Health, 2022, 19, 5847.	2.6	1
3	The Moderating Effect of Educational Background on the Efficacy of a Computer-Based Brief Intervention Addressing the Full Spectrum of Alcohol Use: Randomized Controlled Trial. JMIR Public Health and Surveillance, 2022, 8, e33345.	2.6	3
4	Effect of a 1â€year short message service in detoxified alcoholâ€dependent patients: a multiâ€center, openâ€label randomized controlled trial. Addiction, 2021, 116, 1431-1442.	3.3	5
5	Brief alcohol intervention at a municipal registry office: reach and retention. European Journal of Public Health, 2021, 31, 418-423.	0.3	9
6	Latent alcohol use patterns and their link to depressive symptomatology in medical care patients. Addiction, $2021,116,1063-1073.$	3.3	6
7	Effects of a brief alcohol intervention addressing the full spectrum of drinking in an adult general population sample: a randomized controlled trial. Addiction, 2021, 116, 2056-2066.	3.3	7
8	Alkoholkonsum – Bedeutung für die Präention und Gesundheitsförderung. The Springer Reference Pflegerapie, Gesundheit, 2021, , 441-457.	0.3	0
9	The effect of a videoâ€supported assessment to increase the accuracy of selfâ€reported physical activity. Scandinavian Journal of Medicine and Science in Sports, 2021, 31, 1059-1068.	2.9	1
10	Longitudinal measurement invariance of the patient health questionnaire in a German sample. BMC Psychiatry, 2021, 21, 386.	2.6	2
11	Alcohol abstinence and mortality in a general population sample of adults in Germany: A cohort study. PLoS Medicine, 2021, 18, e1003819.	8.4	13
12	Untreated pathological gamblers: who recovers and who does not?. International Gambling Studies, 2020, 20, 200-213.	2.1	5
13	Use of machine learning to classify adult ADHD and other conditions based on the Conners' Adult ADHD Rating Scales. Scientific Reports, 2020, 10, 18871.	3.3	18
14	Copattern of depression and alcohol use in medical care patients: cross-sectional study in Germany. BMJ Open, 2020, 10, e032826.	1.9	2
15	Mental disorders and total mortality after 20Âyears in an adult general population sample. European Psychiatry, 2020, 63, e30.	0.2	10
16	Stability of Atâ€risk Alcohol Use Screening Results in a General Population Sample. Alcoholism: Clinical and Experimental Research, 2020, 44, 1312-1320.	2.4	4
17	Context matters: On the importance of context-specific analysis for the use of screening questionnaires. Drug and Alcohol Dependence, 2019, 205, 107663.	3.2	0
18	Diagnostic performance of the Alcohol Use Disorders Identification Test (AUDIT) in detecting DSM-5 alcohol use disorders in the General population. Drug and Alcohol Dependence, 2019, 204, 107530.	3.2	42

#	Article	IF	CITATIONS
19	Proactive multipurpose health risk screening in health care settings: Methods, design, and reach. International Journal of Methods in Psychiatric Research, 2019, 28, e1760.	2.1	13
20	Does prior recall of past week alcohol use affect screening results for at-risk drinking? Findings from a randomized study. PLoS ONE, 2019, 14, e0217595.	2.5	8
21	Feasibility and Acceptability of an Intervention Providing Computer-Generated Tailored Feedback to Target Alcohol Consumption and Depressive Symptoms in Proactively Recruited Health Care Patients and Reactively Recruited Media Volunteers: Results of a Pilot Study. European Addiction Research, 2019, 25, 119-131.	2.4	12
22	Can brief alcohol interventions in general hospital inpatients improve mental and general health over 2 years? Results from a randomized controlled trial. Psychological Medicine, 2019, 49, 1722-1730.	4.5	15
23	Alkoholkonsum – Bedeutung für Präention und Gesundheitsförderung. The Springer Reference Pflegerapie, Gesundheit, 2019, , 1-17.	0.3	0
24	The effect of online gambling on gambling problems and resulting economic health costs in Germany. European Journal of Health Economics, 2018, 19, 967-978.	2.8	41
25	Effect of blood pressure and total cholesterol measurement on risk prediction using the Systematic COronary Risk Evaluation (SCORE). BMC Cardiovascular Disorders, 2018, 18, 84.	1.7	1
26	The role of self-esteem in Internet addiction within the context of comorbid mental disorders: Findings from a general population-based sample. Journal of Behavioral Addictions, 2018, 7, 976-984.	3.7	18
27	Short-term stability of different drinking patterns over the course of four weeks among adults. A latent transition analysis. Drug and Alcohol Dependence, 2018, 191, 181-186.	3.2	11
28	Measurement invariance of the alcohol use disorders identification test: Establishing its factor structure in different settings and across gender. Drug and Alcohol Dependence, 2018, 189, 55-61.	3.2	28
29	Explaining the relation between pathological gambling and depression: Rumination as an underlying common cause. Journal of Behavioral Addictions, 2018, 7, 384-391.	3.7	11
30	Associations of specific psychiatric disorders with isolated focal dystonia, and monogenic and idiopathic Parkinson's disease. Journal of Neurology, 2017, 264, 1076-1084.	3.6	10
31	Reach of Individuals at Risk for Cardiovascular Disease by Proactive Recruitment Strategies in General Practices, Job Centers, and Health Insurance. International Journal of Behavioral Medicine, 2017, 24, 153-160.	1.7	14
32	Does impaired mental health interfere with the outcome of brief alcohol intervention at general hospitals?. Journal of Consulting and Clinical Psychology, 2017, 85, 562-573.	2.0	11
33	A Mobile Phone-Based Life Skills Training Program for Substance Use Prevention Among Adolescents: Pre-Post Study on the Acceptance and Potential Effectiveness of the Program, Ready4life. JMIR MHealth and UHealth, 2017, 5, e143.	3.7	30
34	The association between Internet addiction and personality disorders in a general population-based sample. Journal of Behavioral Addictions, 2016, 5, 691-699.	3.7	55
35	Dopamine-receptor 2 gene-methylation and gambling behavior in relation to impulsivity. Psychiatry Research, 2016, 239, 154-155.	3.3	8
36	Motivating smokers to quit using computer-generated letters that target either reduction or cessation: A population-based randomized controlled trial among smokers who do not intend to quit. Drug and Alcohol Dependence, 2016, 166, 177-186.	3.2	14

#	Article	IF	Citations
37	Genome-wide association study of pathological gambling. European Psychiatry, 2016, 36, 38-46.	0.2	82
38	Type of gambling as an independent risk factor for suicidal events in pathological gamblers Psychology of Addictive Behaviors, 2016, 30, 263-269.	2.1	26
39	Characteristics associated with non-participation in 7-day accelerometry. Preventive Medicine Reports, 2015, 2, 413-418.	1.8	15
40	A Pilot Study on the Feasibility and Acceptability of a Text Message-Based Aftercare Treatment Programme Among Alcohol Outpatients. Alcohol and Alcoholism, 2015, 50, 188-194.	1.6	31
41	Suicidal events among pathological gamblers: The role of comorbidity of axis I and axis II disorders. Psychiatry Research, 2015, 225, 413-419.	3.3	48
42	The "Pathological Gambling and Epidemiology―(PAGE) study program: design and fieldwork. International Journal of Methods in Psychiatric Research, 2015, 24, 11-31.	2.1	36
43	Alterations in DNA-methylation of the dopamine-receptor 2 gene are associated with abstinence and health care utilization in individuals with a lifetime history of pathologic gambling. Progress in Neuro-Psychopharmacology and Biological Psychiatry, 2015, 63, 30-34.	4.8	38
44	Factors Associated with Secondhand Smoke Exposure Prevalence and Secondhand Smoke Level of Children Living with Parental Smokers: A Cross Sectional Study. Journal of Community Health, 2015, 40, 501-507.	3.8	8
45	Demographic Risk Factors and Gambling Preference May Not Explain the High Prevalence of Gambling Problems Among the Population with Migration Background: Results from a German Nationwide Survey. Journal of Gambling Studies, 2015, 31, 741-757.	1.6	19
46	Thresholds of probable problematic gambling involvement for the German population: Results of the Pathological Gambling and Epidemiology (PAGE) Study Psychology of Addictive Behaviors, 2015, 29, 794-804.	2.1	23
47	A Surveillance Tool Using Mobile Phone Short Message Service to Reduce Alcohol Consumption Among Alcohol-Dependent Patients. Alcoholism: Clinical and Experimental Research, 2014, 38, 1728-1736.	2.4	36
48	Occurence of Internet Addiction in a General Population Sample: A Latent Class Analysis. European Addiction Research, 2014, 20, 159-166.	2.4	155
49	Assessment of Problematic Internet Use by the Compulsive Internet Use Scale and the Internet Addiction Test: A Sample of Problematic and Pathological Gamblers. European Addiction Research, 2014, 20, 75-81.	2.4	69
50	Association of Sociodemographic, Psychopathological and Gambling-Related Factors with Treatment Utilization for Pathological Gambling. European Addiction Research, 2014, 20, 167-173.	2.4	16
51	Factor Structure of the Compulsive Internet Use Scale. Cyberpsychology, Behavior, and Social Networking, 2014, 17, 46-51.	3.9	23
52	Reducing tobacco smoke exposure in children aged below 4years â€" A randomized controlled trial. Preventive Medicine, 2014, 69, 208-213.	3.4	2
53	Predictors of indoor smoking at young children's homesâ€"a cross-sectional study. European Journal of Pediatrics, 2014, 173, 1187-1191.	2.7	10
54	Predictors of hazardous drinking, tobacco smoking and physical inactivity in vocational school students. BMC Public Health, 2013, 13, 475.	2.9	37

#	Article	IF	CITATIONS
55	Sub-threshold panic attacks and agoraphobic avoidance increase comorbidity of mental disorders: Results from an adult general population sample. Journal of Anxiety Disorders, 2013, 27, 485-493.	3.2	15
56	Comorbid Axis I-disorders among subjects with pathological, problem, or at-risk gambling recruited from the general population in Germany: Results of the PAGE study. Psychiatry Research, 2013, 210, 1065-1070.	3.3	55
57	Excess Mortality of Alcoholâ€Dependent Individuals After 14ÂYears and Mortality Predictors Based on Treatment Participation and Severity of Alcohol Dependence. Alcoholism: Clinical and Experimental Research, 2013, 37, 156-163.	2.4	53
58	Efficacy of a Text Message-Based Smoking Cessation Intervention for Young People: A Cluster Randomized Controlled Trial. Journal of Medical Internet Research, 2013, 15, e171.	4.3	110
59	A Pre-Post Study on the Appropriateness and Effectiveness of a Web- and Text Messaging-Based Intervention to Reduce Problem Drinking in Emerging Adults. Journal of Medical Internet Research, 2013, 15, e196.	4.3	59
60	Adoption, reach and effectiveness of computer-based, practitioner delivered and combined smoking interventions in general medical practices: A three-arm cluster randomized trial. Drug and Alcohol Dependence, 2012, 121, 124-132.	3.2	24
61	Changes in drinking behavior among control group participants in early intervention studies targeting unhealthy alcohol use recruited in general hospitals and general practices. Drug and Alcohol Dependence, 2012, 125, 81-88.	3.2	16
62	Efficacy of a text messaging (SMS) based smoking cessation intervention for adolescents and young adults: Study protocol of a cluster randomised controlled trial. BMC Public Health, 2012, 12, 51.	2.9	53
63	Smokers' Expectations Toward the Engagement of Their General Practitioner in Discussing Lifestyle Behaviors. Journal of Health Communication, 2011, 16, 135-147.	2.4	13
64	Efficacy of an internet program for smoking cessation during and after inpatient rehabilitation treatment: a quasi-randomized controlled trial. Addictive Behaviors, 2011, 36, 1369-1372.	3.0	18
65	Overestimation of Drinking Norms and its Association with Alcohol Consumption in Apprentices. Alcohol and Alcoholism, 2011, 46, 204-209.	1.6	35
66	Multiple dimensions of health locus of control in a representative population sample: ordinal factor analysis and cross-validation of an existing three and a new four factor model. BMC Medical Research Methodology, 2011, 11, 114.	3.1	16
67	Need for cognition as a predictor and a moderator of outcome in a tailored letters smoking cessation intervention Health Psychology, 2010, 29, 367-373.	1.6	18
68	Workplace smoking restrictions: smoking behavior and the intention to change among continuing smokers. International Journal of Public Health, 2010, 55, 599-608.	2.3	8
69	Severity of unhealthy alcohol consumption in medical inpatients and the general population: is the general hospital a suitable place for brief interventions?. International Journal of Public Health, 2010, 55, 637-643.	2.3	11
70	Predictors and moderators of outcome in different brief interventions for smoking cessation in general medical practice. Patient Education and Counseling, 2010, 78, 57-64.	2.2	33
71	Effects of practitioner-delivered brief counseling and computer-generated tailored letters on cigarettes per day among smokers who do not quit—A quasi-randomized controlled trialâ~†. Drug and Alcohol Dependence, 2010, 112, 81-89.	3.2	5
72	Problematischer Alkoholkonsum: Prâalenz und Frühintervention. Public Health Forum, 2010, 18, 29-30.	0.2	0

#	Article	IF	Citations
73	Primary focal dystonia: evidence for distinct neuropsychiatric and personality profiles. Journal of Neurology, Neurosurgery and Psychiatry, 2009, 80, 1176-1179.	1.9	105
74	Continuous individual support of smoking cessation using text messaging: A pilot experimental study. Nicotine and Tobacco Research, 2009, 11, 915-923.	2.6	107
75	Reported Beverage Consumed and Alcohol-Related Diseases among Male Hospital Inpatients with Problem Drinking. Alcohol and Alcoholism, 2009, 44, 216-221.	1.6	2
76	Mental Health and Readiness to Change Smoking Behavior in Daily Smoking Primary Care Patients. International Journal of Behavioral Medicine, 2009, 16, 347-354.	1.7	3
77	Nicotine dependence criteria and nicotine withdrawal symptoms in relation to pain among an adult general population sample. European Journal of Pain, 2009, 13, 82-88.	2.8	15
78	Does the smoking status of general practitioners affect the efficacy of smoking cessation counselling?. Patient Education and Counseling, 2009, 74, 23-28.	2.2	17
79	Differences in help seeking rates after brief intervention for alcohol use disorders in general practice patients with and without comorbid anxiety or depressive disorders. International Journal of Methods in Psychiatric Research, 2008, 17, S74-S77.	2.1	18
80	Intention to quit smoking: Is the partner's smoking status associated with the smoker's intention to quit?. International Journal of Behavioral Medicine, 2008, 15, 328-335.	1.7	32
81	Proactive interventions for smoking cessation in general medical practice: a quasiâ€randomized controlled trial to examine the efficacy of computerâ€tailored letters and physicianâ€delivered brief advice*. Addiction, 2008, 103, 294-304.	3.3	74
82	[Commentary] SUPPORTING GPS IN THE PROVISION OF SMOKING CESSATION INTERVENTIONâ€"FROM RESEARCH TO DAILY ROUTINE. Addiction, 2008, 103, 309-309.	3.3	0
83	Reasons for not using smoking cessation aids. BMC Public Health, 2008, 8, 129.	2.9	36
84	Computer-generated tailored feedback letters for smoking cessation: Theoretical and empirical variability of tailoring. International Journal of Medical Informatics, 2008, 77, 715-722.	3.3	16
85	Evaluation of a telephone-based stepped care intervention for alcohol-related disorders: A randomized controlled trial. Drug and Alcohol Dependence, 2008, 93, 244-251.	3.2	97
86	Effectiveness of brief alcohol interventions for general practice patients with problematic drinking behavior and comorbid anxiety or depressive disorders. Drug and Alcohol Dependence, 2008, 94, 214-220.	3.2	39
87	Computer-tailored smoking cessation intervention in a general population setting in Germany: Outcome of a randomized controlled trial. Nicotine and Tobacco Research, 2008, 10, 371-379.	2.6	16
88	Gender differences in the efficacy of brief interventions with a stepped care approach in general practice patients with alcohol-related disorders. Alcohol and Alcoholism, 2008, 43, 334-340.	1.6	24
89	Does precontemplation represent a homogeneous stage category? A latent class analysis on German smokers Journal of Consulting and Clinical Psychology, 2008, 76, 840-851.	2.0	9
90	Patterns and predictors of change in a smoking intervention study: Latent growth analysis of a multivariate outcome model Health Psychology, 2008, 27, S233-S242.	1.6	8

#	Article	IF	CITATIONS
91	Alcohol Screening in General Practices Using the AUDIT: How Many Response Categories Are Necessary?. European Addiction Research, 2007, 13, 25-30.	2.4	13
92	Development and Evaluation of a Screening Instrument for Alcohol-Use Disorders and At-Risk Drinking: The Brief Alcohol Screening Instrument for Medical Care (BASIC). Journal of Studies on Alcohol and Drugs, 2007, 68, 607-614.	1.0	28
93	Variability of tailoring of a smoking cessation intervention based on the transtheoretical model. Addictive Behaviors, 2007, 32, 3083-3087.	3.0	10
94	Effects of smoking cessation on health care use: Is elevated risk of hospitalization among former smokers attributable to smoking-related morbidity?. Drug and Alcohol Dependence, 2007, 88, 197-203.	3.2	12
95	Stability of subtypes of natural recovery from alcohol dependence after two years. Addiction, 2007, 102, 904-908.	3.3	32
96	Sedative, hypnotic, anxiolytic and opioid medicament use and its co-occurrence with tobacco smoking and alcohol risk drinking in a community sample. BMC Public Health, 2007, 7, 337.	2.9	16
97	Psychometric properties of the †Processes of Change' scale for alcohol misuse and its short form (POC-20). Addictive Behaviors, 2006, 31, 821-832.	3.0	17
98	Tobacco smoking in relation to analgesic drug use in a national adult population sample. Drug and Alcohol Dependence, 2006, 85, 49-55.	3.2	41
99	Psychiatric comorbidity including nicotine dependence among individuals with eating disorder criteria in an adult general population sample. Psychiatry Research, 2006, 141, 71-79.	3.3	19
100	Changes in the "stages of change―as outcome measures of a smoking cessation intervention: A randomized controlled trial. Preventive Medicine, 2006, 43, 101-106.	3.4	23
101	Tobacco smoking in relation to pain in a national general population survey. Preventive Medicine, 2006, 43, 477-481.	3.4	109
102	Alcohol Consumption and Health-Services Utilizationin Germany. Journal of Studies on Alcohol and Drugs, 2006, 67, 429-435.	2.3	17
103	STABILITY OF REMISSION FROM ALCOHOL DEPENDENCE WITHOUT FORMAL HELP. Alcohol and Alcoholism, 2006, 41, 311-314.	1.6	22
104	The Reliability and Validity of the Alcohol Use Disorders Identification Test (AUDIT) in a German General Practice Population Sample. Journal of Studies on Alcohol and Drugs, 2006, 67, 473-481.	2.3	130
105	Predictors of Increased Body Mass Index Following Cessation of Smoking. American Journal on Addictions, 2006, 15, 192-197.	1.4	16
106	Supporting the intention to change health risk behaviors. Zeitschrift Fur Gesundheitswissenschaften, 2006, 14, 377-383.	1.6	1
107	Post-traumatic stress disorder. European Archives of Psychiatry and Clinical Neuroscience, 2006, 256, 299-306.	3.2	136
108	Provision of smoking cessation counseling by general practitioners assisted by training and screening procedure. Patient Education and Counseling, 2006, 63, 232-238.	2,2	39

#	Article	IF	Citations
109	Relation between awareness of circulatory disorders and smoking in a general population health examination. BMC Public Health, 2006, 6, 48.	2.9	17
110	Attitudes towards smoking policies and tobacco control measures in relation to smoking status and smoking behaviour. European Journal of Public Health, 2006, 16, 513-519.	0.3	41
111	Smoking status, obesity and hypertension in a general population sample: a cross-sectional study. QJM - Monthly Journal of the Association of Physicians, 2006, 99, 407-415.	0.5	25
112	Associations of perceived work strain with nicotine dependence in a community sample. Occupational and Environmental Medicine, 2006, 63, 207-211.	2.8	25
113	Psychiatric disorders among at-risk consumers of alcohol in the general population Journal of Studies on Alcohol and Drugs, 2005, 66, 246-253.	2.3	37
114	Association of Smoking and Nicotine Dependence With Trauma and Posttraumatic Stress Disorder in a General Population Sample. Journal of Nervous and Mental Disease, 2005, 193, 843-846.	1.0	126
115	No considerable long-term weight gain after smoking cessation: evidence from a prospective study. European Journal of Cancer Prevention, 2005, 14, 289-295.	1.3	19
116	Stage of Change Transitions and Processes of Change, Decisional Balance, and Self-Efficacy in Smokers: A Transtheoretical Model Validation Using Longitudinal Data Psychology of Addictive Behaviors, 2005, 19, 3-9.	2.1	34
117	Longitudinal relationships among transtheoretical model constructs for smokers in the precontemplation and contemplation stages of change. Annals of Behavioral Medicine, 2005, 30, 12-20.	2.9	5
118	Influence of psychiatric comorbidity in alcoholâ€dependent subjects in a representative population survey on treatment utilization and natural recovery. Addiction, 2005, 100, 405-413.	3.3	55
119	Relationships of Psychiatric Disorders with Overweight and Obesity in an Adult General Population ^{**} . Obesity, 2005, 13, 101-109.	4.0	82
120	Relationships of psychiatric disorders with sleep duration in an adult general population sample. Journal of Psychiatric Research, 2005, 39, 577-583.	3.1	53
121	Gender and age differences among current smokers in a general population survey. BMC Public Health, 2005, 5, 57.	2.9	14
122	INTENTION TO CHANGE DRINKING BEHAVIOUR IN GENERAL PRACTICE PATIENTS WITH PROBLEMATIC DRINKING AND COMORBID DEPRESSION OR ANXIETY. Alcohol and Alcoholism, 2005, 40, 394-400.	1.6	27
123	Consistency or Change in Nicotine Dependence According to the Fagerstrom Test for Nicotine Dependence Over Three Years in a Population Sample. Journal of Addictive Diseases, 2005, 24, 85-100.	1.3	12
124	Gender differences in temptation to drink, self-efficacy to abstain and coping behavior in treated alcohol-dependent individuals: Controlling for severity of dependence. Addiction Research and Theory, 2005, 13, 129-136.	1.9	15
125	Stages of change and other factors in â€`light' cigarette smokers. European Journal of Public Health, 2005, 15, 146-151.	0.3	7
126	Self-rated general health and psychiatric disorders in a general population sample. European Psychiatry, 2005, 20, 223-228.	0.2	7

#	Article	IF	CITATIONS
127	Comparison of a Population-Based Sample of "Risky Drinking―Smokers and Groups Consuming Just One Substance. Substance Use and Misuse, 2005, 40, 1721-1732.	1.4	0
128	Effects of item sequence on the performance of the AUDIT in general practices. Drug and Alcohol Dependence, 2005, 79, 373-377.	3.2	23
129	Autoevaluación de la salud general y trastornos psiquiátricos en una muestra de la población general. European Psychiatry (Ed Española), 2005, 12, 349-354.	0.0	0
130	Prevalence, Characteristics, Associated Mental Disorders and Predictors of DSM-IV Nicotine Dependence. European Addiction Research, 2004, 10, 29-34.	2.4	25
131	Subtyping general population smokers not intending to quit by stages to reduce smoking. Nicotine and Tobacco Research, 2004, 6, 1043-1050.	2.6	13
132	Impact of psychiatric disorders in the general population: satisfaction with life and the influence of comorbidity and disorder duration. Social Psychiatry and Psychiatric Epidemiology, 2004, 39, 435-41.	3.1	43
133	Smoking, nicotine dependence and psychiatric comorbidity—a population-based study including smoking cessation after three years. Drug and Alcohol Dependence, 2004, 76, 287-295.	3.2	197
134	Nicotine dependence, quit attempts, and quitting among smokers in a regional population sample from a country with a high prevalence of tobacco smoking. Preventive Medicine, 2004, 38, 350-358.	3.4	81
135	Self-efficacy to refrain from smoking predicted by major depression and nicotine dependence. Addictive Behaviors, 2004, 29, 857-866.	3.0	45
136	A short form of the Fagerström Test for Nicotine Dependence and the Heaviness of Smoking Index in two adult population samples. Addictive Behaviors, 2004, 29, 1207-1212.	3.0	60
137	Nicotine dependence and lifetime amount of smoking in a population sample. European Journal of Public Health, 2004, 14, 182-185.	0.3	22
138	PrÃ♥ention tabak- und alkoholbezogener Erkrankungen. Public Health Forum, 2004, 12, 29-30.	0.2	1
139	Depressive Disorders Are Related to Nicotine Dependence in the Population But Do Not Necessarily Hamper Smoking Cessation. Journal of Clinical Psychiatry, 2004, 65, 169-176.	2.2	57
140	Depressive disorders were the fourth leading cause of global disease burden in the year 2000. Evidence-Based Mental Health, 2004, 7, 123.	4.5	7
141	Measuring Sense of Coherence with only three items: A useful tool for population surveys. British Journal of Health Psychology, 2003, 8, 409-421.	3.5	62
142	Gender differences in unipolar depression: a general population survey of adults between age 18 to 64 of German nationality. Journal of Affective Disorders, 2003, 77, 203-211.	4.1	50
143	Probabilities of alcohol high-risk drinking, abuse or dependence estimated on grounds of tobacco smoking and nicotine dependence. Addiction, 2003, 98, 805-814.	3.3	99
144	Intentionally reduced smoking among untreated general population smokers: prevalence, stability, prediction of smoking behaviour change and differences between subjects choosing either reduction or abstinence. Addiction, 2003, 98, 1101-1110.	3.3	40

#	Article	IF	CITATIONS
145	Types of natural recovery from alcohol dependence: a cluster analytic approach. Addiction, 2003, 98, 1737-1746.	3.3	106
146	The Fagerström test for nicotine dependence in two adult population samplesâ€"potential influence of lifetime amount of tobacco smoked on the degree of dependence. Drug and Alcohol Dependence, 2003, 71, 1-6.	3.2	37
147	Relation among stage of change, demographic characteristics, smoking history, and nicotine dependence in an adult German population. Preventive Medicine, 2003, 37, 368-374.	3.4	47
148	Alcohol high risk drinking, abuse and dependence among tobacco smoking medical care patients and the general population. Drug and Alcohol Dependence, 2003, 69, 189-195.	3.2	48
149	Specific Somatoform Disorder in the General Population. Psychosomatics, 2003, 44, 304-311.	2.5	48
150	STRENGTH OF THE RELATIONSHIP BETWEEN TOBACCO SMOKING, NICOTINE DEPENDENCE AND THE SEVERITY OF ALCOHOL DEPENDENCE SYNDROME CRITERIA IN A POPULATION-BASED SAMPLE. Alcohol and Alcoholism, 2003, 38, 606-612.	1.6	97
151	Somatoform Pain Disorder in the General Population. Psychotherapy and Psychosomatics, 2003, 72, 88-94.	8.8	48
152	The Role of Family and Partnership in Recovery from Alcohol Dependence: Comparison of Individuals Remitting with and without Formal Help. European Addiction Research, 2002, 8, 122-127.	2.4	26
153	THE READINESS TO CHANGE QUESTIONNAIRE IN SUBJECTS WITH HAZARDOUS ALCOHOL CONSUMPTION, ALCOHOL MISUSE AND DEPENDENCE IN A GENERAL POPULATION SURVEY. Alcohol and Alcoholism, 2002, 37, 362-369.	1.6	38
154	SCREENING FOR ALCOHOL USE DISORDERS AND AT-RISK DRINKING IN THE GENERAL POPULATION: PSYCHOMETRIC PERFORMANCE OF THREE QUESTIONNAIRES. Alcohol and Alcoholism, 2002, 37, 261-268.	1.6	273
155	Remission from alcohol dependence without help: how restrictive should our definition of treatment be?. Journal of Studies on Alcohol and Drugs, 2002, 63, 229-236.	2.3	24
156	Naturalistic Changes in the Readiness to Quit Tobacco Smoking in a German General Population Sample. Preventive Medicine, 2002, 35, 326-333.	3.4	25
157	Screening for mental health: validity of the MHI-5 using DSM-IV Axis I psychiatric disorders as gold standard. Psychiatry Research, 2001, 105, 243-253.	3.3	474
158	Lifetime-comorbidity of obsessive-compulsive disorder and subclinical obsessive-compulsive disorder in northern Germany. European Archives of Psychiatry and Clinical Neuroscience, 2001, 251, 130-135.	3.2	78
159	Factors influencing remission from alcohol dependence without formal help in a representative population sample. Addiction, 2001, 96, 1327-1336.	3.3	43
160	The association between degree of nicotine dependence and other health behaviours: Findings from a German general population study. European Journal of Public Health, 2001, 11, 450-452.	0.3	29
161	Gender differences in natural recovery from alcohol dependence Journal of Studies on Alcohol and Drugs, 2000, 61, 783-786.	2.3	34
162	The Composite International Diagnostic Interview: feasibility and necessity of editing and interviewer training in general population surveys. International Journal of Methods in Psychiatric Research, 2000, 9, 32-42.	2.1	28

#	Article	IF	CITATION
163	Studies on natural recovery from alcohol dependence: sample selection bias by media solicitation?. Addiction, 2000, 95, 765-775.	3.3	65
164	Maintenance Factors of Recovery From Alcohol Dependence in Treated and Untreated Individuals. Alcoholism: Clinical and Experimental Research, 2000, 24, 1773-1777.	2.4	37
165	Prevalence of alcohol consumption, abuse and dependence in a country with high per capita consumption: findings from the German TACOS study. Social Psychiatry and Psychiatric Epidemiology, 2000, 35, 539-547.	3.1	89
166	Prevalence, quality of life and psychosocial function in obsessive-compulsive disorder and subclinical obsessive-compulsive disorder in northern Germany. European Archives of Psychiatry and Clinical Neuroscience, 2000, 250, 262-268.	3.2	119
167	Maintenance Factors of Recovery From Alcohol Dependence in Treated and Untreated Individuals. Alcoholism: Clinical and Experimental Research, 2000, 24, 1773-1777.	2.4	0
168	Motivation to change drinking behavior: comparison of alcohol-dependent individuals in a general hospital and a general population sample. General Hospital Psychiatry, 1999, 21, 348-353.	2.4	54
169	Gender and Age in Gambling Participation, Gambling Onset, and Problematic Gambling in a General Population Sample: Empirical Findings from Germany. International Journal of Mental Health and Addiction, 0 , 1 .	7.4	O