Scott B Reeder

List of Publications by Year in descending order

Source: https://exaly.com/author-pdf/9445281/publications.pdf

Version: 2024-02-01

13099 20,933 346 68 citations h-index papers

131 g-index 353 353 353 15229 docs citations times ranked citing authors all docs

12946

| # | Article | IF | CITATIONS |
|----|---|-----|-----------|
| 1 | Abdominal fellowship-trained versus generalist radiologist accuracy when interpreting MR and CT for the diagnosis of appendicitis. European Radiology, 2022, 32, 533-541. | 4.5 | O |
| 2 | Spectroscopyâ€based multiâ€parametric quantification in subjects with liver iron overload at 1.5T and 3T. Magnetic Resonance in Medicine, 2022, 87, 597-613. | 3.0 | 6 |
| 3 | Editorial for "Effects of <scp>B₁</scp> Heterogeneity on Spin <scp>Echoâ€Based</scp> Liver Iron Estimates― Journal of Magnetic Resonance Imaging, 2022, 55, 1426-1427. | 3.4 | O |
| 4 | Cardiovascular Magnetic Resonance for Patients With COVID-19. JACC: Cardiovascular Imaging, 2022, 15, 685-699. | 5.3 | 79 |
| 5 | Abdominal applications of quantitative 4D flow MRI. Abdominal Radiology, 2022, 47, 3229-3250. | 2.1 | 10 |
| 6 | Proton density water fraction as a reproducible MRâ€based measurement of breast density. Magnetic Resonance in Medicine, 2022, 87, 1742-1757. | 3.0 | 6 |
| 7 | Clinical Applications of 4D Flow MRI in the Portal Venous System. Magnetic Resonance in Medical Sciences, 2022, 21, 340-353. | 2.0 | 11 |
| 8 | Addressing concomitant gradient phase errors in timeâ€interleaved chemical shiftâ€encoded MRI fat fraction and R 2 * mapping with a passâ€specific phase fitting method. Magnetic Resonance in Medicine, 2022, , . | 3.0 | 2 |
| 9 | <scp>Magnetic Resonance</scp> Imaging During a Pandemic: Recommendations by the <scp>ISMRM</scp> Safety Committee. Journal of Magnetic Resonance Imaging, 2022, 55, 1322-1339. | 3.4 | 3 |
| 10 | Editorial for "Bias, Repeatability and Reproducibility of Liver <scp>T1</scp> Mapping With Variable Flip Anglesâ€. Journal of Magnetic Resonance Imaging, 2022, 56, 1053-1054. | 3.4 | 0 |
| 11 | Improved free-breathing liver fat and iron quantification using a 2D chemical shift–encoded MRI with flip angle modulation and motion-corrected averaging. European Radiology, 2022, 32, 5458-5467. | 4.5 | 1 |
| 12 | Characterization of mesenteric and portal hemodynamics using 4D flow MRI: the effects of meals and diurnal variation. Abdominal Radiology, 2022, 47, 2106-2114. | 2.1 | 6 |
| 13 | Determining Biomarkers of Myosteatosis for Sarcopenia and Cachexia Using MRI and Ultrasound. FASEB Journal, 2022, 36, . | 0.5 | 0 |
| 14 | Myosteatosis as a Shared Biomarker for Sarcopenia and Cachexia Using MRI and Ultrasound. Frontiers in Rehabilitation Sciences, 2022, 3, . | 1.2 | 1 |
| 15 | Magnetic Resonance Imaging as an Alternative to <scp>Contrastâ€Enhanced</scp> Computed Tomography to Mitigate Iodinated Contrast Shortages in the United States: Recommendations From the International Society for Magnetic Resonance in Medicine. Journal of Magnetic Resonance Imaging, 2022. 56, 655-656. | 3.4 | 4 |
| 16 | Magnetic Resonance Imaging of Liver Fibrosis, Fat, and Iron. Radiologic Clinics of North America, 2022, 60, 705-716. | 1.8 | 4 |
| 17 | Recommendations for Imaging Patients With Cardiac Implantable Electronic Devices (<scp>CIEDs</scp>). Journal of Magnetic Resonance Imaging, 2021, 53, 1311-1317. | 3.4 | 12 |
| 18 | Feasibility and optimization of ultra-short echo time MRI for improved imaging of IVC-filters at 3.0ÂT. Abdominal Radiology, 2021, 46, 362-372. | 2.1 | 1 |

| # | Article | IF | CITATIONS |
|----|---|--------------|-----------|
| 19 | Complex confounder-corrected R2* mapping for liver iron quantification with MRI. European Radiology, 2021, 31, 264-275. | 4.5 | 20 |
| 20 | Freeâ€breathing liver fat and quantification using motionâ€corrected averaging based on a nonlocal means algorithm. Magnetic Resonance in Medicine, 2021, 85, 653-666. | 3.0 | 2 |
| 21 | A Phase 1 Dose Escalation Study of Neoadjuvant SBRT Plus Elective Nodal Radiation with Concurrent Capecitabine for Resectable Pancreatic Cancer. International Journal of Radiation Oncology Biology Physics, 2021, 109, 458-463. | 0.8 | 8 |
| 22 | B ₀ and B ₁ inhomogeneities in the liver at 1.5 T and 3.0 T. Magnetic Resonance in Medicine, 2021, 85, 2212-2220. | 3.0 | 17 |
| 23 | Design and evaluation of quantitative MRI phantoms to mimic the simultaneous presence of fat, iron, and fibrosis in the liver. Magnetic Resonance in Medicine, 2021, 85, 734-747. | 3.0 | 10 |
| 24 | Diagnosis of Coronavirus Disease 2019 Pneumonia by Using Chest Radiography: Value of Artificial Intelligence. Radiology, 2021, 298, E88-E97. | 7.3 | 102 |
| 25 | Accuracies of Chemical Shift In/Opposed Phase and Chemical Shift Encoded Magnetic Resonance Imaging to Detect Intratumoral Fat in Hepatocellular Carcinoma. Journal of Magnetic Resonance Imaging, 2021, 53, 1791-1802. | 3.4 | 5 |
| 26 | Temperatureâ€corrected proton density fat fraction estimation using chemical shiftâ€encoded MRI in phantoms. Magnetic Resonance in Medicine, 2021, 86, 69-81. | 3.0 | 11 |
| 27 | Linearity and Bias of Proton Density Fat Fraction as a Quantitative Imaging Biomarker: A Multicenter, Multiplatform, Multivendor Phantom Study. Radiology, 2021, 298, 640-651. | 7.3 | 39 |
| 28 | Limits of Fat Quantification in the Presence of Iron Overload. Journal of Magnetic Resonance Imaging, 2021, 54, 1166-1174. | 3 . 4 | 10 |
| 29 | Magnetic resonance imaging versus computed tomography and ultrasound for the diagnosis of female pelvic pathology. Emergency Radiology, 2021, 28, 789-796. | 1.8 | 4 |
| 30 | Portosystemic Shunts: Should We Pay Closer Attention with Cross-Sectional Imaging?. Radiology, 2021, 299, 141-142. | 7.3 | 1 |
| 31 | Magnetic resonance elastography biomarkers for detection of histologic alterations in nonalcoholic fatty liver disease in the absence of fibrosis. European Radiology, 2021, 31, 8408-8419. | 4.5 | 6 |
| 32 | Reproducibility of liver R2* quantification for liver iron quantification from cardiac R2* acquisitions. Abdominal Radiology, 2021, 46, 4200-4209. | 2.1 | 4 |
| 33 | Dual contrast liver MRI: a pictorial illustration. Abdominal Radiology, 2021, 46, 4588-4600. | 2.1 | 3 |
| 34 | Diagnostic Performance of MRI for Esophageal Carcinoma: A Systematic Review and Meta-Analysis. Radiology, 2021, 299, 583-594. | 7.3 | 21 |
| 35 | Multisite multivendor validation of a quantitative MRI and CT compatible fat phantom. Medical Physics, 2021, 48, 4375-4386. | 3.0 | 10 |
| 36 | Myocarditis Associated with mRNA COVID-19 Vaccination. Radiology, 2021, 301, E409-E411. | 7.3 | 48 |

| # | Article | IF | Citations |
|----|--|-------------|-----------|
| 37 | Evaluation for Myocarditis in Competitive Student Athletes Recovering From Coronavirus Disease 2019 With Cardiac Magnetic Resonance Imaging. JAMA Cardiology, 2021, 6, 945. | 6.1 | 161 |
| 38 | Relaxivityâ€iron calibration in hepatic iron overload: Reproducibility and extension of a Monte Carlo model. NMR in Biomedicine, 2021, 34, e4604. | 2.8 | 7 |
| 39 | Quantification of Liver Fat Content with CT and MRI: State of the Art. Radiology, 2021, 301, 250-262. | 7.3 | 77 |
| 40 | Emergence of 3D MR Elastography–based Quantitative Markers for Diffuse Liver Disease. Radiology, 2021, 301, 163-165. | 7.3 | 6 |
| 41 | Ferumoxytol-enhanced MR imaging for differentiating intrapancreatic splenules from other tumors. Abdominal Radiology, 2021, 46, 2003-2013. | 2.1 | 1 |
| 42 | Simultaneous T 1 â€weighted and T 2 â€weighted 3D MRI using RF phaseâ€modulated gradient echo imaging. Magnetic Resonance in Medicine, 2021, 87, 1758. | 3.0 | 0 |
| 43 | Impact of ferumoxytol magnetic resonance imaging on the rhesus macaque maternal–fetal interfaceâ€. Biology of Reproduction, 2020, 102, 434-444. | 2.7 | 5 |
| 44 | Diurnal Variation of Proton Density Fat Fraction in the Liver Using Quantitative Chemical Shift Encoded MRI. Journal of Magnetic Resonance Imaging, 2020, 51, 407-414. | 3.4 | 11 |
| 45 | Combined gadoxetic acid and gadobenate dimeglumine enhanced liver MRI: a parameter optimization study. Abdominal Radiology, 2020, 45, 220-231. | 2.1 | 2 |
| 46 | Sensitivity of quantitative relaxometry and susceptibility mapping to microscopic iron distribution. Magnetic Resonance in Medicine, 2020, 83, 673-680. | 3.0 | 16 |
| 47 | Evaluation of a motionâ€robust 2D chemical shiftâ€encoded technique for R2* and field map quantification in ferumoxytolâ€enhanced MRI of the placenta in pregnant rhesus macaques. Journal of Magnetic Resonance Imaging, 2020, 51, 580-592. | 3.4 | 8 |
| 48 | ACR guidance document on MR safe practices: Updates and critical information 2019. Journal of Magnetic Resonance Imaging, 2020, 51, 331-338. | 3.4 | 61 |
| 49 | Quantitative ferumoxytol-enhanced MRI in pregnancy: A feasibility study in the nonhuman primate. Magnetic Resonance Imaging, 2020, 65, 100-108. | 1.8 | 13 |
| 50 | Pharmacokinetics of Ferumoxytol in the Abdomen and Pelvis: A Dosing Study with 1.5- and 3.0-T MRI Relaxometry. Radiology, 2020, 294, 108-116. | 7. 3 | 13 |
| 51 | T ₁ â€corrected quantitative chemical shiftâ€encoded MRI. Magnetic Resonance in Medicine, 2020, 83, 2051-2063. | 3.0 | 10 |
| 52 | Phaseâ€based T ₂ mapping with gradient echo imaging. Magnetic Resonance in Medicine, 2020, 84, 609-619. | 3.0 | 10 |
| 53 | Accuracy of common proton density fat fraction thresholds for magnitude- and complex-based chemical shift-encoded MRI for assessing hepatic steatosis in patients with obesity. Abdominal Radiology, 2020, 45, 661-671. | 2.1 | 16 |
| 54 | No Cases of Nephrogenic Systemic Fibrosis after Administration of Gadoxetic Acid. Radiology, 2020, 297, 556-562. | 7.3 | 8 |

| # | Article | IF | Citations |
|----|---|-----|-----------|
| 55 | Liver fat quantification: where do we stand?. Abdominal Radiology, 2020, 45, 3386-3399. | 2.1 | 41 |
| 56 | Effect of noise and estimator type on bias for analysis of liver proton density fat fraction. Magnetic Resonance Imaging, 2020, 74, 244-249. | 1.8 | 0 |
| 57 | Motionâ€robust, high‧NR liver fat quantification using a 2D sequential acquisition with a variable flip angle approach. Magnetic Resonance in Medicine, 2020, 84, 2004-2017. | 3.0 | 5 |
| 58 | Gadoxetate-enhanced abbreviated MRI is highly accurate for hepatocellular carcinoma screening. European Radiology, 2020, 30, 6003-6013. | 4.5 | 43 |
| 59 | Prospective comparison of longitudinal change in hepatic proton density fat fraction (PDFF) estimated by magnitude-based MRI (MRI-M) and complex-based MRI (MRI-C). European Radiology, 2020, 30, 5120-5129. | 4.5 | 2 |
| 60 | The Impact of the COVID-19 Pandemic on the Radiology Research Enterprise: Radiology Scientific Expert Panel. Radiology, 2020, 296, E134-E140. | 7.3 | 29 |
| 61 | Interobserver agreement for the direct and indirect signs of pulmonary embolism evaluated using contrast enhanced magnetic angiography. European Journal of Radiology Open, 2020, 7, 100256. | 1.6 | 2 |
| 62 | Value of MRI in medicine: More than just another test?. Journal of Magnetic Resonance Imaging, 2019, 49, e14-e25. | 3.4 | 78 |
| 63 | Comparison of gadoliniumâ€enhanced and ferumoxytolâ€enhanced conventional and UTEâ€MRA for the depiction of the pulmonary vasculature. Magnetic Resonance in Medicine, 2019, 82, 1660-1670. | 3.0 | 14 |
| 64 | Characterizing a short T ₂ * signal component in the liver using ultrashort TE chemical shiftâ€encoded MRI at 1.5T and 3.0T. Magnetic Resonance in Medicine, 2019, 82, 2032-2045. | 3.0 | 7 |
| 65 | Measurement of spleen fat on MRI-proton density fat fraction arises from reconstruction of noise. Abdominal Radiology, 2019, 44, 3295-3303. | 2.1 | 7 |
| 66 | Letter to the Editor: Intrapancreatic Accessory Spleen Masquerading as a Pancreatic Neuroendocrine Tumor. Journal of Gastrointestinal Surgery, 2019, 23, 1717-1718. | 1.7 | 0 |
| 67 | Prospective evaluation of MRI compared with CT for the etiology of abdominal pain in emergency department patients with concern for appendicitis. Journal of Magnetic Resonance Imaging, 2019, 50, 1651-1658. | 3.4 | 4 |
| 68 | Clinical Implementation of a Focused MRI Protocol for Hepatic Fat and Iron Quantification. American Journal of Roentgenology, 2019, 213, 90-95. | 2.2 | 13 |
| 69 | Pilot study on longitudinal change in pancreatic proton density fat fraction during a weightâ€loss surgery program in adults with obesity. Journal of Magnetic Resonance Imaging, 2019, 50, 1092-1102. | 3.4 | 16 |
| 70 | Quantitative MRI Biomarkers of Diffuse Liver Disease. Advances in Clinical Radiology, 2019, 1, 55-69. | 0.2 | 2 |
| 71 | Noncontrast Chest Computed Tomographic Imaging of Obesity and the Metabolic Syndrome. Journal of Thoracic Imaging, 2019, 34, 126-135. | 1.5 | 10 |
| 72 | Monitoring Fatty Liver Disease with MRI Following Bariatric Surgery: A Prospective, Dual-Center Study. Radiology, 2019, 290, 682-690. | 7.3 | 22 |

| # | Article | IF | CITATIONS |
|----|---|-----|-----------|
| 73 | Perfusion of the placenta assessed using arterial spin labeling and ferumoxytol dynamic contrast enhanced magnetic resonance imaging in the rhesus macaque. Magnetic Resonance in Medicine, 2019, 81, 1964-1978. | 3.0 | 23 |
| 74 | Quantification of Liver Function with MRI: Is It Ready?. Radiology, 2019, 290, 134-135. | 7.3 | 3 |
| 75 | Four-dimensional Flow MRI as a Marker for Risk Stratification of Gastroesophageal Varices in Patients with Liver Cirrhosis. Radiology, 2019, 290, 101-107. | 7.3 | 38 |
| 76 | Hepatic steatosis and reduction in steatosis following bariatric weight loss surgery differs between segments and lobes. European Radiology, 2019, 29, 2474-2480. | 4.5 | 11 |
| 77 | Assessment of a high‧NR chemicalâ€shiftâ€encoded MRI with complex reconstruction for proton density fat fraction (PDFF) estimation overall and in the lowâ€fat range. Journal of Magnetic Resonance Imaging, 2019, 49, 229-238. | 3.4 | 9 |
| 78 | Feasibility of high spatiotemporal resolution for an abbreviated 3 <scp>D</scp> radial breast <scp>MRI</scp> protocol. Magnetic Resonance in Medicine, 2018, 80, 1452-1466. | 3.0 | 17 |
| 79 | Prospective Comparison of the Diagnostic Accuracy of MR Imaging versus CT for Acute Appendicitis. Radiology, 2018, 288, 467-475. | 7.3 | 36 |
| 80 | Validation of a motionâ€robust 2D sequential technique for quantification of hepatic proton density fat fraction during free breathing. Journal of Magnetic Resonance Imaging, 2018, 48, 1578-1585. | 3.4 | 16 |
| 81 | Noise properties of proton density fat fraction estimated using chemical shift–encoded MRI. Magnetic Resonance in Medicine, 2018, 80, 685-695. | 3.0 | 17 |
| 82 | Noninvasive, Quantitative Assessment of Liver Fat by MRIâ€PDFF as an Endpoint in NASH Trials. Hepatology, 2018, 68, 763-772. | 7.3 | 299 |
| 83 | Relaxivity of Ferumoxytol at 1.5 T and 3.0 T. Investigative Radiology, 2018, 53, 257-263. | 6.2 | 61 |
| 84 | Surgical planning for living donor liver transplant using 4D flow MRI, computational fluid dynamics and in vitro experiments. Computer Methods in Biomechanics and Biomedical Engineering: Imaging and Visualization, 2018, 6, 545-555. | 1.9 | 24 |
| 85 | Comparison of ferumoxytolâ€based cerebral blood volume estimates using quantitative R ₁ and relaxometry. Magnetic Resonance in Medicine, 2018, 79, 3072-3081. | 3.0 | 7 |
| 86 | MRI proton density fat fraction is robust across the biologically plausible range of triglyceride spectra in adults with nonalcoholic steatohepatitis. Journal of Magnetic Resonance Imaging, 2018, 47, 995-1002. | 3.4 | 27 |
| 87 | Optimization of regionâ€ofâ€interest sampling strategies for hepatic MRI proton density fat fraction quantification. Journal of Magnetic Resonance Imaging, 2018, 47, 988-994. | 3.4 | 20 |
| 88 | Fully phaseâ€encoded MRI near metallic implants using ultrashort echo times and broadband excitation. Magnetic Resonance in Medicine, 2018, 79, 2156-2163. | 3.0 | 9 |
| 89 | Linearity, Bias, and Precision of Hepatic Proton Density Fat Fraction Measurements by Using MR Imaging: A Meta-Analysis. Radiology, 2018, 286, 486-498. | 7.3 | 225 |
| 90 | How bariatric surgery affects liver volume and fat density in NAFLD patients. Surgical Endoscopy and Other Interventional Techniques, 2018, 32, 1675-1682. | 2.4 | 46 |

| # | Article | IF | CITATIONS |
|-----|---|-------------|-----------|
| 91 | Deep Brain Nuclei T1 Shortening after Gadobenate Dimeglumine in Children: Influence of Radiation and Chemotherapy. American Journal of Neuroradiology, 2018, 39, 24-30. | 2.4 | 13 |
| 92 | Vascular input function correction of inflow enhancement for improved pharmacokinetic modeling of liver <scp>DCE</scp> â€ <scp>MRI</scp> . Magnetic Resonance in Medicine, 2018, 79, 3093-3102. | 3.0 | 7 |
| 93 | IDEAL-IQ in an oncologic population: meeting the challenge of concomitant liver fat and liver iron. Cancer Imaging, 2018, 18, 51. | 2.8 | 36 |
| 94 | Interâ€method reproducibility of biexponential <scp>R</scp> ₂ MR relaxometry for estimation of liver iron concentration. Magnetic Resonance in Medicine, 2018, 80, 2691-2701. | 3.0 | 11 |
| 95 | Quantification of Liver Fat Content With Unenhanced MDCT: Phantom and Clinical Correlation With MRI Proton Density Fat Fraction. American Journal of Roentgenology, 2018, 211, W151-W157. | 2.2 | 73 |
| 96 | MRI of the Nontraumatic Acute Abdomen. Gastroenterology Clinics of North America, 2018, 47, 667-690. | 2.2 | 8 |
| 97 | Clinical outcomes after magnetic resonance angiography (MRA) versus computed tomographic angiography (CTA) for pulmonary embolism evaluation. Emergency Radiology, 2018, 25, 469-477. | 1.8 | 15 |
| 98 | MRI liver fat quantification in an oncologic population: the added value of complex chemical shift-encoded MRI. Clinical Imaging, 2018, 52, 193-199. | 1.5 | 14 |
| 99 | Multisite, multivendor validation of the accuracy and reproducibility of proton-density fat-fraction quantification at 1.5T and 3T using a fat-water phantom. Magnetic Resonance in Medicine, 2017, 77, 1516-1524. | 3.0 | 99 |
| 100 | Combined gadoxetic acid and gadofosveset enhanced liver MRI for detection and characterization of liver metastases. European Radiology, 2017, 27, 32-40. | 4.5 | 10 |
| 101 | Guidelines for documentation and consent for nonclinical, nonresearch MRI in human subjects. Journal of Magnetic Resonance Imaging, 2017, 45, 36-41. | 3.4 | 1 |
| 102 | Accelerating fully phaseâ€encoded MRI near metal using multiband radiofrequency excitation. Magnetic Resonance in Medicine, 2017, 77, 1223-1230. | 3.0 | 6 |
| 103 | MR visible localization device for radiographic-pathologic correlation of surgical specimens. Magnetic Resonance Imaging, 2017, 37, 159-163. | 1.8 | 1 |
| 104 | An acetoneâ€based phantom for quantitative diffusion MRI. Journal of Magnetic Resonance Imaging, 2017, 46, 1683-1692. | 3.4 | 13 |
| 105 | Added value of gadoxetic acid-enhanced T1-weighted magnetic resonance cholangiography for the diagnosis of post-transplant biliary complications. European Radiology, 2017, 27, 4415-4425. | 4. 5 | 12 |
| 106 | Prevalence of Fatty Liver Disease and Hepatic Iron Overload in a Northeastern German Population by Using Quantitative MR Imaging. Radiology, 2017, 284, 706-716. | 7.3 | 91 |
| 107 | How to write an original radiological research manuscript. European Radiology, 2017, 27, 4455-4460. | 4.5 | 5 |
| 108 | Crossover comparison of ferumoxytol and gadobenate dimeglumine for abdominal MRâ€angiography at 3.0 tesla: Effects of contrast bolus length and flip angle. Journal of Magnetic Resonance Imaging, 2017, 45, 1617-1626. | 3.4 | 6 |

| # | Article | IF | Citations |
|-----|--|------|-----------|
| 109 | Gadolinium deposition in the brain: summary of evidence and recommendations. Lancet Neurology, The, 2017, 16, 564-570. | 10.2 | 600 |
| 110 | Accuracy of PDFF estimation by magnitudeâ€based and complexâ€based MRI in children with MR spectroscopy as a reference. Journal of Magnetic Resonance Imaging, 2017, 46, 1641-1647. | 3.4 | 19 |
| 111 | Contrast-enhanced pulmonary MRA for the primary diagnosis of pulmonary embolism: current state of the art and future directions. British Journal of Radiology, 2017, 90, 20160901. | 2.2 | 22 |
| 112 | The effects of concomitant gradients on chemical shift encoded MRI. Magnetic Resonance in Medicine, 2017, 78, 730-738. | 3.0 | 22 |
| 113 | Diagnostic Accuracy of MRI Versus CT for the Evaluation of Acute Appendicitis in Children and Young Adults. American Journal of Roentgenology, 2017, 209, 911-919. | 2.2 | 39 |
| 114 | Chelated or dechelated gadolinium deposition – Authors' reply. Lancet Neurology, The, 2017, 16, 955-956. | 10.2 | 5 |
| 115 | Standardized Approach for ROI-Based Measurements of Proton Density Fat Fraction and R2* in the Liver. American Journal of Roentgenology, 2017, 209, 592-603. | 2.2 | 68 |
| 116 | Comparison of radial 4D Flow-MRI with perivascular ultrasound to quantify blood flow in the abdomen and introduction of a porcine model of pre-hepatic portal hypertension. European Radiology, 2017, 27, 5316-5324. | 4.5 | 21 |
| 117 | MRIâ€based quantitative susceptibility mapping (QSM) and R2* mapping of liver iron overload: Comparison with SQUIDâ€based biomagnetic liver susceptometry. Magnetic Resonance in Medicine, 2017, 78, 264-270. | 3.0 | 61 |
| 118 | Intravenous Gadoxetate Disodium Administration Reduces Breath-holding Capacity in the Hepatic Arterial Phase: A Multi-Center Randomized Placebo-controlled Trial. Radiology, 2017, 282, 361-368. | 7.3 | 46 |
| 119 | Quantification of liver fat in the presence of iron overload. Journal of Magnetic Resonance Imaging, 2017, 45, 428-439. | 3.4 | 39 |
| 120 | Externally calibrated parallel imaging for 3D multispectral imaging near metallic implants using broadband ultrashort echo time imaging. Magnetic Resonance in Medicine, 2017, 77, 2303-2309. | 3.0 | 7 |
| 121 | Accuracy of Liver Fat Quantification With Advanced CT, MRI, and Ultrasound Techniques: Prospective Comparison With MR Spectroscopy. American Journal of Roentgenology, 2017, 208, 92-100. | 2.2 | 180 |
| 122 | Thrombus-mimicking artifacts in two-point Dixon MRI: Prevalence, appearance, and severity. Journal of Magnetic Resonance Imaging, 2017, 45, 229-236. | 3.4 | 5 |
| 123 | High SNR Acquisitions Improve the Repeatability of Liver Fat Quantification Using Confounder-corrected Chemical Shift-encoded MR Imaging. Magnetic Resonance in Medical Sciences, 2017, 16, 332-339. | 2.0 | 11 |
| 124 | Intraindividual Crossover Comparison of Gadoxetic Acid Dose for Liver MRI in Normal Volunteers. Magnetic Resonance in Medical Sciences, 2016, 15, 60-72. | 2.0 | 13 |
| 125 | Sensitivity of chemical shiftâ€encoded fat quantification to calibration of fat MR spectrum. Magnetic Resonance in Medicine, 2016, 75, 845-851. | 3.0 | 52 |
| 126 | Systematic review and metaâ€analysis of the accuracy of MRI to diagnose appendicitis in the general population. Journal of Magnetic Resonance Imaging, 2016, 43, 1346-1354. | 3.4 | 34 |

| # | Article | IF | Citations |
|-----|---|-------------|-----------|
| 127 | Quantification of liver proton-density fat fraction in 7.1T preclinical MR systems: Impact of the fitting technique. Journal of Magnetic Resonance Imaging, 2016, 44, 1425-1431. | 3.4 | О |
| 128 | Longitudinal Monitoring of Hepatic Blood Flow before and after TIPS by Using 4D-Flow MR Imaging. Radiology, 2016, 281, 574-582. | 7.3 | 41 |
| 129 | Emerging Applications of Abdominal 4D Flow MRI. American Journal of Roentgenology, 2016, 207, 58-66. | 2.2 | 39 |
| 130 | Contrast-Enhanced Abdominal MRI for Suspected Appendicitis: How We Do It. American Journal of Roentgenology, 2016, 207, 49-57. | 2.2 | 17 |
| 131 | Use of chemical shift encoded magnetic resonance imaging (CSE-MRI) for high resolution fat-suppressed imaging of the brachial and lumbosacral plexuses. European Journal of Radiology, 2016, 85, 1199-1207. | 2.6 | 8 |
| 132 | Trends in the Use of Medical Imaging to Diagnose Appendicitis at an Academic Medical Center. Journal of the American College of Radiology, 2016, 13, 1050-1056. | 1.8 | 14 |
| 133 | Combined gadoxetic acid and gadofosveset enhanced liver MRI: A feasibility and parameter optimization study. Magnetic Resonance in Medicine, 2016, 75, 318-328. | 3.0 | 10 |
| 134 | Safety and technique of ferumoxytol administration for MRI. Magnetic Resonance in Medicine, 2016, 75, 2107-2111. | 3.0 | 171 |
| 135 | Incidence of actionable findings on contrast enhanced magnetic resonance angiography ordered for pulmonary embolism evaluation. European Journal of Radiology, 2016, 85, 1383-1389. | 2.6 | 14 |
| 136 | Mathematical optimization of contrast concentration for t ₁ â€weighted spoiled gradient echo imaging. Magnetic Resonance in Medicine, 2016, 75, 1556-1564. | 3.0 | 9 |
| 137 | Gadolinium Deposition in the Brain: Do We Know Enough to Change Practice?. Radiology, 2016, 279, 323-326. | 7.3 | 26 |
| 138 | Quantitative Imaging Biomarkers of NAFLD. Digestive Diseases and Sciences, 2016, 61, 1337-1347. | 2.3 | 70 |
| 139 | Contrast enhanced pulmonary magnetic resonance angiography for pulmonary embolism: Building a successful program. European Journal of Radiology, 2016, 85, 553-563. | 2.6 | 32 |
| 140 | An Investigation of Transient Severe Motion Related to Gadoxetic Acid–enhanced MR Imaging. Radiology, 2016, 279, 93-102. | 7.3 | 77 |
| 141 | Pulmonary Embolism Detection with Three-dimensional Ultrashort Echo Time MR Imaging: Experimental Study in Canines. Radiology, 2016, 278, 413-421. | 7.3 | 28 |
| 142 | Non-contrast-enhanced MRA of renal artery stenosis: validation against DSA in a porcine model. European Radiology, 2016, 26, 547-555. | 4. 5 | 28 |
| 143 | Thermogenic profiling using magnetic resonance imaging of dermal and other adipose tissues. JCI Insight, 2016, 1, e87146. | 5.0 | 26 |
| 144 | Quantification of liver fat with respiratoryâ€gated quantitative chemical shift encoded MRI. Journal of Magnetic Resonance Imaging, 2015, 42, 1241-1248. | 3.4 | 24 |

| # | Article | IF | CITATIONS |
|-----|--|-------------|-----------|
| 145 | Impaired regulation of portal venous flow in response to a meal challenge as quantified by 4D flow MRI. Journal of Magnetic Resonance Imaging, 2015, 42, spcone-spcone. | 3.4 | O |
| 146 | Flow-induced signal misallocation artifacts in two-point fat-water chemical shift MRI. Magnetic Resonance in Medicine, 2015, 73, 1926-1931. | 3.0 | 6 |
| 147 | Combined dynamic contrast-enhanced liver MRI and MRA using interleaved variable density sampling. Magnetic Resonance in Medicine, 2015, 73, 973-983. | 3.0 | 7 |
| 148 | Quantitative magnetic resonance imaging of hepatic steatosis: Validation in ex vivo human livers. Hepatology, 2015, 62, 1444-1455. | 7. 3 | 128 |
| 149 | On confirmation bias in imaging research. Journal of Magnetic Resonance Imaging, 2015, 41, 1163-1164. | 3.4 | 7 |
| 150 | Reproducibility of MRâ€based liver fat quantification across field strength: Sameâ€day comparison between 1.5T and 3T in obese subjects. Journal of Magnetic Resonance Imaging, 2015, 42, 811-817. | 3.4 | 67 |
| 151 | Impaired regulation of portal venous flow in response to a meal challenge as quantified by 4D flow MRI. Journal of Magnetic Resonance Imaging, 2015, 42, 1009-1017. | 3.4 | 48 |
| 152 | Improving chemical shift encoded water–fat separation using objectâ€based information of the magnetic field inhomogeneity. Magnetic Resonance in Medicine, 2015, 73, 597-604. | 3.0 | 27 |
| 153 | Quantitative susceptibility mapping in the abdomen as an imaging biomarker of hepatic iron overload. Magnetic Resonance in Medicine, 2015, 74, 673-683. | 3.0 | 98 |
| 154 | Characterizing the limits of MRI near metallic prostheses. Magnetic Resonance in Medicine, 2015, 74, 1564-1573. | 3.0 | 19 |
| 155 | In Nonobese Girls, Waist Circumference as a Predictor of Insulin Resistance Is Comparable to MRI Fat Measures and Superior to BMI. Hormone Research in Paediatrics, 2015, 84, 258-265. | 1.8 | 9 |
| 156 | Cytochrome P450 1B1: An unexpected modulator of liver fatty acid homeostasis. Archives of Biochemistry and Biophysics, 2015, 571, 21-39. | 3.0 | 42 |
| 157 | Magnetic Resonance Angiography of the Upper Extremity. Magnetic Resonance Imaging Clinics of North America, 2015, 23, 479-493. | 1.1 | 7 |
| 158 | Pancreatic Steatosis Demonstrated at MR Imaging in the General Population: Clinical Relevance. Radiology, 2015, 276, 129-136. | 7. 3 | 113 |
| 159 | Dermal white adipose tissue: a new component of the thermogenic response. Journal of Lipid Research, 2015, 56, 2061-2069. | 4.2 | 104 |
| 160 | High specificity targeting and detection of human neuroblastoma using multifunctional anti-GD2 iron-oxide nanoparticles. Nanomedicine, 2015, 10, 2973-2988. | 3.3 | 18 |
| 161 | Proton density fat-fraction is an accurate biomarker of hepatic steatosis in adolescent girls and young women. European Radiology, 2015, 25, 2921-2930. | 4.5 | 54 |
| 162 | Longitudinal Changes in Liver Fat Content in Asymptomatic Adults: Hepatic Attenuation on Unenhanced CT as an Imaging Biomarker for Steatosis. American Journal of Roentgenology, 2015, 205, 1167-1172. | 2.2 | 34 |

| # | Article | IF | Citations |
|-----|--|-----|-----------|
| 163 | Effect of hepatocyte-specific gadolinium-based contrast agents on hepatic fat-fraction and R2âŽ. Magnetic Resonance Imaging, 2015, 33, 43-50. | 1.8 | 16 |
| 164 | Hepatobiliary MR contrast agents in hypovascular hepatocellular carcinoma. Journal of Magnetic Resonance Imaging, 2015, 41, 251-265. | 3.4 | 46 |
| 165 | MR system operator: Recommended minimum requirements for performing MRI in human subjects in a research setting. Journal of Magnetic Resonance Imaging, 2015, 41, 899-902. | 3.4 | 10 |
| 166 | Effects of Inhaled Fluticasone on Upper Airway during Sleep and Wakefulness in Asthma: A Pilot Study. Journal of Clinical Sleep Medicine, 2014, 10, 183-193. | 2.6 | 54 |
| 167 | Highâ€spatial and highâ€temporal resolution dynamic contrastâ€enhanced perfusion imaging of the liver with timeâ€resolved threeâ€dimensional radial MRI. Magnetic Resonance in Medicine, 2014, 71, 934-941. | 3.0 | 29 |
| 168 | Highâ€resolution 3D radial bSSFP with IDEAL. Magnetic Resonance in Medicine, 2014, 71, 95-104. | 3.0 | 15 |
| 169 | T1 bias in chemical shiftâ €e ncoded liver fatâ€fraction: Role of the flip angle. Journal of Magnetic Resonance Imaging, 2014, 40, 875-883. | 3.4 | 38 |
| 170 | Navigator flip angle optimization for freeâ€breathing T1â€weighted hepatobiliary phase imaging with gadoxetic acid. Journal of Magnetic Resonance Imaging, 2014, 40, 1129-1136. | 3.4 | 6 |
| 171 | Ethnic differences in the effects of hepatic fat deposition on insulin resistance in nonobese middle school girls. Obesity, 2014, 22, 243-248. | 3.0 | 11 |
| 172 | Effect of flip angle on the accuracy and repeatability of hepatic proton density fat fraction estimation by complex dataâ€based, T1â€independent, T2*â€corrected, spectrumâ€modeled MRI. Journal of Magnetic Resonance Imaging, 2014, 39, 440-447. | 3.4 | 43 |
| 173 | Effect of temporal resolution on 4D flow MRI in the portal circulation. Journal of Magnetic Resonance Imaging, 2014, 39, 819-826. | 3.4 | 28 |
| 174 | On the confounding effect of temperature on chemical shiftâ€encoded fat quantification. Magnetic Resonance in Medicine, 2014, 72, 464-470. | 3.0 | 56 |
| 175 | Effect of temporal resolution on 4D flow MRI in the portal circulation. Journal of Magnetic Resonance Imaging, 2014, 39, spcone-spcone. | 3.4 | 2 |
| 176 | Wholeâ€heart chemical shift encoded water–fat MRI. Magnetic Resonance in Medicine, 2014, 72, 718-725. | 3.0 | 6 |
| 177 | Design of kâ€space channel combination kernels and integration with parallel imaging. Magnetic Resonance in Medicine, 2014, 71, 2139-2154. | 3.0 | 12 |
| 178 | Consensus report from the 6th International forum for liver MRI using gadoxetic acid. Journal of Magnetic Resonance Imaging, 2014, 40, 516-529. | 3.4 | 40 |
| 179 | Non-Contrast Enhanced 3D SSFP MRA of the Renal Allograft Vasculature: A Comparison Between Radial Linear Combination and Cartesian Inflow-Weighted Acquisitions. Magnetic Resonance Imaging, 2014, 32, 190-195. | 1.8 | 9 |
| 180 | Optimal Timing and Diagnostic Adequacy of Hepatocyte Phase Imaging with Gadoxetate-Enhanced Liver MRI. Academic Radiology, 2014, 21, 726-732. | 2.5 | 23 |

| # | Article | IF | Citations |
|-----|---|-----|-----------|
| 181 | New and Improved Imaging Modalities for NAFLD. Current Hepatology Reports, 2014, 13, 88-96. | 0.9 | 1 |
| 182 | CME update: Review articles and commentaries in JMRI. Journal of Magnetic Resonance Imaging, 2014, 40, 778-778. | 3.4 | 1 |
| 183 | Pulmonary MRA: Differentiation of pulmonary embolism from truncation artefact. European Radiology, 2014, 24, 1942-1949. | 4.5 | 16 |
| 184 | Natural History of Hepatic Steatosis: Observed Outcomes for Subsequent Liver and Cardiovascular Complications. American Journal of Roentgenology, 2014, 202, 752-758. | 2.2 | 68 |
| 185 | Quantification of liver iron with MRI: State of the art and remaining challenges. Journal of Magnetic Resonance Imaging, 2014, 40, 1003-1021. | 3.4 | 208 |
| 186 | Gadolinium-based contrast agents: What does "single-dose―mean anymore?. Journal of Magnetic Resonance Imaging, 2014, 39, 1343-1345. | 3.4 | 2 |
| 187 | Reproducibility of Cerebrospinal Venous Blood Flow and Vessel Anatomy with the Use of Phase Contrast-Vastly Undersampled Isotropic Projection Reconstruction and Contrast-Enhanced MRA. American Journal of Neuroradiology, 2014, 35, 999-1006. | 2.4 | 23 |
| 188 | Quantitative chemical shift-encoded MRI is an accurate method to quantify hepatic steatosis. Journal of Magnetic Resonance Imaging, 2014, 39, 1494-1501. | 3.4 | 78 |
| 189 | Quantitative hepatic perfusion modeling using DCEâ€MRI with sequential breathholds. Journal of Magnetic Resonance Imaging, 2014, 39, 853-865. | 3.4 | 13 |
| 190 | Predicting Hepatic Steatosis in a Racially and Ethnically Diverse Cohort of Adolescent Girls. Journal of Pediatrics, 2014, 165, 319-325.e1. | 1.8 | 30 |
| 191 | Spectrally resolved fully phaseâ€encoded threeâ€dimensional fast spinâ€echo imaging. Magnetic Resonance in Medicine, 2014, 71, 681-690. | 3.0 | 13 |
| 192 | Accelerating sequences in the presence of metal by exploiting the spatial distribution of offâ€resonance. Magnetic Resonance in Medicine, 2014, 72, 1658-1667. | 3.0 | 11 |
| 193 | Application of direct virtual coil to dynamic contrastâ€enhanced MRI and MR angiography with dataâ€driven parallel imaging. Magnetic Resonance in Medicine, 2014, 71, 783-789. | 3.0 | 2 |
| 194 | Primer on magnetic resonance imaging of the liver. Clinical Liver Disease, 2014, 4, 120-123. | 2.1 | 1 |
| 195 | Emerging quantitative MRI biomarkers of diffuse liver disease. Clinical Liver Disease, 2014, 4, 129-132. | 2.1 | 4 |
| 196 | Reduction of image noise in low tube current dynamic CT myocardial perfusion imaging using HYPR processing: A timeâ€attenuation curve analysis. Medical Physics, 2013, 40, 011904. | 3.0 | 16 |
| 197 | Proton-density fat fraction and simultaneous R2* estimation as an MRI tool for assessment of osteoporosis. European Radiology, 2013, 23, 3432-3439. | 4.5 | 106 |
| 198 | Clinical Implications of Non–Contrast-Enhanced Computed Tomography for Follow-Up After Endovascular Abdominal Aortic Aneurysm Repair. Annals of Vascular Surgery, 2013, 27, 1042-1048. | 0.9 | 11 |

| # | Article | IF | Citations |
|-----|--|-----|-----------|
| 199 | R2* estimation using "inâ€phase―echoes in the presence of fat: The effects of complex spectrum of fat. Journal of Magnetic Resonance Imaging, 2013, 37, 717-726. | 3.4 | 40 |
| 200 | Comparison of <i>R</i> ₂ * correction methods for accurate fat quantification in fatty liver. Journal of Magnetic Resonance Imaging, 2013, 37, 414-422. | 3.4 | 47 |
| 201 | Fat confounds the observed apparent diffusion coefficient in patients with hepatic steatosis. Magnetic Resonance in Medicine, 2013, 69, 545-552. | 3.0 | 39 |
| 202 | Successful Computed Tomography Angiogram Through Tibial Intraosseous Access: A Case Report. Journal of Emergency Medicine, 2013, 45, 182-185. | 0.7 | 15 |
| 203 | Load-dependent variations in knee kinematics measured with dynamic MRI. Journal of Biomechanics, 2013, 46, 2045-2052. | 2.1 | 35 |
| 204 | In vivo validation of 4D flow MRI for assessing the hemodynamics of portal hypertension. Journal of Magnetic Resonance Imaging, 2013, 37, 1100-1108. | 3.4 | 93 |
| 205 | Improved fat water separation with water selective inversion pulse for inversion recovery imaging in cardiac MRI. Journal of Magnetic Resonance Imaging, 2013, 37, 484-490. | 3.4 | 7 |
| 206 | Quantification of Thoracic Blood Flow Using Volumetric Magnetic Resonance Imaging With Radial Velocity Encoding. Investigative Radiology, 2013, 48, 819-825. | 6.2 | 44 |
| 207 | Adipose tissue MRI for quantitative measurement of central obesity. Journal of Magnetic Resonance Imaging, 2013, 37, 707-716. | 3.4 | 41 |
| 208 | The evolving landscape of self-assessment continuing medical education (SA-CME). Journal of Magnetic Resonance Imaging, 2013, 38, 509-510. | 3.4 | 1 |
| 209 | MRI for acute chest pain: Current state of the Art. Journal of Magnetic Resonance Imaging, 2013, 37, 1290-1300. | 3.4 | 16 |
| 210 | Effectiveness of MR angiography for the primary diagnosis of acute pulmonary embolism: Clinical outcomes at 3 months and 1 year. Journal of Magnetic Resonance Imaging, 2013, 38, 914-925. | 3.4 | 61 |
| 211 | In vivo validation of 4D flow MRI for assessing the hemodynamics of portal hypertension. Journal of Magnetic Resonance Imaging, 2013, 37, spcone-spcone. | 3.4 | 0 |
| 212 | Magnetic susceptibility as a $\langle i\rangle B\langle i\rangle \langle sub\rangle 0\langle sub\rangle$ field strength independent MRI biomarker of liver iron overload. Magnetic Resonance in Medicine, 2013, 70, 648-656. | 3.0 | 36 |
| 213 | Multipeak fatâ€corrected complex R2* relaxometry: Theory, optimization, and clinical validation. Magnetic Resonance in Medicine, 2013, 70, 1319-1331. | 3.0 | 115 |
| 214 | Emerging quantitative magnetic resonance imaging biomarkers of hepatic steatosis. Hepatology, 2013, 58, 1877-1880. | 7.3 | 42 |
| 215 | Quantification of Hepatic Steatosis With Dual-Energy Computed Tomography. Investigative Radiology, 2012, 47, 603-610. | 6.2 | 72 |
| 216 | Effect of Multipeak Spectral Modeling of Fat for Liver Iron and Fat Quantification: Correlation of Biopsy with MR Imaging Results. Radiology, 2012, 265, 133-142. | 7.3 | 169 |

| # | Article | IF | Citations |
|-----|--|-----|-----------|
| 217 | 4D cardiovascular magnetic resonance velocity mapping of alterations of right heart flow patterns and main pulmonary artery hemodynamics in tetralogy of Fallot. Journal of Cardiovascular Magnetic Resonance, 2012, 14, 16. | 3.3 | 129 |
| 218 | Presurgical Localization of Parathyroid Adenomas with Magnetic Resonance Imaging at 3.0 T: An Adjunct Method to Supplement Traditional Imaging. Annals of Surgical Oncology, 2012, 19, 981-989. | 1.5 | 56 |
| 219 | Hepatobiliary MR imaging with gadoliniumâ€based contrast agents. Journal of Magnetic Resonance Imaging, 2012, 35, 492-511. | 3.4 | 121 |
| 220 | Waterâ€silicone separated volumetric MR acquisition for rapid assessment of breast implants. Journal of Magnetic Resonance Imaging, 2012, 35, 1216-1221. | 3.4 | 11 |
| 221 | Validation of MRI biomarkers of hepatic steatosis in the presence of iron overload in the ob/ob mouse. Journal of Magnetic Resonance Imaging, 2012, 35, 844-851. | 3.4 | 41 |
| 222 | "MR Physics for Clinicians―Series: Enhancement for the JMRI CME Program. Journal of Magnetic Resonance Imaging, 2012, 35, 997-997. | 3.4 | 1 |
| 223 | Gadoxetic acid–enhanced T1â€weighted MR cholangiography in primary sclerosing cholangitis. Journal of Magnetic Resonance Imaging, 2012, 36, 632-640. | 3.4 | 34 |
| 224 | Characterization of hepatic adenoma and focal nodular hyperplasia with gadoxetic acid. Journal of Magnetic Resonance Imaging, 2012, 36, 686-696. | 3.4 | 70 |
| 225 | High resolution navigated threeâ€dimensional T ₁ â€weighted hepatobiliary MRI using gadoxetic acid optimized for 1.5 tesla. Journal of Magnetic Resonance Imaging, 2012, 36, 890-899. | 3.4 | 51 |
| 226 | Proton density fatâ€fraction: A standardized mrâ€based biomarker of tissue fat concentration. Journal of Magnetic Resonance Imaging, 2012, 36, 1011-1014. | 3.4 | 385 |
| 227 | Addressing phase errors in fatâ€water imaging using a mixed magnitude/complex fitting method. Magnetic Resonance in Medicine, 2012, 67, 638-644. | 3.0 | 105 |
| 228 | Robust multipoint waterâ€fat separation using fat likelihood analysis. Magnetic Resonance in Medicine, 2012, 67, 1065-1076. | 3.0 | 23 |
| 229 | Improved least squares MR image reconstruction using estimates of <i>kâ€</i> Space data consistency. Magnetic Resonance in Medicine, 2012, 67, 1600-1608. | 3.0 | 42 |
| 230 | R mapping in the presence of macroscopic $\langle i \rangle B \langle i \rangle \langle sub \rangle 0 \langle sub \rangle$ field variations. Magnetic Resonance in Medicine, 2012, 68, 830-840. | 3.0 | 80 |
| 231 | Variations in T2* and fat content of murine brown and white adipose tissues by chemical-shift MRI. Magnetic Resonance Imaging, 2012, 30, 323-329. | 1.8 | 42 |
| 232 | Time-Resolved Interventional Cardiac C-arm Cone-Beam CT: An Application of the PICCS Algorithm. IEEE Transactions on Medical Imaging, 2012, 31, 907-923. | 8.9 | 66 |
| 233 | On the performance of <i>T</i> ₂ * correction methods for quantification of hepatic fat content. Magnetic Resonance in Medicine, 2012, 67, 389-404. | 3.0 | 44 |
| 234 | Quantification of Hepatic Steatosis with T1-independent, T2*-corrected MR Imaging with Spectral Modeling of Fat: Blinded Comparison with MR Spectroscopy. Radiology, 2011, 258, 767-775. | 7.3 | 345 |

| # | Article | IF | CITATIONS |
|-----|---|-----|-----------|
| 235 | PULMONARY ANGIOGRAPHY WITH 64â€MULTIDETECTORâ€ROW COMPUTED TOMOGRAPHY IN NORMAL DOGS. Veterinary Radiology and Ultrasound, 2011, 52, 362-367. | 0.9 | 13 |
| 236 | 64â€MULTIDETECTOR COMPUTED TOMOGRAPHIC ANGIOGRAPHY OF THE CANINE CORONARY ARTERIES. Veterinary Radiology and Ultrasound, 2011, 52, 507-515. | 0.9 | 19 |
| 237 | Effects of postprandial state and mesenteric blood flow on the repeatability of MR elastography in asymptomatic subjects. Journal of Magnetic Resonance Imaging, 2011, 33, 239-244. | 3.4 | 46 |
| 238 | T ₁ independent, T ₂ [*] corrected chemical shift based fat–water separation with multiâ€peak fat spectral modeling is an accurate and precise measure of hepatic steatosis. Journal of Magnetic Resonance Imaging, 2011, 33, 873-881. | 3.4 | 183 |
| 239 | Quantitative assessment of liver fat with magnetic resonance imaging and spectroscopy. Journal of Magnetic Resonance Imaging, 2011, 34, 729-749. | 3.4 | 613 |
| 240 | Fourâ€dimensional velocity mapping of the hepatic and splanchnic vasculature with radial sampling at 3 tesla: A feasibility study in portal hypertension. Journal of Magnetic Resonance Imaging, 2011, 34, 577-584. | 3.4 | 50 |
| 241 | Optimized highâ€resolution contrastâ€enhanced hepatobiliary imaging at 3 tesla: A crossâ€over comparison of gadobenate dimeglumine and gadoxetic acid. Journal of Magnetic Resonance Imaging, 2011, 34, 585-594. | 3.4 | 55 |
| 242 | Quantitative assessment of liver fat with magnetic resonance imaging and spectroscopy. Journal of Magnetic Resonance Imaging, 2011, 34, 729-749. | 3.4 | 359 |
| 243 | Interleaved variable density sampling with a constrained parallel imaging reconstruction for dynamic contrastâ€enhanced MR angiography. Magnetic Resonance in Medicine, 2011, 66, 428-436. | 3.0 | 19 |
| 244 | Combination of complexâ€based and magnitudeâ€based multiecho waterâ€fat separation for accurate quantification of fatâ€fraction. Magnetic Resonance in Medicine, 2011, 66, 199-206. | 3.0 | 166 |
| 245 | Constraining the initial phase in water–fat separation. Magnetic Resonance Imaging, 2011, 29, 216-221. | 1.8 | 34 |
| 246 | Clinical Usefulness of Adding 3D Cartilage Imaging Sequences to a Routine Knee MR Protocol. American Journal of Roentgenology, 2011, 196, 159-167. | 2.2 | 45 |
| 247 | Noninvasive Assessment of Transstenotic Pressure Gradients in Porcine Renal Artery Stenoses by Using Vastly Undersampled Phase-Contrast MR Angiography. Radiology, 2011, 261, 266-273. | 7.3 | 56 |
| 248 | Renal Arteries: Isotropic, High-Spatial-Resolution, Unenhanced MR Angiography with Three-dimensional Radial Phase Contrast. Radiology, 2011, 258, 254-260. | 7.3 | 51 |
| 249 | Quantitative assessment of liver fat with magnetic resonance imaging and spectroscopy., 2011, 34, 729. | | 1 |
| 250 | Fat and water magnetic resonance imaging. Journal of Magnetic Resonance Imaging, 2010, 31, 4-18. | 3.4 | 291 |
| 251 | Flowâ€independent T ₂ â€prepared inversion recovery blackâ€blood MR imaging. Journal of Magnetic Resonance Imaging, 2010, 31, 248-254. | 3.4 | 20 |
| 252 | Cardiac MRI evaluation of nonischemic cardiomyopathies. Journal of Magnetic Resonance Imaging, 2010, 31, 518-530. | 3.4 | 14 |

| # | Article | IF | Citations |
|-----|---|-----|-----------|
| 253 | Repeatability of magnetic resonance elastography for quantification of hepatic stiffness. Journal of Magnetic Resonance Imaging, 2010, 31, 725-731. | 3.4 | 145 |
| 254 | Phase and amplitude correction for multiâ€echo water–fat separation with bipolar acquisitions. Journal of Magnetic Resonance Imaging, 2010, 31, 1264-1271. | 3.4 | 63 |
| 255 | <i>k</i> å€space waterâ€fat decomposition with T ₂ * estimation and multifrequency fat spectrum modeling for ultrashort echo time imaging. Journal of Magnetic Resonance Imaging, 2010, 31, 1027-1034. | 3.4 | 24 |
| 256 | Cartilage morphology at 3.0T: Assessment of threeâ€dimensional magnetic resonance imaging techniques. Journal of Magnetic Resonance Imaging, 2010, 32, 173-183. | 3.4 | 35 |
| 257 | Noise analysis for 3â€point chemical shiftâ€based waterâ€fat separation with spectral modeling of fat. Journal of Magnetic Resonance Imaging, 2010, 32, 493-500. | 3.4 | 16 |
| 258 | Single breathhold cardiac CINE imaging with multiâ€echo threeâ€dimensional hybrid radial SSFP acquisition. Journal of Magnetic Resonance Imaging, 2010, 32, 434-440. | 3.4 | 19 |
| 259 | T ₂ â€weighted 3D fast spin echo imaging with water–fat separation in a single acquisition. Journal of Magnetic Resonance Imaging, 2010, 32, 745-751. | 3.4 | 28 |
| 260 | Frequency response of multipoint chemical shiftâ€based spectral decomposition. Journal of Magnetic Resonance Imaging, 2010, 32, 943-952. | 3.4 | 3 |
| 261 | Independent estimation of <i>T</i> * ₂ for water and fat for improved accuracy of fat quantification. Magnetic Resonance in Medicine, 2010, 63, 849-857. | 3.0 | 78 |
| 262 | Noninvasive temperature mapping with MRI using chemical shift waterâ€fat separation. Magnetic Resonance in Medicine, 2010, 63, 1238-1246. | 3.0 | 63 |
| 263 | The influence of prior hamstring injury on lengthening muscle tissue mechanics. Journal of Biomechanics, 2010, 43, 2254-2260. | 2.1 | 79 |
| 264 | Whole chest MRA and velocimetry for congenital heart disease in less than 10 minutes with 3D radial phase contrast. Journal of Cardiovascular Magnetic Resonance, 2010, 12, . | 3.3 | 0 |
| 265 | 3.0-T Evaluation of Knee Cartilage by Using Three-Dimensional IDEAL GRASS Imaging: Comparison with Fast Spin-Echo Imaging. Radiology, 2010, 255, 117-127. | 7.3 | 55 |
| 266 | Quantification of Hepatic Steatosis with 3-T MR Imaging: Validation in <i>ob/ob</i> Mice. Radiology, 2010, 254, 119-128. | 7.3 | 71 |
| 267 | Presurgical Localization of the Artery of Adamkiewicz with Time-resolved 3.0-T MR Angiography. Radiology, 2010, 255, 873-881. | 7.3 | 62 |
| 268 | Magnetic Resonance Imaging Quantification of Liver Iron. Magnetic Resonance Imaging Clinics of North America, 2010, 18, 359-381. | 1.1 | 170 |
| 269 | Quantification of Liver Fat with Magnetic Resonance Imaging. Magnetic Resonance Imaging Clinics of North America, 2010, 18, 337-357. | 1.1 | 260 |
| 270 | Rapid comprehensive evaluation of luminography and hemodynamic function with 3d radially undersampled phase contrast imaging MRI., 2009, 2009, 4057-60. | | 3 |

| # | Article | IF | Citations |
|-----|---|-----|-----------|
| 271 | Multiecho IDEAL Gradient-Echo Water-Fat Separation for Rapid Assessment of Cartilage Volume at 1.5 T: Initial Experience. Radiology, 2009, 252, 561-567. | 7.3 | 31 |
| 272 | Endovascular Abdominal Aortic Aneurysm Repair: Nonenhanced Volumetric CT for Follow-up. Radiology, 2009, 253, 253-262. | 7.3 | 63 |
| 273 | Improved fat suppression using multipeak reconstruction for IDEAL chemical shift fatâ€water separation: Application with fast spin echo imaging. Journal of Magnetic Resonance Imaging, 2009, 29, 436-442. | 3.4 | 28 |
| 274 | Quantification of hepatic steatosis with MRI: The effects of accurate fat spectral modeling. Journal of Magnetic Resonance Imaging, 2009, 29, 1332-1339. | 3.4 | 221 |
| 275 | Improved timeâ€ofâ€flight magnetic resonance angiography with IDEAL waterâ€fat separation. Journal of Magnetic Resonance Imaging, 2009, 29, 1367-1374. | 3.4 | 17 |
| 276 | T ₁ independent, T ₂ * corrected MRI with accurate spectral modeling for quantification of fat: Validation in a fatâ€waterâ€\$PIO phantom. Journal of Magnetic Resonance Imaging, 2009, 30, 1215-1222. | 3.4 | 191 |
| 277 | Increased volume of coverage for abdominal contrastâ€enhanced MR angiography with twoâ€dimensional autocalibrating parallel imaging: Initial experience at 3.0 Tesla. Journal of Magnetic Resonance Imaging, 2009, 30, 1093-1100. | 3.4 | 30 |
| 278 | Advanced MRI Methods for Assessment of Chronic Liver Disease. American Journal of Roentgenology, 2009, 193, 14-27. | 2.2 | 169 |
| 279 | A New Intercostal Artery Management Strategy for Thoracoabdominal Aortic Aneurysm Repair. Journal of Surgical Research, 2009, 154, 99-104. | 1.6 | 14 |
| 280 | Preface. Magnetic Resonance Imaging Clinics of North America, 2009, 17, xi-xii. | 1.1 | 5 |
| 281 | High temporal resolution cardiac cone-beam CT using a slowly rotating C-arm gantry. Proceedings of SPIE, 2009, , . | 0.8 | 7 |
| 282 | Cartilage imaging at 3.0T with gradient refocused acquisition in the steadyâ€state (GRASS) and IDEAL fatâ€water separation. Journal of Magnetic Resonance Imaging, 2008, 28, 167-174. | 3.4 | 23 |
| 283 | Simultaneous estimation of tongue volume and fat fraction using IDEALâ€FSE. Journal of Magnetic Resonance Imaging, 2008, 28, 504-508. | 3.4 | 27 |
| 284 | Improved delayed enhanced myocardial imaging with T ₂ â€Prep inversion recovery magnetization preparation. Journal of Magnetic Resonance Imaging, 2008, 28, 1280-1286. | 3.4 | 38 |
| 285 | Generalized <i>k</i> å€space decomposition with chemical shift correction for nonâ€cartesian waterâ€fat imaging. Magnetic Resonance in Medicine, 2008, 59, 1151-1164. | 3.0 | 81 |
| 286 | Water–fat separation with bipolar multiecho sequences. Magnetic Resonance in Medicine, 2008, 60, 198-209. | 3.0 | 73 |
| 287 | Effects of refocusing flip angle modulation and view ordering in 3D fast spin echo. Magnetic Resonance in Medicine, 2008, 60, 640-649. | 3.0 | 239 |
| 288 | Multiecho waterâ€fat separation and simultaneous <i>R</i> estimation with multifrequency fat spectrum modeling. Magnetic Resonance in Medicine, 2008, 60, 1122-1134. | 3.0 | 590 |

| # | Article | IF | CITATIONS |
|-----|--|-----|-----------|
| 289 | Influence of multichannel combination, parallel imaging and other reconstruction techniques on MRI noise characteristics. Magnetic Resonance Imaging, 2008, 26, 754-762. | 1.8 | 199 |
| 290 | Temporally Targeted Imaging Method Applied to ECG-Gated Computed Tomography. Academic Radiology, 2008, 15, 93-106. | 2.5 | 6 |
| 291 | Cardiac MRI of ischemic heart disease at 3T: Potential and challenges. European Journal of Radiology, 2008, 65, 15-28. | 2.6 | 83 |
| 292 | Body MRI Using IDEAL. American Journal of Roentgenology, 2008, 190, 1076-1084. | 2.2 | 70 |
| 293 | Noise considerations of three-point water-fat separation imaging methods. Medical Physics, 2008, 35, 3597-3606. | 3.0 | 6 |
| 294 | High-Resolution 3D Cartilage Imaging with IDEAL–SPGR at 3 T. American Journal of Roentgenology, 2007, 189, 1510-1515. | 2.2 | 41 |
| 295 | IDEAL Imaging of the Musculoskeletal System: Robust Water–Fat Separation for Uniform Fat Suppression, Marrow Evaluation, and Cartilage Imaging. American Journal of Roentgenology, 2007, 189, W284-W291. | 2.2 | 67 |
| 296 | Fat quantification with IDEAL gradient echo imaging: Correction of bias from <i>T</i> ₁ and noise. Magnetic Resonance in Medicine, 2007, 58, 354-364. | 3.0 | 418 |
| 297 | Balanced SSFP imaging of the musculoskeletal system. Journal of Magnetic Resonance Imaging, 2007, 25, 270-278. | 3.4 | 27 |
| 298 | Water–fat separation with IDEAL gradient-echo imaging. Journal of Magnetic Resonance Imaging, 2007, 25, 644-652. | 3.4 | 300 |
| 299 | Measurement of signalâ€toâ€noise ratios in MR images: Influence of multichannel coils, parallel imaging, and reconstruction filters. Journal of Magnetic Resonance Imaging, 2007, 26, 375-385. | 3.4 | 809 |
| 300 | Leastâ€squares chemical shift separation for ¹³ C metabolic imaging. Journal of Magnetic Resonance Imaging, 2007, 26, 1145-1152. | 3.4 | 91 |
| 301 | Multiecho reconstruction for simultaneous waterâ€fat decomposition and T2* estimation. Journal of Magnetic Resonance Imaging, 2007, 26, 1153-1161. | 3.4 | 366 |
| 302 | Effects of injection rate and dose on image quality in time-resolved magnetic resonance angiography (MRA) by using 1.0M contrast agents. European Radiology, 2007, 17, 1394-1402. | 4.5 | 24 |
| 303 | Measurement of Signal-to-Noise Ratio and Parallel Imaging. , 2007, , 49-61. | | 17 |
| 304 | ECG-gated HYPR reconstruction for undersampled CT myocardial perfusion imaging., 2007,,. | | 6 |
| 305 | Cardiac Steady-State Free Precession CINE Magnetic Resonance Imaging at 3.0 Tesla. Investigative Radiology, 2006, 41, 141-147. | 6.2 | 42 |
| 306 | Phase-Sensitive Inversion Recovery (PSIR) Single-Shot TrueFISP for Assessment of Myocardial Infarction at 3 Tesla. Investigative Radiology, 2006, 41, 148-153. | 6.2 | 59 |

| # | Article | IF | CITATIONS |
|-----|--|-----|-----------|
| 307 | Single acquisition water-fat separation: Feasibility study for dynamic imaging. Magnetic Resonance in Medicine, 2006, 55, 413-422. | 3.0 | 39 |
| 308 | Cardiac CINE MR imaging with a 32-channel cardiac coil and parallel imaging: Impact of acceleration factors on image quality and volumetric accuracy. Journal of Magnetic Resonance Imaging, 2006, 23, 222-227. | 3.4 | 71 |
| 309 | T1- and T2-weighted fast spin-echo imaging of the brachial plexus and cervical spine with IDEAL water–fat separation. Journal of Magnetic Resonance Imaging, 2006, 24, 825-832. | 3.4 | 50 |
| 310 | Articular Cartilage of the Knee: Rapid Three-dimensional MR Imaging at 3.0 T with IDEAL Balanced Steady-State Free Precession—Initial Experience. Radiology, 2006, 240, 546-551. | 7.3 | 70 |
| 311 | Iterative Decomposition of Water and Fat with Echo Asymmetry and Least-Squares Estimation (IDEAL) Fast Spin-Echo Imaging of the Ankle: Initial Clinical Experience. American Journal of Roentgenology, 2006, 187, 1442-1447. | 2.2 | 42 |
| 312 | MR imaging of articular cartilage at 1.5T and 3.0T: Comparison of SPGR and SSFP sequences. Osteoarthritis and Cartilage, 2005, 13, 338-344. | 1.3 | 124 |
| 313 | Cardiac CINE imaging with IDEAL water-fat separation and steady-state free precession. Journal of Magnetic Resonance Imaging, 2005, 22, 44-52. | 3.4 | 61 |
| 314 | Homodyne reconstruction and IDEAL water-fat decomposition. Magnetic Resonance in Medicine, 2005, 54, 586-593. | 3.0 | 71 |
| 315 | Cram \tilde{A} @r-Rao bounds for three-point decomposition of water and fat. Magnetic Resonance in Medicine, 2005, 54, 625-635. | 3.0 | 194 |
| 316 | Iterative decomposition of water and fat with echo asymmetry and least-squares estimation (IDEAL): Application with fast spin-echo imaging. Magnetic Resonance in Medicine, 2005, 54, 636-644. | 3.0 | 615 |
| 317 | Practical approaches to the evaluation of signal-to-noise ratio performance with parallel imaging: Application with cardiac imaging and a 32-channel cardiac coil. Magnetic Resonance in Medicine, 2005, 54, 748-754. | 3.0 | 274 |
| 318 | Field map estimation with a region growing scheme for iterative 3-point water-fat decomposition. Magnetic Resonance in Medicine, 2005, 54, 1032-1039. | 3.0 | 195 |
| 319 | Controversies in Protocol Selection in the Imaging of Articular Cartilage. Seminars in Musculoskeletal Radiology, 2005, 9, 161-172. | 0.7 | 24 |
| 320 | Steady-State Free Precession MR Imaging: Improved Myocardial Tag Persistence and Signal-to-Noise Ratio for Analysis of Myocardial Motion. Radiology, 2004, 230, 852-861. | 7.3 | 26 |
| 321 | Multicoil Dixon chemical species separation with an iterative least-squares estimation method. Magnetic Resonance in Medicine, 2004, 51, 35-45. | 3.0 | 449 |
| 322 | Signal-to-noise ratio behavior of steady-state free precession. Magnetic Resonance in Medicine, 2004, 52, 123-130. | 3.0 | 48 |
| 323 | Advanced MR imaging of the shoulder: dedicated cartilage techniques. Magnetic Resonance Imaging Clinics of North America, 2004, 12, 143-159. | 1.1 | 16 |
| 324 | Rapid MR Imaging of Articular Cartilage with Steady-State Free Precession and Multipoint Fat-Water Separation. American Journal of Roentgenology, 2003, 180, 357-362. | 2.2 | 74 |

| # | Article | IF | Citations |
|-----|---|-------------|-----------|
| 325 | Sonography in Primary Hyperparathyroidism. Journal of Ultrasound in Medicine, 2002, 21, 539-552. | 1.7 | 90 |
| 326 | Advanced Cardiac MR Imaging of Ischemic Heart Disease. Radiographics, 2001, 21, 1047-1074. | 3.3 | 49 |
| 327 | Ultrafast Pulse Sequence Techniques for Cardiac Magnetic Resonance Imaging. Topics in Magnetic Resonance Imaging, 2000, 11, 312-330. | 1.2 | 16 |
| 328 | Simultaneous Noninvasive Determination of Regional Myocardial Perfusion and Oxygen Content in Rabbits: Toward Direct Measurement of Myocardial Oxygen Consumption at MR Imaging. Radiology, 1999, 212, 739-747. | 7.3 | 23 |
| 329 | A novel object-independent ?balanced? reference scan for echo-planar imaging. Journal of Magnetic Resonance Imaging, 1999, 9, 847-852. | 3.4 | 28 |
| 330 | Referenceless interleaved echo-planar imaging. Magnetic Resonance in Medicine, 1999, 41, 87-94. | 3.0 | 55 |
| 331 | Effects of water exchange on the measurement of myocardial perfusion using paramagnetic contrast agents. Magnetic Resonance in Medicine, 1999, 41, 334-342. | 3.0 | 68 |
| 332 | Multi-echo segmented k-space imaging: An optimized hybrid sequence for ultrafast cardiac imaging. Magnetic Resonance in Medicine, 1999, 41, 375-385. | 3.0 | 74 |
| 333 | In vivo measurement of T*2 and field inhomogeneity maps in the human heart at 1.5 T. Magnetic Resonance in Medicine, 1998, 39, 988-998. | 3.0 | 183 |
| 334 | Single-shot, variable flip-angle slice-selective excitation with four gradient-modulated adiabatic half-passage segments. Magnetic Resonance in Medicine, 1998, 40, 334-340. | 3.0 | 6 |
| 335 | Magnitude and Time Course of Microvascular Obstruction and Tissue Injury After Acute Myocardial Infarction. Circulation, 1998, 98, 1006-1014. | 1.6 | 453 |
| 336 | Techniques for high-speed cardiac magnetic resonance imaging in rats and rabbits. Magnetic Resonance in Medicine, 1997, 37, 124-130. | 3.0 | 20 |
| 337 | Quantification and reduction of ghosting artifacts in interleaved echo-planar imaging. Magnetic Resonance in Medicine, 1997, 38, 429-439. | 3.0 | 56 |
| 338 | Fast ²³ Na Magnetic Resonance Imaging of Acute Reperfused Myocardial Infarction. Circulation, 1997, 95, 1877-1885. | 1.6 | 109 |
| 339 | Quantitative cardiac perfusion: a noninvasive spin-labeling method that exploits coronary vessel geometry Radiology, 1996, 200, 177-184. | 7. 3 | 39 |
| 340 | A magnetization-driven gradient echo pulse sequence for the study of myocardial perfusion. Magnetic Resonance in Medicine, 1995, 34, 276-282. | 3.0 | 51 |
| 341 | Blood oxygenation dependence oft1 andt2 in the isolated, perfused rabbit heart at 4.7t. Magnetic Resonance in Medicine, 1995, 34, 623-627. | 3.0 | 58 |
| 342 | Tag contrast in breath-hold CINE cardiac MRI. Magnetic Resonance in Medicine, 1994, 31, 521-525. | 3.0 | 37 |

SCOTT B REEDER

| # | Article | IF | CITATIONS |
|-----|--|-----|-----------|
| 343 | The effect of high performance gradients on fast gradient echo imaging. Magnetic Resonance in Medicine, 1994, 32, 612-621. | 3.0 | 52 |
| 344 | Tagged MR imaging in a deforming phantom: photographic validation Radiology, 1994, 190, 765-769. | 7.3 | 459 |
| 345 | Cardiac MR imaging., 0,, 34-46. | | 0 |
| 346 | CE-MRA in the primary diagnosis of pulmonary embolism: Building a team to start a clinically relevant program., 0,, 31-36. | | 1 |