

Mei-Zhen Cui

List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/9453251/publications.pdf>

Version: 2024-02-01

52
papers

1,877
citations

236925

25
h-index

254184

43
g-index

53
all docs

53
docs citations

53
times ranked

2544
citing authors

#	ARTICLE	IF	CITATIONS
1	LPA1-mediated PKD2 activation promotes LPA-induced tissue factor expression via the p38 β and JNK2 MAPK pathways in smooth muscle cells. <i>Journal of Biological Chemistry</i> , 2021, 297, 101152.	3.4	3
2	Proapoptotic Mitochondrial Carrier Homolog Protein PSAP Mediates Death Receptor 6 Induced Apoptosis. <i>Journal of Alzheimer's Disease</i> , 2020, 74, 1097-1106.	2.6	8
3	Platelet CD40 Mediates Leukocyte Recruitment and Neointima Formation after Arterial Denudation Injury in Atherosclerosis-Prone Mice. <i>American Journal of Pathology</i> , 2018, 188, 252-263.	3.8	10
4	CRE and SRE mediate LPA-induced CCN1 transcription in mouse aortic smooth muscle cells. <i>Canadian Journal of Physiology and Pharmacology</i> , 2017, 95, 275-280.	1.4	4
5	Pen-2 and Presenilin are Sufficient to Catalyze Notch Processing. <i>Journal of Alzheimer's Disease</i> , 2017, 56, 1263-1269.	2.6	9
6	CD14 is a key mediator of both lysophosphatidic acid and lipopolysaccharide induction of foam cell formation. <i>Journal of Biological Chemistry</i> , 2017, 292, 14391-14400.	3.4	22
7	Lysophosphatidic Acid Triggers Apoptosis in HeLa Cells through the Upregulation of Tumor Necrosis Factor Receptor Superfamily Member 21. <i>Mediators of Inflammation</i> , 2017, 2017, 1-12.	3.0	6
8	JNK1 Mediates Lipopolysaccharide-Induced CD14 and SR-AI Expression and Macrophage Foam Cell Formation. <i>Frontiers in Physiology</i> , 2017, 8, 1075.	2.8	11
9	Nicastrin is required for amyloid precursor protein (<sc>APP</sc>) but not Notch processing, while anterior pharynx-defective 1 is dispensable for processing of both <sc>APP</sc> and Notch. <i>Journal of Neurochemistry</i> , 2016, 136, 1246-1258.	3.9	10
10	Nanospherical arabinogalactan proteins are a key component of the high-strength adhesive secreted by English ivy. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 2016, 113, E3193-202.	7.1	62
11	Lysophosphatidic acid-induced vascular neointimal formation in mouse carotid arteries is mediated by the matricellular protein CCN1/Cyr61. <i>American Journal of Physiology - Cell Physiology</i> , 2016, 311, C975-C984.	4.6	16
12	<i>Potential therapeutics for myocardial ischemia-reperfusion injury</i>. Focus on "Induction of cardioprotection by small netrin-1-derived peptides": <i>American Journal of Physiology - Cell Physiology</i> , 2015, 309, C97-C99.	4.6	4
13	Cellular FLICE-like Inhibitory Protein (c-FLIP) and PS1-associated Protein (PSAP) Mediate Presenilin 1-induced γ -Secretase-dependent and -independent Apoptosis, Respectively. <i>Journal of Biological Chemistry</i> , 2015, 290, 18269-18280.	3.4	14
14	The Matricellular Protein Cyr61 Is a Key Mediator of Platelet-derived Growth Factor-induced Cell Migration. <i>Journal of Biological Chemistry</i> , 2015, 290, 8232-8242.	3.4	22
15	Matricellular Protein Cyr61 Bridges Lysophosphatidic Acid and Integrin Pathways Leading to Cell Migration. <i>Journal of Biological Chemistry</i> , 2014, 289, 5774-5783.	3.4	20
16	Inhibition of Egr1 expression underlies the anti-mitogenic effects of cAMP in vascular smooth muscle cells. <i>Journal of Molecular and Cellular Cardiology</i> , 2014, 72, 9-19.	1.9	26
17	PSAP induces a unique Apaf-1 and Smac-dependent mitochondrial apoptotic pathway independent of Bcl-2 family proteins. <i>Biochimica Et Biophysica Acta - Molecular Basis of Disease</i> , 2013, 1832, 453-474.	3.8	19
18	Lysophosphatidic acid induces increased BACE1 expression and A β 2 formation. <i>Biochimica Et Biophysica Acta - Molecular Basis of Disease</i> , 2013, 1832, 29-38.	3.8	30

#	ARTICLE	IF	CITATIONS
19	Histamine induces activation of protein kinase D that mediates tissue factor expression and activity in human aortic smooth muscle cells. <i>American Journal of Physiology - Heart and Circulatory Physiology</i> , 2012, 303, H1344-H1352.	3.2	13
20	Death Receptor 6 Induces Apoptosis Not through Type I or Type II Pathways, but via a Unique Mitochondria-dependent Pathway by Interacting with Bax Protein. <i>Journal of Biological Chemistry</i> , 2012, 287, 29125-29133.	3.4	79
21	Lysophosphatidic Acid Induces Early Growth Response-1 (Egr-1) Protein Expression via Protein Kinase C β -regulated Extracellular Signal-regulated Kinase (ERK) and c-Jun N-terminal Kinase (JNK) Activation in Vascular Smooth Muscle Cells. <i>Journal of Biological Chemistry</i> , 2012, 287, 22635-22642.	3.4	35
22	Pen β 2 is dispensable for endoproteolysis of presenilin 1, and nicastrin β subcomplex is important for both β -secretase assembly and substrate recruitment. <i>Journal of Neurochemistry</i> , 2012, 123, 837-844.	3.9	13
23	Lysophosphatidic acid signaling in vascular smooth muscle cells. <i>Journal of Clinical & Experimental Cardiology</i> , 2012, 01, .	0.0	0
24	Lysophosphatidic acid effects on atherosclerosis and thrombosis. <i>Clinical Lipidology</i> , 2011, 6, 413-426.	0.4	49
25	LPA induces IL-6 secretion from aortic smooth muscle cells via an LPA β -regulated, PKC-dependent, and p38 β -mediated pathway. <i>American Journal of Physiology - Heart and Circulatory Physiology</i> , 2010, 298, H974-H983.	3.2	30
26	Distinct Cell Responses to Substrates Consisting of Poly(μ -caprolactone) and Poly(propylene fumarate) in the Presence or Absence of Cross-Links. <i>Biomacromolecules</i> , 2010, 11, 2748-2759.	5.4	45
27	The GxxxG Motif in the Transmembrane Domain of A β PP Plays an Essential Role in the Interaction of CTF β with the β -secretase Complex and the Formation of Amyloid- β . <i>Journal of Alzheimer's Disease</i> , 2009, 18, 167-176.	2.6	6
28	Lysophosphatidylcholine Activates a Novel PKD2-Mediated Signaling Pathway That Controls Monocyte Migration. <i>Arteriosclerosis, Thrombosis, and Vascular Biology</i> , 2009, 29, 1376-1382.	2.4	45
29	Residues at P2-P1 positions of β - and γ -cleavage sites are important in formation of β -amyloid peptide. <i>Neurobiology of Disease</i> , 2009, 36, 453-460.	4.4	2
30	Effects of β -secretase cleavage β region mutations on APP processing and A β formation: interpretation with sequential cleavage and β -helical model. <i>Journal of Neurochemistry</i> , 2008, 107, 722-733.	3.9	18
31	Both the N-terminal fragment and the protein β protein interaction domain (PDZ domain) are required for the pro-apoptotic activity of presenilin-associated protein PSAP. <i>Biochimica Et Biophysica Acta - General Subjects</i> , 2008, 1780, 696-708.	2.4	9
32	Histamine Induces Egr-1 Expression in Human Aortic Endothelial Cells via the H1 Receptor-mediated Protein Kinase C β -dependent ERK Activation Pathway. <i>Journal of Biological Chemistry</i> , 2008, 283, 26928-26936.	3.4	34
33	Lysophosphatidic acid induces prostate cancer PC3 cell migration via activation of LPA1, p42 and p38 β . <i>Biochimica Et Biophysica Acta - Molecular and Cell Biology of Lipids</i> , 2007, 1771, 883-892.	2.4	76
34	The same β -secretase accounts for the multiple intramembrane cleavages of APP. <i>Journal of Neurochemistry</i> , 2007, 100, 1234-1246.	3.9	42
35	Lysophosphatidic Acid Induces Early Growth Response Gene 1 Expression in Vascular Smooth Muscle Cells. <i>Arteriosclerosis, Thrombosis, and Vascular Biology</i> , 2006, 26, 1029-1035.	2.4	35
36	Calpain inhibitor MDL28170 modulates A β formation by inhibiting the formation of intermediate A β 46 and protecting A β from degradation. <i>FASEB Journal</i> , 2006, 20, 331-333.	0.5	24

#	ARTICLE	IF	CITATIONS
37	Activation of Protein Kinase C δ by Peroxynitrite Regulates LKB1-dependent AMP-activated Protein Kinase in Cultured Endothelial Cells. <i>Journal of Biological Chemistry</i> , 2006, 281, 6366-6375.	3.4	161
38	β -Cleavage Is Dependent on η -Cleavage during the Proteolytic Processing of Amyloid Precursor Protein within Its Transmembrane Domain. <i>Journal of Biological Chemistry</i> , 2005, 280, 37689-37697.	3.4	120
39	Angiotensin II-Induced Protein Kinase D Activation Is Regulated by Protein Kinase C δ and Mediated via the Angiotensin II Type 1 Receptor in Vascular Smooth Muscle Cells. <i>Arteriosclerosis, Thrombosis, and Vascular Biology</i> , 2004, 24, 2271-2276.	2.4	26
40	Identification of a New Presenilin-dependent η -Cleavage Site within the Transmembrane Domain of Amyloid Precursor Protein. <i>Journal of Biological Chemistry</i> , 2004, 279, 50647-50650.	3.4	126
41	Oxidized lipoprotein regulation of tissue factor in smooth muscle cells. <i>International Congress Series</i> , 2004, 1262, 83-86.	0.2	0
42	Thrombin Rapidly Induces Protein Kinase D Phosphorylation, and Protein Kinase C δ Mediates the Activation. <i>Journal of Biological Chemistry</i> , 2003, 278, 2824-2828.	3.4	51
43	Lysophosphatidic Acid Induction of Tissue Factor Expression in Aortic Smooth Muscle Cells. <i>Arteriosclerosis, Thrombosis, and Vascular Biology</i> , 2003, 23, 224-230.	2.4	45
44	The Novel Presenilin-1-associated Protein Is a Proapoptotic Mitochondrial Protein. <i>Journal of Biological Chemistry</i> , 2002, 277, 48913-48922.	3.4	54
45	Smooth muscle cell surface tissue factor pathway activation by oxidized low-density lipoprotein requires cellular lipid peroxidation. <i>Blood</i> , 2000, 96, 3056-3063.	1.4	50
46	Smooth muscle cell surface tissue factor pathway activation by oxidized low-density lipoprotein requires cellular lipid peroxidation. <i>Blood</i> , 2000, 96, 3056-3063.	1.4	0
47	Identification of a Novel PSD-95/Dlg/ZO-1 (PDZ)-like Protein Interacting with the C Terminus of Presenilin-1. <i>Journal of Biological Chemistry</i> , 1999, 274, 32543-32546.	3.4	44
48	LDL Increases Inactive Tissue Factor on Vascular Smooth Muscle Cell Surfaces. <i>Circulation</i> , 1999, 99, 1753-1759.	1.6	64
49	Native and Oxidized Low Density Lipoprotein Induction of Tissue Factor Gene Expression in Smooth Muscle Cells Is Mediated by Both Egr-1 and Sp1. <i>Journal of Biological Chemistry</i> , 1999, 274, 32795-32802.	3.4	76
50	Transcriptional Regulation of the Tissue Factor Gene in Human Epithelial Cells Is Mediated by Sp1 and EGR-1. <i>Journal of Biological Chemistry</i> , 1996, 271, 2731-2739.	3.4	143
51	The LIF response element of the β 2-microglobulin gene confers LIF-induced transcriptional activation in embryonal stem cells. <i>Cytokine</i> , 1995, 7, 491-502.	3.2	25
52	Nucleotide Sequence of the Gene Encoding NADH Dehydrogenase from an Alkalophile, <i>Bacillus</i> sp. Strain YN-1. <i>Journal of Biochemistry</i> , 1991, 109, 678-683.	1.7	41