Hakan TerzÄ^o

List of Publications by Year in descending order

Source: https://exaly.com/author-pdf/9487506/publications.pdf Version: 2024-02-01



Ηλκάνι Τερζάο

| # | Article | IF | CITATIONS |
|----|--|-----|-----------|
| 1 | Proteomic analysis reveals the role of exogenous cysteine in alleviating chromium stress in maize seedlings. Ecotoxicology and Environmental Safety, 2021, 209, 111784. | 6.0 | 22 |
| 2 | Comparative analysis of salt-induced changes in the root physiology and proteome of the xero-halophyte Salsola crassa. Revista Brasileira De Botanica, 2021, 44, 33-42. | 1.3 | 4 |
| 3 | Alterations in the root proteomes of Brassica napus cultivars under salt stress. Botanica Serbica, 2021, 45, 87-96. | 1.0 | 1 |
| 4 | Proteomic responses of maize roots to the combined stress of sulphur deficiency and chromium toxicity. Biologia (Poland), 2021, 76, 1887-1899. | 1.5 | 0 |
| 5 | Exogenous cysteine alleviates chromium stress via reducing its uptake and regulating proteome in roots of Brassica napus L. seedlings. South African Journal of Botany, 2021, 139, 114-121. | 2.5 | 15 |
| 6 | Growth responses and essential oil profile of Salvia officinalis L. Influenced by water deficit and various nutrient sources in the greenhouse. Saudi Journal of Biological Sciences, 2021, 28, 7327-7335. | 3.8 | 15 |
| 7 | Proteomic analysis of the anticancer effect of various extracts of endemicThermopsisturcica in human cervical cancer cells. Turkish Journal of Medical Sciences, 2020, 50, 1993-2004. | 0.9 | 1 |
| 8 | Proteomic analysis of chromium stress and sulfur deficiency responses in leaves of two canola (Brassica napus L.) cultivars differing in Cr(VI) tolerance. Ecotoxicology and Environmental Safety, 2016, 124, 255-266. | 6.0 | 28 |
| 9 | Proteomic and biochemical responses of canola (Brassica napus L.) exposed to salinity stress and exogenous lipoic acid. Journal of Plant Physiology, 2015, 179, 90-99. | 3.5 | 21 |
| 10 | Interactive effects of sulfur and chromium on antioxidative defense systems and BnMP1 gene expression in canola (Brassica napus L.) cultivars differing in Cr(VI) tolerance. Ecotoxicology, 2015, 24, 1171-1182. | 2.4 | 25 |
| 11 | Variations in Chromium Tolerance and Accumulation among Canola (Brassica napus L.) Cultivars. Bulletin of Environmental Contamination and Toxicology, 2014, 93, 113-119. | 2.7 | 8 |
| 12 | Protective role of hydrogen peroxide pretreatment on defense systems and BnMP1 gene expression in Cr(VI)-stressed canola seedlings. Ecotoxicology, 2013, 22, 1303-1312. | 2.4 | 30 |
| 13 | Lead contamination reduces chlorophyll biosynthesis and genomic template stability in Brassica rapa L Environmental and Experimental Botany, 2010, 67, 467-473. | 4.2 | 195 |
| 14 | Evaluation of 2,4-D and Dicamba genotoxicity in bean seedlings using comet and RAPD assays. Ecotoxicology and Environmental Safety, 2010, 73, 1558-1564. | 6.0 | 53 |
| 15 | Determination of genotoxic effects of copper sulphate and cobalt chloride in Allium cepa root cells by chromosome aberration and comet assays. Chemosphere, 2009, 75, 934-938. | 8.2 | 154 |
| 16 | Small heat shock protein responses in leaf tissues of wheat cultivars with different heat susceptibility. Biologia (Poland), 2008, 63, 521-525. | 1.5 | 12 |
| 17 | Proteomic analysis reveals different responses to drought between the Cleome spinosa (C3) and Cleome gynandra (C4). Turkish Journal of Botany, 0, , . | 1.2 | 2 |