

Hiroyasu Nakano

List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/9499915/publications.pdf>

Version: 2024-02-01

117
papers

19,908
citations

23567

58
h-index

20358

116
g-index

122
all docs

122
docs citations

122
times ranked

32517
citing authors

#	ARTICLE	IF	CITATIONS
1	The scaffold-dependent function of RIPK1 in dendritic cells promotes injury-induced colitis. <i>Mucosal Immunology</i> , 2022, 15, 84-95.	6.0	7
2	Regulation of the release of damage-associated molecular patterns from necroptotic cells. <i>Biochemical Journal</i> , 2022, 479, 677-685.	3.7	17
3	Proscillaridin A Sensitizes Human Colon Cancer Cells to TRAIL-Induced Cell Death. <i>International Journal of Molecular Sciences</i> , 2022, 23, 6973.	4.1	0
4	Interleukin-11-expressing fibroblasts have a unique gene signature correlated with poor prognosis of colorectal cancer. <i>Nature Communications</i> , 2021, 12, 2281.	12.8	60
5	Time-Lapse Imaging of Necroptosis and DAMP Release at Single-Cell Resolution. <i>Methods in Molecular Biology</i> , 2021, 2274, 353-363.	0.9	2
6	MIND bomb 2 prevents RIPK1 kinase activity-dependent and -independent apoptosis through ubiquitylation of cFLIPL. <i>Communications Biology</i> , 2021, 4, 80.	4.4	13
7	Regulation of T cell differentiation by the AP-1 transcription factor JunB. <i>Immunological Medicine</i> , 2021, 44, 197-203.	2.6	25
8	A missense mutation in the MLKL brace region promotes lethal neonatal inflammation and hematopoietic dysfunction. <i>Nature Communications</i> , 2020, 11, 3150.	12.8	75
9	Regenerating islet-derived protein (Reg)3 β plays a crucial role in attenuation of ileitis and colitis in mice. <i>Biochemistry and Biophysics Reports</i> , 2020, 21, 100738.	1.3	11
10	Identification of a phosphorylation site on Ulk1 required for genotoxic stress-induced alternative autophagy. <i>Nature Communications</i> , 2020, 11, 1754.	12.8	46
11	Inhibition of Importin β 1 Augments the Anticancer Effect of Agonistic Anti-Death Receptor 5 Antibody in TRAIL-resistant Tumor Cells. <i>Molecular Cancer Therapeutics</i> , 2020, 19, 1123-1133.	4.1	6
12	Identification of the hallmarks of necroptosis and ferroptosis by transmission electron microscopy. <i>Biochemical and Biophysical Research Communications</i> , 2020, 527, 839-844.	2.1	39
13	JunB plays a crucial role in development of regulatory T cells by promoting IL-2 signaling. <i>Mucosal Immunology</i> , 2019, 12, 1104-1117.	6.0	34
14	Hepatic ferroptosis plays an important role as the trigger for initiating inflammation in nonalcoholic steatohepatitis. <i>Cell Death and Disease</i> , 2019, 10, 449.	6.3	267
15	Necroptosis of Intestinal Epithelial Cells Induces Type 3 Innate Lymphoid Cell-Dependent Lethal Ileitis. <i>IScience</i> , 2019, 15, 536-551.	4.1	21
16	Regulation of membrane phospholipid asymmetry by Notch-mediated flippase expression controls the number of intraepithelial TCR β ^{hi} CD8 α ^{hi} T cells. <i>PLoS Biology</i> , 2019, 17, e3000262.	5.6	3
17	Addendum: A FRET biosensor for necroptosis uncovers two different modes of the release of DAMPs. <i>Nature Communications</i> , 2019, 10, 1923.	12.8	2
18	A murine model of acute lung injury identifies growth factors to promote tissue repair and their biomarkers. <i>Genes To Cells</i> , 2019, 24, 112-125.	1.2	5

#	ARTICLE	IF	CITATIONS
19	Blockade of TNF receptor superfamily 1 (TNFR1)â€“dependent and TNFR1-independent cell death is crucial for normal epidermal differentiation. <i>Journal of Allergy and Clinical Immunology</i> , 2019, 143, 213-228.e10.	2.9	11
20	Development of novel methods that monitor necroptosis and the release of DAMPs at the single cell resolution. <i>Cell Stress</i> , 2019, 3, 66-69.	3.2	10
21	Generation of and characterization of anti-IL-11 antibodies using newly established Il11-deficient mice. <i>Biochemical and Biophysical Research Communications</i> , 2018, 505, 453-459.	2.1	11
22	A FRET biosensor for necroptosis uncovers two different modes of the release of DAMPs. <i>Nature Communications</i> , 2018, 9, 4457.	12.8	65
23	The AP-1 transcription factor JunB is required for Th17 cell differentiation. <i>Scientific Reports</i> , 2017, 7, 17402.	3.3	47
24	Depletion of myeloid cells exacerbates hepatitis and induces an aberrant increase in histone H3 in mouse serum. <i>Hepatology</i> , 2017, 65, 237-252.	7.3	12
25	Critical Contribution of Nuclear Factor Erythroid 2-related Factor 2 (NRF2) to Electrophile-induced Interleukin-11 Production. <i>Journal of Biological Chemistry</i> , 2017, 292, 205-216.	3.4	22
26	HTLV-1 Tax Induces Formation of the Active Macromolecular IKK Complex by Generating Lys63- and Met1-Linked Hybrid Polyubiquitin Chains. <i>PLoS Pathogens</i> , 2017, 13, e1006162.	4.7	30
27	Novel method to rescue a lethal phenotype through integration of target gene onto the X-chromosome. <i>Scientific Reports</i> , 2016, 6, 37200.	3.3	11
28	Short form FLICE-inhibitory protein promotes TNFÎ±-induced necroptosis in fibroblasts derived from CFLARs transgenic mice. <i>Biochemical and Biophysical Research Communications</i> , 2016, 480, 23-28.	2.1	6
29	Guidelines for the use and interpretation of assays for monitoring autophagy (3rd edition). <i>Autophagy</i> , 2016, 12, 1-222.	9.1	4,701
30	FLIP the Switch: Regulation of Apoptosis and Necroptosis by cFLIP. <i>International Journal of Molecular Sciences</i> , 2015, 16, 30321-30341.	4.1	118
31	Cellular FLICE-Inhibitory Protein Regulates Tissue Homeostasis. <i>Current Topics in Microbiology and Immunology</i> , 2015, 403, 119-141.	1.1	14
32	The adaptor TRAF5 limits the differentiation of inflammatory CD4+ T cells by antagonizing signaling via the receptor for IL-6. <i>Nature Immunology</i> , 2014, 15, 449-456.	14.5	38
33	Critical contribution of oxidative stress to TNFÎ±-induced necroptosis downstream of RIPK1 activation. <i>Biochemical and Biophysical Research Communications</i> , 2013, 436, 212-216.	2.1	76
34	Tumor necrosis factor receptorâ€“associated factor 5 is an essential mediator of ischemic brain infarction. <i>Journal of Neurochemistry</i> , 2013, 126, 400-414.	3.9	36
35	Differential topical susceptibility to TGFÎ² ² in intact and injured regions of the epithelium: key role in myofibroblast transition. <i>Molecular Biology of the Cell</i> , 2013, 24, 3326-3336.	2.1	45
36	Hyperosmotic stress regulates the distribution and stability of myocardin-related transcription factor, a key modulator of the cytoskeleton. <i>American Journal of Physiology - Cell Physiology</i> , 2013, 304, C115-C127.	4.6	30

#	ARTICLE	IF	CITATIONS
37	Shigella IpaH0722 E3 Ubiquitin Ligase Effector Targets TRAF2 to Inhibit PKC ϵ -NF κ B Activity in Invaded Epithelial Cells. <i>PLoS Pathogens</i> , 2013, 9, e1003409.	4.7	58
38	Reciprocal expression of MRTF-A and myocardin is crucial for pathological vascular remodelling in mice. <i>EMBO Journal</i> , 2012, 31, 4428-4440.	7.8	83
39	Interleukin-11 Links Oxidative Stress and Compensatory Proliferation. <i>Science Signaling</i> , 2012, 5, ra5.	3.6	87
40	c-FLIP Maintains Tissue Homeostasis by Preventing Apoptosis and Programmed Necrosis. <i>Science Signaling</i> , 2012, 5, ra93.	3.6	66
41	Aberrant accumulation of interleukin-10-secreting neutrophils in TRAF2-deficient mice. <i>Immunology and Cell Biology</i> , 2012, 90, 881-888.	2.3	6
42	Guidelines for the use and interpretation of assays for monitoring autophagy. <i>Autophagy</i> , 2012, 8, 445-544.	9.1	3,122
43	Crucial role for autophagy in degranulation of mast cells. <i>Journal of Allergy and Clinical Immunology</i> , 2011, 127, 1267-1276.e6.	2.9	120
44	SHARPIN is a component of the NF κ B-activating linear ubiquitin chain assembly complex. <i>Nature</i> , 2011, 471, 633-636.	27.8	557
45	β -Catenin and Smad3 regulate the activity and stability of myocardin-related transcription factor during epithelial \rightarrow myofibroblast transition. <i>Molecular Biology of the Cell</i> , 2011, 22, 4472-4485.	2.1	76
46	An unexpected role for autophagy in degranulation of mast cells. <i>Autophagy</i> , 2011, 7, 657-659.	9.1	24
47	Tumor Necrosis Factor Receptor-associated Factor (TRAF) 2 Controls Homeostasis of the Colon to Prevent Spontaneous Development of Murine Inflammatory Bowel Disease. <i>Journal of Biological Chemistry</i> , 2011, 286, 17879-17888.	3.4	31
48	Importin β 1 Protein-mediated Nuclear Localization of Death Receptor 5 (DR5) Limits DR5/Tumor Necrosis Factor (TNF)-related Apoptosis-inducing Ligand (TRAIL)-induced Cell Death of Human Tumor Cells. <i>Journal of Biological Chemistry</i> , 2011, 286, 43383-43393.	3.4	58
49	Human lactoferrin activates NF κ B through the Toll-like receptor 4 pathway while it interferes with the lipopolysaccharide-stimulated TLR4 signaling. <i>FEBS Journal</i> , 2010, 277, 2051-2066.	4.7	95
50	Fate-determining mechanisms in epithelial \rightarrow myofibroblast transition: major inhibitory role for Smad3. <i>Journal of Cell Biology</i> , 2010, 188, 383-399.	5.2	113
51	TRAF5 Deficiency Accelerates Atherogenesis in Mice by Increasing Inflammatory Cell Recruitment and Foam Cell Formation. <i>Circulation Research</i> , 2010, 107, 757-766.	4.5	48
52	Fate-determining mechanisms in epithelial \rightarrow myofibroblast transition: major inhibitory role for Smad3. <i>Journal of Experimental Medicine</i> , 2010, 207, i5-i5.	8.5	1
53	TRAF5 is a critical mediator of in vitro signals and in vivo functions of LMP1, the viral oncogenic mimic of CD40. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 2009, 106, 17140-17145.	7.1	37
54	TRAF2 Phosphorylation Modulates Tumor Necrosis Factor Alpha-Induced Gene Expression and Cell Resistance to Apoptosis. <i>Molecular and Cellular Biology</i> , 2009, 29, 303-314.	2.3	43

#	ARTICLE	IF	CITATIONS
55	NF- κ B2 (p100) limits TNF- α -induced osteoclastogenesis. <i>Journal of Clinical Investigation</i> , 2009, 119, 2879-2881.	8.2	11
56	Downregulation of c-FLIP promotes caspase-dependent JNK activation and reactive oxygen species accumulation in tumor cells. <i>Oncogene</i> , 2008, 27, 76-84.	5.9	50
57	A critical role of RICK/RIP2 polyubiquitination in Nod-induced NF- κ B activation. <i>EMBO Journal</i> , 2008, 27, 373-383.	7.8	469
58	Rac, PAK and p38 regulate cell contact-dependent nuclear translocation of myocardium-related transcription factor. <i>FEBS Letters</i> , 2008, 582, 291-298.	2.8	49
59	Fusion of OTT to BSAC Results in Aberrant Up-regulation of Transcriptional Activity. <i>Journal of Biological Chemistry</i> , 2008, 283, 26820-26828.	3.4	13
60	Mitochondrial Extrusion through the Cytoplasmic Vacuoles during Cell Death. <i>Journal of Biological Chemistry</i> , 2008, 283, 24128-24135.	3.4	70
61	Low shear stress preferentially enhances IKK activity through selective sources of ROS for persistent activation of NF- κ B in endothelial cells. <i>American Journal of Physiology - Cell Physiology</i> , 2007, 292, C362-C371.	4.6	64
62	Cell Contact-dependent Regulation of Epithelial-Mesenchymal Transition via the Rho-Rho Kinase-Phospho-Myosin Pathway. <i>Molecular Biology of the Cell</i> , 2007, 18, 1083-1097.	2.1	161
63	TRAF-1, -2, -3, -5, and -6 Are Induced in Atherosclerotic Plaques and Differentially Mediate Proinflammatory Functions of CD40L in Endothelial Cells. <i>Arteriosclerosis, Thrombosis, and Vascular Biology</i> , 2007, 27, 1101-1107.	2.4	97
64	TNF Receptor-Associated Factor 2-Dependent Canonical Pathway Is Crucial for the Development of Peyer's Patches. <i>Journal of Immunology</i> , 2007, 178, 2272-2277.	0.8	14
65	Inflammatory Reactive Oxygen Species-Mediated Hemopoietic Suppression in <i>Fancc</i> -Deficient Mice. <i>Journal of Immunology</i> , 2007, 178, 5277-5287.	0.8	67
66	Identification of TNF- α -responsive NF- κ B p65-binding element in the distal promoter of the mouse serine protease inhibitor SerpinE2. <i>FEBS Letters</i> , 2006, 580, 3257-3262.	2.8	8
67	FOG-1 represses GATA-1-dependent Fc γ RII β -chain transcription: transcriptional mechanism of mast-cell-specific gene expression in mice. <i>Blood</i> , 2006, 108, 262-269.	1.4	55
68	Reactive oxygen species mediate crosstalk between NF- κ B and JNK. <i>Cell Death and Differentiation</i> , 2006, 13, 730-737.	11.2	332
69	An antiapoptotic protein, c-FLIPL, directly binds to MKK7 and inhibits the JNK pathway. <i>EMBO Journal</i> , 2006, 25, 5549-5559.	7.8	90
70	The C-terminal Activating Region 2 of the Epstein-Barr Virus-encoded Latent Membrane Protein 1 Activates NF- κ B through TRAF6 and TAK1. <i>Journal of Biological Chemistry</i> , 2006, 281, 2162-2169.	3.4	68
71	Nuclear translocation of the SRF co-activator MAL in cortical neurons: role of RhoA signalling. <i>Journal of Neurochemistry</i> , 2005, 94, 169-180.	3.9	36
72	A revival of old players. <i>EMBO Reports</i> , 2005, 6, 126-127.	4.5	3

#	ARTICLE	IF	CITATIONS
73	NF- κ B RelA Phosphorylation Regulates RelA Acetylation. <i>Molecular and Cellular Biology</i> , 2005, 25, 7966-7975.	2.3	402
74	Osteoclast differentiation independent of the TRANCE \rightarrow RANK \rightarrow TRAF6 axis. <i>Journal of Experimental Medicine</i> , 2005, 202, 589-595.	8.5	335
75	Tumor Necrosis Factor κ (TNF κ) Induces the Unfolded Protein Response (UPR) in a Reactive Oxygen Species (ROS)-dependent Fashion, and the UPR Counteracts ROS Accumulation by TNF κ . <i>Journal of Biological Chemistry</i> , 2005, 280, 33917-33925.	3.4	346
76	Recruitment of Tumor Necrosis Factor Receptor-associated Factor Family Proteins to Apoptosis Signal-regulating Kinase 1 Signalosome Is Essential for Oxidative Stress-induced Cell Death. <i>Journal of Biological Chemistry</i> , 2005, 280, 37033-37040.	3.4	196
77	Transient and Selective NF- κ B p65 Serine 536 Phosphorylation Induced by T Cell Costimulation Is Mediated by I κ B Kinase β and Controls the Kinetics of p65 Nuclear Import. <i>Journal of Immunology</i> , 2004, 172, 6336-6344.	0.8	205
78	TNF Receptor-Associated Factor 5 Limits the Induction of Th2 Immune Responses. <i>Journal of Immunology</i> , 2004, 172, 4292-4297.	0.8	54
79	TRAF Family Proteins Link PKR with NF- κ B Activation. <i>Molecular and Cellular Biology</i> , 2004, 24, 4502-4512.	2.3	147
80	Insufficient p65 phosphorylation at S536 specifically contributes to the lack of NF- κ B activation and transformation in resistant JB6 cells. <i>Carcinogenesis</i> , 2004, 25, 1991-2003.	2.8	117
81	Effects of PU.1-induced mouse calcium \rightarrow calmodulin-dependent kinase I-like kinase (CKLiK) on apoptosis of murine erythroleukemia cells. <i>Experimental Cell Research</i> , 2004, 294, 39-50.	2.6	12
82	Signaling crosstalk between NF- κ B and JNK. <i>Trends in Immunology</i> , 2004, 25, 402-405.	6.8	67
83	Genome wide analysis of TNF-inducible genes reveals that antioxidant enzymes are induced by TNF and responsible for elimination of ROS. <i>Molecular Immunology</i> , 2004, 41, 547-551.	2.2	26
84	NF- κ B inhibits TNF-induced accumulation of ROS that mediate prolonged MAPK activation and necrotic cell death. <i>EMBO Journal</i> , 2003, 22, 3898-3909.	7.8	460
85	TRAF5 Functions in Both RANKL- and TNF κ -Induced Osteoclastogenesis. <i>Journal of Bone and Mineral Research</i> , 2003, 18, 443-450.	2.8	63
86	Phosphorylation of serine 276 is essential for p65 NF- κ B subunit-dependent cellular responses. <i>Biochemical and Biophysical Research Communications</i> , 2003, 300, 807-812.	2.1	145
87	The death domain kinase RIP has an essential role in DNA damage-induced NF- κ B activation. <i>Genes and Development</i> , 2003, 17, 873-882.	5.9	126
88	The Role of Apoptosis Signal-regulating Kinase 1 in Lymphotoxin- β Receptor-mediated Cell Death. <i>Journal of Biological Chemistry</i> , 2003, 278, 16073-16081.	3.4	52
89	Tumor Necrosis Factor- κ -induced IKK Phosphorylation of NF- κ B p65 on Serine 536 Is Mediated through the TRAF2, TRAF5, and TAK1 Signaling Pathway. <i>Journal of Biological Chemistry</i> , 2003, 278, 36916-36923.	3.4	316
90	TWEAK Induces NF- κ B2 p100 Processing and Long Lasting NF- κ B Activation. <i>Journal of Biological Chemistry</i> , 2003, 278, 36005-36012.	3.4	279

#	ARTICLE	IF	CITATIONS
91	Epstein-Barr virus latent membrane protein 1 activation of NF- κ B through IRAK1 and TRAF6. Proceedings of the National Academy of Sciences of the United States of America, 2003, 100, 15595-15600.	7.1	120
92	Multiple Pathways of TWEAK-Induced Cell Death. Journal of Immunology, 2002, 168, 734-743.	0.8	160
93	Identification of a Novel Transcriptional Activator, BSAC, by a Functional Cloning to Inhibit Tumor Necrosis Factor-induced Cell Death. Journal of Biological Chemistry, 2002, 277, 28853-28860.	3.4	81
94	Pro-inflammatory effect of TWEAK/Fn14 interaction on human umbilical vein endothelial cells. Biochemical and Biophysical Research Communications, 2002, 299, 488-493.	2.1	163
95	Lymphotoxin- β receptor mediates NEMO-independent NF- κ B activation. FEBS Letters, 2002, 532, 45-51.	2.8	75
96	Protection Against Fas-Mediated and Tumor Necrosis Factor Receptor 1-Mediated Liver Injury by Blockade of FADD Without Loss of Nuclear Factor- κ B Activation. Annals of Surgery, 2001, 234, 681-688.	4.2	13
97	Critical Roles of TRAF2 and TRAF5 in Tumor Necrosis Factor-induced NF- κ B Activation and Protection from Cell Death. Journal of Biological Chemistry, 2001, 276, 36530-36534.	3.4	277
98	Essential Role of Nuclear Factor (NF)- κ B-Inducing Kinase and Inhibitor of κ B ($\text{I}\kappa$ B) Kinase β in NF- κ B Activation through Lymphotoxin β Receptor, but Not through Tumor Necrosis Factor Receptor I. Journal of Experimental Medicine, 2001, 193, 631-636.	8.5	193
99	Molecular Basis for Hematopoietic/Mesenchymal Interaction during Initiation of Peyer's Patch Organogenesis. Journal of Experimental Medicine, 2001, 193, 621-630.	8.5	217
100	Role of Adapter Function in Oncoprotein-mediated Activation of NF- κ B. Journal of Biological Chemistry, 1999, 274, 17402-17405.	3.4	204
101	CAST, a Novel CD3 μ -binding Protein Transducing Activation Signal for Interleukin-2 Production in T Cells. Journal of Biological Chemistry, 1999, 274, 18173-18180.	3.4	43
102	Ubiquitin-dependent degradation of I κ B is mediated by a ubiquitin ligase Skp1/Cul 1/F-box protein FWD1. Proceedings of the National Academy of Sciences of the United States of America, 1999, 96, 3859-3863.	7.1	192
103	Targeted disruption of Traf5 gene causes defects in CD40- and CD27-mediated lymphocyte activation. Proceedings of the National Academy of Sciences of the United States of America, 1999, 96, 9803-9808.	7.1	183
104	Ku in the Cytoplasm Associates with CD40 in Human B Cells and Translocates into the Nucleus following Incubation with IL-4 and Anti-CD40 mAb. Immunity, 1999, 11, 339-348.	14.3	61
105	ASK1 Is Essential for JNK/SAPK Activation by TRAF2. Molecular Cell, 1998, 2, 389-395.	9.7	625
106	cDNA Cloning, Expression, Subcellular Localization, and Chromosomal Assignment of Mammalian Aurora Homologues, Aurora-Related Kinase (ARK) 1 and 2. Biochemical and Biophysical Research Communications, 1998, 244, 285-292.	2.1	73
107	CD27, a Member of the Tumor Necrosis Factor Receptor Superfamily, Activates NF- κ B and Stress-activated Protein Kinase/c-Jun N-terminal Kinase via TRAF2, TRAF5, and NF- κ B-inducing Kinase. Journal of Biological Chemistry, 1998, 273, 13353-13358.	3.4	223
108	Tumor Necrosis Factor Receptor-associated Factor 6 (TRAF6) Stimulates Extracellular Signal-regulated Kinase (ERK) Activity in CD40 Signaling Along a Ras-independent Pathway. Journal of Experimental Medicine, 1998, 187, 237-244.	8.5	116

#	ARTICLE	IF	CITATIONS
109	Characterization of murine CD70 by molecular cloning and mAb. <i>International Immunology</i> , 1998, 10, 517-526.	4.0	103
110	Differential regulation of I κ B kinase \hat{A} and \hat{A} by two upstream kinases, NF- \hat{A} B-inducing kinase and mitogen-activated protein kinase/ERK kinase kinase-1. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 1998, 95, 3537-3542.	7.1	512
111	Human TNF Receptor-Associated Factor 5 (TRAF5): cDNA Cloning, Expression and Assignment of the TRAF5 Gene to Chromosome 1q32. <i>Genomics</i> , 1997, 42, 26-32.	2.9	15
112	Tumor Necrosis Factor Receptor-associated Factor (TRAF) 5 and TRAF2 Are Involved in CD30-mediated NF \hat{A} B Activation. <i>Journal of Biological Chemistry</i> , 1997, 272, 2042-2045.	3.4	193
113	Specific Interaction of Topoisomerase II $\hat{2}$ and the CD3 $\hat{1}$ μ Chain of the T Cell Receptor Complex. <i>Journal of Biological Chemistry</i> , 1996, 271, 6483-6489.	3.4	111
114	TRAF5, an Activator of NF- \hat{A} B and Putative Signal Transducer for the Lymphotoxin- $\hat{1}$ $\hat{2}$ Receptor. <i>Journal of Biological Chemistry</i> , 1996, 271, 14661-14664.	3.4	311
115	Expansion of Circulating $\hat{3}$ $\hat{1}$ T Cells in Active Sarcoidosis Closely Correlates with Defects in Cellular Immunity. <i>Clinical Immunology and Immunopathology</i> , 1995, 74, 217-222.	2.0	8
116	Induction of G1 arrest by down-regulation of cyclin D3 in T cell hybridomas.. <i>Journal of Experimental Medicine</i> , 1995, 182, 401-408.	8.5	33
117	Purification of glutathione S-transferase fusion proteins as a non-degraded form by using a protease-negative E.coli strain, AD202. <i>Nucleic Acids Research</i> , 1994, 22, 543-544.	14.5	52