

Chris Hawes

List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/9506369/publications.pdf>

Version: 2024-02-01

74
papers

8,600
citations

53794

45
h-index

79698

73
g-index

81
all docs

81
docs citations

81
times ranked

7126
citing authors

#	ARTICLE	IF	CITATIONS
1	Functional characterization of <i>Schistosoma mansoni</i> fucosyltransferases in <i>Nicotiana benthamiana</i> plants. <i>Scientific Reports</i> , 2020, 10, 18528.	3.3	14
2	p24 Family Proteins Are Involved in Transport to the Plasma Membrane of GPI-Anchored Proteins in Plants. <i>Plant Physiology</i> , 2020, 184, 1333-1347.	4.8	19
3	Differences in intracellular localisation of ANKH mutants that relate to mechanisms of calcium pyrophosphate deposition disease and craniometaphyseal dysplasia. <i>Scientific Reports</i> , 2020, 10, 7408.	3.3	6
4	Glycaemic index, glycaemic load and dietary fibre characteristics of two commercially available fruit smoothies. <i>International Journal of Food Sciences and Nutrition</i> , 2019, 70, 116-123.	2.8	2
5	A signal motif retains Arabidopsis ER- α -mannosidase I in the cis-Golgi and prevents enhanced glycoprotein ERAD. <i>Nature Communications</i> , 2019, 10, 3701.	12.8	25
6	Plants Neither Possess nor Require Consciousness. <i>Trends in Plant Science</i> , 2019, 24, 677-687.	8.8	75
7	Optimization of Sample Preparation Methods and SEM Imaging Conditions Enables High Resolution X-ray Mapping of Essential Elements in Biological Specimens. <i>Microscopy and Microanalysis</i> , 2019, 25, 1090-1091.	0.4	0
8	Quantitative analysis of plant ER architecture and dynamics. <i>Nature Communications</i> , 2019, 10, 984.	12.8	56
9	In memoriam " Ian Moore. <i>Journal of Cell Science</i> , 2019, 132, .	2.0	2
10	The odd one out: Arabidopsis reticulon 20 does not bend ER membranes but has a role in lipid regulation. <i>Scientific Reports</i> , 2018, 8, 2310.	3.3	18
11	Protein Storage Vacuoles Originate from Remodeled Preexisting Vacuoles in <i>Arabidopsis thaliana</i> . <i>Plant Physiology</i> , 2018, 177, 241-254.	4.8	52
12	Labeling the ER for Light and Fluorescence Microscopy. <i>Methods in Molecular Biology</i> , 2018, 1691, 1-14.	0.9	6
13	Characterization of Proteins Localized to Plant ER-PM Contact Sites. <i>Methods in Molecular Biology</i> , 2018, 1691, 23-31.	0.9	4
14	Predominant Golgi Residency of the Plant K/HDEL Receptor Is Essential for Its Function in Mediating ER Retention. <i>Plant Cell</i> , 2018, 30, 2174-2196.	6.6	19
15	<i>Arabidopsis</i> Lunapark proteins are involved in ER cisternae formation. <i>New Phytologist</i> , 2018, 219, 990-1004.	7.3	29
16	Auxin and Vesicle Traffic. <i>Plant Physiology</i> , 2018, 176, 1884-1888.	4.8	8
17	Stacks off tracks: a role for the golgin AtCASP in plant endoplasmic reticulum-Golgi apparatus tethering. <i>Journal of Experimental Botany</i> , 2017, 68, 3339-3350.	4.8	36
18	Plant Endoplasmic Reticulum-Plasma Membrane Contact Sites. <i>Trends in Plant Science</i> , 2017, 22, 289-297.	8.8	122

#	ARTICLE	IF	CITATIONS
19	Plant <sc>VAP</sc>27 proteins: domain characterization, intracellular localization and role in plant development. <i>New Phytologist</i> , 2016, 210, 1311-1326.	7.3	110
20	Localization and interactions between Arabidopsis auxin biosynthetic enzymes in the TAA/YUC-dependent pathway. <i>Journal of Experimental Botany</i> , 2016, 67, 4195-4207.	4.8	48
21	A C-terminal amphipathic helix is necessary for the in vivo tubule-shaping function of a plant reticulon. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 2016, 113, 10902-10907.	7.1	49
22	Arabidopsis NAP1 Regulates the Formation of Autophagosomes. <i>Current Biology</i> , 2016, 26, 2060-2069.	3.9	83
23	Actin-dependent vacuolar occupancy of the cell determines auxin-induced growth repression. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 2016, 113, 452-457.	7.1	130
24	Vesicles versus Tubes: Is Endoplasmic Reticulum-Golgi Transport in Plants Fundamentally Different from Other Eukaryotes?. <i>Plant Physiology</i> , 2015, 168, 393-406.	4.8	80
25	Putting the Squeeze on Plasmodesmata: A Role for Reticulons in Primary Plasmodesmata Formation. <i>Plant Physiology</i> , 2015, 168, 1563-1572.	4.8	89
26	Reticulomics: Protein-protein interaction studies with two plasmodesmata-localised reticulon family proteins identify binding partners enriched at plasmodesmata, ER and the plasma membrane. <i>Plant Physiology</i> , 2015, 169, pp.01153.2015.	4.8	76
27	Endoplasmic reticulum localization and activity of maize auxin biosynthetic enzymes. <i>Journal of Experimental Botany</i> , 2015, 66, 6009-6020.	4.8	24
28	The endoplasmic reticulum: A dynamic and well-connected organelle. <i>Journal of Integrative Plant Biology</i> , 2015, 57, 50-62.	8.5	72
29	The transmembrane domain of <i>N</i>-acetylglucosaminyltransferase is the key determinant for its Golgi subcompartmentation. <i>Plant Journal</i> , 2014, 80, 809-822.	5.7	22
30	ER – the key to the highway. <i>Current Opinion in Plant Biology</i> , 2014, 22, 30-38.	7.1	60
31	Fluorescent labelling of the actin cytoskeleton in plants using a cameloid antibody. <i>Plant Methods</i> , 2014, 10, 12.	4.3	57
32	The Plant Cytoskeleton, NET3C, and VAP27 Mediate the Link between the Plasma Membrane and Endoplasmic Reticulum. <i>Current Biology</i> , 2014, 24, 1397-1405.	3.9	180
33	An <sc>Arabidopsis</sc> reticulon and the atlastin homologue <sc>RHD3</sc> act together in shaping the tubular endoplasmic reticulum. <i>New Phytologist</i> , 2013, 197, 481-489.	7.3	50
34	An inhibitor of oil body mobilization in Arabidopsis. <i>New Phytologist</i> , 2013, 200, 641-649.	7.3	25
35	Time-Resolved Fluorescence Imaging Reveals Differential Interactions of <i>N</i>-Glycan Processing Enzymes across the Golgi Stack in <i>Planta</i> . <i>Plant Physiology</i> , 2013, 161, 1737-1754.	4.8	51
36	Secretory Pathway Research: The More Experimental Systems the Better. <i>Plant Cell</i> , 2012, 24, 1316-1326.	6.6	39

#	ARTICLE	IF	CITATIONS
37	The ER/Golgi Interface – Is There Anything in-between?. <i>Frontiers in Plant Science</i> , 2012, 3, 73.	3.6	23
38	A Recycling-Defective Vacuolar Sorting Receptor Reveals an Intermediate Compartment Situated between Prevacuoles and Vacuoles in Tobacco. <i>Plant Cell</i> , 2011, 22, 3992-4008.	6.6	77
39	Optical tweezers for the micromanipulation of plant cytoplasm and organelles. <i>Current Opinion in Plant Biology</i> , 2010, 13, 731-735.	7.1	35
40	Sequential Depletion and Acquisition of Proteins during Golgi Stack Disassembly and Reformation. <i>Traffic</i> , 2010, 11, 1429-1444.	2.7	40
41	Transmembrane domain length is responsible for the ability of a plant reticulon to shape endoplasmic reticulum tubules in vivo. <i>Plant Journal</i> , 2010, 64, 411-418.	5.7	78
42	Golgi membrane dynamics after induction of a dominant-negative mutant Sar1 GTPase in tobacco. <i>Journal of Experimental Botany</i> , 2010, 61, 405-422.	4.8	42
43	Biogenesis of the plant Golgi apparatus. <i>Biochemical Society Transactions</i> , 2010, 38, 761-767.	3.4	25
44	Five <i>Arabidopsis</i> Reticulon Isoforms Share Endoplasmic Reticulum Location, Topology, and Membrane-Shaping Properties. <i>Plant Cell</i> , 2010, 22, 1333-1343.	6.6	173
45	A Missense Mutation in the <i>Arabidopsis</i> COPII Coat Protein Sec24A Induces the Formation of Clusters of the Endoplasmic Reticulum and Golgi Apparatus. <i>Plant Cell</i> , 2009, 21, 3655-3671.	6.6	103
46	A Comparative Study of the Involvement of 17 <i>Arabidopsis</i> Myosin Family Members on the Motility of Golgi and Other Organelles. <i>Plant Physiology</i> , 2009, 150, 700-709.	4.8	163
47	Arginine/Lysine Residues in the Cytoplasmic Tail Promote ER Export of Plant Glycosylation Enzymes. <i>Traffic</i> , 2009, 10, 101-115.	2.7	84
48	Grab a Golgi: Laser Trapping of Golgi Bodies Reveals <i>in vivo</i> Interactions with the Endoplasmic Reticulum. <i>Traffic</i> , 2009, 10, 567-571.	2.7	150
49	Fluorescence Lifetime Imaging of Interactions between Golgi Tethering Factors and Small GTPases in Plants. <i>Traffic</i> , 2009, 10, 1034-1046.	2.7	45
50	The plant endoplasmic reticulum: a cell-wide web. <i>Biochemical Journal</i> , 2009, 423, 145-155.	3.7	107
51	Overexpression of a Plant Reticulon Remodels the Lumen of the Cortical Endoplasmic Reticulum but Does not Perturb Protein Transport. <i>Traffic</i> , 2008, 9, 94-102.	2.7	124
52	The Plant ER–Golgi Interface. <i>Traffic</i> , 2008, 9, 1571-1580.	2.7	60
53	BFA effects are tissue and not just plant specific. <i>Trends in Plant Science</i> , 2008, 13, 405-408.	8.8	116
54	Truncated myosin XI tail fusions inhibit peroxisome, Golgi, and mitochondrial movement in tobacco leaf epidermal cells: a genetic tool for the next generation. <i>Journal of Experimental Botany</i> , 2008, 59, 2499-2512.	4.8	140

#	ARTICLE	IF	CITATIONS
55	Localization and domain characterization of Arabidopsis golgin candidates. <i>Journal of Experimental Botany</i> , 2007, 58, 4373-4386.	4.8	69
56	Golgi Regeneration after Brefeldin A Treatment in BY-2 Cells Entails Stack Enlargement and Cisternal Growth followed by Division. <i>Plant Physiology</i> , 2007, 145, 527-538.	4.8	43
57	Plant neurobiology: no brain, no gain?. <i>Trends in Plant Science</i> , 2007, 12, 135-136.	8.8	146
58	Photoactivation of GFP reveals protein dynamics within the endoplasmic reticulum membrane. <i>Journal of Experimental Botany</i> , 2006, 57, 43-50.	4.8	190
59	In tobacco leaf epidermal cells, the integrity of protein export from the endoplasmic reticulum and of ER export sites depends on active COPI machinery. <i>Plant Journal</i> , 2006, 46, 95-110.	5.7	93
60	Rapid, transient expression of fluorescent fusion proteins in tobacco plants and generation of stably transformed plants. <i>Nature Protocols</i> , 2006, 1, 2019-2025.	12.0	1,534
61	Plant N-Glycan Processing Enzymes Employ Different Targeting Mechanisms for Their Spatial Arrangement along the Secretory Pathway. <i>Plant Cell</i> , 2006, 18, 3182-3200.	6.6	201
62	An Arabidopsis GRIP domain protein locates to the trans-Golgi and binds the small GTPase ARL1. <i>Plant Journal</i> , 2005, 44, 459-470.	5.7	66
63	Holding it all together? Candidate proteins for the plant Golgi matrix. <i>Current Opinion in Plant Biology</i> , 2005, 8, 632-639.	7.1	65
64	AtPEX2 and AtPEX10 Are Targeted to Peroxisomes Independently of Known Endoplasmic Reticulum Trafficking Routes. <i>Plant Physiology</i> , 2005, 139, 690-700.	4.8	67
65	Sec22 and Memb11 Are v-SNAREs of the Anterograde Endoplasmic Reticulum-Golgi Pathway in Tobacco Leaf Epidermal Cells. <i>Plant Physiology</i> , 2005, 139, 1244-1254.	4.8	79
66	Endoplasmic Reticulum Export Sites and Golgi Bodies Behave as Single Mobile Secretory Units in Plant Cells[W]. <i>Plant Cell</i> , 2004, 16, 1753-1771.	6.6	258
67	A GFP-based assay reveals a role for RHD3 in transport between the endoplasmic reticulum and Golgi apparatus. <i>Plant Journal</i> , 2004, 37, 398-414.	5.7	148
68	ER quality control can lead to retrograde transport from the ER lumen to the cytosol and the nucleoplasm in plants. <i>Plant Journal</i> , 2003, 34, 269-281.	5.7	118
69	The Destination for Single-Pass Membrane Proteins Is Influenced Markedly by the Length of the Hydrophobic Domain. <i>Plant Cell</i> , 2002, 14, 1077-1092.	6.6	207
70	Membrane Protein Transport between the Endoplasmic Reticulum and the Golgi in Tobacco Leaves Is Energy Dependent but Cytoskeleton Independent. <i>Plant Cell</i> , 2002, 14, 1293-1309.	6.6	303
71	Redistribution of membrane proteins between the Golgi apparatus and endoplasmic reticulum in plants is reversible and not dependent on cytoskeletal networks. <i>Plant Journal</i> , 2002, 29, 661-678.	5.7	247
72	ER confirmed as the location of mystery organelles in Arabidopsis plants expressing GFP!. <i>Trends in Plant Science</i> , 2001, 6, 245-246.	8.8	71

#	ARTICLE	IF	CITATIONS
73	A Rab1 GTPase Is Required for Transport between the Endoplasmic Reticulum and Golgi Apparatus and for Normal Golgi Movement in Plants. <i>Plant Cell</i> , 2000, 12, 2201-2217.	6.6	550
74	Stacks on tracks: the plant Golgi apparatus traffics on an actin/ER network. <i>Plant Journal</i> , 1998, 15, 441-447.	5.7	818