Clarissa M Comim

List of Publications by Year in descending order

Source: https://exaly.com/author-pdf/9570610/publications.pdf Version: 2024-02-01



#	Article	IF	CITATIONS
1	Effects of low-intensity training on the brain and muscle in the congenital muscular dystrophy 1D model. Neurological Sciences, 2022, , 1.	1.9	Ο
2	Swimming Improves Memory and Antioxidant Defense in an Animal Model of Duchenne Muscular Dystrophy. Molecular Neurobiology, 2021, 58, 5067-5077.	4.0	4
3	Effects of chronic treatment with gold nanoparticles on inflammatory responses and oxidative stress in Mdx mice. Journal of Drug Targeting, 2020, 28, 46-54.	4.4	20
4	Involvement of NLRP3 inflammasome in schizophrenia-like behaviour in young animals after maternal immune activation. Acta Neuropsychiatrica, 2020, 32, 321-327.	2.1	11
5	Effect of Aerobic Physical Exercise in an Animal Model of Duchenne Muscular Dystrophy. Journal of Molecular Neuroscience, 2020, 70, 1552-1564.	2.3	3
6	Late Brain Involvement after Neonatal Immune Activation. BioMed Research International, 2019, 2019, 1-11.	1.9	5
7	Neurocognitive Impairment in mdx Mice. Molecular Neurobiology, 2019, 56, 7608-7616.	4.0	15
8	Dermatoglyphical impressions are different between children and adolescents with normal weight, overweight and obesity: a cross-sectional study. F1000Research, 2019, 8, 964.	1.6	5
9	Neonatal Immune Activation May Provoke Long-term Depressive Attributes. Current Neurovascular Research, 2019, 16, 358-364.	1.1	3
10	Dextran Sulphate of Sodium-induced colitis in mice: antihyperalgesic effects of ethanolic extract of Citrus reticulata and potential damage to the central nervous system. Anais Da Academia Brasileira De Ciencias, 2018, 90, 3139-3145.	0.8	7
11	Early fragmentation of polyester urethane sheet neither causes persistent oxidative stress nor alters the outcome of normal tissue healing in rat skin. Anais Da Academia Brasileira De Ciencias, 2018, 90, 2211-2222.	0.8	1
12	Inhibition of indoleamine 2,3-dioxygenase 1/2 prevented cognitive impairment and energetic metabolism changes in the hippocampus of adult rats subjected to polymicrobial sepsis. Journal of Neuroimmunology, 2017, 305, 167-171.	2.3	21
13	Congenital Muscular Dystrophy 1D Causes Matrix Metalloproteinase Activation And Blood-Brain Barrier Impairment. Current Neurovascular Research, 2017, 14, 60-64.	1.1	3
14	Activity of Krebs cycle enzymes in <i>mdx</i> mice. Muscle and Nerve, 2016, 53, 91-95.	2.2	8
15	Experimental Neonatal Sepsis Causes Long-Term Cognitive Impairment. Molecular Neurobiology, 2016, 53, 5928-5934.	4.0	28
16	Behavioral Responses in Animal Model of Congenital Muscular Dystrophy 1D. Molecular Neurobiology, 2016, 53, 402-407.	4.0	5
17	Neurotrophins, cytokines, oxidative parameters and funcionality in Progressive Muscular Dystrophies. Anais Da Academia Brasileira De Ciencias, 2015, 87, 1809-1818.	0.8	13
18	Effects of sodium butyrate on aversive memory in rats submitted to sepsis. Neuroscience Letters, 2015, 595, 134-138.	2.1	28

#	Article	IF	CITATIONS
19	Effect of sepsis on behavioral changes on the ketamine-induced animal model of schizophrenia. Journal of Neuroimmunology, 2015, 281, 78-82.	2.3	5
20	Interleukin-1β Receptor Antagonism Prevents Cognitive Impairment Following Experimental Bacterial Meningitis. Current Neurovascular Research, 2015, 12, 253-261.	1.1	13
21	Environmental enrichment restores cognitive deficits induced by experimental childhood meningitis. Revista Brasileira De Psiquiatria, 2014, 36, 322-329.	1.7	12
22	Methylphenidate treatment causes oxidative stress and alters energetic metabolism in an animal model of attention-deficit hyperactivity disorder. Acta Neuropsychiatrica, 2014, 26, 96-103.	2.1	31
23	Evaluation of NCS-1, DARPP-32, and neurotrophins in hippocampus and prefrontal cortex in rats submitted to sepsis. Synapse, 2014, 68, 474-479.	1.2	10
24	Acute Brain Inflammation and Oxidative Damage Are Related to Long-Term Cognitive Deficits and Markers of Neurodegeneration in Sepsis-Survivor Rats. Molecular Neurobiology, 2014, 49, 380-385.	4.0	72
25	Erythropoietin prevents cognitive impairment and oxidative parameters in Wistar rats subjected to pneumococcal meningitis. Translational Research, 2014, 163, 503-513.	5.0	21
26	NCS-1 deficiency causes anxiety and depressive-like behavior with impaired non-aversive memory in mice. Physiology and Behavior, 2014, 130, 91-98.	2.1	33
27	Inhibition of matrix metalloproteinases-2 and -9 prevents cognitive impairment induced by pneumococcal meningitis in Wistar rats. Experimental Biology and Medicine, 2014, 239, 225-231.	2.4	33
28	Inflammation biomarkers and delirium in critically ill patients. Critical Care, 2014, 18, R106.	5.8	79
29	ll1-β Involvement in Cognitive Impairment after Sepsis. Molecular Neurobiology, 2014, 49, 1069-1076.	4.0	87
30	Neonatal Escherichia coli K1 meningitis causes learning and memory impairments in adulthood. Journal of Neuroimmunology, 2014, 272, 35-41.	2.3	20
31	Attenuation of cognitive impairment by the nonbacteriolytic antibiotic daptomycin in Wistar rats submitted to pneumococcal meningitis. BMC Neuroscience, 2013, 14, 42.	1.9	20
32	Matrix Metalloproteinase-2 and Metalloproteinase-9 Activities are Associated with Blood–Brain Barrier Dysfunction in an Animal Model of Severe Sepsis. Molecular Neurobiology, 2013, 48, 62-70.	4.0	91
33	Central Nervous System Involvement in the Animal Model of Myodystrophy. Molecular Neurobiology, 2013, 48, 71-77.	4.0	5
34	Inhibition of indoleamine 2,3-dioxygenase prevented cognitive impairment in adult Wistar rats subjected to pneumococcal meningitis. Translational Research, 2013, 162, 390-397.	5.0	26
35	Late brain alterations in sepsisâ€survivor rats. Synapse, 2013, 67, 786-793.	1.2	22
36	Caspase-3 Mediates In Part Hippocampal Apoptosis in Sepsis. Molecular Neurobiology, 2013, 47, 394-398.	4.0	48

#	Article	lF	CITATIONS
37	Chronic administration of branchedâ€chain amino acids impairs spatial memory and increases brainâ€derived neurotrophic factor in a rat model. Journal of Inherited Metabolic Disease, 2013, 36, 721-730.	3.6	27
38	Behavioral Responses in Rats Submitted to Chronic Administration of Branched-Chain Amino Acids. JIMD Reports, 2013, 13, 159-167.	1.5	14
39	Methylphenidate treatment affects mitogen-activated protein kinase activation in the striatum of young rats. Acta Neuropsychiatrica, 2013, 25, 235-239.	2.1	0
40	Correlation of Acute Phase Inflammatory and Oxidative Markers With Long-term Cognitive Impairment in Sepsis Survivors Rats. Shock, 2013, 40, 45-48.	2.1	34
41	Brain-derived neurotrophic factor plasma levels are associated with mortality in critically ill patients even in the absence of brain injury. Critical Care, 2012, 16, R234.	5.8	16
42	Cannabidiol reduces host immune response and prevents cognitive impairments in Wistar rats submitted to pneumococcal meningitis. European Journal of Pharmacology, 2012, 697, 158-164.	3.5	61
43	Erythropoietin reverts cognitive impairment and alters the oxidative parameters and energetic metabolism in sepsis animal model. Journal of Neural Transmission, 2012, 119, 1267-1274.	2.8	16
44	Antioxidant treatment prevents cognitive impairment and oxidative damage in pneumococcal meningitis survivor rats. Metabolic Brain Disease, 2012, 27, 587-593.	2.9	14
45	Effects of experimental cerebral malaria in memory, brain-derived neurotrophic factor and acetylcholinesterase acitivity in the hippocampus of survivor mice. Neuroscience Letters, 2012, 523, 104-107.	2.1	22
46	Antioxidant administration prevents memory impairment in an animal model of maple syrup urine disease. Behavioural Brain Research, 2012, 231, 92-96.	2.2	23
47	Protective effects of guanosine against sepsis-induced damage in rat brain and cognitive impairment. Brain, Behavior, and Immunity, 2012, 26, 904-910.	4.1	61
48	Imipramine reverses depressive-like parameters in pneumococcal meningitis survivor rats. Journal of Neural Transmission, 2012, 119, 653-660.	2.8	12
49	NMDA preconditioning prevents object recognition memory impairment and increases brain viability in mice exposed to traumatic brain injury. Brain Research, 2012, 1466, 82-90.	2.2	29
50	Reduction of acethylcolinesterase activity in the brain of mdx mice. Neuromuscular Disorders, 2011, 21, 359-362.	0.6	7
51	Evaluation of light/dark cycle in anxiety- and depressive-like behaviors after regular treatment with methylphenidate hydrochloride in rats of different ages. Revista Brasileira De Psiquiatria, 2011, 33, 55-58.	1.7	13
52	Dexamethasone Treatment Reverses Cognitive Impairment but Increases Brain Oxidative Stress in Rats Submitted to Pneumococcal Meningitis. Oxidative Medicine and Cellular Longevity, 2011, 2011, 1-7.	4.0	8
53	Serum Heat Shock Protein 70 Levels, Oxidant Status, and Mortality in Sepsis. Shock, 2011, 35, 466-470.	2.1	65
54	Acute low dose of MK-801 prevents memory deficits without altering hippocampal DARPP-32 expression and BDNF levels in sepsis survivor rats. Journal of Neuroimmunology, 2011, 230, 48-51.	2.3	19

#	Article	IF	CITATIONS
55	Alterations in Inflammatory Mediators, Oxidative Stress Parameters and Energetic Metabolism in the Brain of Sepsis Survivor Rats. Neurochemical Research, 2011, 36, 304-311.	3.3	53
56	Aversive memory in sepsis survivor rats. Journal of Neural Transmission, 2011, 118, 213-217.	2.8	15
57	Traffic of leukocytes and cytokine up-regulation in the central nervous system in sepsis. Intensive Care Medicine, 2011, 37, 711-718.	8.2	78
58	Diurnal differences in memory and learning in young and adult rats treated with methylphenidate. Journal of Neural Transmission, 2010, 117, 457-462.	2.8	15
59	Time-dependent behavioral recovery after pneumococcal meningitis in rats. Journal of Neural Transmission, 2010, 117, 819-826.	2.8	23
60	Depressive-Like Parameters in Sepsis Survivor Rats. Neurotoxicity Research, 2010, 17, 279-286.	2.7	28
61	TNF-α, IL-1β, IL-6, and cinc-1 levels in rat brain after meningitis induced by Streptococcus pneumoniae. Journal of Neuroimmunology, 2010, 221, 42-45.	2.3	56
62	Correlation between behavioral deficits and decreased brain-derived neurotrofic factor in neonatal meningitis. Journal of Neuroimmunology, 2010, 223, 73-76.	2.3	32
63	Treatment with cannabidiol reverses oxidative stress parameters, cognitive impairment and mortality in rats submitted to sepsis by cecal ligation and puncture. Brain Research, 2010, 1348, 128-138.	2.2	72
64	Mitochondrial respiratory chain and creatine kinase activities in <i>mdx</i> mouse brain. Muscle and Nerve, 2010, 41, 257-260.	2.2	12
65	Effect of chronic administration of ketamine on the mitochondrial respiratory chain activity caused by chronic mild stress. Acta Neuropsychiatrica, 2010, 22, 292-299.	2.1	12
66	Cognitive Dysfunction Is Sustained after Rescue Therapy in Experimental Cerebral Malaria, and Is Reduced by Additive Antioxidant Therapy. PLoS Pathogens, 2010, 6, e1000963.	4.7	91
67	Low dose dexamethasone reverses depressive-like parameters and memory impairment in rats submitted to sepsis. Neuroscience Letters, 2010, 473, 126-130.	2.1	22
68	Antibiotic therapy prevents, in part, the oxidative stress in the rat brain after meningitis induced by Streptococcus pneumoniae. Neuroscience Letters, 2010, 478, 93-96.	2.1	29
69	Depressive-like-behavior and proinflamatory interleukine levels in the brain of rats submitted to pneumococcal meningitis. Brain Research Bulletin, 2010, 82, 243-246.	3.0	22
70	Cognitive Impairment in the Septic Brain. Current Neurovascular Research, 2009, 6, 194-203.	1.1	44
71	Ketamine induces rapid onset of antidepressant action: neurophysiological biomarkers as predictors of effect. Biomarkers in Medicine, 2009, 3, 5-8.	1.4	3
72	Effects of acute treatment with amphetamine in locomotor activity in sepsis survivor rats. Journal of Neuroimmunology, 2009, 212, 145-147.	2.3	4

#	Article	IF	CITATIONS
73	Increased oxidative stress in submitochondrial particles into the brain of rats submitted to the chronic mild stress paradigm. Journal of Psychiatric Research, 2009, 43, 864-869.	3.1	120
74	Methylphenidate treatment increases Na+, K+-ATPase activity in the cerebrum of young and adult rats. Journal of Neural Transmission, 2009, 116, 1681-1687.	2.8	13
75	Acute administration of ketamine reverses the inhibition of mitochondrial respiratory chain induced by chronic mild stress. Brain Research Bulletin, 2009, 79, 418-421.	3.0	54
76	Ketamine treatment reverses behavioral and physiological alterations induced by chronic mild stress in rats. Progress in Neuro-Psychopharmacology and Biological Psychiatry, 2009, 33, 450-455.	4.8	214
77	Effects of chronic mild stress on the oxidative parameters in the rat brain. Neurochemistry International, 2009, 54, 358-362.	3.8	217
78	Striatum brain-derived neurotrophic factor levels are decreased in dystrophin-deficient mice. Neuroscience Letters, 2009, 459, 66-68.	2.1	9
79	Early antibiotic administration prevents cognitive impairment induced by meningitis in rats. Neuroscience Letters, 2009, 465, 71-73.	2.1	8
80	Tumor necrosis factor alpha (TNF-α) levels in the brain and cerebrospinal fluid after meningitis induced by Streptococcus pneumoniae. Neuroscience Letters, 2009, 467, 217-219.	2.1	44
81	Oxidative variables and antioxidant enzymes activities in the mdx mouse brain. Neurochemistry International, 2009, 55, 802-805.	3.8	13
82	RIVASTIGMINE REVERSES HABITUATION MEMORY IMPAIRMENT OBSERVED IN SEPSIS SURVIVOR RATS. Shock, 2009, 32, 270-271.	2.1	25
83	Effect of acute administration of ketamine and imipramine on creatine kinase activity in the brain of rats. Revista Brasileira De Psiquiatria, 2009, 31, 247-252.	1.7	28
84	Chronic Methylphenidate-Effects Over Circadian Cycle of Young and Adult Rats Submitted to Open-Field and Object Recognition Tests. Current Neurovascular Research, 2009, 6, 259-266.	1.1	11
85	DARPP-32 and NCS-1 Expression is not Altered in Brains of Rats Treated with Typical or Atypical Antipsychotics. Neurochemical Research, 2008, 33, 533-538.	3.3	27
86	The Septic Brain. Neurochemical Research, 2008, 33, 2171-2177.	3.3	65
87	Time-dependent behavioral recovery after sepsis in rats. Intensive Care Medicine, 2008, 34, 1724-1731.	8.2	93
88	Chronic Administration of Ketamine Elicits Antidepressant‣ike Effects in Rats without Affecting Hippocampal Brainâ€Đerived Neurotrophic Factor Protein Levels. Basic and Clinical Pharmacology and Toxicology, 2008, 103, 502-506.	2.5	101
89	DNA Damage after Acute and Chronic Treatment with Malathion in Rats. Journal of Agricultural and Food Chemistry, 2008, 56, 7560-7565.	5.2	36
90	Mitochondrial respiratory chain and creatine kinase activities in rat brain after sepsis induced by cecal ligation and perforation. Mitochondrion, 2008, 8, 313-318.	3.4	74

#	Article	IF	CITATIONS
91	Inhibition of mitochondrial respiratory chain in brain of rats subjected to an experimental model of depression. Neurochemistry International, 2008, 53, 395-400.	3.8	172
92	Acute administration of ketamine induces antidepressant-like effects in the forced swimming test and increases BDNF levels in the rat hippocampus. Progress in Neuro-Psychopharmacology and Biological Psychiatry, 2008, 32, 140-144.	4.8	377
93	Memory-enhancing treatments reverse the impairment of inhibitory avoidance retention in sepsis-surviving rats. Critical Care, 2008, 12, R133.	5.8	17
94	Chronic Mild Stress Paradigm Reduces Sweet Food Intake in Rats without Affecting Brain Derived Neurotrophic Factor Protein Levels. Current Neurovascular Research, 2008, 5, 207-213.	1.1	45
95	DNA damage in rats after treatment with methylphenidate. Progress in Neuro-Psychopharmacology and Biological Psychiatry, 2007, 31, 1282-1288.	4.8	64
96	Effect of Antipsychotics on Creatine Kinase Activity in Rat Brain. Basic and Clinical Pharmacology and Toxicology, 2007, 101, 315-319.	2.5	20
97	Imipramine reverses the depressive symptoms in sepsis survivor rats. Intensive Care Medicine, 2007, 33, 2165-2167.	8.2	23
98	Dermatoglyphics and abdominal resistance in female children and adolescents: a cross-sectional study. F1000Research, 0, 10, 945.	1.6	0