

# Leonardo Velasco

## List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/9574443/publications.pdf>

Version: 2024-02-01

177  
papers

3,367  
citations

136950

32  
h-index

243625

44  
g-index

179  
all docs

179  
docs citations

179  
times ranked

2240  
citing authors

#	ARTICLE	IF	CITATIONS
1	Genetic and physiological characterization of sunflower resistance provided by the wild-derived OrDeb2 gene against highly virulent races of <i>Orobanche cumana</i> Wallr. <i>Theoretical and Applied Genetics</i> , 2022, 135, 501-525.	3.6	9
2	Characterization of press and solvent extraction oils from new sunflower seeds with modified phytosterol compositions. <i>Journal of the Science of Food and Agriculture</i> , 2021, 101, 101-109.	3.5	6
3	Chemical components influencing oxidative stability and sensorial properties of extra virgin olive oil and effect of genotype and location on their expression. <i>LWT - Food Science and Technology</i> , 2021, 136, 110257.	5.2	21
4	Attempts of Physical Refining of Sterol-Rich Sunflower Press Oil to Obtain Minimally Processed Edible Oil. <i>Foods</i> , 2021, 10, 1901.	4.3	3
5	Genotype by environment interaction for oil quality components in olive tree. <i>European Journal of Agronomy</i> , 2020, 119, 126115.	4.1	20
6	Gene banks for wild and cultivated sunflower genetic resources. <i>OCL - Oilseeds and Fats, Crops and Lipids</i> , 2020, 27, 9.	1.4	20
7	Characterization of post-haustorial resistance to sunflower broomrape. <i>Crop Science</i> , 2020, 60, 1188-1198.	1.8	15
8	Comparative Study of Fatty Acid Composition, Total Phenolics, and Antioxidant Capacity in Rapeseed Mutant Lines. <i>JAOCS, Journal of the American Oil Chemists' Society</i> , 2020, 97, 397-407.	1.9	3
9	Oil Phytosterol Concentration in Sunflower Presents a Dilution Response with Oil Weight per Grain. <i>JAOCS, Journal of the American Oil Chemists' Society</i> , 2019, 96, 1115-1123.	1.9	5
10	An SSR-SNP Linkage Map of the Parasitic Weed <i>Orobanche cumana</i> Wallr. Including a Gene for Plant Pigmentation. <i>Frontiers in Plant Science</i> , 2019, 10, 797.	3.6	9
11	A receptor-like kinase enhances sunflower resistance to <i>Orobanche cumana</i> . <i>Nature Plants</i> , 2019, 5, 1211-1215.	9.3	53
12	First Report of Sunflower Broomrape ( <i>Orobanche cumana</i> ) in Portugal. <i>Plant Disease</i> , 2019, 103, 2143-2143.	1.4	4
13	An Analytical Simplification for Faster Determination of Fatty Acid Composition and Phytosterols in Seed Oils. <i>Food Analytical Methods</i> , 2018, 11, 1234-1242.	2.6	9
14	First Report of Sunflower Broomrape, <i>Orobanche cumana</i> Wallr., in Morocco. <i>Plant Disease</i> , 2018, 102, 457-457.	1.4	15
15	Genetic Diversity of a Germplasm Collection of Confectionery Sunflower Landraces from Spain. <i>Crop Science</i> , 2018, 58, 1972-1981.	1.8	4
16	Using Wild Olives in Breeding Programs: Implications on Oil Quality Composition. <i>Frontiers in Plant Science</i> , 2018, 9, 232.	3.6	33
17	Dynamics of phytosterols content and concentration in sunflower grains. <i>Crop and Pasture Science</i> , 2018, 69, 724.	1.5	6
18	Research on resistance to sunflower broomrape: an integrated vision. <i>OCL - Oilseeds and Fats, Crops and Lipids</i> , 2016, 23, D203.	1.4	13

#	ARTICLE	IF	CITATIONS
19	Genetic Analysis of Reduced $\gamma$ -Tocopherol Content in Ethiopian Mustard Seeds. Scientific World Journal, The, 2016, 2016, 1-7.	2.1	1
20	Increased Virulence in Sunflower Broomrape ( <i>Orobanche cumana</i> Wallr.) Populations from Southern Spain Is Associated with Greater Genetic Diversity. Frontiers in Plant Science, 2016, 7, 589.	3.6	28
21	Sunflower Resistance to Broomrape ( <i>Orobanche cumana</i> ) Is Controlled by Specific QTLs for Different Parasitism Stages. Frontiers in Plant Science, 2016, 7, 590.	3.6	45
22	Early selection for oil quality components in olive breeding progenies. European Journal of Lipid Science and Technology, 2016, 118, 1160-1167.	1.5	38
23	Erucic acid in feed and food. EFSA Journal, 2016, 14, e04593.	1.8	45
24	Molecular basis of the high-palmitic acid trait in sunflower seed oil. Molecular Breeding, 2016, 36, 1.	2.1	9
25	Genetic study of recessive broomrape resistance in sunflower. Euphytica, 2016, 209, 419-428.	1.2	18
26	Tocopherols in Sunflower Seedlings under Light and Dark Conditions. Scientific World Journal, The, 2015, 2015, 1-11.	2.1	5
27	Sunflower Broomrape ( <i>Orobanche cumana</i> Wallr.), , 2015, , 129-155.		12
28	Sunflower Production in the European Union**This chapter does not consider sunflower producing countries such as Bulgaria, Croatia, Greece, and Romania, which are included in the chapter on Eastern European zone, except when the global EU figures are calculated. Part of the information provided in this chapter was obtained from technical personnel of several seed and oil companies, who preferred to remain anonymous.. , 2015, , 555-573.		1
29	Sunflower Oil Minor Constituents. , 2015, , 297-329.		8
30	Association mapping for kernel phytosterol content in almond. Frontiers in Plant Science, 2015, 6, 530.	3.6	20
31	Natural variability in phytosterols in almond ( <i>Prunus amygdalus</i> ) trees growing under a southern Mediterranean climate. Journal of Horticultural Science and Biotechnology, 2015, 90, 543-549.	1.9	4
32	Variability of seed quality traits in wild and semi-wild accessions of castor collected in Spain. Industrial Crops and Products, 2015, 65, 203-209.	5.2	13
33	Initial selection steps in olive breeding programs. Euphytica, 2015, 201, 453-462.	1.2	16
34	Advanced olive selections with enhanced quality for minor constituents. Grasas Y Aceites, 2015, 66, e100.	0.9	7
35	Variability for Seed Phytosterols in Sunflower Germplasm. Crop Science, 2014, 54, 190-197.	1.8	16
36	The Genetic Structure of Wild <i>Orobanche cumana</i> Wallr. ( <i>Orobanchaceae</i> ) Populations in Eastern Bulgaria Reflects Introgressions from Weedy Populations. Scientific World Journal, The, 2014, 2014, 1-15.	2.1	12

#	ARTICLE	IF	CITATIONS
37	New sunflower seeds with high contents of phytosterols. OCL - Oilseeds and Fats, Crops and Lipids, 2014, 21, D604.	1.4	8
38	Selection for Some Olive Oil Quality Components Through the Analysis of Fruit Flesh. JAOCS, Journal of the American Oil Chemists' Society, 2014, 91, 1731-1736.	1.9	20
39	Genetic Studies in Sunflower Broomrape. Helia, 2014, 37, .	0.4	1
40	Phylogenetic Relationships and Genetic Diversity among <i>Orobanche cumana</i> Wallr. and <i>O. cernua</i> L. (Orobanchaceae) Populations in the Iberian Peninsula. Helia, 2014, 37, .	0.4	3
41	Novel safflower oil with high $\alpha$ -tocopherol content has a high oxidative stability. European Journal of Lipid Science and Technology, 2014, 116, 832-836.	1.5	24
42	Changes in plastochromanol-8 and tocopherols during germination in Ethiopian mustard lines with contrasting tocopherol levels. Seed Science Research, 2014, 24, 101-112.	1.7	5
43	Identification, characterisation and discriminatory power of microsatellite markers in the parasitic weed <i>Orobanche cumana</i> . Weed Research, 2014, 54, 120-132.	1.7	18
44	Characterization of a $\beta$ -tocopherol methyltransferase mutant gene in wild ( <i>Carthamus oxyacanthus</i> M.) Tj ETQq0 0,0,rgBT /Oyerlock 10	1.2	0
45	Variability of seed quality traits in a collection of Spanish landraces of confectionery sunflower. Crop and Pasture Science, 2014, 65, 242.	1.5	5
46	Broomrape ( <i>Orobanche Cumana</i> Wallr.) Resistance Breeding Utilizing Wild <i>Helianthus</i> Species. Helia, 2014, 37, .	0.4	8
47	Extent of cross-fertilization in <i>Orobanche cumana</i> Wallr.. Biologia Plantarum, 2013, 57, 559-562.	1.9	14
48	Accumulation dynamics of seed tocopherols in sunflower lines with modified tocopherol levels. Acta Physiologiae Plantarum, 2013, 35, 3157-3165.	2.1	3
49	Variability of Phytosterols in <i>Jatropha curcas</i> Germplasm. JAOCS, Journal of the American Oil Chemists' Society, 2013, 90, 1713-1718.	1.9	1
50	A dominant avirulence gene in <i>Orobanche cumana</i> triggers <i>O</i> r5 resistance in sunflower. Weed Research, 2013, 53, 322-327.	1.7	25
51	Fruit characteristics and fatty acid composition in advanced olive breeding selections along the ripening period. Food Research International, 2013, 54, 1890-1896.	6.2	40
52	Changes in squalene and sterols associated with olive maturation. Food Research International, 2013, 54, 1885-1889.	6.2	64
53	Genetic diversity of <i>Orobanche cumana</i> populations from Spain assessed using SSR markers. Weed Research, 2013, 53, 279-289.	1.7	34
54	Near-infrared spectroscopy for analysis of oil content and fatty acid profile in almond flour. European Journal of Lipid Science and Technology, 2013, 115, 211-216.	1.5	13

#	ARTICLE	IF	CITATIONS
55	Selection for contrasting tocopherol content and profile in Ethiopian mustard. <i>Plant Breeding</i> , 2013, 132, 694-700.	1.9	7
56	Genetic Variation and Genotype × Environment Interactions for Seed Phytosterols in Sunflower. <i>Crop Science</i> , 2013, 53, 1589-1593.	1.8	14
57	Marker-Assisted and Physiology-Based Breeding for Resistance to Root Parasitic Orobanchaceae. , 2013, , 369-391.		23
58	FRUIT AND OIL CHARACTERISTICS OF ADVANCED SELECTIONS FROM AN OLIVE BREEDING PROGRAM. <i>Acta Horticulturae</i> , 2013, , 415-419.	0.2	2
59	Environmental Stability of Contrasting Seed Tocopherol Profiles in Sunflower. <i>Crop Science</i> , 2012, 52, 2446-2452.	1.8	6
60	Quantitative Trait Loci for Seed Tocopherol Content in Sunflower. <i>Crop Science</i> , 2012, 52, 786-794.	1.8	6
61	Progress in research on breeding for resistance to sunflower broomrape. <i>Helia</i> , 2012, 35, 47-56.	0.4	12
62	Tocopherols and phytosterols in sunflower seeds for the human food market. <i>Grasas Y Aceites</i> , 2012, 63, 321-327.	0.9	13
63	Mapping of major and modifying genes for high oleic acid content in safflower. <i>Molecular Breeding</i> , 2012, 30, 1279-1293.	2.1	31
64	Genetic basis of unstable expression of high gamma-tocopherol content in sunflower seeds. <i>BMC Plant Biology</i> , 2012, 12, 71.	3.6	16
65	Castor. , 2012, , 237-265.		2
66	Validation of a method for the analysis of phytosterols in sunflower seeds. <i>European Journal of Lipid Science and Technology</i> , 2012, 114, 325-331.	1.5	36
67	Free sterols and steryl glycosides in sunflower seeds with high phytosterol contents. <i>European Journal of Lipid Science and Technology</i> , 2012, 114, 1212-1216.	1.5	12
68	Identification of High Oleic Castor Seeds by Near Infrared Reflectance Spectroscopy. <i>JAOCS, Journal of the American Oil Chemists' Society</i> , 2012, 89, 431-435.	1.9	3
69	Inheritance of resistance to sunflower broomrape ( <i>Orobanche cumana</i> Wallr.) in an interspecific cross between <i>Helianthus annuus</i> and <i>Helianthus debilis</i> subsp <i> . tardiflorus</i> . <i>Plant Breeding</i> , 2012, 131, 220-221.	1.9	49
70	Tocochromanol content and composition in <i>Jatropha curcas</i> seeds. <i>Industrial Crops and Products</i> , 2012, 36, 304-307.	5.2	6
71	Expression of modified tocopherol content and profile in sunflower tissues. <i>Journal of the Science of Food and Agriculture</i> , 2012, 92, 351-357.	3.5	5
72	Phytosterol Variability in Almond Germplasm. <i>Journal of the American Society for Horticultural Science</i> , 2012, 137, 343-348.	1.0	19

#	ARTICLE	IF	CITATIONS
73	Short communication. Estimation of cross-fertilization rate in safflower ( <i>Carthamus tinctorius</i> L.). Spanish Journal of Agricultural Research, 2012, 10, 155.	0.6	2
74	Performance and seed quality of Moroccan sunflower varieties and Spanish landraces used for confectionery and snack food. <i>Helia</i> , 2011, 34, 75-82.	0.4	3
75	Development and characterization of genomic microsatellite markers in safflower ( <i>Carthamus tinctorius</i> L.). <i>Plant Breeding</i> , 2011, 130, 540-543.	1.9	2
76	Inheritance of increased seed tocopherol content in sunflower line IASTA413. <i>Plant Breeding</i> , 2011, 130, 540-543.	1.9	2
77	Inheritance of the unpigmented plant trait in <i>Orobanche cumana</i> . <i>Weed Research</i> , 2011, 51, 151-156.	1.7	10
78	Inheritance of deficient tocopherol accumulation in sunflower seeds. <i>Journal of Genetics</i> , 2011, 90, 489-491.	0.7	0
79	Molecular tagging and candidate gene analysis of the high gamma-tocopherol trait in safflower ( <i>Carthamus tinctorius</i> L.). <i>Molecular Breeding</i> , 2011, 28, 367-379.	2.1	16
80	Phytosterols in the seeds of wild sunflower species. <i>Helia</i> , 2011, 34, 31-38.	0.4	8
81	Update on breeding for resistance to sunflower broomrape. <i>Helia</i> , 2010, 33, 1-11.	0.4	16
82	Transferability, amplification quality, and genome specificity of microsatellites in <i>Brassica carinata</i> and related species. <i>Journal of Applied Genetics</i> , 2010, 51, 123-131.	1.9	10
83	Transferability of non-genic microsatellite and gene-based sunflower markers to safflower. <i>Euphytica</i> , 2010, 175, 145-150.	1.2	13
84	Selection for contrasting seed tocopherol content in sunflower seeds. <i>Journal of Agricultural Science</i> , 2010, 148, 393-400.	1.3	19
85	Inheritance of Low Linolenic Acid Content in Zero-Erucic Acid Ethiopian Mustard. <i>Crop Science</i> , 2009, 49, 549-553.	1.8	1
86	Current research strategies for sunflower broomrape control in Spain. <i>Helia</i> , 2009, 32, 47-55.	0.4	9
87	Inheritance of high oleic acid content in safflower. <i>Euphytica</i> , 2009, 168, 61-69.	1.2	31
88	Inheritance of very high glucosinolate content in Ethiopian mustard seeds. <i>Plant Breeding</i> , 2009, 128, 278-281.	1.9	5
89	Sunflower. , 2009, , 155-232.		24
90	Novel Safflower Germplasm with Increased Saturated Fatty Acid Content. <i>Crop Science</i> , 2009, 49, 127-132.	1.8	12

#	ARTICLE	IF	CITATIONS
91	Other Brassicas. , 2009, , 127-153.		4
92	Development of SCAR markers linked to male sterility and very high linoleic acid content in safflower. <i>Molecular Breeding</i> , 2008, 22, 385-393.	2.1	15
93	Development and characterisation of a Brassica carinata inbred line incorporating genes for low glucosinolate content from B. juncea. <i>Euphytica</i> , 2008, 164, 365-375.	1.2	16
94	Thermostability of genetically modified sunflower oils differing in fatty acid and tocopherol compositions. <i>European Journal of Lipid Science and Technology</i> , 2008, 110, 776-782.	1.5	35
95	Inheritance of very high linoleic acid content and its relationship with nuclear male sterility in safflower. <i>Plant Breeding</i> , 2008, 127, 507-509.	1.9	19
96	Novel seed oil types of Ethiopian mustard with high levels of polyunsaturated fatty acids. <i>Industrial Crops and Products</i> , 2008, 27, 359-363.	5.2	12
97	Update on breeding for resistance to sunflower broomrape. <i>Helia</i> , 2008, 31, 73-84.	0.4	32
98	A new sunflower mutant with increased levels of palmitic acid in seed oil. <i>Helia</i> , 2008, 31, 55-60.	0.4	12
99	Breeding for specialty oil types in sunflower. <i>Helia</i> , 2007, 30, 75-84.	0.4	22
100	Inheritance of resistance to broomrape ( <i>Orobanche cumana</i> Wallr.) race F in a sunflower line derived from wild sunflower species. <i>Plant Breeding</i> , 2007, 126, 67-71.	1.9	50
101	Relationships between seed oil content and fatty acid composition in high stearic acid sunflower. <i>Plant Breeding</i> , 2007, 126, 503-508.	1.9	10
102	Inheritance of Mid and High Oleic Acid Content in Ethiopian Mustard. <i>Crop Science</i> , 2006, 46, 2361-2367.	1.8	2
103	Registration of Three Sunflower Germplasms with Quantitative Resistance to Race F of Broomrape. <i>Crop Science</i> , 2006, 46, 1406-1407.	1.8	12
104	Genetic and Molecular Analysis of High Gamma-Tocopherol Content in Sunflower. <i>Crop Science</i> , 2006, 46, 2015-2021.	1.8	22
105	Inheritance of High Stearic Acid Content in the Sunflower Mutant CAS-14. <i>Crop Science</i> , 2006, 46, 22-29.	1.8	13
106	Transgressive segregation for reduced glucosinolate content in Brassica carinata A. Braun. <i>Plant Breeding</i> , 2006, 125, 400-402.	1.9	6
107	Genetic Mapping of the Tph1 Gene Controlling Beta-tocopherol Accumulation in Sunflower Seeds. <i>Molecular Breeding</i> , 2006, 17, 291-296.	2.1	25
108	Molecular analysis of the high stearic acid content in sunflower mutant CAS-14. <i>Theoretical and Applied Genetics</i> , 2006, 112, 867-875.	3.6	19

#	ARTICLE	IF	CITATIONS
109	Molecular Mapping of Nuclear Male Sterility Genes in Sunflower. <i>Crop Science</i> , 2005, 45, 1851-1857.	1.8	28
110	Identification and genetic characterization of a safflower mutant with a modified tocopherol profile. <i>Plant Breeding</i> , 2005, 124, 459-463.	1.9	33
111	Fatty acid and tocopherol accumulation in the seeds of a high oleic acid castor mutant. <i>Industrial Crops and Products</i> , 2005, 22, 201-206.	5.2	31
112	Spatial and Temporal Expression of Mutations for High Oleic Acid and Low Linolenic Acid Concentration in Ethiopian Mustard. <i>Crop Science</i> , 2005, 45, crops2005.0202.	1.8	8
113	Comparative genetic study of two sources of beta-tocopherol in sunflower. <i>Helia</i> , 2005, 28, 1-8.	0.4	11
114	Evaluation of wild sunflower species for tocopherol content and composition. <i>Helia</i> , 2004, 27, 107-112.	0.4	5
115	Use of Near-Infrared Reflectance Spectroscopy for Selecting for High Stearic Acid Concentration in Single Husked Achenes of Sunflower. <i>Crop Science</i> , 2004, 44, 93-97.	1.8	22
116	Registration of Four Sunflower Germplasms Resistant to Race F of Broomrape. <i>Crop Science</i> , 2004, 44, 1033-1034.	1.8	36
117	Registration of T589 and T2100 Sunflower Germplasms with Modified Tocopherols. <i>Crop Science</i> , 2004, 44, 362-363.	1.8	14
118	Registration of CR34 and CR81 Safflower Germplasms with Increased Tocopherol. <i>Crop Science</i> , 2004, 44, 2278-2278.	1.8	17
119	Novel variation for the tocopherol profile in a sunflower created by mutagenesis and recombination. <i>Plant Breeding</i> , 2004, 123, 490-492.	1.9	44
120	Quantitative trait loci for broomrape ( <i>Orobanche cumana</i> Wallr.) resistance in sunflower. <i>Theoretical and Applied Genetics</i> , 2004, 109, 92-102.	3.6	85
121	Allelic variation in linolenic acid content of high erucic acid Ethiopian mustard and incorporation of the low linolenic acid trait into zero erucic acid germplasm. <i>Plant Breeding</i> , 2004, 123, 137-140.	1.9	8
122	Use of Near-Infrared Reflectance Spectroscopy for Selecting for High Stearic Acid Concentration in Single Husked Achenes of Sunflower. <i>Crop Science</i> , 2004, 44, 93.	1.8	11
123	Registration of T589 and T2100 Sunflower Germplasms with Modified Tocopherols. <i>Crop Science</i> , 2004, 44, 362.	1.8	12
124	Development of sunflower germplasm with high delta-tocopherol content. <i>Helia</i> , 2004, 27, 99-106.	0.4	9
125	Dominance relationships for genes conferring resistance to broomrape ( <i>Orobanche cumana wallr</i> ) in sunflower. <i>Helia</i> , 2004, 27, 183-192.	0.4	11
126	Development of high-oleic, low-linolenic acid Ethiopian-mustard ( <i>Brassica carinata</i> ) germplasm. <i>Theoretical and Applied Genetics</i> , 2003, 107, 823-830.	3.6	25



#	ARTICLE	IF	CITATIONS
127	Inheritance of reduced plant height in the sunflower line Dw 89. <i>Plant Breeding</i> , 2003, 122, 441-443.	1.9	6
128	Registration of Dw 89 and Dw 271 Dwarf Parental Lines of Sunflower. <i>Crop Science</i> , 2003, 43, 1140-1141.	1.8	7
129	Inheritance of Increased Oleic Acid Concentration in High-Erucic Acid Ethiopian Mustard. <i>Crop Science</i> , 2003, 43, 106.	1.8	13
130	Inheritance of Increased Oleic Acid Concentration in High-Erucic Acid Ethiopian Mustard. <i>Crop Science</i> , 2003, 43, 106.	1.8	7
131	Identification and genetic characterization of new sources of beta- and gamma-tocopherol in sunflower germplasm. <i>Helia</i> , 2003, 26, 17-23.	0.4	15
132	Breeding Oilseed Crops for Improved Oil Quality. <i>The Journal of Crop Improvement: Innovations in Practiceory and Research</i> , 2002, 5, 309-344.	0.4	40
133	Genetic and environmental variation for tocopherol content and composition in sunflower commercial hybrids. <i>Journal of Agricultural Science</i> , 2002, 139, 425-429.	1.3	56
134	Inheritance of reduced linolenic acid content in the Ethiopian mustard mutant N2-4961. <i>Plant Breeding</i> , 2002, 121, 263-265.	1.9	11
135	Title is missing!. <i>Euphytica</i> , 2002, 123, 89-93.	1.2	32
136	Registration of One Low, Two Medium, and One High Erucic Acid Ethiopian Mustard Genetic Stocks. <i>Crop Science</i> , 2001, 41, 281-282.	1.8	2
137	Registration of Zero Erucic Acid Ethiopian Mustard Genetic Stock 25Xâ€¹. <i>Crop Science</i> , 2001, 41, 282-282.	1.8	13
138	Inheritance of trichome density in Ethiopian mustard leaves. <i>Euphytica</i> , 2001, 117, 241-244.	1.2	0
139	Inheritance of plant height in the dwarf mutant 'Enana' of safflower. <i>Plant Breeding</i> , 2000, 119, 525-527.	1.9	3
140	Fatty acids and tocochromanols in seeds of Orobanche. <i>Phytochemistry</i> , 2000, 54, 295-300.	2.9	37
141	Tocopherol, plastochromanol and fatty acid patterns in the genus <i>Linum</i> . <i>Plant Systematics and Evolution</i> , 2000, 221, 77-88.	0.9	46
142	Comparative studies on <i>Orobanche cernua</i> L. and <i>O. cumana</i> Wallr. (Orobanchaceae) in the Iberian Peninsula. <i>Botanical Journal of the Linnean Society</i> , 2000, 134, 513-527.	1.6	23
143	Variability for seed glucosinolates in a germplasm collection of the genus <i>Brassica</i> . , 2000, 47, 231-238.		37
144	Registration of the Dwarf Safflower Genetic Stock Enana. <i>Crop Science</i> , 2000, 40, 1207-1208.	1.8	1

#	ARTICLE	IF	CITATIONS
145	Use of near-infrared reflectance spectroscopy to assess nitrogen concentration in different plant tissues of rapeseed. <i>Communications in Soil Science and Plant Analysis</i> , 2000, 31, 2987-2995.	1.4	9
146	Nondestructive Screening for Oleic and Linoleic Acid in Single Sunflower Achenes by Near-Infrared Reflectance Spectroscopy. <i>Crop Science</i> , 1999, 39, 219-222.	1.8	34
147	Chemotaxonomic value of tocopherols in Brassicaceae. <i>Phytochemistry</i> , 1999, 50, 793-798.	2.9	55
148	Chemotaxonomic significance of fatty acids and tocopherols in Boraginaceae. <i>Phytochemistry</i> , 1999, 52, 423-426.	2.9	73
149	Tocopherol and fatty acid composition of twenty-five species of Onagraceae Juss.. <i>Botanical Journal of the Linnean Society</i> , 1999, 129, 359-366.	1.6	17
150	Title is missing!. <i>Euphytica</i> , 1999, 106, 125-130.	1.2	16
151	Title is missing!. <i>Euphytica</i> , 1999, 106, 79-85.	1.2	78
152	Development of calibration equations to predict oil content and fatty acid composition in brassicaceae germplasm by near-infrared reflectance spectroscopy. <i>JAOCS, Journal of the American Oil Chemists' Society</i> , 1999, 76, 25-30.	1.9	34
153	Characterization of yam bean ( <i>Pachyrhizus</i> spp.) Seeds as potential sources of high palmitic acid oil. <i>JAOCS, Journal of the American Oil Chemists' Society</i> , 1999, 76, 1309-1312.	1.9	26
154	Seed yield, oil and phytate concentration in the seeds of two oilseed rape cultivars as affected by different phosphorus supply. <i>European Journal of Agronomy</i> , 1999, 11, 293-299.	4.1	43
155	Quantitative determination of tocopherols in single seeds of rapeseed ( <i>Brassica napus</i> L.). <i>Lipid - Fett</i> , 1999, 101, 142-145.	0.4	34
156	Tocopherols accumulation in developing seeds and pods of rapeseed ( <i>Brassica napus</i> L.). <i>Lipid - Fett</i> , 1999, 101, 400-403.	0.4	21
157	Analysis of dry matter and protein contents in fresh yam bean tubers by near-infrared reflectance spectroscopy. <i>Communications in Soil Science and Plant Analysis</i> , 1999, 30, 1797-1805.	1.4	6
158	Tocopherol and fatty acid composition of twenty-five species of Onagraceae Juss.. <i>Botanical Journal of the Linnean Society</i> , 1999, 129, 359-366.	1.6	1
159	Title is missing!. <i>Euphytica</i> , 1998, 101, 221-230.	1.2	68
160	Title is missing!. <i>Genetic Resources and Crop Evolution</i> , 1998, 45, 371-382.	1.6	73
161	Increasing erucic acid content in Ethiopian mustard through mutation breeding. <i>Plant Breeding</i> , 1998, 117, 85-87.	1.9	22
162	Analysis of total glucosinolate content and individual glucosinolates in Brassica spp. by near-infrared reflectance spectroscopy. <i>Plant Breeding</i> , 1998, 117, 97-102.	1.9	40

#	ARTICLE	IF	CITATIONS
163	Performance of near-infrared reflectance spectroscopy (NIRS) in routine analysis of C18 unsaturated fatty acids in intact rapeseed. <i>Lipid - Fett</i> , 1998, 100, 44-48.	0.4	10
164	Application of near-infrared reflectance spectroscopy to estimate the bulk density of Ethiopian mustard seeds. , 1998, 77, 312-318.		10
165	Determination of seed oil content and fatty acid composition in sunflower through the analysis of intact seeds, husked seeds, meal and oil by near-infrared reflectance spectroscopy. <i>JAOCS, Journal of the American Oil Chemists' Society</i> , 1998, 75, 547-555.	1.9	89
166	A rapid and simple approach to identify different sunflower oil types by means of near-infrared reflectance spectroscopy. <i>JAOCS, Journal of the American Oil Chemists' Society</i> , 1998, 75, 1883-1888.	1.9	13
167	Nondestructive Assessment of Sinapic Acid Esters in Brassica Species: I. Analysis by Near Infrared Reflectance Spectroscopy. <i>Crop Science</i> , 1998, 38, 1645-1650.	1.8	13
168	Registration of Six Ethiopian Mustard Germplasm Lines. <i>Crop Science</i> , 1998, 38, 558-558.	1.8	9
169	Nondestructive Assessment of Sinapic Acid Esters in Brassica Species: II. Evaluation of Germplasm and Identification of Phenotypes with Reduced Levels. <i>Crop Science</i> , 1998, 38, 1650-1654.	1.8	29
170	Within-plant variation for seed weight and seed quality traits in white lupin ( <i>Lupinus albus</i> L.). <i>Australian Journal of Agricultural Research</i> , 1998, 49, 59.	1.5	3
171	Induced variability for C18 unsaturated fatty acids in Ethiopian mustard. <i>Canadian Journal of Plant Science</i> , 1997, 77, 91-95.	0.9	35
172	Use of Near Infrared Reflectance Spectroscopy to Screen Ethiopian Mustard for Seed Weight. <i>Agronomy Journal</i> , 1997, 89, 151-153.	1.8	5
173	Selection for reduced linolenic acid content in Ethiopian mustard ( <i>Brassica carinata</i> Braun). <i>Plant Breeding</i> , 1997, 116, 396-397.	1.9	11
174	Determination of the fatty acid composition of the oil in intact-seed mustard by near-infrared reflectance spectroscopy. <i>JAOCS, Journal of the American Oil Chemists' Society</i> , 1997, 74, 1595-1602.	1.9	43
175	Screening Ethiopian Mustard for Erucic Acid By Near Infrared Reflectance Spectroscopy. <i>Crop Science</i> , 1996, 36, 1068-1071.	1.8	16
176	An efficient method for screening seed colour in Ethiopian mustard using visible reflectance spectroscopy and multivariate analysis. <i>Euphytica</i> , 1996, 90, 359-363.	1.2	1
177	Isolation of induced mutants in Ethiopian mustard ( <i>Brassica carinata</i> Braun) with low levels of erucic acid. <i>Plant Breeding</i> , 1995, 114, 454-456.	1.9	25