## Sharon Li Ting Pek

List of Publications by Year in descending order

Source: https://exaly.com/author-pdf/979642/publications.pdf Version: 2024-02-01



#	Article	IF	CITATIONS
1	Association of Genetic Variants for Plasma LRG1 With Rapid Decline in Kidney Function in Patients With Type 2 Diabetes. Journal of Clinical Endocrinology and Metabolism, 2021, 106, 2384-2394.	3.6	9
2	Persistent hypercholesterolemia in child with homozygous autosomal recessive hypercholesterolemia: A decade of lipid management. Journal of Clinical Lipidology, 2021, 15, 441-446.	1.5	0
3	Successful pharmacological management of a child with compound heterozygous familial hypercholesterolemia and review of the recent literature. Journal of Clinical Lipidology, 2020, 14, 639-645.	1.5	1
4	A Multifunctional Role of Leucine-Rich α-2-Glycoprotein 1 in Cutaneous Wound Healing Under Normal and Diabetic Conditions. Diabetes, 2020, 69, 2467-2480.	0.6	41
5	Plasma osteoprotegerin as a biomarker of poor glycaemic control that predicts progression of albuminuria in type 2 diabetes mellitus: A 3-year longitudinal cohort study. Diabetes Research and Clinical Practice, 2020, 161, 107992.	2.8	5
6	Endothelin-1 predicts incident diabetic peripheral neuropathy in Type 2 Diabetes: a cohort study. European Journal of Endocrinology, 2020, 182, 429-438.	3.7	9
7	Spectrum of mutations in index patients with familial hypercholesterolemia in Singapore: Single center study. Atherosclerosis, 2018, 269, 106-116.	0.8	22
8	Association of circulating proinflammatory marker, leucineâ€richâ€Î±2â€glycoprotein (LRG1), following metabolic/bariatric surgery. Diabetes/Metabolism Research and Reviews, 2018, 34, e3029.	4.0	25
9	Association of the anti-angiogenic factor secreted protein and rich in cysteine (SPARC) with vascular complications among Chinese type 2 diabetic patients in Singapore. Journal of Diabetes and Its Complications, 2017, 31, 1222-1227.	2.3	2
10	Association of apolipoprotein-CIII (apoC-III), endothelium-dependent vasodilation and peripheral neuropathy in a multi-ethnic population with type 2 diabetes. Metabolism: Clinical and Experimental, 2017, 72, 75-82.	3.4	9
11	Plasma Leucine-Rich α-2-Glycoprotein 1 Predicts Rapid eGFR Decline and Albuminuria Progression in Type 2 Diabetes Mellitus. Journal of Clinical Endocrinology and Metabolism, 2017, 102, 3683-3691.	3.6	43
12	Circulating and visceral adipose miR-100 is down-regulated in patients with obesity and Type 2 diabetes. Molecular and Cellular Endocrinology, 2016, 427, 112-123.	3.2	47