## Dirk Eick

## List of Publications by Year in descending order

Source: https://exaly.com/author-pdf/996536/publications.pdf

Version: 2024-02-01

53 papers

5,875 citations

36 h-index 53 g-index

55 all docs

55 docs citations

55 times ranked 7596 citing authors

#	Article	IF	Citations
1	Extension of the minimal functional unit of the RNA polymerase II CTD from yeast to mammalian cells. Biology Letters, 2019, 15, 20190068.	2.3	1
2	MIR sequences recruit zinc finger protein ZNF768 to expressed genes. Nucleic Acids Research, 2019, 47, 700-715.	14.5	14
3	Arginine Citrullination at the C-Terminal Domain Controls RNA Polymerase II Transcription. Molecular Cell, 2019, 73, 84-96.e7.	9.7	50
4	Tyrosine-1 of RNA Polymerase II CTD Controls Global Termination of Gene Transcription in Mammals. Molecular Cell, 2018, 69, 48-61.e6.	9.7	66
5	Getting to grips with c-Myc. ELife, 2018, 7, .	6.0	1
6	Different phosphoisoforms of RNA polymerase II engage the Rtt103 termination factor in a structurally analogous manner. Proceedings of the National Academy of Sciences of the United States of America, 2017, 114, E3944-E3953.	7.1	24
7	Transcriptome analysis of dominant-negative Brd4 mutants identifies Brd4-specific target genes of small molecule inhibitor JQ1. Scientific Reports, 2017, 7, 1684.	3.3	17
8	CDK9-dependent RNA polymerase II pausing controls transcription initiation. ELife, 2017, 6, .	6.0	179
9	A Nonradioactive Assay to Measure Production and Processing of Ribosomal RNA by 4sU-Tagging. Methods in Molecular Biology, 2016, 1455, 121-131.	0.9	2
10	Specific threonine-4 phosphorylation and function of RNA polymerase II CTD during M phase progression. Scientific Reports, 2016, 6, 27401.	3.3	17
11	Getting Access to Low-Complexity Domain Modifications. Trends in Biochemical Sciences, 2016, 41, 894-897.	7.5	11
12	Heptad-Specific Phosphorylation of RNA PolymeraseÂII CTD. Molecular Cell, 2016, 61, 305-314.	9.7	118
13	DEAD-box helicase DDX27 regulates 3′ end formation of ribosomal 47S RNA and stably associates with the PeBoW-complex. Experimental Cell Research, 2015, 334, 146-159.	2.6	26
14	Site-specific methylation and acetylation of lysine residues in the C-terminal domain (CTD) of RNA polymerase II. Transcription, 2015, 6, 91-101.	3.1	22
15	The structure and substrate specificity of human Cdk12/Cyclin K. Nature Communications, 2014, 5, 3505.	12.8	141
16	Tyrosine phosphorylation of RNA polymerase II CTD is associated with antisense promoter transcription and active enhancers in mammalian cells. ELife, 2014, 3, e02105.	6.0	76
17	The RNA Polymerase II Carboxy-Terminal Domain (CTD) Code. Chemical Reviews, 2013, 113, 8456-8490.	47.7	368
18	Dynamic phosphorylation patterns of RNA polymerase II CTD during transcription. Biochimica Et Biophysica Acta - Gene Regulatory Mechanisms, 2013, 1829, 55-62.	1.9	225

#	Article	IF	Citations
19	4-thiouridine inhibits rRNA synthesis and causes a nucleolar stress response. RNA Biology, 2013, 10, 1623-1630.	3.1	117
20	Cyclin-dependent Kinase 9 Links RNA Polymerase II Transcription to Processing of Ribosomal RNA. Journal of Biological Chemistry, 2013, 288, 21173-21183.	3.4	27
21	Dose-dependent regulation of target gene expression and cell proliferation by c-Myc levels. Transcription, 2013, 4, 192-197.	3.1	31
22	CTD serine-2 plays a critical role in splicing and termination factor recruitment to RNA polymerase II in vivo. Nucleic Acids Research, 2013, 41, 1591-1603.	14.5	87
23	Functional ribosome biogenesis is a prerequisite for p53 destabilization: impact of chemotherapy on nucleolar functions and RNA metabolism. Biological Chemistry, 2013, 394, 1133-1143.	2.5	27
24	Tyrosine-1 and threonine-4 phosphorylation marks complete the RNA polymerase II CTD phospho-code. RNA Biology, 2012, 9, 1144-1146.	3.1	16
25	CTD Tyrosine Phosphorylation Impairs Termination Factor Recruitment to RNA Polymerase II. Science, 2012, 336, 1723-1725.	12.6	215
26	Threonine-4 of mammalian RNA polymerase II CTD is targeted by Polo-like kinase 3 and required for transcriptional elongation. EMBO Journal, 2012, 31, 2784-2797.	7.8	123
27	Ultrashort and progressive 4sU-tagging reveals key characteristics of RNA processing at nucleotide resolution. Genome Research, 2012, 22, 2031-2042.	5.5	132
28	Splicing enhances recruitment of methyltransferase HYPB/Setd2 and methylation of histone H3 Lys36. Nature Structural and Molecular Biology, 2011, 18, 977-983.	8.2	204
29	The C-Terminal Domain of RNA Polymerase II Is Modified by Site-Specific Methylation. Science, 2011, 332, 99-103.	12.6	190
30	Transcription initiation platforms and GTF recruitment at tissue-specific enhancers and promoters. Nature Structural and Molecular Biology, 2011, 18, 956-963.	8.2	296
31	Chemotherapeutic Drugs Inhibit Ribosome Biogenesis at Various Levels. Journal of Biological Chemistry, 2010, 285, 12416-12425.	3.4	356
32	RNA Polymerase II C-terminal Heptarepeat Domain Ser-7 Phosphorylation Is Established in a Mediator-dependent Fashion. Journal of Biological Chemistry, 2010, 285, 188-196.	3.4	49
33	The tumor suppressor p53 connects ribosome biogenesis to cell cycle control: a double-edged sword. Oncotarget, 2010, 1, 43-47.	1.8	21
34	CDK9 directs H2B monoubiquitination and controls replicationâ€dependent histone mRNA 3′â€end processing. EMBO Reports, 2009, 10, 894-900.	4.5	142
35	TFIIH Kinase Places Bivalent Marks on the Carboxy-Terminal Domain of RNA Polymerase II. Molecular Cell, 2009, 34, 387-393.	9.7	235
36	The nucleolar SUMOâ€specific protease SENP3 reverses SUMO modification of nucleophosmin and is required for rRNA processing. EMBO Reports, 2008, 9, 273-279.	4.5	141

#	Article	IF	CITATIONS
37	Molecular evolution of the RNA polymerase II CTD. Trends in Genetics, 2008, 24, 289-296.	6.7	129
38	The CALM and CALM/AF10 interactor CATS is a marker for proliferation. Molecular Oncology, 2008, 2, 356-367.	4.6	36
39	The BRCT domain of mammalian Pes1 is crucial for nucleolar localization and rRNA processing. Nucleic Acids Research, 2007, 35, 789-800.	14.5	41
40	Rapid conditional knock-down–knock-in system for mammalian cells. Nucleic Acids Research, 2007, 35, e17-e17.	14.5	12
41	Interdependence of Pes1, Bop1, and WDR12 Controls Nucleolar Localization and Assembly of the PeBoW Complex Required for Maturation of the 60S Ribosomal Subunit. Molecular and Cellular Biology, 2007, 27, 3682-3694.	2.3	116
42	H2B Ubiquitylation Acts as a Barrier to Ctk1 Nucleosomal Recruitment Prior to Removal by Ubp8 within a SAGA-Related Complex. Molecular Cell, 2007, 27, 275-288.	9.7	196
43	Serine-7 of the RNA Polymerase II CTD Is Specifically Required for snRNA Gene Expression. Science, 2007, 318, 1777-1779.	12.6	221
44	Transcribing RNA Polymerase II Is Phosphorylated at CTD Residue Serine-7. Science, 2007, 318, 1780-1782.	12.6	258
45	Dominant-negative Pes1 mutants inhibit ribosomal RNA processing and cell proliferation via incorporation into the PeBoW-complex. Nucleic Acids Research, 2006, 34, 3030-3043.	14.5	79
46	Dissection of transcriptional programmes in response to serum and c-Myc in a human B-cell line. Oncogene, 2005, 24, 520-524.	5.9	68
47	Mammalian WDR12 is a novel member of the Pes1–Bop1 complex and is required for ribosome biogenesis and cell proliferation. Journal of Cell Biology, 2005, 170, 367-378.	5.2	166
48	Role of the Mammalian RNA Polymerase II C-Terminal Domain (CTD) Nonconsensus Repeats in CTD Stability and Cell Proliferation. Molecular and Cellular Biology, 2005, 25, 7665-7674.	2.3	49
49	The last CTD repeat of the mammalian RNA polymerase II large subunit is important for its stability. Nucleic Acids Research, 2004, 32, 35-44.	14.5	58
50	A role for c-Myc in the regulation of ribosomal RNA processing. Nucleic Acids Research, 2003, 31, 6148-6156.	14.5	160
51	Cell cycle activation by c-myc in a Burkitt lymphoma model cell line. International Journal of Cancer, 2000, 87, 787-793.	5.1	178
52	Conditional Expression of RNA Polymerase II in Mammalian Cells. Journal of Biological Chemistry, 2000, 275, 24375-24382.	3 <b>.</b> 4	74
53	Control of cell growth by c-Myc in the absence of cell division. Current Biology, 1999, 9, 1255-1258.	3.9	267