
Eduardo Carasek da Rocha

List of Publications by Year in descending order

Source: https://exaly.com/author-pdf/999981/publications.pdf Version: 2024-02-01



#	Article	IF	CITATIONS
1	Ultrasound-assisted and disposable pipette extraction for the determination of faecal contaminants in sediment samples by GC-MS. International Journal of Environmental Analytical Chemistry, 2022, 102, 4288-4299.	1.8	2
2	Application of Homogeneous Liquid–Liquid Microextraction With Switchable Hydrophilicity Solvents to the Determination of MDMA, MDA and NBOMes in Postmortem Blood Samples. Journal of Analytical Toxicology, 2022, 46, 776-782.	1.7	8
3	A semi-quantitative model through PLS-DA in the evaluation of carbendazim in grape juices. Food Chemistry, 2022, 368, 130742.	4.2	11
4	Determination of five 3-alkyl-2-methoxypyrazines employing HS-SPME-GC-NPD: Application in evaluation of off-flavor of South American wines. Journal of Food Composition and Analysis, 2022, 105, 104237.	1.9	6
5	Disposable pipette extraction: A critical review of concepts, applications, and directions. Analytica Chimica Acta, 2022, 1192, 339383.	2.6	27
6	Rotating sorptive extraction (RSE) as a green approach for determining pesticides from fruit and vegetable wash-water using GC–MS. Advances in Sample Preparation, 2022, 1, 100003.	1.1	2
7	Multiclass determination of endocrine disruptors in urine by hollow fiber microporous membrane and liquid chromatography. Analytical Biochemistry, 2022, 652, 114725.	1.1	2
8	Novel strategy for disposable pipette extraction (DPX): Low-cost Parallel-DPX for determination of phthalate migration from common plastic materials to saliva simulant with GC-MS. Talanta, 2021, 221, 121443.	2.9	10
9	Determination of bisphenol A: Old problem, recent creative solutions based on novel materials. Journal of Separation Science, 2021, 44, 1148-1173.	1.3	13
10	High-throughput approach for the in situ generation of magnetic ionic liquids in parallel-dispersive droplet extraction of organic micropollutants in aqueous environmental samples. Talanta, 2021, 223, 121759.	2.9	19
11	Exploring the Use of Switchable Hydrophilicity Solvents as Extraction Phase for the Determination of Food-Packaging Contaminants in Coconut Water Samples by Gas Chromatography-Mass Spectrometry. Food Analytical Methods, 2021, 14, 319-330.	1.3	12
12	A green approach to DDT degradation and metabolite monitoring in water comparing the hydrodechlorination efficiency of Pd, Au-on-Pd and Cu-on-Pd nanoparticle catalysis. Science of the Total Environment, 2021, 760, 143403.	3.9	4
13	Photocatalytic Cellulose-Paper: Deepening in the Sustainable and Synergic Combination of Sorption and Photodegradation. ACS Omega, 2021, 6, 9577-9586.	1.6	11
14	Sustainable green solvents for microextraction techniques: Recent developments and applications. Journal of Chromatography A, 2021, 1640, 461944.	1.8	46
15	A proofâ€ofâ€concept of parallel singleâ€drop microextraction for the rapid and sensitive biomonitoring of pesticides in urine. Journal of Separation Science, 2021, 44, 1961-1968.	1.3	17
16	Exploring the use of cork pellets in bar adsorptive microextraction for the determination of organochloride pesticides in water samples with gas chromatography/electron capture detection quantification. Journal of Chromatography A, 2021, 1645, 462099.	1.8	12
17	The use of microwave-induced plasma optical emission spectrometry for fluorine determination and its application to tea infusions. Talanta, 2021, 227, 122190.	2.9	7
18	Assessment of a natural extraction phase in disposable pipette extraction coupled with the sub-minute determination of 11-nor-1"9-tetrahydrocannabinol-9-carboxylic acid in human urine by fast-GC-FID. Sustainable Chemistry and Pharmacy, 2021, 20, 100390.	1.6	4

#	Article	IF	CITATIONS
19	Adsorption of hazardous and noxious 4-nitrophenol by a silsesquioxane organic-inorganic hybrid material. Journal of Sol-Gel Science and Technology, 2021, 99, 402-412.	1.1	7
20	A Rapid Analytical Approach for Monitoring Pharmaceuticals in Hospital Wastewater—A DPX-Based Procedure with Environmentally-Friendly Extraction Phase Coupled to High Performance Liquid Chromatography–Diode Array/Fluorescence Detectors. Separations, 2021, 8, 109.	1.1	1
21	Ionic liquids. , 2021, , 427-451.		5
22	A low-voltage paper spray ionization QTOF-MS method for the qualitative analysis of NPS in street drug blotter samples. Forensic Toxicology, 2020, 38, 227-231.	1.4	13
23	A high throughput approach to rotating-disk sorptive extraction (RDSE) using laminar cork for the simultaneous determination of multiclass organic micro-pollutants in aqueous sample by GC-MS. Talanta, 2020, 208, 120459.	2.9	21
24	Determination of pesticides of different chemical classes in drinking water of the state of Santa Catarina (Brazil) using solid-phase microextraction coupled to chromatographic determinations. Environmental Science and Pollution Research, 2020, 27, 43870-43883.	2.7	5
25	Estrogens determination through disposable pipette extraction coupled to ultraviolet spectroscopy and nonlinear pseudoâ€univariate calibration: Solving rank deficiency problems. Journal of Chemometrics, 2020, 34, e3276.	0.7	1
26	Emerging micropollutants determination by NIR spectroscopy using pseudo-univariate calibration and TF-SPME coupled with 96-well plate system. Microchemical Journal, 2020, 155, 104789.	2.3	4
27	A green and low-cost method employing switchable hydrophilicity solvent for the simultaneous determination of antidepressants in human urine by gas chromatography - mass spectrometry detection. Journal of Chromatography B: Analytical Technologies in the Biomedical and Life Sciences, 2020, 1143, 122069.	1.2	26
28	Application of bar adsorptive microextraction (BAµE) for the determination of pesticides and emerging contaminants in water used for rice cultivation in southern Brazil. SN Applied Sciences, 2020, 2, 1.	1.5	2
29	Designing a green device to BAμE: Recycled cork pellet as extraction phase for the determination of parabens in river water samples. Talanta, 2020, 219, 121369.	2.9	18
30	A green - high throughput –extraction method based on hydrophobic natural deep eutectic solvent for the determination of emerging contaminants in water by high performance liquid chromatography – diode array detection. Journal of Chromatography A, 2020, 1626, 461377.	1.8	23
31	Returning to Nature for the Design of Sorptive Phases in Solid-Phase Microextraction. Separations, 2020, 7, 2.	1.1	39
32	Simple and straightforward analysis of cannabinoids in medicinal products by fast-GC–FID. Forensic Toxicology, 2020, 38, 531-535.	1.4	9
33	Expanding the applicability of magnetic ionic liquids for multiclass determination in biological matrices based on dispersive liquid–liquid microextraction and HPLC with diode array detector analysis. Journal of Separation Science, 2020, 43, 2657-2665.	1.3	17
34	A straightforward and semiautomated membrane-based method as efficient tool for the determination of cocaine and its metabolites in urine samples using liquid chromatography coupled to quadrupole time-of-flight-mass spectrometry. Journal of Chromatography A, 2020, 1621, 461088.	1.8	10
35	Alternative Green Extraction Phases Applied to Microextraction Techniques for Organic Compound Determination. Separations, 2019, 6, 35.	1.1	20
36	Cork sheet as a sorptive phase to extract hormones from water by rotating-disk sorptive extraction (RDSE). Analytica Chimica Acta, 2019, 1087, 1-10.	2.6	30

#	Article	IF	CITATIONS
37	The use of Ca + Pd + Zr as modifiers in the determination of sulfur by HR-CS GF MAS with solid sampling. Journal of Analytical Atomic Spectrometry, 2019, 34, 498-503.	1.6	5
38	lodine determination by high-resolution continuum source molecular absorption spectrometry – A comparison between potential molecules. Spectrochimica Acta, Part B: Atomic Spectroscopy, 2019, 153, 42-49.	1.5	4
39	Histamine functionalized magnetic nanoparticles (HIS-MNP) as a sorbent for thin film microextraction of endocrine disrupting compounds in aqueous samples and determination by high performance liquid chromatography-fluorescence detection. Journal of Chromatography A, 2019, 1602, 41-47.	1.8	17
40	A rapid and environmentally friendly analytical method based on conductive polymer as extraction phase for disposable pipette extraction for the determination of hormones and polycyclic aromatic hydrocarbons in river water samples using high-performance liquid chromatography/diode array detection. Journal of Environmental Chemical Engineering, 2019, 7, 103156.	3.3	15
41	Bract as a novel extraction phase in thin-film SPME combined with 96-well plate system for the high-throughput determination of estrogens in human urine by liquid chromatography coupled to fluorescence detection. Journal of Chromatography B: Analytical Technologies in the Biomedical and Life Sciences. 2019. 1118-1119. 17-24.	1.2	27
42	Magnetic ionic liquids as an efficient tool for the multiresidue screening of organic contaminants in river water samples. Separation Science Plus, 2019, 2, 51-58.	0.3	12
43	Effective and High-Throughput Analytical Methodology for the Determination of Lead and Cadmium in Water Samples by Disposable Pipette Extraction Coupled with High-Resolution Continuum Source Graphite Furnace Atomic Absorption Spectrometry (HR-CS GF AAS). Analytical Letters, 2019, 52, 2133-2149.	1.0	19
44	A green and simple sample preparation method to determine pesticides in rice using a combination of SPME and rotating disk sorption devices. Analytica Chimica Acta, 2019, 1069, 57-65.	2.6	25
45	A natural and renewable biosorbent phase as a lowâ€cost approach in disposable pipette extraction technique for the determination of emerging contaminants in lake water samples. Journal of Separation Science, 2019, 42, 1404-1411.	1.3	20
46	Single drop microextraction in a 96-well plate format: A step toward automated and high-throughput analysis. Analytica Chimica Acta, 2019, 1063, 159-166.	2.6	67
47	Magnetic ionic liquids as versatile extraction phases for the rapid determination of estrogens in human urine by dispersive liquid-liquid microextraction coupled with high-performance liquid chromatography-diode array detection. Analytical and Bioanalytical Chemistry, 2018, 410, 4689-4699.	1.9	58
48	Low-cost approach to increase the analysis throughput of bar adsorptive microextraction (BAµE) combined with environmentally-friendly renewable sorbent phase of recycled diatomaceous earth. Talanta, 2018, 178, 886-893.	2.9	22
49	Development of a highâ€throughput method based on thinâ€film microextraction using a 96â€well plate system with a cork coating for the extraction of emerging contaminants in river water samples. Journal of Separation Science, 2018, 41, 697-703.	1.3	24
50	Use of a Natural Sorbent as Alternative Solid-Phase Microextraction Coating for the Determination of Polycyclic Aromatic Hydrocarbons in Water Samples by Gas Chromatography-Mass Spectrometry. Journal of the Brazilian Chemical Society, 2018, , .	0.6	3
51	A Low-Cost Approach Using Diatomaceous Earth Biosorbent as Alternative SPME Coating for the Determination of PAHs in Water Samples by GC-MS. Separations, 2018, 5, 55.	1.1	9
52	Improvement of dispersive liquidâ€liquid microextraction robustness by performing consecutive extractions: Determination of polycyclic aromatic hydrocarbons in Brazilian sugar cane spirits by GCâ€MS. Separation Science Plus, 2018, 1, 564-573.	0.3	3
53	Expanding the applicability of cork as extraction phase for disposable pipette extraction in multiresidue analysis of pharmaceuticals in urine samples. Journal of Chromatography B: Analytical Technologies in the Biomedical and Life Sciences, 2018, 1102-1103, 159-166.	1.2	36
54	Hollow-fiber renewal liquid membrane extraction coupled with 96-well plate system as innovative high-throughput configuration forÂthe determination of endocrine disrupting compounds by high-performance liquid chromatography-fluorescence and diode array detection. Analytica Chimica Acta, 2018, 1040, 33-40.	2.6	23

#	Article	IF	CITATIONS
55	A recent overview of the application of liquid-phase microextraction to the determination of organic micro-pollutants. TrAC - Trends in Analytical Chemistry, 2018, 108, 203-209.	5.8	55
56	Exploiting green sorbents in rotatingâ€disk sorptive extraction for the determination of parabens by highâ€performance liquid chromatography with tandem electrospray ionization triple quadrupole mass spectrometry. Journal of Separation Science, 2018, 41, 4047-4054.	1.3	21
57	Median preoptic nucleus excitatory neurotransmitters in the maintenance of hypertensive state. Brain Research Bulletin, 2018, 142, 207-215.	1.4	5
58	An effective and high-throughput analytical methodology for pesticide screening in human urine by disposable pipette extraction and gas chromatography – mass spectrometry. Journal of Chromatography B: Analytical Technologies in the Biomedical and Life Sciences, 2018, 1092, 459-465.	1.2	33
59	Development of an eco-friendly method for the determination of total polychlorinated biphenyls in solid waste by gas chromatography-mass spectrometry. Journal of Environmental Chemical Engineering, 2018, 6, 5042-5048.	3.3	7
60	Basic principles, recent trends and future directions of microextraction techniques for the analysis of aqueous environmental samples. Trends in Environmental Analytical Chemistry, 2018, 19, e00060.	5.3	53
61	Determination of arsenic in agricultural soil samples using High-resolution continuum source graphite furnace atomic absorption spectrometry and direct solid sample analysis. Talanta, 2018, 188, 722-728.	2.9	37
62	A new configuration for bar adsorptive microextraction (BAμE) for the quantification of biomarkers (hexanal and heptanal) in human urine by HPLC providing an alternative for early lung cancer diagnosis. Analytica Chimica Acta, 2017, 965, 54-62.	2.6	35
63	Investigation of chemical modifiers for the determination of cadmium and chromium in fish oil and lipoid matrices using HR-CS GF AAS and a simple â€~dilute-and-shoot' approach. Microchemical Journal, 2017, 133, 175-181.	2.3	19
64	A low-cost biosorbent-based coating for the highly sensitive determination of organochlorine pesticides by solid-phase microextraction and gas chromatography-electron capture detection. Journal of Chromatography A, 2017, 1525, 23-31.	1.8	34
65	Novel approach to high-throughput determination of endocrine disruptors using recycled diatomaceous earth as a green sorbent phase for thin-film solid-phase microextraction combined with 96-well plate system. Analytica Chimica Acta, 2017, 996, 29-37.	2.6	50
66	A novel extraction-based procedure for the determination of cadmium in marine macro-algae using HR-CS GF AAS. Analytical Methods, 2017, 9, 5400-5406.	1.3	3
67	Evaluation of two membrane-based microextraction techniques for the determination of endocrine disruptors in aqueous samples by HPLC with diode array detection. Journal of Separation Science, 2017, 40, 4431-4438.	1.3	10
68	A hybrid material as a sorbent phase for the disposable pipette extraction technique enhances efficiency in the determination of phenolic endocrine-disrupting compounds. Journal of Chromatography A, 2017, 1513, 42-50.	1.8	33
69	Determination of emerging contaminants in aqueous matrices with hollow fiber-supported dispersive liquid-liquid microextraction (HF-DLLME) and separation/detection by liquid chromatography – Diode array detection. Microchemical Journal, 2017, 130, 371-376.	2.3	21
70	Evaluation of volatile profiles obtained for minimally-processed pineapple fruit samples during storage by headspace-solid phase microextraction gas chromatography-mass spectrometry. Food Science and Technology, 2017, 37, 663-672.	0.8	9
71	Exploiting Cork as Biosorbent Extraction Phase for Solid-Phase Microextraction to Determine 3-(4-Methylbenzylidene)camphor and 2-Ethylhexyl 4-(Dimethylamino)benzoate in River Water by Gas Chromatography-Mass Spectrometry. Journal of the Brazilian Chemical Society, 2017, , .	0.6	2
72	Application of a robust solid-phase microextraction fiber consisting of NiTi wires coated with polypyrrole for the determination of haloanisoles in water and wine. Analytical Methods, 2016, 8, 5503-5510.	1.3	6

#	Article	IF	CITATIONS
73	Determination of compounds with varied volatilities from aqueous samples using a polymeric ionic liquid sorbent coating by direct immersion-headspace solid-phase microextraction. Analytical Methods, 2016, 8, 4108-4118.	1.3	16
74	Arsenic containing medium and long chain fatty acids in marine fish oil identified as degradation products using reversed-phase HPLC-ICP-MS/ESI-MS. Journal of Analytical Atomic Spectrometry, 2016, 31, 1836-1845.	1.6	27
75	Coupling solid phase microextraction to complementary separation platforms for metabotyping of E. coli metabolome in response to natural antibacterial agents. Metabolomics, 2016, 12, 1.	1.4	20
76	Investigation of chemical modifiers for the direct determination of arsenic in fish oil using high-resolution continuum source graphite furnace atomic absorption spectrometry. Talanta, 2016, 150, 142-147.	2.9	24
77	A simple sample preparation procedure for the fast screening of selenium species in soil samples using alkaline extraction and hydride-generation graphite furnace atomic absorption spectrometry. Microchemical Journal, 2016, 125, 50-55.	2.3	21
78	Novel analytical procedure using a combination of hollow fiber supported liquid membrane and dispersive liquid–liquid microextraction for the determination of aflatoxins in soybean juice by high performance liquid chromatography – Fluorescence detector. Food Chemistry, 2016, 196, 292-300.	4.2	35
79	Determination of sulfur in crude oil using high-resolution continuum source molecular absorption spectrometry of the SnS molecule in a graphite furnace. Talanta, 2016, 146, 203-208.	2.9	24
80	Combination of hollow-fiber-supported liquid membrane and dispersive liquid-liquid microextraction as a fast and sensitive technique for the extraction of pesticides from grape juice followed by high-performance liquid chromatography. Journal of Separation Science, 2015, 38, 1959-1968.	1.3	19
81	A novel approach to bar adsorptive microextraction: Cork as extractor phase for determination of benzophenone, triclocarban and parabens in aqueous samples. Analytica Chimica Acta, 2015, 888, 59-66.	2.6	52
82	Fluorine determination in coal using high-resolution graphite furnace molecular absorption spectrometry and direct solid sample analysis. Spectrochimica Acta, Part B: Atomic Spectroscopy, 2015, 105, 18-24.	1.5	22
83	Application of disposable starch-based platforms for sample introduction and determination of refractory elements using graphite furnace atomic absorption spectrometry and direct solid sample analysis. Journal of Analytical Atomic Spectrometry, 2015, 30, 381-388.	1.6	9
84	Membrane-based microextraction techniques in analytical chemistry: A review. Analytica Chimica Acta, 2015, 880, 8-25.	2.6	134
85	Frog Volatile Compounds: Application of in vivo SPME for the Characterization of the Odorous Secretions from Two Species of Hypsiboas Treefrogs. Journal of Chemical Ecology, 2015, 41, 360-372.	0.9	22
86	Determination of chlorine in coal via the SrCl molecule using high-resolution graphite furnace molecular absorption spectrometry and direct solid sample analysis. Spectrochimica Acta, Part B: Atomic Spectroscopy, 2015, 114, 46-50.	1.5	22
87	Determination of aluminum in moisturizing body lotions using graphite furnace atomic absorption spectrometry. Analytical Methods, 2015, 7, 9636-9640.	1.3	9
88	Use of green coating (cork) in solid-phase microextraction for the determination of organochlorine pesticides in water by gas chromatography-electron capture detection. Talanta, 2015, 134, 409-414.	2.9	55
89	Simultaneous Determination of Environmental Contaminants with Different Volatilities in Tap Water Samples Using a New Approach to Single-Drop Microextraction Procedure. Journal of the Brazilian Chemical Society, 2015, , .	0.6	0
90	Determination of fluorine in plant materials via calcium mono-fluoride using high-resolution graphite furnace molecular absorption spectrometry with direct solid sample introduction. Journal of Analytical Atomic Spectrometry, 2014, 29, 1564-1569.	1.6	31

#	Article	IF	CITATIONS
91	Screening of volatile compounds in honey using a new sampling strategy combining multiple extraction temperatures in a single assay by HS-SPME–GC–MS. Food Chemistry, 2014, 145, 1061-1065.	4.2	37
92	Use of Doehlert design in the optimization of extraction conditions in the determination of organochlorine pesticides in bovine milk samples by HS-SPME. Analytical Methods, 2014, 6, 3254-3260.	1.3	20
93	Strontium mono-chloride — A new molecule for the determination of chlorine using high-resolution graphite furnace molecular absorption spectrometry and direct solid sample analysis. Spectrochimica Acta, Part B: Atomic Spectroscopy, 2014, 102, 1-6.	1.5	26
94	Method development for the determination of bromine in coal using high-resolution continuum source graphite furnace molecular absorption spectrometry and direct solid sample analysis. Spectrochimica Acta, Part B: Atomic Spectroscopy, 2014, 96, 33-39.	1.5	23
95	Determination of selenium in soil samples using high-resolution continuum source graphite furnace atomic absorption spectrometry and direct solid sample analysis. Analytical Methods, 2014, 6, 2870-2875.	1.3	18
96	DESENVOLVIMENTO DE METODOLOGIA EM SISTEMA EM FLUXO PARA DETERMINAÇÃO DE CD USANDO ERVA MATE E CHÕPRETO COMO ADSORVENTE E ESPECTROMETRIA DE ABSORÇÃO ATà "MICA EM CHAMA. Ecletica Quimica, 2014, 39, 68.	0.2	0
97	Determination of volatile profile of citrus fruit by HS-SPME/GC-MS with oxidized NiTi fibers using two temperatures in the same extraction procedure. Microchemical Journal, 2013, 109, 128-133.	2.3	45
98	Asymmetric reduction of (4S)-(+)-carvone catalyzed by baker's yeast: A green method for monitoring the conversion based on liquid–liquid–liquid microextraction with polypropylene hollow fiber membranes. Process Biochemistry, 2013, 48, 1159-1165.	1.8	15
99	Simultaneous determination of trihalomethanes and organochlorine pesticides in water samples by direct immersion-headspace-solid phase microextraction. Journal of Chromatography A, 2013, 1321, 30-37.	1.8	33
100	Detection of extraction artifacts in the analysis of honey volatiles using comprehensive two-dimensional gas chromatography. Food Chemistry, 2013, 141, 1828-1833.	4.2	35
101	Determination of sulfur in coal using direct solid sampling and high-resolution continuum source molecular absorption spectrometry of the CS molecule in a graphite furnace. Talanta, 2013, 106, 368-374.	2.9	39
102	Use of two different coating temperatures for a cold fiber headspace solidâ€phase microextraction system to determine the volatile profile of Brazilian medicinal herbs. Journal of Separation Science, 2013, 36, 1410-1417.	1.3	22
103	Cork as a new (green) coating for solid-phase microextraction: Determination of polycyclic aromatic hydrocarbons in water samples by gas chromatography–mass spectrometry. Analytica Chimica Acta, 2013, 772, 33-39.	2.6	75
104	Effect of magnesium acetylacetonate on the signal of organic forms of vanadium in graphite furnace atomic absorption spectrometry. Talanta, 2013, 103, 66-74.	2.9	4
105	Development of a simple analytical method for determining trihalomethanes in beer using a headspace solid-phase microextraction technique. Quimica Nova, 2013, 36, 1052-1056.	0.3	6
106	Desenvolvimento de um método analÃŧico baseado em microextração lÃquido-lÃquido para a determinação de cromo (VI) em amostras aquosas com detecção por espectrometria de absorção atÃ′mica em chama. Quimica Nova, 2013, 36, 942-946.	0.3	1
107	Microextração em fase lÃquida suportada com fibra oca (HF-LPME): Fundamentos e aplicações recentes. Scientia Chromatographica, 2013, 5, 249-262.	0.2	4
108	Pendimethalin in surface waters of rivers in the proximity of irrigated paddy fields by solid phase microextraction and gas chromatography. International Journal of Environmental Analytical Chemistry, 2012, 92, 313-323.	1.8	8

#	Article	IF	CITATIONS
109	Multiresidue determination of pesticides in industrial and fresh orange juice by hollow fiber microporous membrane liquid–liquid extraction and detection by liquid chromatography–electrospray-tandem mass spectrometry. Talanta, 2012, 88, 573-580.	2.9	64
110	Comparison of three different sample preparation procedures for the determination of traffic-related elements in airborne particulate matter collected on glass fiber filters. Talanta, 2012, 88, 689-695.	2.9	30
111	Evaluation of a completely automated cold fiber device using compounds with varying volatility and polarity. Analytica Chimica Acta, 2012, 742, 22-29.	2.6	24
112	Determination of Zn(II) in alcohol fuel by flame atomic absorption spectrometry after on-line preconcentration using a solid phase extraction system. Journal of Analytical Chemistry, 2012, 67, 448-454.	0.4	10
113	Simultaneous determination of polycyclic aromatic hydrocarbons and benzene, toluene, ethylbenzene and xylene in water samples using a new sampling strategy combining different extraction modes and temperatures in a single extraction solid-phase microextraction-gas chromatography–mass spectrometry procedure, lournal of Chromatography A. 2012. 1233. 22-29.	1.8	71
114	Application of solidâ€phase microextraction and gas chromatographyâ€mass spectrometry for the determination of chlorophenols in leather. Journal of Separation Science, 2012, 35, 602-607.	1.3	12
115	Extraction and on-fiber derivatization of chlorophenols in leather by internally cooled solid phase microextraction. Journal of the Brazilian Chemical Society, 2012, 23, 2232-2236.	0.6	3
116	DETERMINATION OF BENZENE IN BEVERAGES BY SOLID-PHASE MICROEXTRACTION AND GAS CHROMATOGRAPHY. Scientia Chromatographica, 2012, 4, 209-216.	0.2	2
117	Hollow-fiber liquid–liquid–solid micro-extraction of lead in soft drinks and determination by graphite furnace atomic absorption spectrometry. Talanta, 2011, 84, 989-994.	2.9	25
118	Method development and optimization for the determination of selenium in bean and soil samples using hydride generation electrothermal atomic absorption spectrometry. Talanta, 2011, 85, 1350-1356.	2.9	42
119	Determination of fluorine in tea using high-resolution molecular absorption spectrometry with electrothermal vaporization of the calcium mono-fluoride CaF. Talanta, 2011, 85, 2681-2685.	2.9	77
120	A fast and accurate method for the determination of total and soluble fluorine in toothpaste using high-resolution graphite furnace molecular absorption spectrometry and its comparison with established techniques. Journal of Pharmaceutical and Biomedical Analysis, 2011, 54, 1040-1046.	1.4	61
121	A new optimization strategy for gaseous phase sampling by an internally cooled solid-phase microextraction technique. Journal of Chromatography A, 2011, 1218, 367-372.	1.8	8
122	A new approach based on a combination of direct and headspace cold-fiber solid-phase microextraction modes in the same procedure for the determination of polycyclic aromatic hydrocarbons and phthalate esters in soil samples. Journal of Chromatography A, 2011, 1218, 1707-1714.	1.8	42
123	Determination of THMs in soft drink by solid-phase microextraction and gas chromatography. Food Chemistry, 2011, 127, 290-295.	4.2	22
124	Use of different sample temperatures in a single extraction procedure for the screening of the aroma profile of plant matrices by headspace solid-phase microextraction. Journal of Chromatography A, 2011, 1218, 3731-3736.	1.8	15
125	Volatile compounds of leaves and fruits of Mangifera indica var. coquinho (Anacardiaceae) obtained using solid phase microextraction and hydrodistillation. Food Chemistry, 2011, 127, 689-693.	4.2	36
126	Determination of Trace Silver in Water Samples by Online Column Preconcentration Flame Atomic Absorption Spectrometry Using Termite Digestion Product. Journal of Automated Methods and Management in Chemistry, 2011, 2011, 1-7.	0.5	6

#	Article	IF	CITATIONS
127	Produção de gás de sÃntese por plasma térmico via pirólise de metano e dióxido de carbono. Quimica Nova, 2011, 34, 1491-1495.	0.3	3
128	Fibras altamente robustas de microextração em fase sólida e recobrimento via reação sol-gel. Scientia Chromatographica, 2011, 3, 133-143.	0.2	1
129	High-Resolution Continuum Source Atomic and Molecular Absorption Spectrometry—A Review. Applied Spectroscopy Reviews, 2010, 45, 327-354.	3.4	87
130	A simple hollow fiber renewal liquid membrane extraction method for analysis of sulfonamides in honey samples with determination by liquid chromatography–tandem mass spectrometry. Journal of Chromatography A, 2010, 1217, 6449-6454.	1.8	52
131	Isolation and preconcentration of Cd(II) from environmental samples using polypropylene porous membrane in a hollow fiber renewal liquid membrane extraction procedure and determination by FAAS. Journal of Hazardous Materials, 2010, 177, 567-572.	6.5	30
132	Simultaneous liquid–liquid microextraction and polypropylene microporous membrane solid-phase extraction of organochlorine pesticides in water, tomato and strawberry samples. Journal of Chromatography A, 2010, 1217, 7-13.	1.8	47
133	New sorbents for extraction and microextraction techniques. Journal of Chromatography A, 2010, 1217, 2533-2542.	1.8	224
134	Estudo da eficiência de degradação de tetracloreto de carbono por plasma térmico e caracterização dos produtos formados. Quimica Nova, 2010, 33, 398-403.	0.3	0
135	Determination of sulfur in biological samples using high-resolution molecular absorption spectrometry in a graphite furnace with direct solid sampling. Journal of Analytical Atomic Spectrometry, 2010, 25, 1039.	1.6	35
136	Determination of cadmium in alcohol fuel using Moringa oleifera seeds as a biosorbent in an on-line system coupled to FAAS. Talanta, 2010, 80, 1133-1138.	2.9	67
137	Application of an NiTi alloy coated with ZrO2 solid-phase microextraction fiber for determination of haloanisoles in red wine samples. Mikrochimica Acta, 2009, 164, 197-202.	2.5	20
138	Speciation of Cr(III) and Cr(VI) in environmental samples determined by selective separation and preconcentration on silica gel chemically modified with niobium(V) oxide. Journal of Hazardous Materials, 2009, 161, 450-456.	6.5	53
139	Simple hollow fiber renewal liquid membrane extraction method for pre-concentration of Cd(II) in environmental samples and detection by Flame Atomic Absorption Spectrometry. Analytica Chimica Acta, 2009, 638, 45-50.	2.6	44
140	Development of a flow system for the determination of cadmium in fuel alcohol using vermicompost as biosorbent and flame atomic absorption spectrometry. Talanta, 2009, 78, 333-336.	2.9	68
141	Analysis of Volatile Compounds Released From Flowers and Roots of <i>Polygala cyparissias</i> and <i>Polygala paniculata</i> by Headspace/SPME. Journal of Essential Oil Research, 2009, 21, 255-258.	1.3	13
142	Determination of Phthalates and Adipate in Physiological Saline Solutions by Solid-Phase Microextraction and Gas Chromatography. Analytical Sciences, 2009, 25, 865-868.	0.8	7
143	Application of poly(dimethylsiloxane) fiber sol–gel coated onto NiTi alloy electrodeposited with zirconium oxide for the determination of organochlorine pesticides in herbal infusions. Journal of Separation Science, 2008, 31, 2875-2881.	1.3	14
144	Use of Al2O3 in an automated on-line pre-concentration system for determination of cadmium(II) by FAAS. Journal of Hazardous Materials, 2008, 150, 328-334.	6.5	9

#	Article	IF	CITATIONS
145	Use of 8-hydroxyquinoline-chitosan chelating resin in an automated on-line preconcentration system for determination of zinc(II) by F AAS. Journal of Hazardous Materials, 2008, 157, 88-93.	6.5	47
146	A combination of statistical and analytical evaluation methods as a new optimization strategy for the quantification of pharmaceutical residues in sewage effluent. Analytica Chimica Acta, 2008, 613, 169-176.	2.6	22
147	Headspace solid-phase microextraction–gas chromatographic–time-of-flight mass spectrometric methodology for geographical origin verification of coffee. Analytica Chimica Acta, 2008, 617, 72-84.	2.6	128
148	Application of robust NiTi–ZrO2–PEG SPME fiber in the determination of haloanisoles in cork stopper samples. Analytica Chimica Acta, 2008, 629, 92-97.	2.6	26
149	Preparation and characterization of new solid-phase microextraction fibers obtained by sol–gel technology and zirconium oxide electrodeposited on NiTi alloy. Journal of Chromatography A, 2008, 1187, 34-39.	1.8	61
150	New poly(ethylene glycol) solid-phase microextraction fiber employing zirconium oxide electrolytically deposited onto a NiTi alloy as substrate for sol–gel reactions. Journal of Chromatography A, 2008, 1198-1199, 54-58.	1.8	44
151	Simultaneous on-line pre-concentration and determination of trace metals in environmental samples by flow injection combined with inductively coupled plasma mass spectrometry using silica gel modified with niobium(V) oxide. Talanta, 2008, 74, 586-592.	2.9	19
152	Single drop micro-extraction with O,O-diethyl dithiophosphate for the determination of lead by electrothermal atomic absorption spectrometry. Talanta, 2008, 74, 800-805.	2.9	56
153	Use of Addition Calibration Technique for Determination of Acetaminophen and Hydrochlorothiazide in Human Urine by High-Performance Liquid Chromatography. Journal of Chromatographic Science, 2008, 46, 804-808.	0.7	10
154	Application of Factorial Design and Doehlert Matrix for Determination of Trace Lead in Environmental Samples by On-line Column Preconcentration FAAS Using Silica Gel Chemically Modified with Niobium(V) Oxide. Analytical Sciences, 2008, 24, 365-370.	0.8	7
155	Use of the SPME-GC-MS technique to study the thermal degradation of isotactic polypropylene: Effects of temperature and reaction time, and analysis of the reaction mechanism. E-Polymers, 2008, 8, .	1.3	8
156	Determinação de herbicidas usados no cultivo de arroz irrigado na região sul do estado de Santa Catarina através da SPME-GC-ECD. Quimica Nova, 2008, 31, 79-83.	0.3	9
157	Otimização multivariada e aplicação do sorvente SiO2-Nb2O5 para determinação em linha de Ni(II) em matriz aquosa. Ecletica Quimica, 2008, 33, 25-32.	0.2	4
158	Determination of trihalomethanes in drinking water from three different water sources in Florianopolis-Brazil using purge and trap and gas chromatography. Journal of the Brazilian Chemical Society, 2007, 18, 741-747.	0.6	12
159	Fast and sensitive method to determine chloroanisoles in cork using an internally cooled solid-phase microextraction fiber. Journal of Chromatography A, 2007, 1138, 10-17.	1.8	77
160	Application of fractional factorial experimental and Box-Behnken designs for optimization of single-drop microextraction of 2,4,6-trichloroanisole and 2,4,6-tribromoanisole from wine samples. Journal of Chromatography A, 2007, 1148, 131-136.	1.8	120
161	Preparation and application of NiTi alloy coated with ZrO2 as a new fiber for solid-phase microextraction. Journal of Chromatography A, 2007, 1164, 18-24.	1.8	71
162	Application of NiTi alloy coated with ZrO2 as a new fiber for solid-phase microextraction for determination of halophenols in water samples. Analytica Chimica Acta, 2007, 598, 254-260.	2.6	61

#	Article	IF	CITATIONS
163	Cloud point extraction for the determination of lead and cadmium in urine by graphite furnace atomic absorption spectrometry with multivariate optimization using Box–Behnken design. Spectrochimica Acta, Part B: Atomic Spectroscopy, 2007, 62, 1019-1027.	1.5	51
164	Monitoring the formation of trihalomethanes in the effluents from a shrimp hatchery. Environmental Monitoring and Assessment, 2007, 127, 435-444.	1.3	6
165	Determination of haloanisoles in paper samples for food packaging by solid-phase microextraction and gas chromatography. Mikrochimica Acta, 2007, 159, 229-234.	2.5	11
166	Screening of Tropical Fruit Volatile Compounds Using Solid-Phase Microextraction (SPME) Fibers and Internally Cooled SPME Fiber. Journal of Agricultural and Food Chemistry, 2006, 54, 8688-8696.	2.4	139
167	Development of an on-line preconcentration system for zinc determination in biological samples. Talanta, 2006, 69, 488-493.	2.9	46
168	Analysis of Zinc in Biological Samples by Flame Atomic Absorption Spectrometry: Use of Addition Calibration Technique. Biological Trace Element Research, 2006, 111, 265-280.	1.9	10
169	Thermal Immobilization of Poly(Butylacrylate) on Glass-Ceramic Rod for Preparation of Solid Phase Microextraction Fibers. Mikrochimica Acta, 2006, 155, 465-469.	2.5	3
170	Solid phase microextraction to concentrate volatile products from thermal degradation of polymers. Polymer Degradation and Stability, 2005, 89, 33-37.	2.7	20
171	Class fibers coated with Nb2O5 for use in SPME. Colloids and Surfaces A: Physicochemical and Engineering Aspects, 2005, 259, 15-22.	2.3	10
172	The Use of a Thin Glass-Ceramic Rod as a Surface for Sol-Gel Coating in the Preparation of SPME Fibers. Chromatographia, 2005, 61, 277-283.	0.7	13
173	Development of a DC-plasma torch constructed with graphite electrodes and an integrated nebulization system for decomposition of CCl4. Journal of the Brazilian Chemical Society, 2005, 16, 531-534.	0.6	9
174	Chromium speciation and preconcentration using zirconium(IV) and zirconium(IV) phosphate chemically immobilized onto silica gel surface using a flow system and F AAS. Talanta, 2005, 65, 537-542.	2.9	88
175	Niobium(V) oxide coated on thin glass–ceramic rod as a solid phase microextraction fiber. Talanta, 2005, 66, 74-79.	2.9	26
176	A Comparative Study of Homemade C 18 and Commercial C 18 Sorbents for Preconcentration of Lead by Minicolumn Solid Phase Extraction. Mikrochimica Acta, 2004, 144, 17-21.	2.5	9
177	Chelating resin from functionalization of chitosan with complexing agent 8-hydroxyquinoline: application for metal ions on line preconcentration system. Analytica Chimica Acta, 2004, 521, 157-162.	2.6	69
178	Determination of Trace Metals in Fuel Alcohol by FAAS Using Nb2O5–SiO2as Sorbent Material in a Flow Injection On‣ine Preconcentration System. Analytical Letters, 2004, 37, 1909-1924.	1.0	24
179	Use of NbO?SiO in an automated on-line preconcentration system for determination of copper and cadmium by FAAS. Talanta, 2004, 62, 727-733.	2.9	52
180	Sulphoxine immobilized onto chitosan microspheres by spray drying: application for metal ions preconcentration by flow injection analysis. Talanta, 2004, 63, 397-403.	2.9	40

#	Article	IF	CITATIONS
181	Application of silica gel organofunctionalized with 3(1-imidazolyl)propyl in an on-line preconcentration system for the determination of copper by FAAS. Talanta, 2004, 64, 181-189.	2.9	53
182	Application of Nb 2 O 5 -SiO 2 in Pre-Concentration and Determination of Copper and Cadmium by Flow System with Flame Atomic Absorption Spectrometry. Mikrochimica Acta, 2003, 141, 169-174.	2.5	18
183	A new procedure for solid-phase microextraction using a flow-through system with air segmentation. Analyst, The, 2002, 127, 603-607.	1.7	3
184	A new method of microvolume back-extraction procedure for enrichment of Pb and Cd and determination by flame atomic absorption spectrometry. Talanta, 2002, 56, 185-191.	2.9	67
185	Pré-concentração de chumbo e cádmio em um sistema de micro extração lÃquido-lÃquido e determinação por espectrometria de absorção atÃ′mica com chama. Quimica Nova, 2002, 25, 748-752.	0.3	15
186	A low-cost flame atomic absorption spectrometry method for determination of trace metals in aqueous samples. Talanta, 2000, 51, 173-178.	2.9	44
187	Aplicação de SPME (Solid Phase Micro-Extraction) na análise de águas potáveis de três localidades do estado de São Paulo. Quimica Nova, 1998, 21, 804-806.	0.3	4
188	Assessment of a Fully Optimized DPX-Based Procedure for the Multiclass Determination of Pesticides in Drinking Water Using High-Performance Liquid Chromatography with Diode Array Detection. Journal of the Brazilian Chemical Society, 0, , .	0.6	4
189	A Green Procedure Using Disposable Pipette Extraction to Determine Polycyclic Aromatic Sulfur Heterocycles in Water Samples and Solid Petrochemical Residues. Journal of the Brazilian Chemical Society, 0, , .	0.6	1
190	Electrode Modified with the Ionic Liquid [P6,6,6,14+]2[MnCl42-] for the Determination of Bisphenol A in Plastic Samples. Journal of the Brazilian Chemical Society, 0, , .	0.6	2
191	Exploring Polypyrrole as Extraction Phase for Disposable Pipette Extraction Method for Multiclass Organic Micro-Pollutant Determination in River and Tap Water Using Gas Chromatography-Mass Spectrometry. Journal of the Brazilian Chemical Society, 0, , .	0.6	3
192	An Eco-Friendly Proposal by Integrating Chromatographic Fingerprinting and Multivariate Control Chart in a Non-Target Analysis to Evaluate Grape Juices from Different Farming Practices. Journal of the Brazilian Chemical Society, 0, , .	0.6	1
193	A New Approach for Histamine Functionalized Nanoparticles as Sorbent Phase for Bar Adsorptive Microextraction (BAÂμE) and Determination of Parabens in Water by HPLC-DAD. Journal of the Brazilian Chemical Society, 0, , .	0.6	0