

Transformation of intact yeast cells treated with alkali c

Journal of Bacteriology

153, 163-168

DOI: [10.1128/jb.153.1.163-168.1983](https://doi.org/10.1128/jb.153.1.163-168.1983)

Citation Report

| # | ARTICLE | IF | CITATIONS |
|----|--|------|-----------|
| 1 | Mouse Î±-amylase synthesized by <i>Saccharomyces cerevisiae</i> is released into the culture medium. Carlsberg Research Communications, 1983, 48, 545-555. | 1.7 | 54 |
| 2 | Mutagenesis and genetic transformation of meiotic segregants of lager yeast. Carlsberg Research Communications, 1983, 48, 557-565. | 1.7 | 17 |
| 3 | Molecular cloning and characterization of the threonine deaminase (ILV1) gene of <i>Saccharomyces cerevisiae</i> . Carlsberg Research Communications, 1983, 48, 149-159. | 1.7 | 29 |
| 4 | Isolation and characterization of a maltose transport mutant in the yeast <i>Saccharomyces cerevisiae</i> . Current Genetics, 1983, 7, 195-199. | 0.8 | 37 |
| 5 | Evidence for the biochemical role of an internal sequence in yeast nuclear mRNA introns: Implications for U1 RNA and metazoan mRNA splicing. Cell, 1983, 34, 395-403. | 13.5 | 276 |
| 6 | Molecular cloning and characterization of the RAD1 gene of <i>Saccharomyces cerevisiae</i> . Gene, 1983, 26, 119-126. | 1.0 | 50 |
| 7 | Plasmid-encoded hygromycin B resistance: the sequence of hygromycin B phosphotransferase gene and its expression in <i>Escherichia coli</i> and <i>Saccharomyces cerevisiae</i> . Gene, 1983, 25, 179-188. | 1.0 | 870 |
| 8 | A general method for polyethylene-glycol-induced genetic transformation of bacteria and yeast. Gene, 1983, 25, 333-341. | 1.0 | 493 |
| 9 | Cloning of <i>Saccharomyces cerevisiae</i> DNA replication genes: isolation of the CDC8 gene and two genes that compensate for the cdc8-1 mutation.. Molecular and Cellular Biology, 1983, 3, 1730-1737. | 1.1 | 216 |
| 10 | Isolation and characterization of the RAD3 gene of <i>Saccharomyces cerevisiae</i> and inviability of rad3 deletion mutants. Proceedings of the National Academy of Sciences of the United States of America, 1983, 80, 5680-5684. | 3.3 | 119 |
| 11 | Yeast mutants deficient in protein glycosylation.. Proceedings of the National Academy of Sciences of the United States of America, 1983, 80, 7466-7470. | 3.3 | 245 |
| 12 | [22] Cloning of yeast STE genes in 2 Î¼m vectors. Methods in Enzymology, 1983, 101, 325-343. | 0.4 | 64 |
| 13 | Import of proteins into mitochondria: a 70 kilodalton outer membrane protein with a large carboxy-terminal deletion is still transported to the outer membrane.. EMBO Journal, 1983, 2, 2161-2168. | 3.5 | 150 |
| 14 | Localization and sequence analysis of chloroplast DNA sequences of <i>Chlamydomonas reinhardtii</i> that promote autonomous replication in yeast. EMBO Journal, 1984, 3, 415-421. | 3.5 | 64 |
| 15 | Expression of plasmid R388-encoded type II dihydrofolate reductase as a dominant selective marker in <i>Saccharomyces cerevisiae</i> .. Molecular and Cellular Biology, 1984, 4, 407-414. | 1.1 | 55 |
| 16 | The expression of cDNA clones of yeast M1 double-stranded RNA in yeast confers both killer and immunity phenotypes.. EMBO Journal, 1984, 3, 1383-1387. | 3.5 | 57 |
| 17 | Analysis of full-length cDNA clones carrying GAL1 of <i>Saccharomyces cerevisiae</i> : a model system for cDNA expression. Nucleic Acids Research, 1984, 12, 6397-6414. | 6.5 | 30 |
| 18 | Transformation of protoplasted yeast cells is directly associated with cell fusion.. Molecular and Cellular Biology, 1984, 4, 771-778. | 1.1 | 69 |

| # | ARTICLE | IF | CITATIONS |
|----|--|------|-----------|
| 19 | Deletion mutations affecting autonomously replicating sequence ARS1 of <i>Saccharomyces cerevisiae</i> . <i>Molecular and Cellular Biology</i> , 1984, 4, 2455-2466. | 1.1 | 235 |
| 20 | Identification of autonomous replication sequences in genomic and mitochondrial DNA of <i>Crithidia fasciculata</i> . <i>Molecular and Biochemical Parasitology</i> , 1984, 10, 151-160. | 0.5 | 7 |
| 21 | Autonomous replication sequences in the maxicircle kinetoplast DNA of <i>Leishmania tarentolae</i> . <i>Molecular and Biochemical Parasitology</i> , 1984, 13, 263-275. | 0.5 | 13 |
| 22 | Mitochondrial membrane biogenesis: Characterization and use of pet mutants to clone the nuclear gene coding for subunit V of yeast cytochrome c oxidase. <i>Journal of Cellular Biochemistry</i> , 1984, 24, 229-242. | 1.2 | 18 |
| 23 | Cloning of the ILV2, ILV3 and ILV5 genes of <i>Saccharomyces cerevisiae</i> . <i>Carlsberg Research Communications</i> , 1984, 49, 577-584. | 1.7 | 27 |
| 24 | Site-specific mutagenesis of the Ti plasmid by transformation of <i>Agrobacterium tumefaciens</i> with mutagenized T-DNA fragments cloned in <i>E. coli</i> plasmids. <i>Molecular Genetics and Genomics</i> , 1984, 194, 188-194. | 2.4 | 22 |
| 25 | A new, rapid and efficient transformation procedure for <i>Neurospora</i> . <i>Current Genetics</i> , 1984, 8, 77-79. | 0.8 | 95 |
| 26 | Cloning by genetic complementation and restriction mapping of the yeast HEM1 gene coding for 5-aminolevulinic acid synthase. <i>Current Genetics</i> , 1984, 8, 327-331. | 0.8 | 17 |
| 27 | Rapid genetic test for discrimination between haploid and polyploid transformants in <i>Saccharomyces</i> . <i>Current Genetics</i> , 1984, 8, 341-344. | 0.8 | 5 |
| 28 | Hygromycin B resistance as dominant selectable marker in yeast. <i>Current Genetics</i> , 1984, 8, 353-358. | 0.8 | 96 |
| 29 | Conformational alterations in the proximal portion of the yeast invertase signal peptide do not block secretion. <i>Molecular Genetics and Genomics</i> , 1984, 197, 351-357. | 2.4 | 22 |
| 30 | Telomeres and P-element of <i>Drosophila melanogaster</i> contain sequences that replicate autonomously in <i>Saccharomyces cerevisiae</i> . <i>Molecular Genetics and Genomics</i> , 1984, 197, 342-344. | 2.4 | 5 |
| 31 | Regulation of expression of the galactose gene cluster in <i>Saccharomyces cerevisiae</i> . <i>Molecular Genetics and Genomics</i> , 1984, 195, 29-34. | 2.4 | 83 |
| 32 | Transfer of yeast telomeres to linear plasmids by recombination. <i>Cell</i> , 1984, 39, 191-201. | 13.5 | 172 |
| 33 | Isolation and characterization of the RAD2 gene of <i>Saccharomyces cerevisiae</i> . <i>Gene</i> , 1984, 30, 121-128. | 1.0 | 42 |
| 34 | A 189-bp fragment of <i>Crithidia fasciculata</i> maxicircle DNA confers autonomous replication in <i>Saccharomyces cerevisiae</i> . <i>Gene</i> , 1984, 29, 103-112. | 1.0 | 11 |
| 35 | The major promoter element of rRNA transcription in yeast lies 2 kb upstream. <i>Cell</i> , 1984, 39, 663-673. | 13.5 | 268 |
| 36 | The yeast cloning vector YEp13 contains a tRNA ^{3Leu} gene that can mutate to an amber suppressor. <i>Gene</i> , 1984, 27, 239-251. | 1.0 | 21 |

| # | ARTICLE | IF | CITATIONS |
|----|---|------|-----------|
| 37 | Cis-acting, recombination-stimulating activity in a fragment of the ribosomal DNA of <i>S. cerevisiae</i> . <i>Cell</i> , 1984, 39, 377-386. | 13.5 | 277 |
| 38 | Lariat structures are in vivo intermediates in yeast pre-mRNA splicing. <i>Cell</i> , 1984, 39, 611-621. | 13.5 | 412 |
| 39 | Two modules from the hypersuppressive rho ⁺ mitochondrial DNA are required for plasmid replication in yeast. <i>Gene</i> , 1984, 30, 47-61. | 1.0 | 29 |
| 40 | Introduction of the plasmid pKM101-associated muc genes into <i>Saccharomyces cerevisiae</i> . <i>Mutation Research - DNA Repair Reports</i> , 1984, 131, 197-204. | 1.9 | 10 |
| 41 | DNA Transformation in Cyanobacteria, Algae and Fungi: Trends and Perspectives. , 1984, , 192-207. | | 0 |
| 42 | Expression of calf prochymosin in <i>Saccharomyces cerevisiae</i> . <i>Gene</i> , 1984, 27, 35-46. | 1.0 | 118 |
| 43 | The SPT3 gene is required for normal transcription of Ty elements in <i>S. cerevisiae</i> . <i>Cell</i> , 1984, 39, 675-682. | 13.5 | 240 |
| 44 | Micronuclear DNA of <i>Oxytricha nova</i> contains sequences with autonomously replicating activity in <i>Saccharomyces cerevisiae</i> .. <i>Molecular and Cellular Biology</i> , 1984, 4, 1725-1729. | 1.1 | 5 |
| 45 | Two genes for ribosomal protein 51 of <i>Saccharomyces cerevisiae</i> complement and contribute to the ribosomes.. <i>Molecular and Cellular Biology</i> , 1984, 4, 1871-1879. | 1.1 | 112 |
| 46 | Yeast gene CDC8 encodes thymidylate kinase and is complemented by herpes thymidine kinase gene TK.. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 1984, 81, 5821-5825. | 3.3 | 113 |
| 47 | Transformation of intact yeast cells treated with alkali cations or thiol compounds.. <i>Agricultural and Biological Chemistry</i> , 1984, 48, 341-347. | 0.3 | 44 |
| 48 | <i>Lactobacillus casei</i> IFO-3831. <i>Nippon Nogeikagaku Kaishi</i> , 1984, 58, 479-482. | | 0 |
| 49 | Invertase beta-galactosidase hybrid proteins fail to be transported from the endoplasmic reticulum in <i>Saccharomyces cerevisiae</i> .. <i>Molecular and Cellular Biology</i> , 1984, 4, 2347-2355. | 1.1 | 89 |
| 50 | Repetitive <i>Dictyostelium</i> heat-shock promotor functions in <i>Saccharomyces cerevisiae</i> .. <i>Molecular and Cellular Biology</i> , 1984, 4, 591-598. | 1.1 | 35 |
| 51 | MAL6 of <i>Saccharomyces</i> : a complex genetic locus containing three genes required for maltose fermentation.. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 1984, 81, 2811-2815. | 3.3 | 131 |
| 52 | Structure and expression of the SNF1 gene of <i>Saccharomyces cerevisiae</i> .. <i>Molecular and Cellular Biology</i> , 1984, 4, 54-60. | 1.1 | 74 |
| 53 | Intrachromosomal movement of genetically marked <i>Saccharomyces cerevisiae</i> transposons by gene conversion.. <i>Molecular and Cellular Biology</i> , 1984, 4, 703-711. | 1.1 | 52 |
| 54 | Two nuclear mutations that block mitochondrial protein import in yeast.. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 1984, 81, 4819-4823. | 3.3 | 509 |

| # | ARTICLE | IF | CITATIONS |
|----|--|-----|-----------|
| 55 | Regulation of basal and induced levels of the MEL1 transcript in <i>Saccharomyces cerevisiae</i> .. <i>Molecular and Cellular Biology</i> , 1984, 4, 1238-1245. | 1.1 | 109 |
| 56 | An immunological method for detecting gene expression in yeast colonies.. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 1984, 81, 7426-7430. | 3.3 | 33 |
| 57 | Intracellular targeting and import of an F1-ATPase beta-subunit-beta-galactosidase hybrid protein into yeast mitochondria.. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 1984, 81, 3983-3987. | 3.3 | 124 |
| 58 | Codon recognition during frameshift suppression in <i>Saccharomyces cerevisiae</i> .. <i>Molecular and Cellular Biology</i> , 1984, 4, 2052-2061. | 1.1 | 42 |
| 59 | Effects of altered 5'-flanking sequences on the in vivo expression of a <i>Saccharomyces cerevisiae</i> tRNATyr gene.. <i>Molecular and Cellular Biology</i> , 1984, 4, 657-665. | 1.1 | 136 |
| 60 | Expression of phaseolin cDNA genes in yeast under control of natural plant DNA sequences. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 1985, 82, 334-338. | 3.3 | 40 |
| 61 | Cloning of yeast TOP1, the gene encoding DNA topoisomerase I, and construction of mutants defective in both DNA topoisomerase I and DNA topoisomerase II.. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 1985, 82, 7178-7182. | 3.3 | 165 |
| 62 | Influenza viral (A/WSN/33) hemagglutinin is expressed and glycosylated in the yeast <i>Saccharomyces cerevisiae</i> .. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 1985, 82, 2019-2023. | 3.3 | 37 |
| 63 | Recombination hot spot in the human beta-globin gene cluster: meiotic recombination of human DNA fragments in <i>Saccharomyces cerevisiae</i> .. <i>Molecular and Cellular Biology</i> , 1985, 5, 2029-2038. | 1.1 | 54 |
| 64 | Identification of autonomously replicating circular subtelomeric Y' elements in <i>Saccharomyces cerevisiae</i> .. <i>Molecular and Cellular Biology</i> , 1985, 5, 2369-2380. | 1.1 | 62 |
| 65 | Isolation and DNA sequence of ADH3, a nuclear gene encoding the mitochondrial isozyme of alcohol dehydrogenase in <i>Saccharomyces cerevisiae</i> .. <i>Molecular and Cellular Biology</i> , 1985, 5, 3024-3034. | 1.1 | 164 |
| 66 | Effect of RP51 gene dosage alterations on ribosome synthesis in <i>Saccharomyces cerevisiae</i> .. <i>Molecular and Cellular Biology</i> , 1985, 5, 3429-3435. | 1.1 | 114 |
| 67 | Structure of the promoter of the <i>Dictyostelium discoideum</i> prespore EB4 gene.. <i>Molecular and Cellular Biology</i> , 1985, 5, 1465-1472. | 1.1 | 33 |
| 68 | Cloning and molecular analysis of the HAP2 locus: a global regulator of respiratory genes in <i>Saccharomyces cerevisiae</i> .. <i>Molecular and Cellular Biology</i> , 1985, 5, 3410-3416. | 1.1 | 155 |
| 69 | Specific protein binding to far upstream activating sequences in polymerase II promoters.. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 1985, 82, 43-47. | 3.3 | 262 |
| 70 | Activation regions in a yeast transposon have homology to mating type control sequences and to mammalian enhancers.. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 1985, 82, 5423-5427. | 3.3 | 110 |
| 71 | Two-micrometer circle site-specific recombination: the minimal substrate and the possible role of flanking sequences.. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 1985, 82, 5875-5879. | 3.3 | 83 |
| 72 | Point mutations implicate repeated sequences as essential elements of the CYC7 negative upstream site in <i>Saccharomyces cerevisiae</i> .. <i>Molecular and Cellular Biology</i> , 1985, 5, 2951-2958. | 1.1 | 30 |

| # | ARTICLE | IF | CITATIONS |
|----|--|------|-----------|
| 73 | RAD3 gene of <i>Saccharomyces cerevisiae</i> : nucleotide sequence of wild-type and mutant alleles, transcript mapping, and aspects of gene regulation.. <i>Molecular and Cellular Biology</i> , 1985, 5, 17-26. | 1.1 | 102 |
| 74 | Suppressible and nonsuppressible +1 G-C base pair insertions induced by ICR-170 at the <i>his4</i> locus in <i>Saccharomyces cerevisiae</i> .. <i>Molecular and Cellular Biology</i> , 1985, 5, 2247-2256. | 1.1 | 31 |
| 75 | A novel method for transformation of intact yeast cells by electroinjection of plasmid DNA. <i>Applied Microbiology and Biotechnology</i> , 1985, 21, 336. | 1.7 | 129 |
| 76 | Mitochondrial DNA of the basidiomycete <i>Polyporus ciliatus</i> . <i>Applied Microbiology and Biotechnology</i> , 1985, 21, 196-199. | 1.7 | 6 |
| 77 | Organization of the MAL loci of <i>Saccharomyces</i> . <i>Molecular Genetics and Genomics</i> , 1985, 200, 1-8. | 2.4 | 56 |
| 78 | The use of plasmid DNA to probe DNA repair functions in the yeast <i>Saccharomyces cerevisiae</i> . <i>Molecular Genetics and Genomics</i> , 1985, 201, 99-106. | 2.4 | 32 |
| 79 | A mutant of <i>Saccharomyces cerevisiae</i> with impaired maintenance of centromeric plasmids. <i>Current Genetics</i> , 1985, 10, 15-20. | 0.8 | 42 |
| 80 | Maintenance and copy number control of ARS1 plasmids in <i>Saccharomyces cerevisiae</i> . <i>Current Genetics</i> , 1985, 10, 21-27. | 0.8 | 10 |
| 81 | An <i>E. coli</i> -yeast shuttle cosmid with positive selection for inserted fragments. <i>Current Genetics</i> , 1985, 10, 29-33. | 0.8 | 2 |
| 82 | Integrative transformation of the yeast <i>Yarrowia lipolytica</i> . <i>Current Genetics</i> , 1985, 10, 39-48. | 0.8 | 113 |
| 83 | Integrative transformation of the yeast <i>Yarrowia lipolytica</i> . <i>Current Genetics</i> , 1985, 10, 49-58. | 0.8 | 77 |
| 84 | Selection and stability of yeast transformants expressing cDNA of an M1 killer toxin-immunity gene. <i>Current Genetics</i> , 1985, 9, 285-291. | 0.8 | 32 |
| 85 | The transformation of brewing yeasts with a plasmid containing the gene for copper resistance. <i>Current Genetics</i> , 1985, 9, 133-138. | 0.8 | 95 |
| 86 | Regulation of the expression of iso 2-cytochrome c gene in <i>S. cerevisiae</i> : cloning of the positive regulatory gene CYP1 and identification of the region of its target sequence on the structural gene CYP3. <i>Molecular Genetics and Genomics</i> , 1985, 199, 524-533. | 2.4 | 42 |
| 87 | Primary structure of the <i>Saccharomyces cerevisiae</i> GAL7 gene. <i>Yeast</i> , 1985, 1, 67-77. | 0.8 | 169 |
| 88 | The PET18 locus of <i>Saccharomyces cerevisiae</i> : A complex locus containing multiple genes. <i>Yeast</i> , 1985, 1, 159-171. | 0.8 | 50 |
| 89 | Transformation of <i>Kluyveromyces lactis</i> cells by plasmid DNA does not require alkaline cations. <i>Biotechnology Letters</i> , 1985, 7, 723-726. | 1.1 | 1 |
| 90 | Site-directed mutagenesis of cytochrome c shows that an invariant Phe is not essential for function. <i>Nature</i> , 1985, 313, 152-154. | 13.7 | 175 |

| # | ARTICLE | IF | CITATIONS |
|-----|---|------|-----------|
| 91 | Site-directed mutagenesis shows that tyrosine 248 of carboxypeptidase A does not play a crucial role in catalysis. <i>Nature</i> , 1985, 317, 551-555. | 13.7 | 130 |
| 92 | Increased hydrophobicity of the Sâ€²1 binding site in carboxypeptidase Y obtained by site-directed mutagenesis. <i>Carlsberg Research Communications</i> , 1985, 50, 273-284. | 1.7 | 22 |
| 93 | Nucleotide sequences of <i>STE2</i> and <i>STE3</i>, cell type-specific sterile genes from <i>Saccharomyces cerevisiae</i>. <i>EMBO Journal</i> , 1985, 4, 2643-2648. | 3.5 | 327 |
| 94 | The first twelve amino acids (less than half of the pre-sequence) of an imported mitochondrial protein can direct mouse cytosolic dihydrofolate reductase into the yeast mitochondrial matrix.. <i>EMBO Journal</i> , 1985, 4, 2061-2068. | 3.5 | 264 |
| 95 | The first twelve amino acids of a yeast mitochondrial outer membrane protein can direct a nuclear-coded cytochrome oxidase subunit to the mitochondrial inner membrane.. <i>EMBO Journal</i> , 1985, 4, 3509-3518. | 3.5 | 132 |
| 96 | Molecular cloning and characterization of the yeast RAD10 gene and expression of RAD10 protein in <i>E. coli</i> .. <i>EMBO Journal</i> , 1985, 4, 1575-1582. | 3.5 | 43 |
| 97 | In vivo modulation of yeast tRNA gene expression by 5â€²-flanking sequences.. <i>EMBO Journal</i> , 1985, 4, 2649-2656. | 3.5 | 63 |
| 98 | Role of conserved sequence elements in yeast centromere DNA.. <i>EMBO Journal</i> , 1985, 4, 1867-1874. | 3.5 | 94 |
| 99 | Yeast mRNA initiation sites are determined primarily by specific sequences, not by the distance from the TATA element.. <i>EMBO Journal</i> , 1985, 4, 3273-3280. | 3.5 | 148 |
| 100 | Structure, expression and regulation of a nuclear gene encoding a mitochondrial protein: the yeast L(+)-lactate cytochrome c oxidoreductase (cytochrome b2).. <i>EMBO Journal</i> , 1985, 4, 3265-3272. | 3.5 | 165 |
| 101 | Effect of ARS1 mutations on chromosome stability in <i>Saccharomyces cerevisiae</i> .. <i>Molecular and Cellular Biology</i> , 1985, 5, 1676-1684. | 1.1 | 96 |
| 102 | Physical and functional structure of a yeast plasmid, pSB3, isolated from <i>Zygosaccharomyces bisporus</i> . <i>Nucleic Acids Research</i> , 1985, 13, 4267-4283. | 6.5 | 55 |
| 103 | Nucleotide sequence of <i>Candida pelliculosa</i> â²-glucosidase gene. <i>Nucleic Acids Research</i> , 1985, 13, 6273-6282. | 6.5 | 64 |
| 104 | The yeast Î±-factor receptor: structural properties deduced from the sequence of the STE2 gene. <i>Nucleic Acids Research</i> , 1985, 13, 8463-8475. | 6.5 | 340 |
| 105 | Invertase signal and mature sequence substitutions that delay intercompartmental transport of active enzyme.. <i>Journal of Cell Biology</i> , 1985, 100, 1664-1675. | 2.3 | 179 |
| 106 | Transcription of a <i>Drosophila</i> tRNA Arg gene in yeast extract: 5â€²-flanking sequence dependence for transcription in a heterologous system. <i>Nucleic Acids Research</i> , 1985, 13, 2803-2814. | 6.5 | 25 |
| 107 | Characterization of a gene product (Sec53p) required for protein assembly in the yeast endoplasmic reticulum.. <i>Journal of Cell Biology</i> , 1985, 101, 2374-2382. | 2.3 | 79 |
| 108 | Expression of the <i>Pseudomonas</i> Gene Coding for Carboxypeptidase G2 in <i>Saccharomyces cerevisiae</i> . <i>Microbiology (United Kingdom)</i> , 1985, 131, 897-903. | 0.7 | 1 |

| # | ARTICLE | IF | CITATIONS |
|-----|--|------|-----------|
| 109 | Nucleotide sequence and transcriptional mapping of the yeast <i>pet56-his3-ded1</i> gene region. <i>Nucleic Acids Research</i> , 1985, 13, 8587-8601. | 6.5 | 307 |
| 110 | A test of clathrin function in protein secretion and cell growth. <i>Science</i> , 1985, 230, 1009-1014. | 6.0 | 191 |
| 111 | A System for Dominant Transformation and Plasmid Amplification in <i>Saccharomyces cerevisiae</i> . <i>Nature Biotechnology</i> , 1985, 3, 451-456. | 9.4 | 22 |
| 112 | Autonomously replicating sequences from the non transcribed spacers of <i>Tetrahymena</i> thermophilus ribosomal DNA. <i>Nucleic Acids Research</i> , 1985, 13, 2647-2659. | 6.5 | 26 |
| 113 | One-step gene replacement in yeast by cotransformation. <i>Gene</i> , 1985, 36, 87-95. | 1.0 | 77 |
| 114 | Expression of a nitrogen-fixation gene encoding a nitrogenase subunit in yeast. <i>Gene</i> , 1985, 35, 1-9. | 1.0 | 22 |
| 115 | New vectors for construction of recombinant high-copy-number yeast acentric-ring plasmids. <i>Gene</i> , 1985, 40, 217-229. | 1.0 | 15 |
| 116 | Transformation of <i>Rhodospiridium toruloides</i> . <i>Gene</i> , 1985, 36, 235-240. | 1.0 | 51 |
| 117 | Ty elements transpose through an RNA intermediate. <i>Cell</i> , 1985, 40, 491-500. | 13.5 | 906 |
| 118 | Heterologous Gene Expression in <i>Saccharomyces cerevisiae</i> . <i>Biotechnology and Genetic Engineering Reviews</i> , 1985, 3, 377-416. | 2.4 | 88 |
| 119 | Mitotic stability of yeast chromosomes: A colony color assay that measures nondisjunction and chromosome loss. <i>Cell</i> , 1985, 40, 381-392. | 13.5 | 537 |
| 120 | Ty element transposition: Reverse transcriptase and virus-like particles. <i>Cell</i> , 1985, 42, 507-517. | 13.5 | 367 |
| 121 | Conservation of a 29-base-pair sequence within maxicircle ARS fragments from six species of trypanosomes. <i>Gene</i> , 1985, 40, 291-299. | 1.0 | 5 |
| 122 | A eukaryotic transcriptional activator bearing the DNA specificity of a prokaryotic repressor. <i>Cell</i> , 1985, 43, 729-736. | 13.5 | 679 |
| 123 | DNA sequence and characterization of the <i>S. cerevisiae</i> gene encoding adenylate cyclase. <i>Cell</i> , 1985, 43, 493-505. | 13.5 | 468 |
| 124 | Transcription of the human adenovirus E1a gene in <i>Saccharomyces cerevisiae</i> . <i>Gene</i> , 1985, 33, 159-168. | 1.0 | 6 |
| 125 | Functional selection and analysis of yeast centromeric DNA. <i>Cell</i> , 1985, 42, 913-921. | 13.5 | 270 |
| 126 | Controlled transcription of the yeast regulatory gene <i>GAL80</i> . <i>Gene</i> , 1985, 39, 1-9. | 1.0 | 53 |

| # | ARTICLE | IF | CITATIONS |
|-----|---|------|-----------|
| 127 | Molecular consequences of specific intron mutations on yeast mRNA splicing in vivo and in vitro. <i>Cell</i> , 1985, 42, 335-344. | 13.5 | 181 |
| 128 | Molecular cloning of the RAD10 gene of <i>Saccharomyces cerevisiae</i> . <i>Gene</i> , 1985, 34, 55-61. | 1.0 | 31 |
| 129 | Total deletion of yeast LEU4: Further evidence for a second Î±-isopropylmalate synthase and evidence for tight LEU4-MET4 linkage. <i>Gene</i> , 1985, 33, 333-339. | 1.0 | 33 |
| 130 | A quantitative analysis of the effects of 5â€² junction and TACTAAC box mutants and mutant combinations on yeast mRNA splicing. <i>Cell</i> , 1985, 43, 423-430. | 13.5 | 123 |
| 131 | A region flanking the GAL7 gene and a binding site for GAL4 protein as upstream activating sequences in yeast. <i>Journal of Molecular Biology</i> , 1985, 186, 821-824. | 2.0 | 45 |
| 132 | Fructose biphosphatase of <i>Saccharomyces cerevisiae</i> . <i>Journal of Molecular Biology</i> , 1985, 186, 307-319. | 2.0 | 65 |
| 133 | Expression of the gene for ornithine decarboxylase of <i>Saccharomyces cerevisiae</i> in <i>Escherichia coli</i> .. <i>Molecular and Cellular Biology</i> , 1985, 5, 161-166. | 1.1 | 42 |
| 134 | Characteristics of hepatitis B surface antigen produced in yeast. <i>Biochemical and Biophysical Research Communications</i> , 1985, 131, 160-166. | 1.0 | 7 |
| 135 | Specific DNA binding of GAL4, a positive regulatory protein of yeast. <i>Cell</i> , 1985, 40, 767-774. | 13.5 | 773 |
| 136 | Interaction of plasmid DNA with yeast protoplasts and a mechanism of genetic transformation. <i>FEBS Letters</i> , 1985, 183, 211-214. | 1.3 | 7 |
| 137 | <i>Saccharomyces cerevisiae</i> nuclear fusion requires prior activation by alpha factor.. <i>Molecular and Cellular Biology</i> , 1986, 6, 3490-3497. | 1.1 | 98 |
| 138 | Bakers' Yeast. <i>Critical Reviews in Biotechnology</i> , 1986, 4, 75-109. | 5.1 | 29 |
| 139 | Expression of hepatitis B surface antigen by yeast actin gene promoter. <i>Biochemical and Biophysical Research Communications</i> , 1986, 136, 1063-1070. | 1.0 | 4 |
| 140 | Effect of vector type, host strains and transcription terminator on heterologous gene expression in yeast. <i>Biochemical and Biophysical Research Communications</i> , 1986, 140, 602-608. | 1.0 | 5 |
| 141 | Yeast centromeric plasmids as shuttle vectors between <i>Escherichia coli</i> , <i>Bacillus subtilis</i> and <i>Saccharomyces cerevisiae</i> . <i>Biochemical and Biophysical Research Communications</i> , 1986, 135, 915-921. | 1.0 | 0 |
| 142 | A temperature-sensitive mutant of the yeast plasma membrane ATPase obtained by in vitro mutagenesis. <i>FEBS Letters</i> , 1986, 208, 143-146. | 1.3 | 18 |
| 143 | Moloney murine sarcoma virus encoded p37mos expressed in yeast has protein kinase activity. <i>Virology</i> , 1986, 152, 502-506. | 1.1 | 22 |
| 144 | An efficient chloramphenicol-resistance marker for <i>Saccharomyces cerevisiae</i> and <i>Escherichia coli</i> . <i>Gene</i> , 1986, 45, 149-158. | 1.0 | 77 |

| # | ARTICLE | IF | CITATIONS |
|-----|--|------|-----------|
| 145 | The HTS1 gene encodes both the cytoplasmic and mitochondrial histidine tRNA synthetases of <i>S. cerevisiae</i> . <i>Cell</i> , 1986, 46, 235-243. | 13.5 | 326 |
| 146 | Identification of a sequence containing the positive regulatory region of <i>Saccharomyces cerevisiae</i> gene ENO1. <i>Gene</i> , 1986, 45, 67-75. | 1.0 | 57 |
| 147 | Resistance to antimycin A in yeast by amplification of ADH4 on a linear, 42 kb palindromic plasmid. <i>Cell</i> , 1986, 46, 857-863. | 13.5 | 60 |
| 148 | Expression of the hepatitis B virus large envelope protein in <i>Saccharomyces cerevisiae</i> . <i>Gene</i> , 1986, 48, 155-163. | 1.0 | 18 |
| 149 | Splicing of a yeast proline tRNA containing a novel suppressor mutation in the anticodon stem. <i>Journal of Molecular Biology</i> , 1986, 192, 49-63. | 2.0 | 39 |
| 150 | Molecular cloning and regulation of the expression of the MET2 gene of <i>Saccharomyces cerevisiae</i> . <i>Gene</i> , 1986, 46, 71-78. | 1.0 | 28 |
| 151 | Synthesis and processing of Semliki forest virus polyprotein in <i>Saccharomyces cerevisiae</i> : a yeast type glycosylation of E1 envelope protein. <i>Gene</i> , 1986, 48, 267-275. | 1.0 | 11 |
| 152 | The <i>S. cerevisiae</i> structural gene for chitin synthase is not required for chitin synthesis in vivo. <i>Cell</i> , 1986, 46, 213-225. | 13.5 | 274 |
| 153 | Molecular Breeding of Yeasts for Production of Useful Compounds: Novel Methods of Transformation and New Vector Systems. <i>Biotechnology and Genetic Engineering Reviews</i> , 1986, 4, 39-58. | 2.4 | 6 |
| 154 | Application of recombinant DNA technology to the production of useful biomaterials. , 1986, 33, 1-27. | | 22 |
| 155 | Plasmid-mediated Transformed <i>ura3 fur1</i> Double-Mutants of <i>S. cerevisiae</i> : An Autoselection System Applicable to the Production of Foreign Proteins. <i>Nature Biotechnology</i> , 1986, 4, 433-437. | 9.4 | 33 |
| 156 | Developmental regulation of <i>Dictyostelium discoideum</i> actin gene fusions carried on low-copy and high-copy transformation vectors.. <i>Molecular and Cellular Biology</i> , 1986, 6, 3973-3983. | 1.1 | 224 |
| 157 | Genetic Manipulation of Commercial Yeast Strains. <i>Biotechnology and Genetic Engineering Reviews</i> , 1986, 4, 1-38. | 2.4 | 9 |
| 158 | Construction of LYS2 cartridges for use in genetic manipulations of <i>Saccharomyces cerevisiae</i> . <i>Gene</i> , 1986, 46, 237-245. | 1.0 | 108 |
| 159 | Expression of synthetic human-lysozyme gene in <i>Saccharomyces cerevisiae</i> : use of a synthetic chicken-lysozyme signal sequence for secretion and processing. <i>Gene</i> , 1986, 43, 273-279. | 1.0 | 81 |
| 160 | Two yeast mutants defective in endocytosis are defective in pheromone response. <i>Cell</i> , 1986, 46, 355-364. | 13.5 | 186 |
| 161 | Protein sorting in yeast: Mutants defective in vacuole biogenesis mislocalize vacuolar proteins into the late secretory pathway. <i>Cell</i> , 1986, 47, 1041-1051. | 13.5 | 413 |
| 162 | A mitochondrial RNA maturase gene transferred to the yeast nucleus can control mitochondrial mRNA splicing. <i>Cell</i> , 1986, 46, 837-844. | 13.5 | 148 |

| # | ARTICLE | IF | CITATIONS |
|-----|---|------|-----------|
| 163 | Normal stoichiometry of histone dimer sets is necessary for high fidelity of mitotic chromosome transmission. <i>Cell</i> , 1986, 44, 43-52. | 13.5 | 354 |
| 164 | Expression of the human salivary α -amylase gene in yeast and characterization of the secreted protein. <i>Gene</i> , 1986, 50, 247-257. | 1.0 | 38 |
| 165 | Direct identification of small sequence changes in chromosomal DNA. <i>Gene</i> , 1986, 44, 151-158. | 1.0 | 48 |
| 166 | A cloned tryptophan-synthesis gene from the Ascomycete <i>Cochliobolus heterostrophus</i> functions in <i>Escherichia coli</i> , yeast and <i>Aspergillus nidulans</i> . <i>Gene</i> , 1986, 42, 79-88. | 1.0 | 29 |
| 167 | Yeast killer toxin: Site-directed mutations implicate the precursor protein as the immunity component. <i>Cell</i> , 1986, 46, 105-113. | 13.5 | 95 |
| 168 | Normal yeast tRNA ^{CAG} can suppress amber codons and is encoded by an essential gene. <i>Journal of Molecular Biology</i> , 1986, 192, 725-735. | 2.0 | 89 |
| 169 | Multiple upstream AUG codons mediate translational control of GCN4. <i>Cell</i> , 1986, 45, 201-207. | 13.5 | 637 |
| 170 | Regulated yeast promoters produced by DNA rearrangements selected in vivo. <i>Journal of Molecular Biology</i> , 1986, 191, 355-365. | 2.0 | 4 |
| 171 | U2 RNA from yeast is unexpectedly large and contains homology to vertebrate U4, U5, and U6 small nuclear RNAs. <i>Cell</i> , 1986, 47, 49-59. | 13.5 | 168 |
| 172 | Regulation of the yeast metallothionein gene. <i>Gene</i> , 1986, 48, 13-22. | 1.0 | 37 |
| 173 | Hsp26 is not required for growth at high temperatures, nor for thermotolerance, spore development, or germination. <i>Cell</i> , 1986, 45, 885-894. | 13.5 | 244 |
| 174 | Isolation of two genes that affect mitotic chromosome transmission in <i>S. cerevisiae</i> . <i>Cell</i> , 1986, 44, 53-63. | 13.5 | 113 |
| 175 | Identification of sequences in a yeast histone promoter involved in periodic transcription. <i>Cell</i> , 1986, 45, 537-544. | 13.5 | 218 |
| 176 | The presequences of two imported mitochondrial proteins contain information for intracellular and intramitochondrial sorting. <i>Cell</i> , 1986, 44, 801-812. | 13.5 | 213 |
| 177 | Association of reciprocal exchange with gene conversion between the repeated segments of 2- μ m circle. <i>Journal of Molecular Biology</i> , 1986, 191, 341-354. | 2.0 | 12 |
| 178 | Construction of stable laboratory and industrial yeast strains expressing a foreign gene by integrative transformation using a dominant selection system. <i>Gene</i> , 1986, 50, 225-237. | 1.0 | 34 |
| 179 | Site-specific recombination promotes plasmid amplification in yeast. <i>Cell</i> , 1986, 46, 541-550. | 13.5 | 227 |
| 180 | Expression of murine and human granulocyte-macrophage colony-stimulating factors in <i>S. cerevisiae</i> : mutagenesis of the potential glycosylation sites. <i>EMBO Journal</i> , 1986, 5, 1193-1197. | 3.5 | 88 |

| # | ARTICLE | IF | CITATIONS |
|-----|---|-----|-----------|
| 181 | Heme regulates the expression in <i>Saccharomyces cerevisiae</i> of chimaeric genes containing 5â€²-flanking soybean leghemoglobin sequences.. EMBO Journal, 1986, 5, 843-847. | 3.5 | 19 |
| 182 | Chromosome walking shows a highly homologous repetitive sequence present in all the centromere regions of fission yeast. EMBO Journal, 1986, 5, 1011-1021. | 3.5 | 182 |
| 183 | Isolation and characterization of <i>Schizosaccharomyces pombe</i> mutants that block nuclear division but not cytokinesis. EMBO Journal, 1986, 5, 2973-2979. | 3.5 | 279 |
| 184 | At least two nuclear gene products are specifically required for translation of a single yeast mitochondrial mRNA.. EMBO Journal, 1986, 5, 3637-3641. | 3.5 | 89 |
| 185 | The nucleotide sequence of the fission yeast DNA topoisomerase II gene: structural and functional relationships to other DNA topoisomerases.. EMBO Journal, 1986, 5, 2355-2361. | 3.5 | 112 |
| 186 | Suppression of defective RAS1 and RAS2 functions in yeast by an adenylate cyclase activated by a single amino acid change.. EMBO Journal, 1986, 5, 3657-3663. | 3.5 | 87 |
| 187 | Molecular cloning and transcriptional analysis of the start gene <i>CDC25</i> of <i>Saccharomyces cerevisiae</i> . EMBO Journal, 1986, 5, 2363-2369. | 3.5 | 48 |
| 188 | Mitotic spindle pulls but fails to separate chromosomes in type II DNA topoisomerase mutants: uncoordinated mitosis. EMBO Journal, 1986, 5, 1003-1010. | 3.5 | 270 |
| 189 | Yeast pre-messenger RNA splicing efficiency depends on critical spacing requirements between the branch point and 3â€² splice site.. EMBO Journal, 1986, 5, 1023-1030. | 3.5 | 48 |
| 190 | Mutations in conserved intron sequences affect multiple steps in the yeast splicing pathway, particularly assembly of the spliceosome.. EMBO Journal, 1986, 5, 1683-1695. | 3.5 | 197 |
| 191 | The cleavable pre-sequence of an imported chloroplast protein directs attached polypeptides into yeast mitochondria. EMBO Journal, 1986, 5, 1343-1350. | 3.5 | 149 |
| 192 | Site-specific mutagenesis of <i>cdc2+</i> , a cell cycle control gene of the fission yeast <i>Schizosaccharomyces pombe</i> .. Molecular and Cellular Biology, 1986, 6, 3523-3530. | 1.1 | 163 |
| 193 | Expression of the <i>denV</i> gene of coliphage T4 in UV-sensitive rad mutants of <i>Saccharomyces cerevisiae</i> .. Molecular and Cellular Biology, 1986, 6, 3559-3562. | 1.1 | 13 |
| 194 | Single base-pair mutations in centromere element III cause aberrant chromosome segregation in <i>Saccharomyces cerevisiae</i> .. Molecular and Cellular Biology, 1986, 6, 530-538. | 1.1 | 128 |
| 195 | Calcium-sensitive <i>cls4</i> mutant of <i>Saccharomyces cerevisiae</i> with a defect in bud formation. Journal of Bacteriology, 1986, 165, 28-33. | 1.0 | 102 |
| 196 | Construction of a host-vector system in <i>Candida maltosa</i> by using an ARS site isolated from its genome. Journal of Bacteriology, 1986, 167, 551-555. | 1.0 | 53 |
| 197 | Cloning in <i>Saccharomyces cerevisiae</i> of a cycloheximide resistance gene from the <i>Candida maltosa</i> genome which modifies ribosomes. Journal of Bacteriology, 1986, 168, 417-419. | 1.0 | 23 |
| 198 | Intracellular sorting of alcohol dehydrogenase isoenzymes in yeast: a cytosolic location reflects absence of an amino-terminal targeting sequence for the mitochondrion.. EMBO Journal, 1986, 5, 161-165. | 3.5 | 56 |

| # | ARTICLE | IF | CITATIONS |
|-----|---|-----|-----------|
| 199 | Studies of the radical species in compound ES of cytochrome c peroxidase altered by site-directed mutagenesis.. Proceedings of the National Academy of Sciences of the United States of America, 1986, 83, 1295-1299. | 3.3 | 59 |
| 200 | Molecular analysis of SNF2 and SNF5, genes required for expression of glucose-repressible genes in <i>Saccharomyces cerevisiae</i> .. Molecular and Cellular Biology, 1986, 6, 3643-3651. | 1.1 | 127 |
| 201 | Multiple GCD genes required for repression of GCN4, a transcriptional activator of amino acid biosynthetic genes in <i>Saccharomyces cerevisiae</i> .. Molecular and Cellular Biology, 1986, 6, 3990-3998. | 1.1 | 110 |
| 202 | A nuclear gene of <i>Saccharomyces cerevisiae</i> needed for stable maintenance of plasmids.. Molecular and Cellular Biology, 1986, 6, 4053-4059. | 1.1 | 13 |
| 203 | In vivo evidence for posttranslational translocation and signal cleavage of the killer preprotoxin of <i>Saccharomyces cerevisiae</i> .. Molecular and Cellular Biology, 1986, 6, 4274-4280. | 1.1 | 38 |
| 204 | Sequences required for delivery and localization of the ADP/ATP translocator to the mitochondrial inner membrane.. Molecular and Cellular Biology, 1986, 6, 626-634. | 1.1 | 215 |
| 205 | Tripartite upstream promoter element essential for expression of <i>Saccharomyces cerevisiae</i> ribosomal protein genes.. Molecular and Cellular Biology, 1986, 6, 674-687. | 1.1 | 174 |
| 206 | Activation of a cryptic TACTAAC box in the <i>Saccharomyces cerevisiae</i> actin intron.. Molecular and Cellular Biology, 1986, 6, 1571-1578. | 1.1 | 33 |
| 207 | Sequence and expression of the dCMP deaminase gene (DCD1) of <i>Saccharomyces cerevisiae</i> .. Molecular and Cellular Biology, 1986, 6, 1711-1721. | 1.1 | 57 |
| 208 | Isolation and functional analysis of sporulation-induced transcribed sequences from <i>Saccharomyces cerevisiae</i> .. Molecular and Cellular Biology, 1986, 6, 2185-2197. | 1.1 | 55 |
| 209 | Effect of intron mutations on processing and function of <i>Saccharomyces cerevisiae</i> SUP53 tRNA in vitro and in vivo.. Molecular and Cellular Biology, 1986, 6, 2663-2673. | 1.1 | 98 |
| 210 | Transcriptional control of glucoamylase synthesis in vegetatively growing and sporulating <i>Saccharomyces</i> species.. Molecular and Cellular Biology, 1986, 6, 3034-3041. | 1.1 | 59 |
| 211 | Artificial mitochondrial presequences.. Proceedings of the National Academy of Sciences of the United States of America, 1986, 83, 9011-9015. | 3.3 | 213 |
| 212 | Shuttle mutagenesis: a method of transposon mutagenesis for <i>Saccharomyces cerevisiae</i> .. Proceedings of the National Academy of Sciences of the United States of America, 1986, 83, 735-739. | 3.3 | 333 |
| 213 | Sequences upstream of the STE6 gene required for its expression and regulation by the mating type locus in <i>Saccharomyces cerevisiae</i> .. Proceedings of the National Academy of Sciences of the United States of America, 1986, 83, 2536-2540. | 3.3 | 75 |
| 214 | RNA splicing and intron turnover are greatly diminished by a mutant yeast branch point.. Proceedings of the National Academy of Sciences of the United States of America, 1986, 83, 5835-5839. | 3.3 | 79 |
| 215 | Isolation of yeast mutants defective in protein targeting to the vacuole.. Proceedings of the National Academy of Sciences of the United States of America, 1986, 83, 9075-9079. | 3.3 | 389 |
| 216 | Genetically essential and nonessential alpha-tubulin genes specify functionally interchangeable proteins.. Molecular and Cellular Biology, 1986, 6, 3722-3733. | 1.1 | 254 |

| # | ARTICLE | IF | CITATIONS |
|-----|--|-----|-----------|
| 217 | Glycolytic gene expression in <i>Saccharomyces cerevisiae</i> : nucleotide sequence of GCR1, null mutants, and evidence for expression.. <i>Molecular and Cellular Biology</i> , 1986, 6, 3774-3784. | 1.1 | 113 |
| 218 | Structural and functional analysis of the MAL1 locus of <i>Saccharomyces cerevisiae</i> .. <i>Molecular and Cellular Biology</i> , 1986, 6, 3891-3899. | 1.1 | 104 |
| 219 | <i>Saccharomyces cerevisiae</i> contains two functional citrate synthase genes.. <i>Molecular and Cellular Biology</i> , 1986, 6, 1936-1942. | 1.1 | 147 |
| 220 | Transcriptional regulation of ribosomal proteins during a nutritional upshift in <i>Saccharomyces cerevisiae</i> .. <i>Molecular and Cellular Biology</i> , 1986, 6, 2429-2435. | 1.1 | 65 |
| 221 | Identification of a second trans-acting gene controlling maltose fermentation in <i>Saccharomyces carlsbergensis</i> .. <i>Molecular and Cellular Biology</i> , 1986, 6, 2757-2765. | 1.1 | 36 |
| 222 | Toxic effects of excess cloned centromeres.. <i>Molecular and Cellular Biology</i> , 1986, 6, 2213-2222. | 1.1 | 104 |
| 223 | Cloning and characterization of the high-affinity cAMP phosphodiesterase of <i>Saccharomyces cerevisiae</i> .. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 1986, 83, 9303-9307. | 3.3 | 300 |
| 224 | Isolation and characterization of PRT1, a gene required for the initiation of protein biosynthesis in <i>Saccharomyces cerevisiae</i> .. <i>Molecular and Cellular Biology</i> , 1986, 6, 4419-4424. | 1.1 | 26 |
| 225 | Identification of a regulatory region that mediates glucose-dependent induction of the <i>Saccharomyces cerevisiae</i> enolase gene ENO2.. <i>Molecular and Cellular Biology</i> , 1986, 6, 2287-2297. | 1.1 | 82 |
| 226 | Transformation of <i>Phycomyces blakesleeana</i> to G-418 resistance by an autonomously replicating plasmid.. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 1986, 83, 7344-7347. | 3.3 | 68 |
| 227 | Effects of null mutations in the hexokinase genes of <i>Saccharomyces cerevisiae</i> on catabolite repression.. <i>Molecular and Cellular Biology</i> , 1986, 6, 4046-4052. | 1.1 | 129 |
| 228 | An RNA polymerase I enhancer in <i>Saccharomyces cerevisiae</i> .. <i>Molecular and Cellular Biology</i> , 1986, 6, 2089-2097. | 1.1 | 178 |
| 229 | Yeast RPO41 gene product is required for transcription and maintenance of the mitochondrial genome.. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 1986, 83, 3391-3394. | 3.3 | 148 |
| 230 | Meiotic disjunction of circular minichromosomes in yeast does not require DNA homology.. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 1986, 83, 6017-6019. | 3.3 | 43 |
| 231 | Molecular analysis of the DNA sequences involved in the transcriptional regulation of the phosphate-repressible acid phosphatase gene (PHO5) of <i>Saccharomyces cerevisiae</i> .. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 1986, 83, 6070-6074. | 3.3 | 29 |
| 232 | Rapid mapping of antigenic coding regions and constructing insertion mutations in yeast genes by mini-Tn10 "transposon" mutagenesis.. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 1986, 83, 730-734. | 3.3 | 34 |
| 233 | Tightly centromere-linked gene (SPO15) essential for meiosis in the yeast <i>Saccharomyces cerevisiae</i> .. <i>Molecular and Cellular Biology</i> , 1986, 6, 158-167. | 1.1 | 39 |
| 234 | Structure and sequence of the centromeric DNA of chromosome 4 in <i>Saccharomyces cerevisiae</i> .. <i>Molecular and Cellular Biology</i> , 1986, 6, 241-245. | 1.1 | 77 |

| # | ARTICLE | IF | CITATIONS |
|-----|--|-----|-----------|
| 235 | Duplicate upstream activating sequences in the promoter region of the <i>Saccharomyces cerevisiae</i> GAL7 gene.. <i>Molecular and Cellular Biology</i> , 1986, 6, 246-256. | 1.1 | 81 |
| 236 | Mating type-like conversion promoted by the 2 micrograms circle site-specific recombinase: implications for the double-strand-gap repair model.. <i>Molecular and Cellular Biology</i> , 1986, 6, 3831-3837. | 1.1 | 12 |
| 237 | Sequence analysis of temperature-sensitive mutations in the <i>Saccharomyces cerevisiae</i> gene CDC28.. <i>Molecular and Cellular Biology</i> , 1986, 6, 4099-4103. | 1.1 | 68 |
| 238 | RAD7 gene of <i>Saccharomyces cerevisiae</i> : transcripts, nucleotide sequence analysis, and functional relationship between the RAD7 and RAD23 gene products.. <i>Molecular and Cellular Biology</i> , 1986, 6, 1497-1507. | 1.1 | 79 |
| 239 | Differential expressions of essential and nonessential alpha-tubulin genes in <i>Schizosaccharomyces pombe</i> .. <i>Molecular and Cellular Biology</i> , 1986, 6, 2168-2178. | 1.1 | 98 |
| 240 | Flow cytometry analysis of recombinant <i>Saccharomyces cerevisiae</i> populations. <i>Cytometry</i> , 1986, 7, 132-141. | 1.8 | 48 |
| 241 | Extrachromosomal genetics of <i>Cephalosporium acremonium</i> . <i>Applied Microbiology and Biotechnology</i> , 1986, 23, 280. | 1.7 | 2 |
| 242 | Characterization of a chloroplast DNA sequence from <i>Chlorella ellipsoidea</i> that promotes autonomous replication in yeast. <i>Plant Molecular Biology</i> , 1986, 6, 245-252. | 2.0 | 7 |
| 243 | Trehalose and maltose metabolism in yeast transformed by a MAL4 regulatory gene cloned from a constitutive donor strain. <i>Current Genetics</i> , 1986, 11, 97-106. | 0.8 | 11 |
| 244 | Comparison of expression of the endo- β -1,3-1,4-glucanase gene from <i>Bacillus subtilis</i> in <i>Saccharomyces cerevisiae</i> from the CYC1 and ADH1 promoters. <i>Current Genetics</i> , 1986, 11, 65-70. | 0.8 | 28 |
| 245 | Cloning of a <i>Candida utilis</i> gene which complements leu2 mutation in <i>Saccharomyces cerevisiae</i> . <i>Current Genetics</i> , 1986, 10, 573-578. | 0.8 | 8 |
| 246 | Plasmids of mitochondrial origin in senescent mycelia of <i>Podospora curvicolla</i> . <i>Current Genetics</i> , 1986, 10, 803-810. | 0.8 | 36 |
| 247 | Construction of a mini-chromosome by deletion and its mitotic and meiotic behaviour in fission yeast. <i>Molecular Genetics and Genomics</i> , 1986, 203, 397-405. | 2.4 | 105 |
| 248 | Mutations in the right boundary of <i>Saccharomyces cerevisiae</i> centromere 6 lead to nonfunctional or partially functional centromeres. <i>Molecular Genetics and Genomics</i> , 1986, 205, 305-311. | 2.4 | 40 |
| 249 | Molecular cloning and characterization of the STA2 glucoamylase gene of <i>Saccharomyces diastaticus</i> . <i>Molecular Genetics and Genomics</i> , 1986, 203, 29-35. | 2.4 | 50 |
| 250 | Cloning of <i>Candida pelliculosa</i> β -glucosidase gene and its expression in <i>Saccharomyces cerevisiae</i> . <i>Molecular Genetics and Genomics</i> , 1986, 203, 89-94. | 2.4 | 27 |
| 251 | Genetic analysis of partitioning functions encoded by the 2 μ m circle of <i>Saccharomyces cerevisiae</i> . <i>Molecular Genetics and Genomics</i> , 1986, 203, 154-162. | 2.4 | 48 |
| 252 | Partial complementation of the UV sensitivity of <i>E. coli</i> and yeast excision repair mutants by the cloned denV gene of bacteriophage T4. <i>Molecular Genetics and Genomics</i> , 1986, 203, 163-171. | 2.4 | 27 |

| # | ARTICLE | IF | CITATIONS |
|-----|---|------|-----------|
| 253 | Primary structure of wild-type and mutant alleles of the PET494 gene of <i>Saccharomyces cerevisiae</i> . <i>Molecular Genetics and Genomics</i> , 1986, 202, 294-301. | 2.4 | 32 |
| 254 | Transformation of the methylotrophic yeast <i>Hansenula polymorpha</i> by autonomous replication and integration vectors. <i>Molecular Genetics and Genomics</i> , 1986, 202, 302-308. | 2.4 | 125 |
| 255 | Cloning and sequencing of the PHO80 gene and CEN15 of <i>Saccharomyces cerevisiae</i> . <i>Yeast</i> , 1986, 2, 129-139. | 0.8 | 63 |
| 256 | Yeast/ <i>E. coli</i> shuttle vectors with multiple unique restriction sites. <i>Yeast</i> , 1986, 2, 163-167. | 0.8 | 1,471 |
| 257 | Mutations in ARS1 increase the rate of simple loss of plasmids in <i>Saccharomyces cerevisiae</i> . <i>Yeast</i> , 1986, 2, 169-178. | 0.8 | 46 |
| 258 | Physical separation and functional interaction of <i>Kluyveromyces lactis</i> and <i>Saccharomyces cerevisiae</i> ARS elements derived from killer plasmid DNA. <i>Yeast</i> , 1986, 2, 179-191. | 0.8 | 20 |
| 259 | Messenger RNA encoding a glutathione-S-transferase responsible for herbicide tolerance in maize is induced in response to safener treatment. <i>Plant Molecular Biology</i> , 1986, 7, 235-243. | 2.0 | 84 |
| 260 | The stability of the yeast plasmid pJDB248 depends on growth rate of the culture. <i>Biotechnology Letters</i> , 1986, 8, 225-230. | 1.1 | 33 |
| 261 | Sequence homology of the yeast regulatory protein ADR1 with <i>Xenopus</i> transcription factor TFIIIA. <i>Nature</i> , 1986, 320, 283-287. | 13.7 | 232 |
| 262 | Fungal cloning vectors. <i>Trends in Biotechnology</i> , 1986, 4, 93-98. | 4.9 | 11 |
| 263 | Yeasts in molecular biology. Spheroplast preparation with <i>Candida utilis</i> , <i>Schizosaccharomyces pombe</i> and <i>Saccharomyces cerevisiae</i> . <i>Bioscience Reports</i> , 1986, 6, 597-602. | 1.1 | 11 |
| 264 | Analysis of chromosome V and the LV1 gene from <i>Saccharomyces carlsbergensis</i> . <i>Carlsberg Research Communications</i> , 1986, 51, 309-326. | 1.7 | 79 |
| 265 | Cloning and analysis of two alleles of the LV3 gene from <i>Saccharomyces carlsbergensis</i> . <i>Carlsberg Research Communications</i> , 1986, 51, 327-341. | 1.7 | 37 |
| 266 | Construction of a yeast vector directing the synthesis and release of barley (α -3, α -4)- β -glucanase. <i>Carlsberg Research Communications</i> , 1986, 51, 445-458. | 1.7 | 38 |
| 267 | The PEP4 gene encodes an aspartyl protease implicated in the posttranslational regulation of <i>Saccharomyces cerevisiae</i> vacuolar hydrolases. <i>Molecular and Cellular Biology</i> , 1986, 6, 2500-2510. | 1.1 | 301 |
| 268 | Analysis of the yeast SPT3 gene and identification of its product, a positive regulator of Ty transcription. <i>Nucleic Acids Research</i> , 1986, 14, 6885-6900. | 6.5 | 41 |
| 269 | Separation of DNA binding from the transcription-activating function of a eukaryotic regulatory protein. <i>Science</i> , 1986, 231, 699-704. | 6.0 | 637 |
| 270 | Part of the human ribosomal RNA locus stabilizes a plasmid in yeast. <i>Nucleic Acids Research</i> , 1986, 14, 5683-5692. | 6.5 | 0 |

| # | ARTICLE | IF | CITATIONS |
|-----|---|-----|-----------|
| 271 | Factors Enhancing Genetic Transformation of Intact Yeast Cells Modify Cell Wall Porosity. <i>Microbiology (United Kingdom)</i> , 1986, 132, 3089-3093. | 0.7 | 24 |
| 272 | High frequency excision of Ty elements during transformation of yeast. <i>Nucleic Acids Research</i> , 1986, 14, 2989-3001. | 6.5 | 14 |
| 273 | Biosynthesis of the Torpedo californica acetylcholine receptor alpha subunit in yeast. <i>Science</i> , 1986, 231, 1284-1287. | 6.0 | 60 |
| 274 | Identification of the crossover site during FLP-mediated recombination in the <i>Saccharomyces cerevisiae</i> plasmid 2 microns circle.. <i>Molecular and Cellular Biology</i> , 1986, 6, 3357-3367. | 1.1 | 119 |
| 275 | Replicating plasmids in <i>Schizosaccharomyces pombe</i> : improvement of symmetric segregation by a new genetic element.. <i>Molecular and Cellular Biology</i> , 1986, 6, 80-89. | 1.1 | 164 |
| 276 | Efficient isolation of the linear DNA killer plasmid of <i>Kluyveromyces lactis</i> : evidence for location and expression in the cytoplasm and characterization of their terminally bound proteins. <i>Nucleic Acids Research</i> , 1986, 14, 6871-6884. | 6.5 | 86 |
| 277 | Expression of the <i>Saccharomyces cerevisiae</i> inositol-1-phosphate synthase (INO1) gene is regulated by factors that affect phospholipid synthesis.. <i>Molecular and Cellular Biology</i> , 1986, 6, 3320-3328. | 1.1 | 222 |
| 278 | Expression of Rat NADPH-Cytochrome P-450 Reductase cDNA in <i>Saccharomyces cerevisiae</i> . <i>DNA and Cell Biology</i> , 1986, 5, 1-10. | 5.1 | 77 |
| 279 | Gene dosage-dependent secretion of yeast vacuolar carboxypeptidase Y.. <i>Journal of Cell Biology</i> , 1986, 102, 1551-1557. | 2.3 | 222 |
| 280 | The amino terminus of the yeast F1-ATPase beta-subunit precursor functions as a mitochondrial import signal.. <i>Journal of Cell Biology</i> , 1986, 102, 523-533. | 2.3 | 292 |
| 281 | Clathrin requirement for normal growth of yeast. <i>Science</i> , 1987, 238, 504-509. | 6.0 | 120 |
| 282 | A single intronless actin gene in the fission yeast <i>Schizosaccharomyces pombe</i> : nucleotide sequence and transcripts formed in homologous and heterologous yeast. <i>Nucleic Acids Research</i> , 1987, 15, 7369-7379. | 6.5 | 82 |
| 283 | [70] Heterologous gene expression in yeast. <i>Methods in Enzymology</i> , 1987, 152, 673-684. | 0.4 | 15 |
| 284 | A yeast mutant defective at an early stage in import of secretory protein precursors into the endoplasmic reticulum.. <i>Journal of Cell Biology</i> , 1987, 105, 633-645. | 2.3 | 410 |
| 285 | [53] Isolation of genes by complementation in yeast. <i>Methods in Enzymology</i> , 1987, 152, 481-504. | 0.4 | 87 |
| 286 | Molecular Cloning and Synthesis of Biologically Active Human Tissue Inhibitor of Metalloproteinases in Yeast. <i>Nature Biotechnology</i> , 1987, 5, 595-598. | 9.4 | 4 |
| 287 | Yeast carboxypeptidase Y can be translocated and glycosylated without its amino-terminal signal sequence.. <i>Journal of Cell Biology</i> , 1987, 104, 1183-1191. | 2.3 | 81 |
| 288 | Many random sequences functionally replace the secretion signal sequence of yeast invertase. <i>Science</i> , 1987, 235, 312-317. | 6.0 | 386 |

| # | ARTICLE | IF | CITATIONS |
|-----|---|-----|-----------|
| 289 | Cloning and sequencing of <i>Schizosaccharomyces pombe</i> DNA topoisomerase I gene, and effect of gene disruption. <i>Nucleic Acids Research</i> , 1987, 15, 9727-9739. | 6.5 | 91 |
| 290 | Negative regulators of the PHO system in <i>Saccharomyces cerevisiae</i> : isolation and structural characterization of PHO85. <i>Nucleic Acids Research</i> , 1987, 15, 10299-10309. | 6.5 | 78 |
| 291 | The plant vacuolar protein, phytohemagglutinin, is transported to the vacuole of transgenic yeast.. <i>Journal of Cell Biology</i> , 1987, 105, 1971-1979. | 2.3 | 70 |
| 292 | A Method for Gene Disruption That Allows Repeated Use of URA3 Selection in the Construction of Multiply Disrupted Yeast Strains. <i>Genetics</i> , 1987, 116, 541-545. | 1.2 | 946 |
| 293 | Metal-binding, nucleic acid-binding finger sequences in the CDC16 gene of <i>Saccharomyces cerevisiae</i> . <i>Nucleic Acids Research</i> , 1987, 15, 8439-8450. | 6.5 | 37 |
| 294 | Identification and isolation of the gene encoding the small subunit of ribonucleotide reductase from <i>Saccharomyces cerevisiae</i> : DNA damage-inducible gene required for mitotic viability.. <i>Molecular and Cellular Biology</i> , 1987, 7, 2783-2793. | 1.1 | 244 |
| 295 | Molecular characterization of the <i>Saccharomyces cerevisiae</i> dihydrofolate reductase gene (DFRI). <i>Nucleic Acids Research</i> , 1987, 15, 10355-10371. | 6.5 | 48 |
| 296 | Expression and identification of immunity determinants on linear DNA killer plasmids pGKL1 and pGKL2 in <i>Kluyveromyces lactis</i> . <i>Nucleic Acids Research</i> , 1987, 15, 1031-1046. | 6.5 | 71 |
| 297 | Cloning and characterization of BCY1, a locus encoding a regulatory subunit of the cyclic AMP-dependent protein kinase in <i>Saccharomyces cerevisiae</i> .. <i>Molecular and Cellular Biology</i> , 1987, 7, 1371-1377. | 1.1 | 526 |
| 298 | The SPT6 gene is essential for growth and is required for delta-mediated transcription in <i>Saccharomyces cerevisiae</i> .. <i>Molecular and Cellular Biology</i> , 1987, 7, 679-686. | 1.1 | 156 |
| 299 | [22] Visual assay for chromosome ploidy. <i>Methods in Enzymology</i> , 1987, 155, 351-372. | 0.4 | 68 |
| 300 | [33] Expression and secretion vectors for yeast. <i>Methods in Enzymology</i> , 1987, 153, 516-544. | 0.4 | 47 |
| 301 | Isolation and expression of the gene encoding yeast mitochondrial malate dehydrogenase. <i>Journal of Bacteriology</i> , 1987, 169, 5157-5166. | 1.0 | 103 |
| 302 | GAL1-GAL10 divergent promoter region of <i>Saccharomyces cerevisiae</i> contains negative control elements in addition to functionally separate and possibly overlapping upstream activating sequences.. <i>Genes and Development</i> , 1987, 1, 1118-1131. | 2.7 | 64 |
| 303 | [22] Cloning vectors of mitochondrial origin for eukaryotes. <i>Methods in Enzymology</i> , 1987, 153, 366-382. | 0.4 | 2 |
| 304 | The sequence of the <i>Saccharomyces cerevisiae</i> gene PHO2 codes for a regulatory protein with unusual amino acid composition. <i>Nucleic Acids Research</i> , 1987, 15, 233-246. | 6.5 | 147 |
| 305 | <i>Saccharomyces cerevisiae</i> mutants that tolerate centromere plasmids at high copy number.. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 1987, 84, 7203-7207. | 3.3 | 11 |
| 306 | The yeast ARD1 gene product is required for repression of cryptic mating-type information at the HML locus.. <i>Molecular and Cellular Biology</i> , 1987, 7, 3713-3722. | 1.1 | 120 |

| # | ARTICLE | IF | CITATIONS |
|-----|--|-----|-----------|
| 307 | The yeast F1-ATPase beta subunit precursor contains functionally redundant mitochondrial protein import information.. <i>Molecular and Cellular Biology</i> , 1987, 7, 4038-4047. | 1.1 | 74 |
| 308 | Amino acid sequences that determine the nuclear localization of yeast histone 2B.. <i>Molecular and Cellular Biology</i> , 1987, 7, 4048-4057. | 1.1 | 215 |
| 309 | mRNA cap-binding protein: cloning of the gene encoding protein synthesis initiation factor eIF-4E from <i>Saccharomyces cerevisiae</i> .. <i>Molecular and Cellular Biology</i> , 1987, 7, 998-1003. | 1.1 | 156 |
| 310 | Transcriptional regulation of an hsp70 heat shock gene in the yeast <i>Saccharomyces cerevisiae</i> .. <i>Molecular and Cellular Biology</i> , 1987, 7, 1906-1916. | 1.1 | 151 |
| 311 | <i>Saccharomyces cerevisiae</i> positive regulatory gene PET111 encodes a mitochondrial protein that is translated from an mRNA with a long 5' leader.. <i>Molecular and Cellular Biology</i> , 1987, 7, 2728-2734. | 1.1 | 63 |
| 312 | The yeast acid phosphatase can enter the secretory pathway without its N-terminal signal sequence.. <i>Molecular and Cellular Biology</i> , 1987, 7, 3306-3314. | 1.1 | 39 |
| 313 | Alterations in the adenine-plus-thymine-rich region of CEN3 affect centromere function in <i>Saccharomyces cerevisiae</i> .. <i>Molecular and Cellular Biology</i> , 1987, 7, 68-75. | 1.1 | 109 |
| 314 | Identification of healed terminal DNA fragments in linear minichromosomes of <i>Schizosaccharomyces pombe</i> .. <i>Molecular and Cellular Biology</i> , 1987, 7, 4424-4430. | 1.1 | 84 |
| 315 | Mutational and in vitro protein-binding studies on centromere DNA from <i>Saccharomyces cerevisiae</i> .. <i>Molecular and Cellular Biology</i> , 1987, 7, 4522-4534. | 1.1 | 115 |
| 316 | The GCR1 gene encodes a positive transcriptional regulator of the enolase and glyceraldehyde-3-phosphate dehydrogenase gene families in <i>Saccharomyces cerevisiae</i> .. <i>Molecular and Cellular Biology</i> , 1987, 7, 813-820. | 1.1 | 102 |
| 317 | Cell-type-dependent gene activation by yeast transposon Ty1 involves multiple regulatory determinants.. <i>Molecular and Cellular Biology</i> , 1987, 7, 3205-3211. | 1.1 | 45 |
| 318 | Comparison of tRNA gene transcription complexes formed in vitro and in nuclei.. <i>Molecular and Cellular Biology</i> , 1987, 7, 3212-3220. | 1.1 | 40 |
| 319 | The RAD3 gene of <i>Saccharomyces cerevisiae</i> encodes a DNA-dependent ATPase.. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 1987, 84, 6045-6049. | 3.3 | 137 |
| 320 | Accurate initiation at RNA polymerase II promoters in extracts from <i>Saccharomyces cerevisiae</i> .. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 1987, 84, 8839-8843. | 3.3 | 152 |
| 321 | Constitutive expression of the yeast HEM1 gene is actually a composite of activation and repression.. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 1987, 84, 9113-9117. | 3.3 | 89 |
| 322 | Viruses in fungi: infection of yeast with the K1 and K2 killer viruses.. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 1987, 84, 4293-4297. | 3.3 | 75 |
| 323 | Primary structure requirements for correct sorting of the yeast mitochondrial protein ADH III to the yeast mitochondrial matrix space.. <i>Molecular and Cellular Biology</i> , 1987, 7, 294-304. | 1.1 | 51 |
| 324 | Guanine nucleotide activation of, and competition between, RAS proteins from <i>Saccharomyces cerevisiae</i> .. <i>Molecular and Cellular Biology</i> , 1987, 7, 2128-2133. | 1.1 | 127 |

| # | ARTICLE | IF | CITATIONS |
|-----|---|-----|-----------|
| 325 | Isolation and characterization of temperature-sensitive RNA polymerase II mutants of <i>Saccharomyces cerevisiae</i> .. <i>Molecular and Cellular Biology</i> , 1987, 7, 2155-2164. | 1.1 | 52 |
| 326 | Regulatory function of the <i>Saccharomyces cerevisiae</i> RAS C-terminus.. <i>Molecular and Cellular Biology</i> , 1987, 7, 2309-2315. | 1.1 | 88 |
| 327 | Sequences from a prokaryotic genome or the mouse dihydrofolate reductase gene can restore the import of a truncated precursor protein into yeast mitochondria.. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 1987, 84, 3117-3121. | 3.3 | 172 |
| 328 | The upstream activation site of a Ty2 element of yeast is necessary but not sufficient to promote maximal transcription of the element.. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 1987, 84, 8520-8524. | 3.3 | 68 |
| 329 | Temperature-sensitive mutations in the yeast DNA polymerase I gene.. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 1987, 84, 2838-2842. | 3.3 | 117 |
| 330 | Cloning and characterization of the low-affinity cyclic AMP phosphodiesterase gene of <i>Saccharomyces cerevisiae</i> .. <i>Molecular and Cellular Biology</i> , 1987, 7, 3629-3636. | 1.1 | 213 |
| 331 | Genetic and biochemical characterization of clathrin-deficient <i>Saccharomyces cerevisiae</i> .. <i>Molecular and Cellular Biology</i> , 1987, 7, 3888-3898. | 1.1 | 90 |
| 332 | Trans-acting regulatory mutations that alter transcription of <i>Saccharomyces cerevisiae</i> histone genes.. <i>Molecular and Cellular Biology</i> , 1987, 7, 4204-4210. | 1.1 | 124 |
| 333 | Cloning and characterization of LOS1, a <i>Saccharomyces cerevisiae</i> gene that affects tRNA splicing.. <i>Molecular and Cellular Biology</i> , 1987, 7, 1208-1216. | 1.1 | 115 |
| 334 | Phosphorylation of the <i>Saccharomyces cerevisiae</i> equivalent of ribosomal protein S6 has no detectable effect on growth.. <i>Molecular and Cellular Biology</i> , 1987, 7, 1338-1345. | 1.1 | 63 |
| 335 | The membrane-associated enzyme phosphatidylserine synthase is regulated at the level of mRNA abundance.. <i>Molecular and Cellular Biology</i> , 1987, 7, 167-176. | 1.1 | 129 |
| 336 | Molecular cloning of chromosome I DNA from <i>Saccharomyces cerevisiae</i> : isolation and analysis of the CEN1-ADE1-CDC15 region.. <i>Molecular and Cellular Biology</i> , 1987, 7, 410-419. | 1.1 | 57 |
| 337 | Differential effectiveness of yeast cytochrome c oxidase subunit genes results from differences in expression not function.. <i>Molecular and Cellular Biology</i> , 1987, 7, 3520-3526. | 1.1 | 84 |
| 338 | Insertions of up to 17 amino acids into a region of alpha-tubulin do not disrupt function in vivo.. <i>Molecular and Cellular Biology</i> , 1987, 7, 3799-3805. | 1.1 | 44 |
| 339 | Yeast pre-mRNA splicing requires a minimum distance between the 5' splice site and the internal branch acceptor site.. <i>Molecular and Cellular Biology</i> , 1987, 7, 4010-4016. | 1.1 | 20 |
| 340 | Sequence and nuclear localization of the <i>Saccharomyces cerevisiae</i> HAP2 protein, a transcriptional activator.. <i>Molecular and Cellular Biology</i> , 1987, 7, 578-585. | 1.1 | 159 |
| 341 | Regulation of RAD54- and RAD52-lacZ gene fusions in <i>Saccharomyces cerevisiae</i> in response to DNA damage.. <i>Molecular and Cellular Biology</i> , 1987, 7, 1078-1084. | 1.1 | 142 |
| 342 | Fatty acylation is important but not essential for <i>Saccharomyces cerevisiae</i> RAS function.. <i>Molecular and Cellular Biology</i> , 1987, 7, 2344-2351. | 1.1 | 123 |

| # | ARTICLE | IF | CITATIONS |
|-----|---|-----|-----------|
| 343 | Genetic manipulation of centromere function.. <i>Molecular and Cellular Biology</i> , 1987, 7, 2397-2405. | 1.1 | 206 |
| 344 | <i>Phycomyces blakesleeanus</i> TRP1 gene: organization and functional complementation in <i>Escherichia coli</i> and <i>Saccharomyces cerevisiae</i> .. <i>Molecular and Cellular Biology</i> , 1987, 7, 2664-2670. | 1.1 | 22 |
| 345 | Transcription of the constitutively expressed yeast enolase gene ENO1 is mediated by positive and negative cis-acting regulatory sequences.. <i>Molecular and Cellular Biology</i> , 1987, 7, 2753-2761. | 1.1 | 101 |
| 346 | Codon replacement in the PGK1 gene of <i>Saccharomyces cerevisiae</i> : experimental approach to study the role of biased codon usage in gene expression.. <i>Molecular and Cellular Biology</i> , 1987, 7, 2914-2924. | 1.1 | 198 |
| 347 | Import and processing of human ornithine transcarbamoylase precursor by mitochondria from <i>Saccharomyces cerevisiae</i> .. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 1987, 84, 4063-4067. | 3.3 | 40 |
| 348 | Complete nucleotide sequence of a gene conferring polymyxin B resistance on yeast: similarity of the predicted polypeptide to protein kinases.. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 1987, 84, 5848-5852. | 3.3 | 116 |
| 349 | Cloning and heterologous expression of glycosidase genes from <i>Saccharomyces cerevisiae</i> .. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 1987, 84, 2585-2589. | 3.3 | 115 |
| 350 | Protein glycosylation in yeast: transcript heterogeneity of the ALG7 gene.. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 1987, 84, 2145-2149. | 3.3 | 59 |
| 351 | A segment of GCN4 mRNA containing the upstream AUG codons confers translational control upon a heterologous yeast transcript.. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 1987, 84, 2863-2867. | 3.3 | 96 |
| 352 | Isolation and characterization of MOD5, a gene required for isopentenylation of cytoplasmic and mitochondrial tRNAs of <i>Saccharomyces cerevisiae</i> .. <i>Molecular and Cellular Biology</i> , 1987, 7, 177-184. | 1.1 | 140 |
| 353 | Ty1 sequence with enhancer and mating-type-dependent regulatory activities.. <i>Molecular and Cellular Biology</i> , 1987, 7, 258-265. | 1.1 | 79 |
| 354 | Sequences required for transcriptional initiation of the <i>Saccharomyces cerevisiae</i> CYC7 genes.. <i>Molecular and Cellular Biology</i> , 1987, 7, 3785-3791. | 1.1 | 45 |
| 355 | Role of transcriptional and posttranscriptional regulation in expression of histone genes in <i>Saccharomyces cerevisiae</i> .. <i>Molecular and Cellular Biology</i> , 1987, 7, 614-621. | 1.1 | 84 |
| 356 | RAD4 gene of <i>Saccharomyces cerevisiae</i> : molecular cloning and partial characterization of a gene that is inactivated in <i>Escherichia coli</i> .. <i>Molecular and Cellular Biology</i> , 1987, 7, 1180-1192. | 1.1 | 42 |
| 357 | Effect of deletion and insertion on double-strand-break repair in <i>Saccharomyces cerevisiae</i> .. <i>Molecular and Cellular Biology</i> , 1987, 7, 1300-1303. | 1.1 | 15 |
| 358 | Structure and expression of the <i>Saccharomyces cerevisiae</i> CRY1 gene: a highly conserved ribosomal protein gene.. <i>Molecular and Cellular Biology</i> , 1987, 7, 1764-1775. | 1.1 | 82 |
| 359 | Upstream regulatory regions controlling the expression of the yeast maltase gene.. <i>Molecular and Cellular Biology</i> , 1987, 7, 2477-2483. | 1.1 | 38 |
| 360 | Characterization of <i>Saccharomyces cerevisiae</i> genes encoding subunits of cyclic AMP-dependent protein kinase.. <i>Molecular and Cellular Biology</i> , 1987, 7, 2653-2663. | 1.1 | 273 |

| # | ARTICLE | IF | CITATIONS |
|-----|--|-----|-----------|
| 361 | Expression and secretion of foreign polypeptides in yeast. , 1987, , 401-429. | | 7 |
| 362 | Frameshift suppressor mutations affecting the major glycine transfer RNAs of <i>Saccharomyces cerevisiae</i> . <i>Journal of Molecular Biology</i> , 1987, 194, 41-58. | 2.0 | 39 |
| 363 | Expression of human antithrombin III in <i>Saccharomyces cerevisiae</i> and <i>Schizosaccharomyces pombe</i> . <i>Biochimica Et Biophysica Acta Gene Regulatory Mechanisms</i> , 1987, 908, 203-213. | 2.4 | 62 |
| 364 | [10] 5-Fluoroorotic acid as a selective agent in yeast molecular genetics. <i>Methods in Enzymology</i> , 1987, 154, 164-175. | 0.4 | 1,384 |
| 365 | Specificity of cotranslational amino-terminal processing of proteins in yeast. <i>Biochemistry</i> , 1987, 26, 8242-8246. | 1.2 | 213 |
| 366 | [7] $\hat{\text{t}}$ gt 11: Gene isolation with antibody probes and other applications. <i>Methods in Enzymology</i> , 1987, 154, 107-128. | 0.4 | 161 |
| 367 | Plasmid construction by homologous recombination in yeast. <i>Gene</i> , 1987, 58, 201-216. | 1.0 | 569 |
| 368 | Structural characteristics of the PHO8 gene encoding repressible alkaline phosphatase in <i>Saccharomyces cerevisiae</i> . <i>Gene</i> , 1987, 58, 137-148. | 1.0 | 95 |
| 369 | Efficient secretion of <i>Bacillus amyloliquefaciens</i> $\hat{\text{t}}$ -amylase cells by its own signal peptide from <i>Saccharomyces cerevisiae</i> host. <i>Gene</i> , 1987, 59, 161-170. | 1.0 | 35 |
| 370 | Repair of UV-irradiated plasmid DNA in <i>Saccharomyces cerevisiae</i> Inability to complement mutational defects in excision repair by in vitro treatment with <i>Micrococcus luteus</i> UV endonuclease. <i>Mutation Research - DNA Repair Reports</i> , 1987, 183, 161-167. | 1.9 | 6 |
| 371 | Construction of a hybrid cDNA encoding a major legumin precursor polypeptide and its expression and localization in <i>Saccharomyces cerevisiae</i> . <i>FEBS Letters</i> , 1987, 222, 175-180. | 1.3 | 5 |
| 372 | Tight control of the amount of yeast plasma membrane ATPase during changes in growth conditions and gene dosage. <i>FEBS Letters</i> , 1987, 224, 193-197. | 1.3 | 78 |
| 373 | A method to isolate DNA sequences that are promoter-active in <i>Escherichia coli</i> and in yeast. <i>Biochemical and Biophysical Research Communications</i> , 1987, 149, 846-851. | 1.0 | 5 |
| 374 | Translational regulation of the expression of zein cloned in yeast under an inducible GAL promoter. <i>Biochemical and Biophysical Research Communications</i> , 1987, 146, 809-814. | 1.0 | 9 |
| 375 | Gene disruption indicates that the only essential function of the SK18 chromosomal gene is to protect <i>Saccharomyces cerevisiae</i> from viral cytopathology. <i>Virology</i> , 1987, 157, 252-256. | 1.1 | 32 |
| 376 | Expression of the lacZ gene from two methanol-regulated promoters in <i>Pichia pastoris</i> . <i>Nucleic Acids Research</i> , 1987, 15, 3859-3876. | 6.5 | 327 |
| 377 | Concerted deletions and inversions are caused by mitotic recombination between delta sequences in <i>Saccharomyces cerevisiae</i> .. <i>Molecular and Cellular Biology</i> , 1987, 7, 1198-1207. | 1.1 | 131 |
| 378 | A novel yeast secretion vector utilizing secretion signal of killer toxin encoded on the yeast linear DNA plasmid pGKL1. <i>Biochemical and Biophysical Research Communications</i> , 1987, 144, 613-619. | 1.0 | 39 |

| # | ARTICLE | IF | CITATIONS |
|-----|---|------|-----------|
| 379 | Enriched autonomously replicating sequences in a nuclear matrix-DNA complex isolated from synchronized HeLa cells. <i>Biochemical and Biophysical Research Communications</i> , 1987, 144, 1018-1024. | 1.0 | 15 |
| 380 | Recognition of the TACTAAC box during mRNA splicing in yeast involves base pairing to the U2-like snRNA. <i>Cell</i> , 1987, 49, 229-239. | 13.5 | 519 |
| 381 | Recombination-stimulating sequences in yeast ribosomal DNA correspond to sequences regulating transcription by RNA polymerase I. <i>Cell</i> , 1987, 48, 1071-1079. | 13.5 | 352 |
| 382 | Yeast KEX1 gene encodes a putative protease with a carboxypeptidase B-like function involved in killer toxin and $\hat{\pm}$ -factor precursor processing. <i>Cell</i> , 1987, 50, 573-584. | 13.5 | 199 |
| 383 | A family of yeast expression vectors containing the phage f1 intergenic region1. <i>Gene</i> , 1987, 52, 225-233. | 1.0 | 536 |
| 384 | The carboxy-terminal 30 amino acids of GAL4 are recognized by GAL80. <i>Cell</i> , 1987, 50, 137-142. | 13.5 | 348 |
| 385 | RPC40, a unique gene for a subunit shared between yeast RNA polymerases A and C. <i>Cell</i> , 1987, 48, 627-637. | 13.5 | 199 |
| 386 | Mutants of GAL4 protein altered in an activation function. <i>Cell</i> , 1987, 51, 121-126. | 13.5 | 309 |
| 387 | A ras-like protein is required for a post-Golgi event in yeast secretion. <i>Cell</i> , 1987, 49, 527-538. | 13.5 | 901 |
| 388 | A new class of yeast transcriptional activators. <i>Cell</i> , 1987, 51, 113-119. | 13.5 | 754 |
| 389 | GPA1, a haploid-specific essential gene, encodes a yeast homolog of mammalian G protein which may be involved in mating factor signal transduction. <i>Cell</i> , 1987, 50, 1011-1019. | 13.5 | 400 |
| 390 | The RNA components of <i>Schizosaccharomyces pombe</i> RNase P are essential for cell viability. <i>Gene</i> , 1987, 60, 157-161. | 1.0 | 18 |
| 391 | Expression of a wheat $\hat{\pm}$ -gliadin gene in <i>Saccharomyces cerevisiae</i> . <i>Gene</i> , 1987, 55, 303-317. | 1.0 | 28 |
| 392 | Three different genes in <i>S. cerevisiae</i> encode the catalytic subunits of the cAMP-dependent protein kinase. <i>Cell</i> , 1987, 50, 277-287. | 13.5 | 705 |
| 393 | The <i>S. cerevisiae</i> CDC25 gene product regulates the RAS/adenylate cyclase pathway. <i>Cell</i> , 1987, 48, 789-799. | 13.5 | 523 |
| 394 | Allele-specific complementation of an <i>Escherichia coli</i> leuB mutation by a <i>Lactobacillus bulgaricus</i> tRNA gene. <i>Gene</i> , 1987, 60, 75-83. | 1.0 | 27 |
| 395 | KAR1, a gene required for function of both intranuclear and extranuclear microtubules in yeast. <i>Cell</i> , 1987, 48, 1047-1060. | 13.5 | 404 |
| 396 | Synthesis and secretion of wheat $\hat{\pm}$ -amylase in <i>Saccharomyces cerevisiae</i> . <i>Gene</i> , 1987, 55, 353-356. | 1.0 | 39 |

| # | ARTICLE | IF | CITATIONS |
|-----|---|------|-----------|
| 397 | A yeast silencer contains sequences that can promote autonomous plasmid replication and transcriptional activation. <i>Cell</i> , 1987, 51, 709-719. | 13.5 | 360 |
| 398 | A genetic analysis of dicentric minichromosomes in <i>Saccharomyces cerevisiae</i> . <i>Cell</i> , 1987, 48, 801-812. | 13.5 | 107 |
| 399 | Deletion analysis of GAL4 defines two transcriptional activating segments. <i>Cell</i> , 1987, 48, 847-853. | 13.5 | 976 |
| 400 | An essential snRNA from <i>S. cerevisiae</i> has properties predicted for U4, including interaction with a U6-like snRNA. <i>Cell</i> , 1987, 50, 585-592. | 13.5 | 112 |
| 401 | Expression of human pancreatic secretory trypsin inhibitor in <i>Saccharomyces cerevisiae</i> . <i>Gene</i> , 1987, 59, 151-159. | 1.0 | 15 |
| 402 | Nucleotide sequence of the CLS4 (CDC24) gene of <i>Saccharomyces cerevisiae</i> . <i>Gene</i> , 1987, 54, 125-132. | 1.0 | 93 |
| 403 | Protein sorting in yeast: The localization determinant of yeast vacuolar carboxypeptidase Y resides in the propeptide. <i>Cell</i> , 1987, 48, 887-897. | 13.5 | 344 |
| 404 | A <i>Saccharomyces cerevisiae</i> genomic plasmid bank based on a centromere-containing shuttle vector. <i>Gene</i> , 1987, 60, 237-243. | 1.0 | 1,273 |
| 405 | RRP1, a <i>Saccharomyces cerevisiae</i> gene affecting rRNA processing and production of mature ribosomal subunits. <i>Journal of Bacteriology</i> , 1987, 169, 1571-1578. | 1.0 | 51 |
| 406 | Mutational inactivation of the <i>Saccharomyces cerevisiae</i> RAD4 gene in <i>Escherichia coli</i> . <i>Journal of Bacteriology</i> , 1987, 169, 4884-4892. | 1.0 | 16 |
| 407 | Common signal transduction system shared by <i>STE2</i> and <i>STE3</i> in haploid cells of <i>Saccharomyces cerevisiae</i> : autocrine cell-cycle arrest results from forced expression of <i>STE2</i> . <i>EMBO Journal</i> , 1987, 6, 249-254. | 3.5 | 77 |
| 408 | Gene-enzyme relationships in the proline biosynthetic pathway of <i>Saccharomyces cerevisiae</i> . <i>Journal of Bacteriology</i> , 1987, 169, 5364-5372. | 1.0 | 71 |
| 409 | Efficient splicing of two yeast mitochondrial introns controlled by a nuclear-encoded maturase.. <i>EMBO Journal</i> , 1987, 6, 1085-1091. | 3.5 | 53 |
| 410 | Elements involved in oxygen regulation of the <i>Saccharomyces cerevisiae</i> CYC7 gene.. <i>Molecular and Cellular Biology</i> , 1987, 7, 2212-2220. | 1.1 | 91 |
| 411 | MKT1, a nonessential <i>Saccharomyces cerevisiae</i> gene with a temperature-dependent effect on replication of M2 double-stranded RNA. <i>Journal of Bacteriology</i> , 1987, 169, 4941-4945. | 1.0 | 37 |
| 412 | Specific accessory sequences in <i>Saccharomyces cerevisiae</i> introns control assembly of pre-mRNAs into spliceosomes.. <i>EMBO Journal</i> , 1987, 6, 3833-3839. | 3.5 | 44 |
| 413 | Cloning and characterization of <i>Saccharomyces cerevisiae</i> genes that confer L-methionine sulfoximine and tabtoxin resistance. <i>Journal of Bacteriology</i> , 1987, 169, 2440-2448. | 1.0 | 18 |
| 414 | Nuclear pre-mRNA splicing in the fission yeast <i>Schizosaccharomyces pombe</i> strictly requires an intron-contained, conserved sequence element.. <i>EMBO Journal</i> , 1987, 6, 1757-1763. | 3.5 | 60 |

| # | ARTICLE | IF | CITATIONS |
|-----|--|-----|-----------|
| 415 | Transport of proteins to the mitochondrial intermembrane space: the "matrix-targeting"™ and the "sorting"™ domains in the cytochrome c1 presequence.. EMBO Journal, 1987, 6, 2433-2439. | 3.5 | 79 |
| 416 | Three suppressor mutations which cure a mitochondrial RNA maturase deficiency occur at the same codon in the open reading frame of the nuclear NAM2 gene.. EMBO Journal, 1987, 6, 713-721. | 3.5 | 67 |
| 417 | Independent mutations at the amino terminus of a protein act as surrogate signals for mitochondrial import.. EMBO Journal, 1987, 6, 705-711. | 3.5 | 56 |
| 418 | The free sulfhydryl group (Cys341) of carboxypeptidase Y: Functional effects of mutational substitutions. Carlsberg Research Communications, 1987, 52, 263-273. | 1.7 | 17 |
| 419 | Yeast cells may use AUC or AAG as initiation codon for protein synthesis. Carlsberg Research Communications, 1987, 52, 83-90. | 1.7 | 5 |
| 420 | Analysis of the effect of dGÂ·dC homopolymer tails on expression of a mouse Î±-amylase cDNA gene in yeast. Carlsberg Research Communications, 1987, 52, 91-97. | 1.7 | 8 |
| 421 | Identification of the domain of Saccharomyces cerevisiae adenylate cyclase associated with the regulatory function of RAS products. Molecular Genetics and Genomics, 1987, 210, 187-194. | 2.4 | 36 |
| 422 | In addition to the ARS core, the ARS box is necessary for autonomously replicating sequences. Molecular Genetics and Genomics, 1987, 206, 60-65. | 2.4 | 21 |
| 423 | Mutant regulatory subunit of 3â€²,5â€²-cAMP-dependent protein kinase of yeast Saccharomyces cerevisiae. Molecular Genetics and Genomics, 1987, 210, 413-418. | 2.4 | 19 |
| 424 | Transcriptional regulation of sporulation genes in yeast. Molecular Genetics and Genomics, 1987, 210, 449-459. | 2.4 | 22 |
| 425 | Secretion of Mucor rennin, a fungal aspartic protease of Mucor pusillus, by recombinant yeast cells. Molecular Genetics and Genomics, 1987, 210, 462-467. | 2.4 | 47 |
| 426 | Strains of Schizosaccharomyces pombe with a disrupted swi1 gene still show some mating-type switching. Molecular Genetics and Genomics, 1987, 210, 485-489. | 2.4 | 10 |
| 427 | Mutations that increase the mitotic stability of minichromosomes in yeast: Characterization of RAR1. Molecular Genetics and Genomics, 1987, 210, 509-517. | 2.4 | 31 |
| 428 | Isolation and expression analysis of two yeast regulatory genes involved in the derepression of glucose-repressible enzymes. Molecular Genetics and Genomics, 1987, 209, 366-373. | 2.4 | 79 |
| 429 | Structure of the yeast HIS5 gene responsive to general control of amino acid biosynthesis. Molecular Genetics and Genomics, 1987, 208, 159-167. | 2.4 | 67 |
| 430 | The RAD3 gene of Saccharomyces cerevisiae: Isolation and characterization of a temperature-sensitive mutant in the essential function and of extragenic suppressors of this mutant. Molecular Genetics and Genomics, 1987, 209, 458-466. | 2.4 | 27 |
| 431 | Regulation of MAL gene expression in yeast: Gene dosage effects. Molecular Genetics and Genomics, 1987, 209, 508-517. | 2.4 | 43 |
| 432 | Phleomycin resistance encoded by the ble gene from transposon Tn 5 as a dominant selectable marker in Saccharomyces cerevisiae. Molecular Genetics and Genomics, 1987, 207, 342-348. | 2.4 | 80 |

| # | ARTICLE | IF | CITATIONS |
|-----|---|-----|-----------|
| 433 | Curing of endogenous 2 micron DNA in yeast by recombinant vectors. <i>Current Genetics</i> , 1987, 11, 315-319. | 0.8 | 22 |
| 434 | Induced cellular resistance to ultraviolet light in <i>Saccharomyces cerevisiae</i> is not accompanied by increased repair of plasmid DNA. <i>Current Genetics</i> , 1987, 11, 321-326. | 0.8 | 8 |
| 435 | ARS activity along the yeast mitochondrial apocytochrome b region: correlation with the location of petite genomes and consensus sequences. <i>Current Genetics</i> , 1987, 12, 583-589. | 0.8 | 3 |
| 436 | Sequence of the bifunctional <i>ade1</i> gene in the purine biosynthetic pathway of the fission yeast <i>Schizosaccharomyces pombe</i> . <i>Current Genetics</i> , 1987, 12, 591-597. | 0.8 | 17 |
| 437 | Phenotypic differences between induced and spontaneous 2?-plasmid-free segregants of <i>Saccharomyces cerevisiae</i> . <i>Current Genetics</i> , 1987, 11, 415-418. | 0.8 | 9 |
| 438 | The stability of chromosomes in yeast. <i>Current Genetics</i> , 1987, 11, 435-443. | 0.8 | 15 |
| 439 | Genetic mapping of two pairs of linked ribosomal protein genes in <i>Saccharomyces cerevisiae</i> . <i>Current Genetics</i> , 1987, 11, 445-450. | 0.8 | 11 |
| 440 | Nucleotide sequencing analysis of a LEU gene of <i>Candida maltosa</i> which complements <i>leuB</i> mutation of <i>Escherichia coli</i> and <i>leu2</i> mutation of <i>Saccharomyces cerevisiae</i> . <i>Current Genetics</i> , 1987, 11, 451-457. | 0.8 | 42 |
| 441 | Studies on stability of miniplasmids comprised of only yeast DNA. <i>Current Genetics</i> , 1987, 12, 15-20. | 0.8 | 6 |
| 442 | Isolation and characterization of the regulatory <i>HEX2</i> gene necessary for glucose repression in yeast. <i>Molecular Genetics and Genomics</i> , 1987, 206, 505-509. | 2.4 | 60 |
| 443 | Terminal segment of <i>Kluyveromyces lactis</i> linear DNA plasmid pGKL2 supports autonomous replication of hybrid plasmids in <i>Saccharomyces cerevisiae</i> . <i>Current Genetics</i> , 1987, 12, 99-104. | 0.8 | 25 |
| 444 | Replacement of the promoter of the yeast plasma membrane ATPase gene by a galactose-dependent promoter and its physiological consequences. <i>Current Genetics</i> , 1987, 12, 105-110. | 0.8 | 129 |
| 445 | The plasmids of the morels: characterization and prerequisites for vector development. <i>Applied Microbiology and Biotechnology</i> , 1987, 27, 276. | 1.7 | 11 |
| 446 | Transformation of <i>Penicillium chrysogenum</i> by a vector containing a mitochondrial origin of replication. <i>Applied Microbiology and Biotechnology</i> , 1987, 26, 237-241. | 1.7 | 31 |
| 447 | PHO5 Upstream sequences confer phosphate control on the constitutive PHO3 gene. <i>Yeast</i> , 1987, 3, 33-42. | 0.8 | 9 |
| 448 | Molecular cloning of chromosome I DNA from <i>Saccharomyces cerevisiae</i> : Isolation of the <i>MAK16</i> gene and analysis of an adjacent gene essential for growth at low temperatures. <i>Yeast</i> , 1987, 3, 51-57. | 0.8 | 76 |
| 449 | Specificity of DNA uptake during whole cell transformation of <i>S. cerevisiae</i> . <i>Yeast</i> , 1987, 3, 131-137. | 0.8 | 43 |
| 450 | Overexpression of the <i>RAD2</i> gene of <i>S. cerevisiae</i> : Identification and preliminary characterization of Rad2 protein. <i>Yeast</i> , 1987, 3, 149-160. | 0.8 | 24 |

| # | ARTICLE | IF | CITATIONS |
|-----|--|-----|-----------|
| 451 | Transformation of yeast spheroplasts without cell fusion. <i>Analytical Biochemistry</i> , 1987, 163, 391-397. | 1.1 | 288 |
| 452 | Expression of the <i>Klebsiella pneumoniae</i> <i>nifH</i> gene in <i>Saccharomyces cerevisiae</i> . <i>Enzyme and Microbial Technology</i> , 1987, 9, 438-443. | 1.6 | 0 |
| 453 | Identification of sequence elements that confer cell-type-specific control of MF alpha 1 expression in <i>Saccharomyces cerevisiae</i> . <i>Molecular and Cellular Biology</i> , 1987, 7, 3185-3193. | 1.1 | 43 |
| 454 | Development of a yeast system to assay mutational specificity. <i>Mutation Research - Environmental Mutagenesis and Related Subjects Including Methodology</i> , 1987, 182, 65-74. | 0.4 | 58 |
| 455 | Nitrogen catabolite regulation of proline permease in <i>Saccharomyces cerevisiae</i> . Cloning of the PUT4 gene and study of PUT4 RNA levels in wild-type and mutant strains. <i>FEBS Journal</i> , 1987, 164, 601-606. | 0.2 | 120 |
| 456 | Isolation of the NPRI gene responsible for the reactivation of ammonia-sensitive amino-acid permeases in <i>Saccharomyces cerevisiae</i> . RNA analysis and gene dosage effects. <i>FEBS Journal</i> , 1987, 164, 607-612. | 0.2 | 61 |
| 457 | Aminopeptidase <i>yscII</i> of yeast. Isolation of mutants and their biochemical and genetic analysis. <i>FEBS Journal</i> , 1988, 173, 589-598. | 0.2 | 28 |
| 458 | Probing the active site of flavocytochrome b2 by site-directed mutagenesis. <i>FEBS Journal</i> , 1988, 178, 329-333. | 0.2 | 46 |
| 459 | Genetic engineering in <i>Yarrowia lipolytica</i> . <i>Journal of Basic Microbiology</i> , 1988, 28, 161-174. | 1.8 | 17 |
| 460 | Replication and recombination in gene establishment in non- <i>Saccharomyces</i> yeasts. <i>Journal of Basic Microbiology</i> , 1988, 28, 197-208. | 1.8 | 2 |
| 461 | A gene-cloning system for <i>Kluyveromyces lactis</i> and isolation of a chromosomal gene required for killer toxin production. <i>Journal of Basic Microbiology</i> , 1988, 28, 211-220. | 1.8 | 28 |
| 462 | Transformation of methylotrophic yeast <i>Hansenula polymorpha</i> : Cloning and expression of genes. <i>Journal of Basic Microbiology</i> , 1988, 28, 343-351. | 1.8 | 23 |
| 463 | Stability and expression of a plasmid-containing killer toxin cDNA in batch and chemostat cultures of <i>Saccharomyces cerevisiae</i> . <i>Biotechnology and Bioengineering</i> , 1988, 31, 783-789. | 1.7 | 15 |
| 464 | Functional analysis of mitochondrial protein import in yeast. <i>Journal of Cellular Biochemistry</i> , 1988, 36, 275-287. | 1.2 | 19 |
| 465 | DNA sequence polymorphisms in the genus <i>Saccharomyces</i> . V. Cloning and characterization of a <i>LEU2</i> gene from <i>S. carlsbergensis</i> . <i>Carlsberg Research Communications</i> , 1988, 53, 209-219. | 1.7 | 23 |
| 466 | Cloning, sequencing and analysis of the yeast <i>S. uvarum</i> <i>ERG10</i> gene encoding acetoacetyl CoA thiolase. <i>Current Genetics</i> , 1988, 13, 471-478. | 0.8 | 36 |
| 467 | Effect of acetoacetyl coa thiolase amplification on sterol synthesis in the yeasts <i>S. cerevisiae</i> and <i>S. uvarum</i> . <i>Biotechnology Letters</i> , 1988, 10, 457-462. | 1.1 | 6 |
| 468 | Construction of a promoter-probe vector with the <i>PHO5</i> gene encoding repressible acid phosphatase in <i>Saccharomyces cerevisiae</i> . <i>Applied Microbiology and Biotechnology</i> , 1988, 28, 155-159. | 1.7 | 5 |

| # | ARTICLE | IF | CITATIONS |
|-----|--|-----|-----------|
| 469 | Physiological and genetic modulation of inducible expression of Escherichia coli β -galactosidase in Saccharomyces cerevisiae. Applied Microbiology and Biotechnology, 1988, 28, 160-165. | 1.7 | 14 |
| 470 | Genomic organization of two families of highly repeated nuclear DNA sequences of maize selected for autonomous replicating activity in yeast. Plant Molecular Biology, 1988, 11, 161-172. | 2.0 | 5 |
| 471 | Liposome-mediated introduction of the chloramphenicol acetyl transferase (CAT) gene and its expression in tobacco protoplasts. Plant Molecular Biology, 1988, 10, 185-191. | 2.0 | 14 |
| 472 | The Escherichia coli proB gene corrects the proline auxotrophy of Saccharomyces cerevisiae pro1 mutants. Molecular Genetics and Genomics, 1988, 212, 124-128. | 2.4 | 12 |
| 473 | Genetic characterization of hyperresistance to formaldehyde and 4-nitroquinoline-N-oxide in the yeast Saccharomyces cerevisiae. Molecular Genetics and Genomics, 1988, 211, 260-265. | 2.4 | 20 |
| 474 | The 2 $\frac{1}{4}$ m D region plays a role in yeast plasmid maintenance. Molecular Genetics and Genomics, 1988, 212, 426-431. | 2.4 | 20 |
| 475 | Yeast ARS function and nuclear matrix association coincide in a short sequence from the human HPRT locus. Molecular Genetics and Genomics, 1988, 212, 301-309. | 2.4 | 71 |
| 476 | A nuclear gene essential for mitochondrial replication suppresses a defect of mitochondrial transcription in Saccharomyces cerevisiae. Molecular Genetics and Genomics, 1988, 214, 218-223. | 2.4 | 90 |
| 477 | Divergence of the mitochondrial leucyl tRNA synthetase genes in two closely related yeasts Saccharomyces cerevisiae and Saccharomyces douglasii: A paradigm of incipient evolution. Molecular Genetics and Genomics, 1988, 213, 297-309. | 2.4 | 52 |
| 478 | Retention of autonomous replicating plasmids in cultured Drosophila cells. Molecular Genetics and Genomics, 1988, 213, 359-363. | 2.4 | 9 |
| 479 | Precise excision of bacterial transposon Tn 5 in yeast. Molecular Genetics and Genomics, 1988, 213, 388-393. | 2.4 | 21 |
| 480 | Repair of double-strand breaks in plasmid DNA in the yeast Saccharomyces cerevisiae. Molecular Genetics and Genomics, 1988, 213, 421-424. | 2.4 | 24 |
| 481 | Expression of MF1 in MATa cells supersensitive to $\hat{\pm}$ -factor leads to self-arrest. Molecular Genetics and Genomics, 1988, 214, 85-88. | 2.4 | 24 |
| 482 | A downstream activator sequence regulates the expression of the yeast transposon. Current Genetics, 1988, 13, 357-362. | 0.8 | 5 |
| 483 | MAL64 c is a global regulator of β -glucoside fermentation: identification of a new gene involved in melezitose fermentation. Current Genetics, 1988, 13, 369-375. | 0.8 | 13 |
| 484 | A colony procedure for transformation of Saccharomyces cerevisiae. Current Genetics, 1988, 13, 21-23. | 0.8 | 78 |
| 485 | Primary structure of the ribosomal protein gene S6 from Schizosaccharomyces pombe. Current Genetics, 1988, 13, 57-63. | 0.8 | 22 |
| 486 | Cloning and expression in Saccharomyces cerevisiae of an endo- β -glucanase gene from a thermophilic cellulolytic anaerobe. Applied Microbiology and Biotechnology, 1988, 28, 57. | 1.7 | 12 |

| # | ARTICLE | IF | CITATIONS |
|-----|--|------|-----------|
| 487 | Identification of an upstream activation site in the pyruvate decarboxylase structural gene (PDC1) of <i>Saccharomyces cerevisiae</i> . <i>Current Genetics</i> , 1988, 14, 405-412. | 0.8 | 43 |
| 488 | Cosmids carrying <i>Aspergillus terreus</i> DNA can integrate into <i>Saccharomyces cerevisiae</i> chromosome XII via recombination between yeast and foreign DNAs. <i>Current Genetics</i> , 1988, 14, 183-189. | 0.8 | 1 |
| 489 | MAL63 codes for a positive regulator of maltose fermentation in <i>Saccharomyces cerevisiae</i> . <i>Current Genetics</i> , 1988, 14, 201-209. | 0.8 | 44 |
| 490 | Comparison of <i>Tetrahymena</i> ARS sequence function in the yeasts <i>Saccharomyces cerevisiae</i> and <i>Schizosaccharomyces pombe</i> . <i>Current Genetics</i> , 1988, 14, 225-233. | 0.8 | 7 |
| 491 | Regulation of isoleucine-valine biosynthesis in <i>Saccharomyces cerevisiae</i> . <i>Current Genetics</i> , 1988, 13, 207-217. | 0.8 | 58 |
| 492 | Cloning of the LYS5 gene encoding saccharopine dehydrogenase from the yeast <i>Yarrowia lipolytica</i> by target integration. <i>Current Genetics</i> , 1988, 14, 15-21. | 0.8 | 126 |
| 493 | Construction and characterization of a haploid strain of <i>Saccharomyces cerevisiae</i> that completely lacks all genomic CYH2 sequences. <i>Current Genetics</i> , 1988, 14, 325-329. | 0.8 | 2 |
| 494 | Multiple elements regulate expression of the cell cycle-regulated thymidylate synthase gene of <i>Saccharomyces cerevisiae</i> . <i>Current Genetics</i> , 1988, 14, 363-373. | 0.8 | 9 |
| 495 | The primary structure of the leu1 + gene of <i>Schizosaccharomyces pombe</i> . <i>Current Genetics</i> , 1988, 14, 375-379. | 0.8 | 63 |
| 496 | Plasmid associations with residual nuclear structures in <i>Saccharomyces cerevisiae</i> . <i>Current Genetics</i> , 1988, 13, 291-297. | 0.8 | 7 |
| 497 | Cloning and biochemical characterization of LYS5 gene of <i>Saccharomyces cerevisiae</i> . <i>Current Genetics</i> , 1988, 13, 299-304. | 0.8 | 13 |
| 498 | A non-conserved sequence in the 5' region of the CYH2 intron from <i>Saccharomyces cerevisiae</i> controls splicing efficiency of the pre-mRNA. <i>Yeast</i> , 1988, 4, 209-217. | 0.8 | 8 |
| 499 | Genetic control of chromosome stability in the yeast <i>Saccharomyces cerevisiae</i> . <i>Yeast</i> , 1988, 4, 257-269. | 0.8 | 37 |
| 500 | Defective <i>Escherichia coli</i> signal peptides function in yeast. <i>Molecular Microbiology</i> , 1988, 2, 209-217. | 1.2 | 13 |
| 501 | Hepatitis B Surface Antigen as Carrier Matrix for the Repetitive Epitope of the Circumsporozoite Protein of <i>Plasmodium falciparum</i> . <i>Nature Biotechnology</i> , 1988, 6, 1065-1070. | 9.4 | 55 |
| 502 | Expression and Secretion in <i>S. Cerevisiae</i> of Biologically Active Leech Hirudin. <i>Nature Biotechnology</i> , 1988, 6, 72-77. | 9.4 | 58 |
| 503 | The human oestrogen receptor functions in yeast. <i>Nature</i> , 1988, 334, 31-36. | 13.7 | 430 |
| 504 | Spliceosomal RNA U6 is remarkably conserved from yeast to mammals. <i>Nature</i> , 1988, 334, 213-218. | 13.7 | 413 |

| # | ARTICLE | IF | CITATIONS |
|-----|---|------|-----------|
| 505 | Negative effect of the transcriptional activator GAL4. <i>Nature</i> , 1988, 334, 721-724. | 13.7 | 687 |
| 506 | A rapid preparation of plasmid DNA from <i>Saccharomyces cerevisiae</i> . <i>Folia Microbiologica</i> , 1988, 33, 34-37. | 1.1 | 4 |
| 507 | Expression of synthetic thaumatin genes in yeast. <i>Biochemistry</i> , 1988, 27, 5101-5107. | 1.2 | 62 |
| 508 | The <i>Giardia lamblia</i> trophozoite contains sets of closely related chromosomes. <i>Nucleic Acids Research</i> , 1988, 16, 4555-4567. | 6.5 | 100 |
| 509 | Processing of preproglycinin expressed from cDNA-encoding A1aB1b subunit in <i>Saccharomyces cerevisiae</i> . <i>FEBS Letters</i> , 1988, 233, 273-276. | 1.3 | 11 |
| 510 | Regulation of yeast fructose-1,6-bisphosphatase in strains containing multicopy plasmids coding for this enzyme. <i>FEBS Letters</i> , 1988, 242, 149-152. | 1.3 | 27 |
| 511 | Molecular cloning of a gene enhancing yeast glyoxalase I activity and its effect on yeast cell size. <i>Journal of Fermentation Technology</i> , 1988, 66, 495-500. | 0.6 | 8 |
| 512 | Strain improvement in industrial microorganisms by recombinant DNA techniques. , 1988, , 129-168. | | 4 |
| 513 | Nonconventional Yeasts: Their Genetics and Biotechnological Applications. <i>Critical Reviews in Biotechnology</i> , 1988, 7, 281-337. | 5.1 | 13 |
| 514 | DNA sequence analysis of the <i>ade6</i> gene of <i>Schizosaccharomyces pombe</i> . <i>Journal of Molecular Biology</i> , 1988, 204, 917-925. | 2.0 | 193 |
| 515 | Three ribonucleases H and a reverse transcriptase from the yeast, <i>Saccharomyces cerevisiae</i> . <i>Biochimica Et Biophysica Acta Gene Regulatory Mechanisms</i> , 1988, 951, 322-329. | 2.4 | 9 |
| 516 | Autoregulation of 27m circle gene expression provides a model for maintenance of stable plasmid copy levels. <i>Cell</i> , 1988, 52, 27-37. | 13.5 | 132 |
| 517 | New yeast- <i>Escherichia coli</i> shuttle vectors constructed with in vitro mutagenized yeast genes lacking six-base pair restriction sites. <i>Gene</i> , 1988, 74, 527-534. | 1.0 | 2,923 |
| 518 | A GTP-binding protein required for secretion rapidly associates with secretory vesicles and the plasma membrane in yeast. <i>Cell</i> , 1988, 53, 753-768. | 13.5 | 604 |
| 519 | IME1, a positive regulator gene of meiosis in <i>S. cerevisiae</i> . <i>Cell</i> , 1988, 52, 853-862. | 13.5 | 316 |
| 520 | The C-terminus of the <i>S. cerevisiae</i> $\hat{\pm}$ -pheromone receptor mediates an adaptive response to pheromone. <i>Cell</i> , 1988, 54, 609-620. | 13.5 | 237 |
| 521 | Stability of recombinant plasmids in yeast. <i>Journal of Biotechnology</i> , 1988, 8, 173-192. | 1.9 | 35 |
| 522 | Testing for intron function in the essential <i>Saccharomyces cerevisiae</i> tRNA ^{Ser} UCC gene. <i>Journal of Molecular Biology</i> , 1988, 202, 667-672. | 2.0 | 10 |

| # | ARTICLE | IF | CITATIONS |
|-----|--|------|-----------|
| 523 | Chromosomal ARS and CEN elements bind specifically to the yeast nuclear scaffold. <i>Cell</i> , 1988, 54, 967-978. | 13.5 | 282 |
| 524 | Yeast KEX2 gene encodes an endopeptidase homologous to subtilisin-like serine proteases. <i>Biochemical and Biophysical Research Communications</i> , 1988, 156, 246-254. | 1.0 | 260 |
| 525 | Expression in yeast of a <i>Bacillus</i> alpha-amylase gene by the ADH1 promoter. <i>Journal of Biotechnology</i> , 1988, 7, 33-47. | 1.9 | 9 |
| 526 | Converting a eukaryotic transcriptional inhibitor into an activator. <i>Cell</i> , 1988, 55, 443-446. | 13.5 | 139 |
| 527 | Statistical positioning of nucleosomes by specific protein-binding to an upstream activating sequence in yeast. <i>Journal of Molecular Biology</i> , 1988, 204, 109-127. | 2.0 | 270 |
| 528 | Structure of yeast glucokinase, a strongly diverged specific aldohexose-phosphorylating isoenzyme. <i>Gene</i> , 1988, 73, 141-152. | 1.0 | 58 |
| 529 | Transcription-dependent DNA supercoiling in yeast DNA topoisomerase mutants. <i>Cell</i> , 1988, 54, 403-411. | 13.5 | 291 |
| 530 | Transcriptional activation by the SV40 AP-1 recognition element in yeast is mediated by a factor similar to AP-1 that is distinct from GCN4. <i>Cell</i> , 1988, 53, 321-330. | 13.5 | 251 |
| 531 | Cloning and characterization of the CYC8 gene mediating glucose repression in yeast. <i>Gene</i> , 1988, 73, 97-111. | 1.0 | 96 |
| 532 | Molecular characterization of yeast regulatory gene CAT3 necessary for glucose derepression and nuclear localization of its product. <i>Gene</i> , 1988, 67, 247-257. | 1.0 | 81 |
| 533 | In vivo functional characterization of a yeast nucleotide sequence: construction of a mini-Mu derivative adapted to yeast. <i>Gene</i> , 1988, 62, 45-54. | 1.0 | 26 |
| 534 | Copy number and stability of yeast 2 μ -based plasmids carrying a transcription-conditional centromere. <i>Gene</i> , 1988, 67, 59-68. | 1.0 | 10 |
| 535 | Cloning and sequence determination of the gene encoding the largest subunit of the fission yeast <i>Schizosaccharomyces pombe</i> RNA polymerase I. <i>Gene</i> , 1988, 74, 503-515. | 1.0 | 25 |
| 536 | Meiotic recombination within the centromere of a yeast chromosome. <i>Cell</i> , 1988, 52, 237-240. | 13.5 | 29 |
| 537 | Cloning and analysis of a yeast genomic DNA sequence capable of directing gene transcription in <i>Escherichia coli</i> as well as in yeast. <i>Gene</i> , 1988, 64, 165-172. | 1.0 | 4 |
| 538 | Isolation of the gene encoding the <i>S. cerevisiae</i> heat shock transcription factor. <i>Cell</i> , 1988, 54, 841-853. | 13.5 | 426 |
| 539 | The carboxy-terminal segment of the yeast $\hat{1}$ -factor receptor is a regulatory domain. <i>Cell</i> , 1988, 55, 221-234. | 13.5 | 329 |
| 540 | SPA1: A gene important for chromosome segregation and other mitotic functions in <i>S. cerevisiae</i> . <i>Cell</i> , 1988, 54, 743-754. | 13.5 | 72 |

| # | ARTICLE | IF | CITATIONS |
|-----|---|------|-----------|
| 541 | DNA-bound Fos proteins activate transcription in yeast. <i>Cell</i> , 1988, 52, 179-184. | 13.5 | 238 |
| 542 | Mutations at a Zn(II) finger motif in the yeast eIF-2 ¹² gene alter ribosomal start-site selection during the scanning process. <i>Cell</i> , 1988, 54, 621-632. | 13.5 | 297 |
| 543 | Study of a temperature-sensitive mutant of the ras-related YPT1 gene product in yeast suggests a role in the regulation of intracellular calcium. <i>Cell</i> , 1988, 53, 635-647. | 13.5 | 263 |
| 544 | SUF12 suppressor protein of yeast. <i>Journal of Molecular Biology</i> , 1988, 199, 559-573. | 2.0 | 133 |
| 545 | Retrovirus-like vectors for <i>Saccharomyces cerevisiae</i> : integration of foreign genes controlled by efficient promoters into yeast chromosomal DNA. <i>Gene</i> , 1988, 67, 259-269. | 1.0 | 23 |
| 546 | The ease of DNA unwinding as a determinant of initiation at yeast replication origins. <i>Cell</i> , 1988, 52, 559-567. | 13.5 | 208 |
| 547 | A series of shuttle vectors using chloramphenicol acetyltransferase as a reporter enzyme in yeast. <i>Gene</i> , 1988, 67, 287-294. | 1.0 | 16 |
| 548 | A membrane glycoprotein, Sec12p, required for protein transport from the endoplasmic reticulum to the Golgi apparatus in yeast.. <i>Journal of Cell Biology</i> , 1988, 107, 851-863. | 2.3 | 267 |
| 549 | Gene transfer in industrial <i>Saccharomyces</i> yeasts. <i>Food Biotechnology</i> , 1988, 2, 1-41. | 0.6 | 17 |
| 550 | DNA-mediated Transformation in the Aquatic Filamentous Fungus <i>Achlya ambisexualis</i> . <i>Microbiology (United Kingdom)</i> , 1988, 134, 2019-2028. | 0.7 | 2 |
| 551 | The RAD6 protein of <i>Saccharomyces cerevisiae</i> polyubiquitinates histones, and its acidic domain mediates this activity.. <i>Genes and Development</i> , 1988, 2, 1476-1485. | 2.7 | 199 |
| 552 | Depletion of <i>Saccharomyces cerevisiae</i> ribosomal protein L16 causes a decrease in 60S ribosomal subunits and formation of half-mer polyribosomes.. <i>Genes and Development</i> , 1988, 2, 160-172. | 2.7 | 154 |
| 553 | SCH9, a gene of <i>Saccharomyces cerevisiae</i> that encodes a protein distinct from, but functionally and structurally related to, cAMP-dependent protein kinase catalytic subunits.. <i>Genes and Development</i> , 1988, 2, 517-527. | 2.7 | 199 |
| 554 | A transcriptional barrier to expression of cloned toxin genes of the linear plasmid k1 of <i>Kluyveromyces lactis</i> : evidence that native k1 has novel promoters. <i>Nucleic Acids Research</i> , 1988, 16, 7333-7350. | 6.5 | 67 |
| 555 | SEC11 is required for signal peptide processing and yeast cell growth.. <i>Journal of Cell Biology</i> , 1988, 106, 1035-1042. | 2.3 | 192 |
| 556 | Changes in histone gene dosage alter transcription in yeast.. <i>Genes and Development</i> , 1988, 2, 150-159. | 2.7 | 308 |
| 557 | Mammalian glucocorticoid receptor derivatives enhance transcription in yeast. <i>Science</i> , 1988, 241, 965-967. | 6.0 | 336 |
| 558 | A dominant trifluoperazine resistance gene from <i>Saccharomyces cerevisiae</i> has homology with FOF1 ATP synthase and confers calcium-sensitive growth.. <i>Molecular and Cellular Biology</i> , 1988, 8, 3094-3103. | 1.1 | 88 |

| # | ARTICLE | IF | CITATIONS |
|-----|---|-----|-----------|
| 559 | tRNAi(met) functions in directing the scanning ribosome to the start site of translation. <i>Science</i> , 1988, 242, 93-97. | 6.0 | 195 |
| 560 | Impairment of yeast pre-mRNA splicing by potential secondary structure-forming sequences near the conserved branchpoint sequence. <i>Nucleic Acids Research</i> , 1988, 16, 10413-10423. | 6.5 | 25 |
| 561 | Structure, evolution and properties of a novel repetitive DNA family in <i>Caenorhabditis elegans</i> . <i>Nucleic Acids Research</i> , 1988, 16, 8213-8231. | 6.5 | 20 |
| 562 | ACE1 regulates expression of the <i>Saccharomyces cerevisiae</i> metallothionein gene.. <i>Molecular and Cellular Biology</i> , 1988, 8, 2745-2752. | 1.1 | 305 |
| 563 | An ARS element from <i>Drosophila melanogaster</i> telomeres contains the yeast ARS core and bent replication enhancer. <i>Nucleic Acids Research</i> , 1988, 16, 1169-1180. | 6.5 | 11 |
| 564 | Foreign transcriptional enhancers in yeast. I. Interactions of papovavirus transcriptional enhancers and a quiescent pseudopromoter on supercoiled plasmids. <i>Nucleic Acids Research</i> , 1988, 16, 8847-8868. | 6.5 | 7 |
| 565 | Sequence-directed bends of DNA helix axis at the upstream activation sites of $\hat{\pm}$ -cell-specific genes in yeast. <i>Nucleic Acids Research</i> , 1988, 16, 6693-6711. | 6.5 | 28 |
| 566 | RAD1, an excision repair gene of <i>Saccharomyces cerevisiae</i> , is also involved in recombination.. <i>Molecular and Cellular Biology</i> , 1988, 8, 3619-3626. | 1.1 | 250 |
| 567 | A novel yeast secretion signal isolated from 28K killer precursor protein encoded on the linear DNA plasmid pGKL1. <i>Nucleic Acids Research</i> , 1988, 16, 7499-7511. | 6.5 | 28 |
| 568 | 5' splice site selection in yeast: genetic alterations in base-pairing with U1 reveal additional requirements.. <i>Genes and Development</i> , 1988, 2, 1258-1267. | 2.7 | 294 |
| 569 | Truncation of the carboxy-terminal domain of yeast beta-tubulin causes temperature-sensitive growth and hypersensitivity to antimetabolic drugs [published erratum appears in <i>J Cell Biol</i> 1989 Feb;108(2):747]. <i>Journal of Cell Biology</i> , 1988, 107, 1427-1435. | 2.3 | 36 |
| 570 | A temperature-sensitive mutation of the <i>Schizosaccharomyces pombe</i> gene <i>nuc2+</i> that encodes a nuclear scaffold-like protein blocks spindle elongation in mitotic anaphase.. <i>Journal of Cell Biology</i> , 1988, 106, 1171-1183. | 2.3 | 238 |
| 571 | Structure and expression of the PHO80 gene of <i>Saccharomyces cerevisiae</i> . <i>Nucleic Acids Research</i> , 1988, 16, 2625-2637. | 6.5 | 34 |
| 572 | Ribosomal protein synthesis is not regulated at the translational level in <i>Saccharomyces cerevisiae</i> : balanced accumulation of ribosomal proteins L16 and rp59 is mediated by turnover of excess protein.. <i>Genes and Development</i> , 1988, 2, 664-676. | 2.7 | 100 |
| 573 | Mutations that alter both localization and production of a yeast nuclear protein.. <i>Genes and Development</i> , 1988, 2, 707-717. | 2.7 | 77 |
| 574 | Transpositional competence and transcription of endogenous Ty elements in <i>Saccharomyces cerevisiae</i> : implications for regulation of transposition.. <i>Molecular and Cellular Biology</i> , 1988, 8, 3571-3581. | 1.1 | 46 |
| 575 | Molecular analysis of GCN3, a translational activator of GCN4: evidence for posttranslational control of GCN3 regulatory function.. <i>Molecular and Cellular Biology</i> , 1988, 8, 4808-4820. | 1.1 | 91 |
| 576 | Gene conversion adjacent to regions of double-strand break repair.. <i>Molecular and Cellular Biology</i> , 1988, 8, 5292-5298. | 1.1 | 62 |

| # | ARTICLE | IF | CITATIONS |
|-----|---|-----|-----------|
| 577 | Bent DNA functions as a replication enhancer in <i>Saccharomyces cerevisiae</i> .. <i>Molecular and Cellular Biology</i> , 1988, 8, 2763-2769. | 1.1 | 95 |
| 578 | Dominant negative protein kinase mutations that confer a G1 arrest phenotype.. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 1988, 85, 4426-4430. | 3.3 | 75 |
| 579 | Physical mapping of large DNA by chromosome fragmentation.. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 1988, 85, 6027-6031. | 3.3 | 196 |
| 580 | Genetic evidence for promoter competition in <i>Saccharomyces cerevisiae</i> .. <i>Molecular and Cellular Biology</i> , 1988, 8, 4608-4615. | 1.1 | 81 |
| 581 | Domain structure and functional analysis of the carboxyl-terminal polyacidic sequence of the RAD6 protein of <i>Saccharomyces cerevisiae</i> .. <i>Molecular and Cellular Biology</i> , 1988, 8, 1179-1185. | 1.1 | 114 |
| 582 | The a-factor pheromone of <i>Saccharomyces cerevisiae</i> is essential for mating.. <i>Molecular and Cellular Biology</i> , 1988, 8, 1309-1318. | 1.1 | 258 |
| 583 | Functional organization of the retrotransposon Ty from <i>Saccharomyces cerevisiae</i> : Ty protease is required for transposition.. <i>Molecular and Cellular Biology</i> , 1988, 8, 1421-1431. | 1.1 | 110 |
| 584 | Identification of functional regions in the yeast transcriptional activator ADR1.. <i>Molecular and Cellular Biology</i> , 1988, 8, 2125-2131. | 1.1 | 65 |
| 585 | Physical monitoring of mating type switching in <i>Saccharomyces cerevisiae</i> .. <i>Molecular and Cellular Biology</i> , 1988, 8, 2342-2349. | 1.1 | 114 |
| 586 | The positive regulatory function of the 5'-proximal open reading frames in GCN4 mRNA can be mimicked by heterologous, short coding sequences.. <i>Molecular and Cellular Biology</i> , 1988, 8, 3827-3836. | 1.1 | 46 |
| 587 | Specific translational activation by nuclear gene products occurs in the 5' untranslated leader of a yeast mitochondrial mRNA.. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 1988, 85, 2677-2681. | 3.3 | 133 |
| 588 | Chitin synthase 2 is essential for septum formation and cell division in <i>Saccharomyces cerevisiae</i> .. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 1988, 85, 4735-4739. | 3.3 | 198 |
| 589 | Plasmids can stably transform yeast mitochondria lacking endogenous mtDNA.. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 1988, 85, 7288-7292. | 3.3 | 149 |
| 590 | Cooperative DNA binding of the yeast transcriptional activator GAL4.. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 1988, 85, 382-386. | 3.3 | 222 |
| 591 | SAM2 encodes the second methionine S-adenosyl transferase in <i>Saccharomyces cerevisiae</i> : physiology and regulation of both enzymes.. <i>Molecular and Cellular Biology</i> , 1988, 8, 5132-5139. | 1.1 | 144 |
| 592 | Nematode repetitive DNA with ARS and segregation function in <i>Saccharomyces cerevisiae</i> .. <i>Molecular and Cellular Biology</i> , 1988, 8, 875-883. | 1.1 | 19 |
| 593 | Constitutive expression of the maltose fermentative enzymes in <i>Saccharomyces carlsbergensis</i> is dependent upon the mutational activation of a nonessential homolog of MAL63.. <i>Molecular and Cellular Biology</i> , 1988, 8, 1027-1035. | 1.1 | 29 |
| 594 | Identification of a Ty1 regulatory sequence responsive to STE7 and STE12.. <i>Molecular and Cellular Biology</i> , 1988, 8, 2545-2554. | 1.1 | 78 |

| # | ARTICLE | IF | CITATIONS |
|-----|--|-----|-----------|
| 595 | Transformation of yeast with synthetic oligonucleotides.. Proceedings of the National Academy of Sciences of the United States of America, 1988, 85, 524-528. | 3.3 | 151 |
| 596 | Role of SGP2, a suppressor of a <i>gpa1</i> mutation, in the mating-factor signaling pathway of <i>Saccharomyces cerevisiae</i> .. Molecular and Cellular Biology, 1988, 8, 5410-5416. | 1.1 | 29 |
| 597 | Single-amino-acid substitutions within the signal sequence of yeast prepro-alpha-factor affect membrane translocation.. Molecular and Cellular Biology, 1988, 8, 1915-1922. | 1.1 | 23 |
| 598 | Mutational analysis of centromere DNA from chromosome VI of <i>Saccharomyces cerevisiae</i> .. Molecular and Cellular Biology, 1988, 8, 2523-2535. | 1.1 | 165 |
| 599 | Potential positive and negative autoregulation of p60c-src by intermolecular autophosphorylation.. Proceedings of the National Academy of Sciences of the United States of America, 1988, 85, 4232-4236. | 3.3 | 173 |
| 600 | Host function of MAK16: G1 arrest by a <i>mak16</i> mutant of <i>Saccharomyces cerevisiae</i> .. Proceedings of the National Academy of Sciences of the United States of America, 1988, 85, 6007-6011. | 3.3 | 33 |
| 601 | DNA topoisomerase-targeting antitumor drugs can be studied in yeast.. Proceedings of the National Academy of Sciences of the United States of America, 1988, 85, 7501-7505. | 3.3 | 451 |
| 602 | Secretion of functional antibody and Fab fragment from yeast cells.. Proceedings of the National Academy of Sciences of the United States of America, 1988, 85, 8678-8682. | 3.3 | 167 |
| 603 | A regulatory region responsible for proline-specific induction of the yeast <i>PUT2</i> gene is adjacent to its TATA box.. Molecular and Cellular Biology, 1988, 8, 4634-4641. | 1.1 | 43 |
| 604 | <i>GAL11</i> protein, an auxiliary transcription activator for genes encoding galactose-metabolizing enzymes in <i>Saccharomyces cerevisiae</i> .. Molecular and Cellular Biology, 1988, 8, 4991-4999. | 1.1 | 160 |
| 605 | <i>Ty3</i> , a yeast retrotransposon associated with tRNA genes, has homology to animal retroviruses.. Molecular and Cellular Biology, 1988, 8, 5245-5256. | 1.1 | 194 |
| 606 | Cloning of the <i>PYR3</i> gene of <i>Ustilago maydis</i> and its use in DNA transformation.. Molecular and Cellular Biology, 1988, 8, 5417-5424. | 1.1 | 36 |
| 607 | The first and fourth upstream open reading frames in <i>GCN4</i> mRNA have similar initiation efficiencies but respond differently in translational control to change in length and sequence.. Molecular and Cellular Biology, 1988, 8, 5439-5447. | 1.1 | 37 |
| 608 | Gene conversion associated with site-specific recombination in yeast plasmid <i>pSR1</i> .. Molecular and Cellular Biology, 1988, 8, 955-962. | 1.1 | 27 |
| 609 | The <i>Saccharomyces cerevisiae</i> <i>ACP2</i> gene encodes an essential HMG1-like protein.. Molecular and Cellular Biology, 1988, 8, 1282-1289. | 1.1 | 43 |
| 610 | Regulation of expression and activity of the yeast transcription factor <i>ADR1</i> .. Molecular and Cellular Biology, 1988, 8, 1868-1876. | 1.1 | 57 |
| 611 | Transcription interferes with elements important for chromosome maintenance in <i>Saccharomyces cerevisiae</i> .. Molecular and Cellular Biology, 1988, 8, 2184-2194. | 1.1 | 158 |
| 612 | A new means of inducibly inactivating a cellular protein.. Molecular and Cellular Biology, 1988, 8, 2638-2646. | 1.1 | 91 |

| # | ARTICLE | IF | CITATIONS |
|-----|---|-----|-----------|
| 613 | Genetic selection for mutations that reduce or abolish ribosomal recognition of the HIS4 translational initiator region.. <i>Molecular and Cellular Biology</i> , 1988, 8, 2955-2963. | 1.1 | 134 |
| 614 | Mutational analysis of the HIS4 translational initiator region in <i>Saccharomyces cerevisiae</i> .. <i>Molecular and Cellular Biology</i> , 1988, 8, 2964-2975. | 1.1 | 140 |
| 615 | Structure and regulation of a nuclear gene in <i>Saccharomyces cerevisiae</i> that specifies MRP7, a protein of the large subunit of the mitochondrial ribosome.. <i>Molecular and Cellular Biology</i> , 1988, 8, 3636-3646. | 1.1 | 81 |
| 616 | Mitotic recombination within the centromere of a yeast chromosome. <i>Science</i> , 1988, 241, 1074-1077. | 6.0 | 23 |
| 617 | Structural organization and functional analysis of centromeric DNA in the fission yeast <i>Schizosaccharomyces pombe</i> .. <i>Molecular and Cellular Biology</i> , 1988, 8, 754-763. | 1.1 | 107 |
| 618 | TRK1 encodes a plasma membrane protein required for high-affinity potassium transport in <i>Saccharomyces cerevisiae</i> .. <i>Molecular and Cellular Biology</i> , 1988, 8, 2848-2859. | 1.1 | 289 |
| 619 | A CONVENIENT DOMINANT SELECTION MARKER FOR GENE TRANSFER IN INDUSTRIAL STRAINS OF <i>SACCHAROMYCES</i> YEAST: SMRI ENCODED RESISTANCE TO THE HERBICIDE SULFOMETURON METHYL. <i>Journal of the Institute of Brewing</i> , 1988, 94, 93-97. | 0.8 | 69 |
| 620 | Construction of the transcriptional and translational signals-cloning vector in <i>Saccharomyces cerevisiae</i> .. <i>Agricultural and Biological Chemistry</i> , 1988, 52, 1597-1598. | 0.3 | 0 |
| 621 | Construction of new ARS1 vectors with increased mitotic stability in <i>Saccharomyces cerevisiae</i> .. <i>Agricultural and Biological Chemistry</i> , 1988, 52, 2021-2025. | 0.3 | 0 |
| 622 | Efficient translational frameshifting occurs within a conserved sequence of the overlap between the two genes of a yeast Ty1 transposon.. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 1988, 85, 6816-6820. | 3.3 | 127 |
| 623 | Expression of diphtheria toxin fragment A and hormone-toxin fusion proteins in toxin-resistant yeast mutants.. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 1988, 85, 8386-8390. | 3.3 | 31 |
| 624 | Isolation and characterization of temperature-sensitive mutations in RPA190, the gene encoding the largest subunit of RNA polymerase I from <i>Saccharomyces cerevisiae</i> .. <i>Molecular and Cellular Biology</i> , 1988, 8, 3997-4008. | 1.1 | 60 |
| 625 | Characterization of two telomeric DNA processing reactions in <i>Saccharomyces cerevisiae</i> .. <i>Molecular and Cellular Biology</i> , 1988, 8, 4642-4650. | 1.1 | 82 |
| 626 | SPT3 is required for normal levels of a-factor and alpha-factor expression in <i>Saccharomyces cerevisiae</i> .. <i>Molecular and Cellular Biology</i> , 1988, 8, 822-827. | 1.1 | 33 |
| 627 | Mutations in a gene encoding the alpha subunit of a <i>Saccharomyces cerevisiae</i> G protein indicate a role in mating pheromone signaling.. <i>Molecular and Cellular Biology</i> , 1988, 8, 2484-2493. | 1.1 | 153 |
| 628 | Role of STE genes in the mating factor signaling pathway mediated by GPA1 in <i>Saccharomyces cerevisiae</i> .. <i>Molecular and Cellular Biology</i> , 1988, 8, 3777-3783. | 1.1 | 113 |
| 629 | Efficient repair of HO-induced chromosomal breaks in <i>Saccharomyces cerevisiae</i> by recombination between flanking homologous sequences.. <i>Molecular and Cellular Biology</i> , 1988, 8, 3918-3928. | 1.1 | 204 |
| 630 | Sequence and genetic analysis of a dispensible 189 nucleotide snRNA from <i>Saccharomyces cerevisiae</i> . <i>Nucleic Acids Research</i> , 1988, 16, 5587-5601. | 6.5 | 14 |

| # | ARTICLE | IF | CITATIONS |
|-----|--|-----|-----------|
| 631 | Molecular cloning and nucleotide sequence of the nuclear PET122 gene required for expression of the mitochondrial COX3 gene in <i>S.cerevisiae</i> . <i>Nucleic Acids Research</i> , 1988, 16, 10783-10802. | 6.5 | 33 |
| 632 | No strict alignment is required between a transcriptional activator binding site and the "TATA box" of a yeast gene.. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 1988, 85, 4262-4266. | 3.3 | 61 |
| 633 | The copper, zinc-superoxide dismutase gene of <i>Saccharomyces cerevisiae</i> : cloning, sequencing, and biological activity.. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 1988, 85, 4789-4793. | 3.3 | 139 |
| 634 | Evidence for regulation of reinitiation in translational control of GCN4 mRNA.. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 1988, 85, 7279-7283. | 3.3 | 26 |
| 635 | An adenylate cyclase from <i>Saccharomyces cerevisiae</i> that is stimulated by RAS proteins with effector mutations.. <i>Molecular and Cellular Biology</i> , 1988, 8, 52-61. | 1.1 | 57 |
| 636 | Properties of promoters cloned randomly from the <i>Saccharomyces cerevisiae</i> genome.. <i>Molecular and Cellular Biology</i> , 1988, 8, 4217-4224. | 1.1 | 48 |
| 637 | Direction of chromosome rearrangements in <i>Saccharomyces cerevisiae</i> by use of his3 recombinational substrates.. <i>Molecular and Cellular Biology</i> , 1988, 8, 4370-4380. | 1.1 | 47 |
| 638 | Copy number and partition of the <i>Saccharomyces cerevisiae</i> 2 micron plasmid controlled by transcription regulators.. <i>Molecular and Cellular Biology</i> , 1988, 8, 4949-4957. | 1.1 | 29 |
| 639 | Connections between transcriptional activators, silencers, and telomeres as revealed by functional analysis of a yeast DNA-binding protein.. <i>Molecular and Cellular Biology</i> , 1988, 8, 5086-5099. | 1.1 | 315 |
| 640 | Isolation of a yeast gene involved in species-specific pre-tRNA processing.. <i>Molecular and Cellular Biology</i> , 1988, 8, 5140-5149. | 1.1 | 76 |
| 641 | A Ty1 cell-type-specific regulatory sequence is a recognition element for a constitutive binding factor.. <i>Molecular and Cellular Biology</i> , 1988, 8, 5299-5309. | 1.1 | 43 |
| 642 | Genetic analysis of small nuclear RNAs in <i>Saccharomyces cerevisiae</i> : viable sextuple mutant.. <i>Molecular and Cellular Biology</i> , 1988, 8, 3150-3159. | 1.1 | 91 |
| 643 | Characterization of an SNR gene locus in <i>Saccharomyces cerevisiae</i> that specifies both dispensible and essential small nuclear RNAs.. <i>Molecular and Cellular Biology</i> , 1988, 8, 3282-3290. | 1.1 | 106 |
| 644 | Yeast mutants temperature-sensitive for growth after random mutagenesis of the chromosomal RAS2 gene and deletion of the RAS1 gene.. <i>EMBO Journal</i> , 1988, 7, 3375-3383. | 3.5 | 37 |
| 645 | The processing peptidase of yeast mitochondria: the two co-operating components MPP and PEP are structurally related.. <i>EMBO Journal</i> , 1988, 7, 3493-3500. | 3.5 | 149 |
| 646 | The WHI1+ gene of <i>Saccharomyces cerevisiae</i> tethers cell division to cell size and is a cyclin homolog.. <i>EMBO Journal</i> , 1988, 7, 4335-4346. | 3.5 | 512 |
| 647 | Expression of double-stranded-RNA-specific RNase III of <i>Escherichia coli</i> is lethal to <i>Saccharomyces cerevisiae</i> . <i>Journal of Bacteriology</i> , 1988, 170, 2989-2993. | 1.0 | 18 |
| 648 | Transcriptional regulation of the cell cycle-dependent thymidylate synthase gene of <i>Saccharomyces cerevisiae</i> .. <i>Molecular and Cellular Biology</i> , 1988, 8, 4616-4624. | 1.1 | 43 |

| # | ARTICLE | IF | CITATIONS |
|-----|---|-----|-----------|
| 649 | Biogenesis of the yeast lysosome (vacuole): biosynthesis and maturation of proteinase yscB.. EMBO Journal, 1988, 7, 1705-1710. | 3.5 | 50 |
| 650 | Dissection of functional domains of the yeast proton-pumping ATPase by directed mutagenesis.. EMBO Journal, 1988, 7, 1793-1798. | 3.5 | 106 |
| 651 | Cold-sensitive and caffeine-supersensitive mutants of the <i>Schizosaccharomyces pombe</i> dis genes implicated in sister chromatid separation during mitosis.. EMBO Journal, 1988, 7, 1465-1473. | 3.5 | 197 |
| 652 | Cysteine biosynthesis in <i>Saccharomyces cerevisiae</i> : mutation that confers cystathionine beta-synthase deficiency. Journal of Bacteriology, 1988, 170, 5883-5889. | 1.0 | 58 |
| 653 | Import of proteins into yeast mitochondria: the nuclear MAS2 gene encodes a component of the processing protease that is homologous to the MAS1-encoded subunit.. EMBO Journal, 1988, 7, 3863-3871. | 3.5 | 145 |
| 654 | Genetic Engineering of Yeast: Construction of Strains That Degrade β -Glucans with the Aid of a Barley Gene. Journal of the American Society of Brewing Chemists, 1988, 46, 31-36. | 0.8 | 9 |
| 655 | 7SL RNA from <i>Schizosaccharomyces pombe</i> is encoded by a single copy essential gene. EMBO Journal, 1988, 7, 231-237. | 3.5 | 66 |
| 656 | Dual regulation of the expression of the polyubiquitin gene by cyclic AMP and heat shock in yeast.. EMBO Journal, 1988, 7, 495-502. | 3.5 | 121 |
| 657 | The NAM2 proteins from <i>S. cerevisiae</i> and <i>S. douglasii</i> are mitochondrial leucyl-tRNA synthetases, and are involved in mRNA splicing.. EMBO Journal, 1988, 7, 473-483. | 3.5 | 180 |
| 658 | A yeast gene required for the G1-to-S transition encodes a protein containing an A-kinase target site and GTPase domain.. EMBO Journal, 1988, 7, 1175-1182. | 3.5 | 112 |
| 659 | A U1 snRNA:pre-mRNA base pairing interaction is required early in yeast spliceosome assembly but does not uniquely define the 5' splice site.. EMBO Journal, 1988, 7, 2533-2538. | 3.5 | 328 |
| 660 | Mutation of lysine-48 to arginine in the yeast RAD3 protein abolishes its ATPase and DNA helicase activities but not the ability to bind ATP.. EMBO Journal, 1988, 7, 3263-3269. | 3.5 | 251 |
| 661 | A novel nucleoskeletal-like protein located at the nuclear periphery is required for the life cycle of <i>Saccharomyces cerevisiae</i> .. EMBO Journal, 1988, 7, 4323-4334. | 3.5 | 143 |
| 662 | Cloning and nucleotide sequence of DNA mismatch repair gene PMS1 from <i>Saccharomyces cerevisiae</i> : homology of PMS1 to prokaryotic MutL and HexB. Journal of Bacteriology, 1989, 171, 5339-5346. | 1.0 | 227 |
| 663 | The <i>Saccharomyces cerevisiae</i> PUT3 activator protein associates with proline-specific upstream activation sequences.. Molecular and Cellular Biology, 1989, 9, 4706-4712. | 1.1 | 77 |
| 664 | Molecular cloning of a cell wall exo-beta-1,3-glucanase from <i>Saccharomyces cerevisiae</i> . Journal of Bacteriology, 1989, 171, 6259-6264. | 1.0 | 127 |
| 665 | Isolation, characterization, and inactivation of the APA1 gene encoding yeast diadenosine 5',5'''-P1,P4-tetraphosphate phosphorylase. Journal of Bacteriology, 1989, 171, 6437-6445. | 1.0 | 35 |
| 666 | Interpathway regulation of the TRP4 gene of yeast.. EMBO Journal, 1989, 8, 939-945. | 3.5 | 56 |

| # | ARTICLE | IF | CITATIONS |
|-----|--|-----|-----------|
| 667 | Isolation and DNA-binding characteristics of a protein involved in transcription activation of two divergently transcribed, essential yeast genes.. EMBO Journal, 1989, 8, 3029-3037. | 3.5 | 71 |
| 668 | A human homologue of the yeast GST1 gene codes for a GTP-binding protein and is expressed in a proliferation-dependent manner in mammalian cells.. EMBO Journal, 1989, 8, 3807-3814. | 3.5 | 107 |
| 669 | Sequence, expression and mutational analysis of BAF1, a transcriptional activator and ARS1-binding protein of the yeast <i>Saccharomyces cerevisiae</i> .. EMBO Journal, 1989, 8, 4265-4272. | 3.5 | 86 |
| 670 | Identification and characterization of the maltose permease in genetically defined <i>Saccharomyces</i> strain. <i>Journal of Bacteriology</i> , 1989, 171, 6148-6154. | 1.0 | 35 |
| 671 | The cloned human oestrogen receptor contains a mutation which alters its hormone binding properties.. EMBO Journal, 1989, 8, 1981-1986. | 3.5 | 391 |
| 672 | Mutants of H-ras that interfere with RAS effector function in <i>Saccharomyces cerevisiae</i> .. EMBO Journal, 1989, 8, 3039-3044. | 3.5 | 47 |
| 673 | A yeast nucleolar protein related to mammalian fibrillarin is associated with small nucleolar RNA and is essential for viability.. EMBO Journal, 1989, 8, 4015-4024. | 3.5 | 283 |
| 674 | Sequence and structural requirements of a mitochondrial protein import signal defined by saturation cassette mutagenesis.. <i>Molecular and Cellular Biology</i> , 1989, 9, 1014-1025. | 1.1 | 66 |
| 675 | Mutational analysis of SEC4 suggests a cyclical mechanism for the regulation of vesicular traffic.. EMBO Journal, 1989, 8, 1685-1693. | 3.5 | 299 |
| 676 | <i>Saccharomyces cerevisiae</i> STE6 gene product: a novel pathway for protein export in eukaryotic cells.. EMBO Journal, 1989, 8, 3973-3984. | 3.5 | 412 |
| 677 | Identification of an upstream activating sequence and an upstream repressible sequence of the pyruvate kinase gene of the yeast <i>Saccharomyces cerevisiae</i> .. <i>Molecular and Cellular Biology</i> , 1989, 9, 442-451. | 1.1 | 107 |
| 678 | Characterization of the END1 gene required for vacuole biogenesis and gluconeogenic growth of budding yeast.. EMBO Journal, 1989, 8, 1349-1359. | 3.5 | 67 |
| 679 | The ras-related mouse ypt1 protein can functionally replace the YPT1 gene product in yeast.. EMBO Journal, 1989, 8, 1427-1432. | 3.5 | 128 |
| 680 | A point mutation in the CYC1 UAS1 creates a new combination of regulatory elements that activate transcription synergistically.. EMBO Journal, 1989, 8, 1801-1808. | 3.5 | 14 |
| 681 | A yeast mitochondrial presequence functions as a signal for targeting to plant mitochondria in vivo.. <i>Plant Cell</i> , 1989, 1, 783-791. | 3.1 | 76 |
| 682 | Regulated phosphorylation and dephosphorylation of GAL4, a transcriptional activator.. <i>Genes and Development</i> , 1989, 3, 1157-1165. | 2.7 | 107 |
| 683 | Functional analysis of the papilloma virus E2 trans-activator in <i>Saccharomyces cerevisiae</i> .. <i>Genes and Development</i> , 1989, 3, 38-48. | 2.7 | 78 |
| 684 | N2,N2-dimethylguanosine-specific tRNA methyltransferase contains both nuclear and mitochondrial targeting signals in <i>Saccharomyces cerevisiae</i> .. <i>Journal of Cell Biology</i> , 1989, 109, 1411-1419. | 2.3 | 60 |

| # | ARTICLE | IF | CITATIONS |
|-----|---|-----|-----------|
| 685 | Loss of Ras activity in <i>Saccharomyces cerevisiae</i> is suppressed by disruptions of a new kinase gene, YAKI, whose product may act downstream of the cAMP-dependent protein kinase.. <i>Genes and Development</i> , 1989, 3, 1336-1348. | 2.7 | 210 |
| 686 | Mutations in the glucocorticoid receptor zinc finger region that distinguish interdigitated DNA binding and transcriptional enhancement activities.. <i>Genes and Development</i> , 1989, 3, 1590-1601. | 2.7 | 226 |
| 687 | The acid phosphatase genes PHO10 and PHO11 in <i>S.cerevisiae</i> are located at the telomeres of chromosomes VIII and I. <i>Nucleic Acids Research</i> , 1989, 17, 1353-1369. | 6.5 | 37 |
| 688 | The GTP-binding protein Ypt1 is required for transport in vitro: the Golgi apparatus is defective in ypt1 mutants.. <i>Journal of Cell Biology</i> , 1989, 109, 1015-1022. | 2.3 | 238 |
| 689 | A yeast gene important for protein assembly into the endoplasmic reticulum and the nucleus has homology to DnaJ, an <i>Escherichia coli</i> heat shock protein.. <i>Journal of Cell Biology</i> , 1989, 109, 2665-2675. | 2.3 | 304 |
| 690 | <i>Saccharomyces cerevisiae</i> and <i>Schizosaccharomyces pombe</i> contain a homologue to the 54-kD subunit of the signal recognition particle that in <i>S. cerevisiae</i> is essential for growth.. <i>Journal of Cell Biology</i> , 1989, 109, 3223-3230. | 2.3 | 118 |
| 691 | Multiple genes are required for proper insertion of secretory proteins into the endoplasmic reticulum in yeast.. <i>Journal of Cell Biology</i> , 1989, 109, 2641-2652. | 2.3 | 341 |
| 692 | Isolation and characterization of STI1, a stress-inducible gene from <i>Saccharomyces cerevisiae</i> .. <i>Molecular and Cellular Biology</i> , 1989, 9, 3638-3646. | 1.1 | 242 |
| 693 | Processing pathway for protease B of <i>Saccharomyces cerevisiae</i> .. <i>Journal of Cell Biology</i> , 1989, 108, 309-325. | 2.3 | 82 |
| 694 | The Effects of Culture Conditions on the Secretion of Human Lysozyme by <i>Saccharomyces cerevisiae</i> A2-1-1A. <i>Agricultural and Biological Chemistry</i> , 1989, 53, 2687-2694. | 0.3 | 0 |
| 695 | The SPA2 protein of yeast localizes to sites of cell growth.. <i>Journal of Cell Biology</i> , 1989, 108, 1419-1429. | 2.3 | 204 |
| 696 | The <i>Saccharomyces cerevisiae</i> SEC14 gene encodes a cytosolic factor that is required for transport of secretory proteins from the yeast Golgi complex.. <i>Journal of Cell Biology</i> , 1989, 108, 1271-1281. | 2.3 | 386 |
| 697 | Higher order chromosome structure is affected by cold-sensitive mutations in a <i>Schizosaccharomyces pombe</i> gene <i>crm1+</i> which encodes a 115-kD protein preferentially localized in the nucleus and its periphery.. <i>Journal of Cell Biology</i> , 1989, 108, 1195-1207. | 2.3 | 232 |
| 698 | Increased Cell Wall Porosity in <i>Saccharomyces cerevisiae</i> after Treatment with Dithiothreitol or EDTA. <i>Microbiology (United Kingdom)</i> , 1989, 135, 2077-2084. | 0.7 | 30 |
| 699 | Essential roles of the RNA polymerase I largest subunit and DNA topoisomerases in the formation of fission yeast nucleolus.. <i>Journal of Cell Biology</i> , 1989, 108, 243-253. | 2.3 | 89 |
| 700 | Mutations in the SAC1 gene suppress defects in yeast Golgi and yeast actin function.. <i>Journal of Cell Biology</i> , 1989, 109, 2939-2950. | 2.3 | 246 |
| 701 | Expression of pGKL killer 28K subunit in <i>Saccharomyces cerevisiae</i> : identification of 28K subunit as a killer protein. <i>Nucleic Acids Research</i> , 1989, 17, 3435-3446. | 6.5 | 79 |
| 702 | Integrative transformation of yeast using electroporation. <i>Nucleic Acids Research</i> , 1989, 17, 8011-8011. | 6.5 | 12 |

| # | ARTICLE | IF | CITATIONS |
|-----|---|-----|-----------|
| 703 | The Segregation of the 2 ^Å -based Yeast Plasmid pJDB248 Breaks Down under Conditions of Slow, Glucose-limited Growth. <i>Microbiology (United Kingdom)</i> , 1989, 135, 2891-2897. | 0.7 | 9 |
| 704 | Mutations in the anticodon stem affect removal of introns from pre-tRNA in <i>Saccharomyces cerevisiae</i> .. <i>Molecular and Cellular Biology</i> , 1989, 9, 4220-4228. | 1.1 | 17 |
| 705 | Control of the <i>Saccharomyces cerevisiae</i> regulatory gene PET494: transcriptional repression by glucose and translational induction by oxygen.. <i>Molecular and Cellular Biology</i> , 1989, 9, 484-491. | 1.1 | 59 |
| 706 | The Sec15 protein responds to the function of the GTP binding protein, Sec4, to control vesicular traffic in yeast.. <i>Journal of Cell Biology</i> , 1989, 109, 1023-1036. | 2.3 | 136 |
| 707 | The residual enzymatic phosphorylation activity of hexokinase II mutants is correlated with glucose repression in <i>Saccharomyces cerevisiae</i> .. <i>Molecular and Cellular Biology</i> , 1989, 9, 5643-5649. | 1.1 | 128 |
| 708 | Mating-type control in <i>Saccharomyces cerevisiae</i> : isolation and characterization of mutants defective in repression by a1-alpha 2.. <i>Molecular and Cellular Biology</i> , 1989, 9, 4523-4530. | 1.1 | 32 |
| 709 | [33] Expression of engineered antibodies and antibody fragments in microorganisms. <i>Methods in Enzymology</i> , 1989, 178, 476-496. | 0.4 | 23 |
| 710 | The yeast gene ERG6 is required for normal membrane function but is not essential for biosynthesis of the cell-cycle-sparking sterol.. <i>Molecular and Cellular Biology</i> , 1989, 9, 3447-3456. | 1.1 | 256 |
| 711 | Structure, biosynthesis, and localization of dipeptidyl aminopeptidase B, an integral membrane glycoprotein of the yeast vacuole.. <i>Journal of Cell Biology</i> , 1989, 108, 1363-1373. | 2.3 | 162 |
| 712 | Construction and Characterization of a Temperature-sensitive Expression System in Recombinant Yeast. <i>Biotechnology Progress</i> , 1989, 5, 18-26. | 1.3 | 25 |
| 713 | Characterization and localization of the sporulation glucoamylase of <i>Saccharomyces cerevisiae</i> . <i>BBA - Proteins and Proteomics</i> , 1989, 994, 200-209. | 2.1 | 22 |
| 714 | Improvement and application of a promoter-probe vector bearing the PHO5 gene as the indicator marker in <i>Saccharomyces cerevisiae</i> . <i>Journal of Bioscience and Bioengineering</i> , 1989, 67, 1-7. | 0.9 | 11 |
| 715 | Expression of the glutathione synthetase gene of <i>Escherichia coli</i> B in <i>Saccharomyces cerevisiae</i> . <i>Journal of Bioscience and Bioengineering</i> , 1989, 68, 390-394. | 0.9 | 33 |
| 716 | Construction of yeast xylulokinase mutant by recombinant dna techniques. <i>Applied Biochemistry and Biotechnology</i> , 1989, 20-21, 327-334. | 1.4 | 7 |
| 717 | Processing and secretion of barley (1 ^Å 3, 1 ^Å 4)- β -glucanase in yeast. <i>Carlsberg Research Communications</i> , 1989, 54, 29-39. | 1.7 | 14 |
| 718 | Three regulatory systems control expression of glutamine synthetase in <i>Saccharomyces cerevisiae</i> at the level of transcription. <i>Molecular Genetics and Genomics</i> , 1989, 217, 370-377. | 2.4 | 40 |
| 719 | Direct Introduction of Fluorescein Isothiocyanate-conjugated Albumin into Intact Macroconidia of <i>Fusarium oxysporum</i> f. sp. <i>lycopersici</i> by Electroporation. <i>Journal of Phytopathology</i> , 1989, 125, 89-96. | 0.5 | 6 |
| 720 | Molecular cloning, primary structure and disruption of the structural gene of aldolase from <i>Saccharomyces cerevisiae</i> . <i>FEBS Journal</i> , 1989, 180, 301-308. | 0.2 | 66 |

| # | ARTICLE | IF | CITATIONS |
|-----|--|-----|-----------|
| 721 | Structural requirements for protein N-glycosylation. Influence of acceptor peptides on cotranslational glycosylation of yeast invertase and site-directed mutagenesis around a sequon sequence. <i>FEBS Journal</i> , 1989, 181, 525-529. | 0.2 | 77 |
| 722 | Site-directed mutagenesis of yeast phosphoglycerate kinase. The 'basic-patch' residue arginine 168. <i>FEBS Journal</i> , 1989, 183, 49-55. | 0.2 | 24 |
| 723 | One- and two-dimensional NMR studies of yeast phosphoglycerate kinase. <i>FEBS Journal</i> , 1989, 184, 617-625. | 0.2 | 20 |
| 724 | The role of subunit 4, a nuclear-encoded protein of the F0 sector of yeast mitochondrial ATP synthase, in the assembly of the whole complex. <i>FEBS Journal</i> , 1989, 185, 163-171. | 0.2 | 90 |
| 725 | Characterization of the methyltransferases in the yeast phosphatidylethanolamine methylation pathway by selective gene disruption. <i>FEBS Journal</i> , 1989, 185, 243-251. | 0.2 | 100 |
| 726 | Transformation of a glucose negative mutant of <i>Pachysolen tannophilus</i> with a plasmid carrying the cloned hexokinase PII gene from <i>Saccharomyces cerevisiae</i> . <i>Biotechnology Letters</i> , 1989, 11, 601-604. | 1.1 | 5 |
| 727 | Meiotic segregation of circular plasmid-minichromosomes from intact chromosomes in <i>Saccharomyces cerevisiae</i> . <i>Current Genetics</i> , 1989, 15, 385-392. | 0.8 | 20 |
| 728 | Molecular transformation of <i>Fusarium solani</i> with an antibiotic resistance marker having no fungal DNA homology. <i>Current Genetics</i> , 1989, 15, 421-428. | 0.8 | 36 |
| 729 | Molecular cloning of chromosome I DNA from <i>Saccharomyces cerevisiae</i> : localization of a repeated sequence containing an acid phosphatase gene near a telomere of chromosome I and chromosome VIII. <i>Current Genetics</i> , 1989, 16, 131-137. | 0.8 | 53 |
| 730 | Cloning and characterization of the SKI3 gene of <i>Saccharomyces cerevisiae</i> demonstrates allelism to SKI5. <i>Current Genetics</i> , 1989, 16, 139-143. | 0.8 | 5 |
| 731 | Molecular characterization of the two genes SNQ and SFA that confer Hyperresistance to 4-nitroquinoline-N-oxide and formaldehyde in <i>Saccharomyces cerevisiae</i> . <i>Current Genetics</i> , 1989, 16, 65-74. | 0.8 | 21 |
| 732 | Cloning and physical characterization of linked lysine genes (LYS4, LYS15) of <i>Saccharomyces cerevisiae</i> . <i>Current Genetics</i> , 1989, 16, 7-12. | 0.8 | 14 |
| 733 | A novel class of vector for yeast transformation. <i>Current Genetics</i> , 1989, 16, 21-25. | 0.8 | 40 |
| 734 | Regulated overproduction of α -amylase by transformation of the amylolytic yeast <i>Schwanniomyces occidentalis</i> . <i>Current Genetics</i> , 1989, 15, 319-325. | 0.8 | 48 |
| 735 | Interactions of the RAD7 and RAD23 excision repair genes of <i>Saccharomyces cerevisiae</i> with DNA repair genes in different epistasis groups. <i>Current Genetics</i> , 1989, 16, 219-223. | 0.8 | 13 |
| 736 | An improved host-vector system for <i>Candida maltosa</i> using a gene isolated from its genome that complements the hiss mutation of <i>Saccharomyces cerevisiae</i> . <i>Current Genetics</i> , 1989, 16, 261-266. | 0.8 | 38 |
| 737 | Dependence of inessential late gene expression on early meiotic events in <i>Saccharomyces cerevisiae</i> . <i>Molecular Genetics and Genomics</i> , 1989, 215, 490-500. | 2.4 | 31 |
| 738 | Novel class of nuclear genes involved in both mRNA splicing and protein synthesis in <i>Saccharomyces cerevisiae</i> mitochondria. <i>Molecular Genetics and Genomics</i> , 1989, 215, 517-528. | 2.4 | 62 |

| # | ARTICLE | IF | CITATIONS |
|-----|---|-----|-----------|
| 739 | Heterologous gene expression of the glyphosate resistance marker and its application in yeast transformation. <i>Current Genetics</i> , 1989, 15, 91-98. | 0.8 | 22 |
| 740 | Enhanced canavanine uptake is associated with nucleotide permeability in a thymidylate auxotroph of <i>Saccharomyces cerevisiae</i> . <i>Current Genetics</i> , 1989, 15, 129-134. | 0.8 | 3 |
| 741 | G418 resistance in the yeast <i>Saccharomyces cerevisiae</i> : comparison of the neomycin resistance genes from Tn5 and Tn903. <i>Applied Microbiology and Biotechnology</i> , 1989, 30, 388-394. | 1.7 | 26 |
| 742 | Molecular cloning of SNM1, a yeast gene responsible for a specific step in the repair of cross-linked DNA. <i>Molecular Genetics and Genomics</i> , 1989, 218, 64-71. | 2.4 | 21 |
| 743 | Mode of expression of the positive regulatory genes PHO2 and PHO4 of the phosphatase regulon in <i>Saccharomyces cerevisiae</i> . <i>Molecular Genetics and Genomics</i> , 1989, 217, 31-39. | 2.4 | 75 |
| 744 | Structure of the multigene family of MAL loci in <i>Saccharomyces</i> . <i>Molecular Genetics and Genomics</i> , 1989, 217, 60-69. | 2.4 | 44 |
| 745 | Structure of the HOM2 gene of <i>Saccharomyces cerevisiae</i> and regulation of its expression. <i>Molecular Genetics and Genomics</i> , 1989, 217, 149-154. | 2.4 | 24 |
| 746 | $\hat{\Gamma}$ -Factor leader sequence-directed transport of <i>Escherichia coli</i> $\hat{\Gamma}$ -galactosidase in the secretory pathway of <i>Saccharomyces cerevisiae</i> . <i>Molecular Genetics and Genomics</i> , 1989, 218, 240-248. | 2.4 | 19 |
| 747 | Ars region in TL-DNA on octopine type Ti plasmids. <i>Molecular Genetics and Genomics</i> , 1989, 218, 284-288. | 2.4 | 2 |
| 748 | A long region upstream of the IME1 gene regulates meiosis in yeast. <i>Molecular Genetics and Genomics</i> , 1989, 218, 308-314. | 2.4 | 36 |
| 749 | Analysis of expression of hybrid yeast genes containing ARS elements. <i>Molecular Genetics and Genomics</i> , 1989, 218, 531-535. | 2.4 | 10 |
| 750 | A functional analysis of the repeated methionine initiator tRNA genes (IMT) in yeast. <i>Molecular Genetics and Genomics</i> , 1989, 216, 276-286. | 2.4 | 31 |
| 751 | Mutations in ADE3 reduce the efficiency of the omnipotent suppressor sup45-2. <i>Current Genetics</i> , 1989, 16, 315-321. | 0.8 | 10 |
| 752 | High efficiency transformation of intact yeast cells using single stranded nucleic acids as a carrier. <i>Current Genetics</i> , 1989, 16, 339-346. | 0.8 | 2,191 |
| 753 | A ribosomal protein gene family from <i>Schizosaccharomyces pombe</i> consisting of three active members. <i>Current Genetics</i> , 1989, 16, 361-367. | 0.8 | 18 |
| 754 | Construction of yeast secretion vectors designed for production of mature proteins using the signal sequence of yeast invertase. <i>Applied Microbiology and Biotechnology</i> , 1989, 32, 317. | 1.7 | 6 |
| 755 | A direct selection procedure for isolating yeast mutants with an impaired segregation of artificial minichromosomes. <i>Current Genetics</i> , 1989, 15, 17-25. | 0.8 | 10 |
| 756 | Plasmid multimerization is dependent on RAD52 activity in <i>Saccharomyces cerevisiae</i> . <i>Molecular Genetics and Genomics</i> , 1989, 219, 495-498. | 2.4 | 8 |

| # | ARTICLE | IF | CITATIONS |
|-----|--|------|-----------|
| 757 | Cauliflower mosaic virus promoters direct efficient expression of a bacterial G418 resistance gene in <i>Schizosaccharomyces pombe</i> . <i>Molecular Genetics and Genomics</i> , 1989, 220, 95-101. | 2.4 | 27 |
| 758 | Yeast mutants with enhanced ability to secrete human lysozyme: Isolation and identification of a protease-deficient mutant. <i>Molecular Genetics and Genomics</i> , 1989, 219, 58-64. | 2.4 | 33 |
| 759 | Identification of a gene conferring resistance to zinc and cadmium ions in the yeast <i>Saccharomyces cerevisiae</i> . <i>Molecular Genetics and Genomics</i> , 1989, 219, 161-167. | 2.4 | 188 |
| 760 | High expression and efficient secretion of <i>Rhizopus oryzae</i> glucoamylase in the yeast <i>Saccharomyces cerevisiae</i> . <i>Applied Microbiology and Biotechnology</i> , 1989, 30, 515. | 1.7 | 50 |
| 761 | Use of molecular markers for monitoring fungi involved in stalk rot of corn. <i>Theoretical and Applied Genetics</i> , 1989, 77, 535-539. | 1.8 | 20 |
| 762 | The 15 N-terminal amino acids of hexokinase II are not required for in vivo function: Analysis of a truncated form of hexokinase II in <i>Saccharomyces cerevisiae</i> . <i>Proteins: Structure, Function and Bioinformatics</i> , 1989, 5, 218-223. | 1.5 | 13 |
| 763 | Cloning and analysis of the <i>Kluyveromyces lactis</i> TRP1 gene: A chromosomal locus flanked by genes encoding inorganic pyrophosphatase and histone H3. <i>Yeast</i> , 1989, 5, 35-50. | 0.8 | 104 |
| 764 | Increase in gene expression by respiratory-deficient mutation. <i>Yeast</i> , 1989, 5, 91-98. | 0.8 | 26 |
| 765 | Functional exploration of the yeast (<i>Saccharomyces cerevisiae</i>) genome: Use of a mini-mu transposon to analyse randomly cloned DNA sequences. <i>Yeast</i> , 1989, 5, 259-269. | 0.8 | 9 |
| 766 | <i>Saccharomyces cerevisiae</i> mutants defective in chromosome segregation. <i>Yeast</i> , 1989, 5, 271-284. | 0.8 | 26 |
| 767 | Physiology of mutants with reduced expression of plasma membrane H ⁺ -ATPase. <i>Yeast</i> , 1989, 5, 307-319. | 0.8 | 128 |
| 768 | High level of expression of a protective antigen of schistosomes in <i>Saccharomyces cerevisiae</i> . <i>Yeast</i> , 1989, 5, 497-507. | 0.8 | 66 |
| 769 | Isolation, DNA sequence and regulation of a new cell division cycle gene from the yeast <i>Saccharomyces cerevisiae</i> . <i>Yeast</i> , 1989, 5, 509-524. | 0.8 | 21 |
| 770 | Screening of Muteins Secreted by Yeast: Random Mutagenesis of Human Interleukin-2. <i>Nature Biotechnology</i> , 1989, 7, 716-720. | 9.4 | 6 |
| 771 | Secretion of Active, Full-length and Half-length Human Secretory Leukocyte Protease Inhibitor by <i>Saccharomyces Cerevisiae</i> . <i>Nature Biotechnology</i> , 1989, 7, 55-60. | 9.4 | 12 |
| 772 | An initiation site for meiotic gene conversion in the yeast <i>Saccharomyces cerevisiae</i> . <i>Nature</i> , 1989, 338, 35-39. | 13.7 | 380 |
| 773 | An autonomously replicating sequence of pSRI plasmid is effective in two yeast species, <i>Zygosaccharomyces rouxii</i> and <i>Saccharomyces cerevisiae</i> . <i>Journal of Molecular Biology</i> , 1989, 207, 757-769. | 2.0 | 16 |
| 774 | A subthreshold level of DNA topoisomerases leads to the excision of yeast rDNA as extrachromosomal rings. <i>Cell</i> , 1989, 57, 975-985. | 13.5 | 204 |

| # | ARTICLE | IF | CITATIONS |
|-----|---|------|-----------|
| 775 | The STE4 and STE18 genes of yeast encode potential \hat{I}^2 and \hat{I}^3 subunits of the mating factor receptor-coupled G protein. <i>Cell</i> , 1989, 56, 467-477. | 13.5 | 548 |
| 776 | An essential G1 function for cyclin-like proteins in yeast. <i>Cell</i> , 1989, 59, 1127-1133. | 13.5 | 601 |
| 777 | KAR2, a karyogamy gene, is the yeast homolog of the mammalian BiP/GRP78 gene. <i>Cell</i> , 1989, 57, 1211-1221. | 13.5 | 721 |
| 778 | The HIV-1 Gag precursor Pr55gag synthesized in yeast is myristoylated and targeted to the plasma membrane. <i>Gene</i> , 1989, 79, 71-81. | 1.0 | 66 |
| 779 | Identification of a DNA binding factor involved in cell-cycle control of the yeast HO gene. <i>Cell</i> , 1989, 57, 21-29. | 13.5 | 179 |
| 780 | Composite motifs and repeat symmetry in <i>S. pombe</i> centromeres: Direct analysis by integration of NotI restriction sites. <i>Cell</i> , 1989, 57, 739-751. | 13.5 | 226 |
| 781 | Disruption of the single tropomyosin gene in yeast results in the disappearance of actin cables from the cytoskeleton. <i>Cell</i> , 1989, 57, 233-242. | 13.5 | 250 |
| 782 | SPT15, the gene encoding the yeast TATA binding factor TFIID, is required for normal transcription initiation in vivo. <i>Cell</i> , 1989, 58, 1183-1191. | 13.5 | 268 |
| 783 | Identification of functional U1 snRNA-pre-mRNA complexes committed to spliceosome assembly and splicing. <i>Cell</i> , 1989, 59, 349-358. | 13.5 | 396 |
| 784 | Domains of the SFL1 protein of yeasts are homologous to Myc oncoproteins or yeast heat-shock transcription factor. <i>Gene</i> , 1989, 85, 321-328. | 1.0 | 79 |
| 785 | <i>S. cerevisiae</i> encodes an essential protein homologous in sequence and function to mammalian BiP. <i>Cell</i> , 1989, 57, 1223-1236. | 13.5 | 415 |
| 786 | Secretion of glycosylated human erythropoietin from yeast directed by the \hat{I}^{\pm} -factor leader region. <i>Gene</i> , 1989, 79, 167-180. | 1.0 | 62 |
| 787 | The conformation of mature human \hat{I}^{\pm} -amylase conditions its secretion from yeast. <i>Gene</i> , 1989, 83, 355-365. | 1.0 | 15 |
| 788 | The C-terminal part of a gene partially homologous to CDC25 gene suppresses the cdc25-5 mutation in <i>Saccharomyces cerevisiae</i> . <i>Gene</i> , 1989, 77, 21-30. | 1.0 | 64 |
| 789 | Some cis- and trans-acting mutants for splicing target pre-mRNA to the cytoplasm. <i>Cell</i> , 1989, 57, 573-583. | 13.5 | 423 |
| 790 | Primary structure of the maltose-permease-encoding gene of <i>Saccharomyces carlsbergensis</i> . <i>Gene</i> , 1989, 79, 189-197. | 1.0 | 32 |
| 791 | The fission yeast dis2+ gene required for chromosome disjoining encodes one of two putative type 1 protein phosphatases. <i>Cell</i> , 1989, 57, 997-1007. | 13.5 | 515 |
| 792 | Cloning of chromosome I DNA from <i>Saccharomyces cerevisiae</i> : mutational analysis of the FUN2 transcribed region. <i>Gene</i> , 1989, 83, 381-385. | 1.0 | 4 |

| # | ARTICLE | IF | CITATIONS |
|-----|---|------|-----------|
| 793 | The poly(A) binding protein is required for poly(A) shortening and 60S ribosomal subunit-dependent translation initiation. <i>Cell</i> , 1989, 58, 857-867. | 13.5 | 559 |
| 794 | Characterization of AMD, the AMP deaminase gene in yeast. Production of amd strain, cloning, nucleotide sequence, and properties of the protein. <i>Biochemistry</i> , 1989, 28, 8734-8743. | 1.2 | 50 |
| 795 | Expression of cloned <i>Saccharomyces diastaticus</i> glucoamylase under natural and inducible promoters. <i>Biochimica Et Biophysica Acta Gene Regulatory Mechanisms</i> , 1989, 1008, 168-176. | 2.4 | 17 |
| 796 | Double-strand breaks stimulate alternative mechanisms of recombination repair. <i>Journal of Molecular Biology</i> , 1989, 207, 527-541. | 2.0 | 172 |
| 797 | <i>Schizosaccharomyces pombe</i> U4 small nuclear RNA closely resembles vertebrate U4 and is required for growth. <i>Journal of Molecular Biology</i> , 1989, 208, 371-379. | 2.0 | 18 |
| 798 | Site-directed mutagenesis of yeast C1-tetrahydrofolate synthase: analysis of an overlapping active site in a multifunctional enzyme. <i>Biochemistry</i> , 1989, 28, 2099-2106. | 1.2 | 20 |
| 799 | A rapid and efficient procedure for transformation of intact <i>Saccharomyces cerevisiae</i> by electroporation. <i>Biochemical and Biophysical Research Communications</i> , 1989, 164, 1157-1164. | 1.0 | 21 |
| 800 | Secretion of <i>Escherichia coli</i> β -galactosidase in <i>Saccharomyces cerevisiae</i> using the signal sequence from the glucoamylase-encoding STA2 gene. <i>Biochemical and Biophysical Research Communications</i> , 1989, 164, 1331-1338. | 1.0 | 19 |
| 801 | Genetic transfer of the pigment bacteriorhodopsin into the eukaryote <i>Schizosaccharomyces pombe</i> . <i>FEBS Letters</i> , 1989, 243, 137-140. | 1.3 | 51 |
| 802 | Retrotransposon transposition intermediates are encapsidated into virus-like particles. <i>FEBS Letters</i> , 1989, 244, 307-310. | 1.3 | 4 |
| 803 | Expression of the cDNA encoding lipocortin-like 39 kDa protein of guinea pig neutrophils in yeast. <i>FEBS Letters</i> , 1989, 255, 231-236. | 1.3 | 12 |
| 804 | Yeast vacuolar aminopeptidase yscI. <i>FEBS Letters</i> , 1989, 259, 125-129. | 1.3 | 56 |
| 805 | Isolation and characterization of the gene encoding phosphoenolpyruvate carboxykinase from <i>Saccharomyces cerevisiae</i> . <i>FEBS Letters</i> , 1989, 258, 313-316. | 1.3 | 43 |
| 806 | The yeast gene, DBF4, essential for entry into S phase is cell cycle regulated. <i>Experimental Cell Research</i> , 1989, 180, 419-428. | 1.2 | 96 |
| 807 | Synthesis and secretion of human nerve growth factor by <i>Saccharomyces cerevisiae</i> . <i>Gene</i> , 1989, 83, 65-74. | 1.0 | 40 |
| 808 | Dominant effects of tubulin overexpression in <i>Saccharomyces cerevisiae</i> . <i>Molecular and Cellular Biology</i> , 1989, 9, 1049-1059. | 1.1 | 195 |
| 809 | A hyper-recombination mutation in <i>S. cerevisiae</i> identifies a novel eukaryotic topoisomerase. <i>Cell</i> , 1989, 58, 409-419. | 13.5 | 542 |
| 810 | Genetic analysis of mammalian GAP expressed in yeast. <i>Cell</i> , 1989, 59, 681-686. | 13.5 | 156 |

| # | ARTICLE | IF | CITATIONS |
|-----|--|------|-----------|
| 811 | A putative protein kinase overcomes pheromone-induced arrest of cell cycling in <i>S. cerevisiae</i> . <i>Cell</i> , 1989, 58, 1107-1119. | 13.5 | 316 |
| 812 | Distribution of introns in frameshift-suppressor proline-tRNA genes of <i>Saccharomyces cerevisiae</i> . <i>Gene</i> , 1989, 76, 89-97. | 1.0 | 8 |
| 813 | Genetic engineering of filamentous fungi. <i>Science</i> , 1989, 244, 1313-1317. | 6.0 | 149 |
| 814 | Characterization of the prephenate dehydrogenase-encoding gene, <i>TYR1</i> , from <i>Saccharomyces cerevisiae</i> . <i>Gene</i> , 1989, 85, 303-311. | 1.0 | 41 |
| 815 | Characterization of a <i>leuA</i> gene and an ARS element from <i>Mucor circinelloides</i> . <i>Gene</i> , 1989, 84, 335-343. | 1.0 | 114 |
| 816 | Effect of deletions in the 5' noncoding region on the translational efficiency of phosphoglycerate kinase mRNA in yeast. <i>Gene</i> , 1989, 79, 83-95. | 1.0 | 49 |
| 817 | DNA specificity of the bicoid activator protein is determined by homeodomain recognition helix residue 9. <i>Cell</i> , 1989, 57, 1275-1283. | 13.5 | 512 |
| 818 | The yeast secretory pathway is perturbed by mutations in <i>PMR1</i> , a member of a Ca ²⁺ ATPase family. <i>Cell</i> , 1989, 58, 133-145. | 13.5 | 557 |
| 819 | Elevated recombination rates in transcriptionally active DNA. <i>Cell</i> , 1989, 56, 619-630. | 13.5 | 1,692 |
| 820 | Expression of Human and Murine Interleukin-5 in Eukaryotic Systems. <i>DNA and Cell Biology</i> , 1989, 8, 491-501. | 5.1 | 59 |
| 821 | Yeast <i>YAP1</i> encodes a novel form of the jun family of transcriptional activator proteins.. <i>Genes and Development</i> , 1989, 3, 283-292. | 2.7 | 315 |
| 822 | Copper-induced binding of cellular factors to yeast metallothionein upstream activation sequences.. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 1989, 86, 65-69. | 3.3 | 87 |
| 823 | Transcription terminates near the poly(A) site in the <i>CYC1</i> gene of the yeast <i>Saccharomyces cerevisiae</i> .. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 1989, 86, 8348-8352. | 3.3 | 86 |
| 824 | Lack of positional requirements for autonomously replicating sequence elements on artificial yeast chromosomes.. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 1989, 86, 973-977. | 3.3 | 44 |
| 825 | A synthetic intron in a naturally intronless yeast pre-tRNA is spliced efficiently in vivo.. <i>Molecular and Cellular Biology</i> , 1989, 9, 329-331. | 1.1 | 8 |
| 826 | Heteroduplex DNA correction in <i>Saccharomyces cerevisiae</i> is mismatch specific and requires functional PMS genes.. <i>Molecular and Cellular Biology</i> , 1989, 9, 4432-4440. | 1.1 | 194 |
| 827 | DNA damage induction of ribonucleotide reductase.. <i>Molecular and Cellular Biology</i> , 1989, 9, 4932-4940. | 1.1 | 120 |
| 828 | A yeast mutation that stabilizes a plasmid bearing a mutated ARS1 element.. <i>Molecular and Cellular Biology</i> , 1989, 9, 809-816. | 1.1 | 60 |

| # | ARTICLE | IF | CITATIONS |
|-----|---|-----|-----------|
| 829 | Intragenic revertants of yeast invertase variants with secretion-defective leader sequences.. Molecular and Cellular Biology, 1989, 9, 1452-1464. | 1.1 | 6 |
| 830 | Introduction of extra telomeric DNA sequences into <i>Saccharomyces cerevisiae</i> results in telomere elongation.. Molecular and Cellular Biology, 1989, 9, 1488-1497. | 1.1 | 108 |
| 831 | Upstream activation sequence-dependent alteration of chromatin structure and transcription activation of the yeast GAL1-GAL10 genes.. Molecular and Cellular Biology, 1989, 9, 1721-1732. | 1.1 | 75 |
| 832 | Cloning and sequence analysis of the <i>Saccharomyces cerevisiae</i> RAD9 gene and further evidence that its product is required for cell cycle arrest induced by DNA damage.. Molecular and Cellular Biology, 1989, 9, 1882-1896. | 1.1 | 140 |
| 833 | Cell size modulation by CDC25 and RAS2 genes in <i>Saccharomyces cerevisiae</i> .. Molecular and Cellular Biology, 1989, 9, 2715-2723. | 1.1 | 72 |
| 834 | Genomic footprinting of a yeast tRNA gene reveals stable complexes over the 5'-flanking region.. Molecular and Cellular Biology, 1989, 9, 3244-3252. | 1.1 | 75 |
| 835 | Failure to induce a DNA repair gene, RAD54, in <i>Saccharomyces cerevisiae</i> does not affect DNA repair or recombination phenotypes.. Molecular and Cellular Biology, 1989, 9, 3314-3322. | 1.1 | 72 |
| 836 | A gene product needed for induction of allantoin system genes in <i>Saccharomyces cerevisiae</i> but not for their transcriptional activation.. Molecular and Cellular Biology, 1989, 9, 3869-3877. | 1.1 | 24 |
| 837 | Adjacent upstream activation sequence elements synergistically regulate transcription of ADH2 in <i>Saccharomyces cerevisiae</i> .. Molecular and Cellular Biology, 1989, 9, 34-42. | 1.1 | 70 |
| 838 | Sec59 encodes a membrane protein required for core glycosylation in <i>Saccharomyces cerevisiae</i> .. Molecular and Cellular Biology, 1989, 9, 1191-1199. | 1.1 | 46 |
| 839 | Sequences within the spacer region of yeast rRNA cistrons that stimulate 35S rRNA synthesis in vivo mediate RNA polymerase I-dependent promoter and terminator activities.. Molecular and Cellular Biology, 1989, 9, 1243-1254. | 1.1 | 49 |
| 840 | The <i>Saccharomyces cerevisiae</i> CKS1 gene, a homolog of the <i>Schizosaccharomyces pombe</i> suc1+ gene, encodes a subunit of the Cdc28 protein kinase complex.. Molecular and Cellular Biology, 1989, 9, 2034-2041. | 1.1 | 211 |
| 841 | <i>Kluyveromyces lactis</i> maintains <i>Saccharomyces cerevisiae</i> intron-encoded splicing signals.. Molecular and Cellular Biology, 1989, 9, 2208-2213. | 1.1 | 58 |
| 842 | Interaction between transcriptional activator protein LAC9 and negative regulatory protein GAL80.. Molecular and Cellular Biology, 1989, 9, 2950-2956. | 1.1 | 38 |
| 843 | Elements involved in S-adenosylmethionine-mediated regulation of the <i>Saccharomyces cerevisiae</i> MET25 gene.. Molecular and Cellular Biology, 1989, 9, 3292-3298. | 1.1 | 129 |
| 844 | Adenylate cyclases in yeast: a comparison of the genes from <i>Schizosaccharomyces pombe</i> and <i>Saccharomyces cerevisiae</i> .. Proceedings of the National Academy of Sciences of the United States of America, 1989, 86, 5693-5697. | 3.3 | 128 |
| 845 | 3-Hydroxy-3-methylglutaryl-coenzyme A reductase from <i>Arabidopsis thaliana</i> is structurally distinct from the yeast and animal enzymes.. Proceedings of the National Academy of Sciences of the United States of America, 1989, 86, 2779-2783. | 3.3 | 107 |
| 846 | The general control activator protein GCN4 is essential for a basal level of ARO3 gene expression in <i>Saccharomyces cerevisiae</i> .. Molecular and Cellular Biology, 1989, 9, 144-151. | 1.1 | 39 |

| # | ARTICLE | IF | CITATIONS |
|-----|---|-----|-----------|
| 847 | Elongation factor EF-1 alpha gene dosage alters translational fidelity in <i>Saccharomyces cerevisiae</i> .. <i>Molecular and Cellular Biology</i> , 1989, 9, 4571-4575. | 1.1 | 86 |
| 848 | Mutations in the signal sequence of prepro-alpha-factor inhibit both translocation into the endoplasmic reticulum and processing by signal peptidase in yeast cells.. <i>Molecular and Cellular Biology</i> , 1989, 9, 4977-4985. | 1.1 | 16 |
| 849 | Molecular analysis of the SNF4 gene of <i>Saccharomyces cerevisiae</i> : evidence for physical association of the SNF4 protein with the SNF1 protein kinase.. <i>Molecular and Cellular Biology</i> , 1989, 9, 5045-5054. | 1.1 | 209 |
| 850 | Identification of an upstream activation sequence and other cis-acting elements required for transcription of COX6 from <i>Saccharomyces cerevisiae</i> .. <i>Molecular and Cellular Biology</i> , 1989, 9, 5350-5358. | 1.1 | 31 |
| 851 | Interaction of the H4 autonomously replicating sequence core consensus sequence and its 3'-flanking domain.. <i>Molecular and Cellular Biology</i> , 1989, 9, 5464-5472. | 1.1 | 40 |
| 852 | Context affects nuclear protein localization in <i>Saccharomyces cerevisiae</i> .. <i>Molecular and Cellular Biology</i> , 1989, 9, 384-389. | 1.1 | 136 |
| 853 | A cysteine-rich nuclear protein activates yeast metallothionein gene transcription.. <i>Molecular and Cellular Biology</i> , 1989, 9, 421-429. | 1.1 | 105 |
| 854 | Introduction of functional artificial introns into the naturally intronless <i>ura4</i> gene of <i>Schizosaccharomyces pombe</i> .. <i>Molecular and Cellular Biology</i> , 1989, 9, 1526-1535. | 1.1 | 39 |
| 855 | Localization of a minimal binding domain and activation regions in yeast regulatory protein ADR1.. <i>Molecular and Cellular Biology</i> , 1989, 9, 2360-2369. | 1.1 | 53 |
| 856 | The OBF1 protein and its DNA-binding site are important for the function of an autonomously replicating sequence in <i>Saccharomyces cerevisiae</i> .. <i>Molecular and Cellular Biology</i> , 1989, 9, 2914-2921. | 1.1 | 62 |
| 857 | <i>hsp82</i> is an essential protein that is required in higher concentrations for growth of cells at higher temperatures.. <i>Molecular and Cellular Biology</i> , 1989, 9, 3919-3930. | 1.1 | 621 |
| 858 | Regulation of alpha-factor production in <i>Saccharomyces cerevisiae</i> : a-factor pheromone-induced expression of the MF alpha 1 and STE13 genes.. <i>Molecular and Cellular Biology</i> , 1989, 9, 4507-4514. | 1.1 | 39 |
| 859 | The HML mating-type cassette of <i>Saccharomyces cerevisiae</i> is regulated by two separate but functionally equivalent silencers.. <i>Molecular and Cellular Biology</i> , 1989, 9, 4621-4630. | 1.1 | 121 |
| 860 | Role of glutamic acid 177 of the ricin toxin A chain in enzymatic inactivation of ribosomes.. <i>Molecular and Cellular Biology</i> , 1989, 9, 5012-5021. | 1.1 | 86 |
| 861 | IRA1, an inhibitory regulator of the RAS-cyclic AMP pathway in <i>Saccharomyces cerevisiae</i> .. <i>Molecular and Cellular Biology</i> , 1989, 9, 757-768. | 1.1 | 275 |
| 862 | GPA1Val-50 mutation in the mating-factor signaling pathway in <i>Saccharomyces cerevisiae</i> .. <i>Molecular and Cellular Biology</i> , 1989, 9, 2289-2297. | 1.1 | 75 |
| 863 | Use of yeast nuclear DNA sequences to define the mitochondrial RNA polymerase promoter in vitro.. <i>Molecular and Cellular Biology</i> , 1989, 9, 3193-3202. | 1.1 | 34 |
| 864 | Juxtaposition of domains homologous to protein kinases and histidyl-tRNA synthetases in GCN2 protein suggests a mechanism for coupling GCN4 expression to amino acid availability.. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 1989, 86, 4579-4583. | 3.3 | 271 |

| # | ARTICLE | IF | CITATIONS |
|-----|---|-----|-----------|
| 865 | Construction of functional artificial minichromosomes in the fission yeast <i>Schizosaccharomyces pombe</i> .. Proceedings of the National Academy of Sciences of the United States of America, 1989, 86, 577-581. | 3.3 | 123 |
| 866 | A transcriptional cascade governs entry into meiosis in <i>Saccharomyces cerevisiae</i> .. Molecular and Cellular Biology, 1989, 9, 2142-2152. | 1.1 | 192 |
| 867 | The yeast G-protein homolog is involved in the mating pheromone signal transduction system.. Molecular and Cellular Biology, 1989, 9, 152-158. | 1.1 | 22 |
| 868 | Basal-level expression of the yeast HSP82 gene requires a heat shock regulatory element.. Molecular and Cellular Biology, 1989, 9, 4789-4798. | 1.1 | 75 |
| 869 | Positive and negative regulation of basal expression of a yeast HSP70 gene.. Molecular and Cellular Biology, 1989, 9, 2025-2033. | 1.1 | 81 |
| 870 | Isolation and characterization of mutations in the HXK2 gene of <i>Saccharomyces cerevisiae</i> .. Molecular and Cellular Biology, 1989, 9, 5630-5642. | 1.1 | 37 |
| 871 | The yeast cell fusion protein FUS1 is O-glycosylated and spans the plasma membrane.. Proceedings of the National Academy of Sciences of the United States of America, 1989, 86, 9916-9920. | 3.3 | 79 |
| 872 | A plant manganese superoxide dismutase is efficiently imported and correctly processed by yeast mitochondria.. Proceedings of the National Academy of Sciences of the United States of America, 1989, 86, 3237-3241. | 3.3 | 72 |
| 873 | Differential repression of GAL4 and adjacent transcription activators by operators in the yeast GAL upstream activating sequence.. Molecular and Cellular Biology, 1989, 9, 4282-4290. | 1.1 | 24 |
| 874 | Identification of a yeast protein with properties similar to those of the immunoglobulin heavy-chain enhancer-binding protein NF- μ E3.. Molecular and Cellular Biology, 1989, 9, 4535-4540. | 1.1 | 10 |
| 875 | Isolation of constitutive mutations affecting the proline utilization pathway in <i>Saccharomyces cerevisiae</i> and molecular analysis of the PUT3 transcriptional activator.. Molecular and Cellular Biology, 1989, 9, 4696-4705. | 1.1 | 34 |
| 876 | Enhancer and silencerlike sites within the transcribed portion of a Ty2 transposable element of <i>Saccharomyces cerevisiae</i> .. Molecular and Cellular Biology, 1989, 9, 4824-4834. | 1.1 | 47 |
| 877 | Unusual enhancer function in yeast rRNA transcription.. Molecular and Cellular Biology, 1989, 9, 4986-4993. | 1.1 | 68 |
| 878 | Mutational analysis of the <i>Saccharomyces cerevisiae</i> SNF1 protein kinase and evidence for functional interaction with the SNF4 protein.. Molecular and Cellular Biology, 1989, 9, 5034-5044. | 1.1 | 238 |
| 879 | Upstream regulatory sequences of the yeast RNR2 gene include a repression sequence and an activation site that binds the RAP1 protein.. Molecular and Cellular Biology, 1989, 9, 5359-5372. | 1.1 | 59 |
| 880 | Identification of the DNA damage-responsive element of RNR2 and evidence that four distinct cellular factors bind it.. Molecular and Cellular Biology, 1989, 9, 5373-5386. | 1.1 | 77 |
| 881 | The <i>Saccharomyces cerevisiae</i> SPT13/GAL11 gene has both positive and negative regulatory roles in transcription.. Molecular and Cellular Biology, 1989, 9, 5602-5609. | 1.1 | 125 |
| 882 | Selection and characterization of ricin toxin A-chain mutations in <i>Saccharomyces cerevisiae</i> .. Molecular and Cellular Biology, 1989, 9, 415-420. | 1.1 | 62 |

| # | ARTICLE | IF | CITATIONS |
|-----|--|-----|-----------|
| 883 | Amino-terminal extension generated from an upstream AUG codon increases the efficiency of mitochondrial import of yeast N2,N2-dimethylguanosine-specific tRNA methyltransferases.. <i>Molecular and Cellular Biology</i> , 1989, 9, 1611-1620. | 1.1 | 48 |
| 884 | Molecular analysis of GPH1, the gene encoding glycogen phosphorylase in <i>Saccharomyces cerevisiae</i> .. <i>Molecular and Cellular Biology</i> , 1989, 9, 1659-1666. | 1.1 | 101 |
| 885 | Structural and functional analyses of <i>Saccharomyces cerevisiae</i> wild-type and mutant RNA1 genes.. <i>Molecular and Cellular Biology</i> , 1989, 9, 2989-2999. | 1.1 | 74 |
| 886 | Functional domains of a negative regulatory protein, GAL80, of <i>Saccharomyces cerevisiae</i> .. <i>Molecular and Cellular Biology</i> , 1989, 9, 3009-3017. | 1.1 | 44 |
| 887 | PRP4 (RNA4) from <i>Saccharomyces cerevisiae</i> : its gene product is associated with the U4/U6 small nuclear ribonucleoprotein particle.. <i>Molecular and Cellular Biology</i> , 1989, 9, 3698-3709. | 1.1 | 105 |
| 888 | Activation of yeast RNA polymerase II transcription by a thymidine-rich upstream element in vitro.. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 1989, 86, 486-490. | 3.3 | 120 |
| 889 | An essential yeast protein, encoded by duplicated genes TIF1 and TIF2 and homologous to the mammalian translation initiation factor eIF-4A, can suppress a mitochondrial missense mutation.. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 1989, 86, 2286-2290. | 3.3 | 135 |
| 890 | Yeast translation initiation suppressor <i>sui2</i> encodes the alpha subunit of eukaryotic initiation factor 2 and shares sequence identity with the human alpha subunit.. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 1989, 86, 2784-2788. | 3.3 | 222 |
| 891 | Isolation and characterization of a mammalian gene encoding a high-affinity cAMP phosphodiesterase.. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 1989, 86, 3599-3603. | 3.3 | 188 |
| 892 | A family of cyclin homologs that control the G1 phase in yeast.. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 1989, 86, 6255-6259. | 3.3 | 422 |
| 893 | Mutations in the structural genes for eukaryotic initiation factors 2 alpha and 2 beta of <i>Saccharomyces cerevisiae</i> disrupt translational control of GCN4 mRNA.. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 1989, 86, 7515-7519. | 3.3 | 105 |
| 894 | Identification of negative regulatory genes that govern the expression of early meiotic genes in yeast.. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 1989, 86, 10018-10022. | 3.3 | 151 |
| 895 | Mutations in RNA polymerase II enhance or suppress mutations in GAL4.. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 1989, 86, 2794-2798. | 3.3 | 126 |
| 896 | Regulation of STA1 gene expression by MAT during the life cycle of <i>Saccharomyces cerevisiae</i> .. <i>Molecular and Cellular Biology</i> , 1989, 9, 3992-3998. | 1.1 | 41 |
| 897 | N alpha acetylation is required for normal growth and mating of <i>Saccharomyces cerevisiae</i> . <i>Journal of Bacteriology</i> , 1989, 171, 5795-5802. | 1.0 | 62 |
| 898 | The REV1 gene of <i>Saccharomyces cerevisiae</i> : isolation, sequence, and functional analysis. <i>Journal of Bacteriology</i> , 1989, 171, 230-237. | 1.0 | 140 |
| 899 | REV3, a <i>Saccharomyces cerevisiae</i> gene whose function is required for induced mutagenesis, is predicted to encode a nonessential DNA polymerase. <i>Journal of Bacteriology</i> , 1989, 171, 5659-5667. | 1.0 | 273 |
| 900 | Expression and transport of <i>Candida tropicalis</i> peroxisomal acyl-coenzyme A oxidase in the yeast <i>Candida maltosa</i> .. <i>Agricultural and Biological Chemistry</i> , 1989, 53, 179-186. | 0.3 | 43 |

| # | ARTICLE | IF | CITATIONS |
|-----|--|-----|-----------|
| 901 | Chromosomal transformation in <i>Saccharomyces cerevisiae</i> with DNA isolated by pulse field gel electrophoresis.. <i>Agricultural and Biological Chemistry</i> , 1990, 54, 1499-1504. | 0.3 | 4 |
| 902 | Lysis protein S of phage lambda functions in <i>Saccharomyces cerevisiae</i> . <i>Journal of Bacteriology</i> , 1990, 172, 7275-7277. | 1.0 | 24 |
| 903 | Expression and secretion of bovine .BETA.-lactoglobulin in <i>Saccharomyces cerevisiae</i> .. <i>Agricultural and Biological Chemistry</i> , 1990, 54, 3111-3116. | 0.3 | 27 |
| 904 | Overexpression of the STE4 gene leads to mating response in haploid <i>Saccharomyces cerevisiae</i> .. <i>Molecular and Cellular Biology</i> , 1990, 10, 217-222. | 1.1 | 158 |
| 905 | Ribosomal protein L30 is dispensable in the yeast <i>Saccharomyces cerevisiae</i> .. <i>Molecular and Cellular Biology</i> , 1990, 10, 5235-5243. | 1.1 | 61 |
| 906 | A suppressor of an RNA polymerase II mutation of <i>Saccharomyces cerevisiae</i> encodes a subunit common to RNA polymerases I, II, and III.. <i>Molecular and Cellular Biology</i> , 1990, 10, 6123-6131. | 1.1 | 50 |
| 907 | Genetic and biochemical evaluation of eucaryotic membrane protein topology: multiple transmembrane domains of <i>Saccharomyces cerevisiae</i> 3-hydroxy-3-methylglutaryl coenzyme A reductase.. <i>Molecular and Cellular Biology</i> , 1990, 10, 672-680. | 1.1 | 76 |
| 908 | A yeast ARS-binding protein activates transcription synergistically in combination with other weak activating factors.. <i>Molecular and Cellular Biology</i> , 1990, 10, 887-897. | 1.1 | 154 |
| 909 | Vanadate-resistant mutants of <i>Saccharomyces cerevisiae</i> show alterations in protein phosphorylation and growth control.. <i>Molecular and Cellular Biology</i> , 1990, 10, 898-909. | 1.1 | 48 |
| 910 | Conditional expression of RPA190, the gene encoding the largest subunit of yeast RNA polymerase I: effects of decreased rRNA synthesis on ribosomal protein synthesis.. <i>Molecular and Cellular Biology</i> , 1990, 10, 2049-2059. | 1.1 | 50 |
| 911 | SNF6 encodes a nuclear protein that is required for expression of many genes in <i>Saccharomyces cerevisiae</i> .. <i>Molecular and Cellular Biology</i> , 1990, 10, 2544-2553. | 1.1 | 79 |
| 912 | IRA2, a second gene of <i>Saccharomyces cerevisiae</i> that encodes a protein with a domain homologous to mammalian ras GTPase-activating protein.. <i>Molecular and Cellular Biology</i> , 1990, 10, 4303-4313. | 1.1 | 300 |
| 913 | Cloning by function: an alternative approach for identifying yeast homologs of genes from other organisms.. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 1990, 87, 6629-6633. | 3.3 | 166 |
| 914 | Functional analysis of a centromere from fission yeast: a role for centromere-specific repeated DNA sequences.. <i>Molecular and Cellular Biology</i> , 1990, 10, 1863-1872. | 1.1 | 109 |
| 915 | <i>Drosophila</i> scaffold-attached regions bind nuclear scaffolds and can function as ARS elements in both budding and fission yeasts.. <i>Molecular and Cellular Biology</i> , 1990, 10, 5442-5454. | 1.1 | 97 |
| 916 | Molecular genetic analysis of <i>Saccharomyces cerevisiae</i> C1-tetrahydrofolate synthase mutants reveals a noncatalytic function of the ADE3 gene product and an additional folate-dependent enzyme.. <i>Molecular and Cellular Biology</i> , 1990, 10, 5679-5687. | 1.1 | 42 |
| 917 | Role of IME1 expression in regulation of meiosis in <i>Saccharomyces cerevisiae</i> .. <i>Molecular and Cellular Biology</i> , 1990, 10, 6103-6113. | 1.1 | 113 |
| 918 | Sequence and functional similarity between a yeast ribosomal protein and the <i>Escherichia coli</i> S5 ram protein.. <i>Molecular and Cellular Biology</i> , 1990, 10, 6544-6553. | 1.1 | 67 |

| # | ARTICLE | IF | CITATIONS |
|-----|---|-----|-----------|
| 919 | Characterization of RAD9 of <i>Saccharomyces cerevisiae</i> and evidence that its function acts posttranslationally in cell cycle arrest after DNA damage.. <i>Molecular and Cellular Biology</i> , 1990, 10, 6554-6564. | 1.1 | 222 |
| 920 | Mutational analysis of the SNF3 glucose transporter of <i>Saccharomyces cerevisiae</i> .. <i>Molecular and Cellular Biology</i> , 1990, 10, 1105-1115. | 1.1 | 78 |
| 921 | Enhancer and promoter elements from simian virus 40 and polyomavirus can substitute for an upstream activation sequence in <i>Saccharomyces cerevisiae</i> .. <i>Molecular and Cellular Biology</i> , 1990, 10, 947-957. | 1.1 | 7 |
| 922 | The product of the <i>Saccharomyces cerevisiae</i> cell cycle gene DBF2 has homology with protein kinases and is periodically expressed in the cell cycle.. <i>Molecular and Cellular Biology</i> , 1990, 10, 1358-1366. | 1.1 | 135 |
| 923 | Mutations in cell division cycle genes CDC36 and CDC39 activate the <i>Saccharomyces cerevisiae</i> mating pheromone response pathway.. <i>Molecular and Cellular Biology</i> , 1990, 10, 2966-2972. | 1.1 | 40 |
| 924 | Modification and transfer into an embryonal carcinoma cell line of a 360-kilobase human-derived yeast artificial chromosome.. <i>Molecular and Cellular Biology</i> , 1990, 10, 4163-4169. | 1.1 | 93 |
| 925 | A conserved sequence in histone H2A which is a ubiquitination site in higher eucaryotes is not required for growth in <i>Saccharomyces cerevisiae</i> .. <i>Molecular and Cellular Biology</i> , 1990, 10, 4905-4911. | 1.1 | 46 |
| 926 | PRP18, a protein required for the second reaction in pre-mRNA splicing.. <i>Molecular and Cellular Biology</i> , 1990, 10, 324-332. | 1.1 | 75 |
| 927 | HPR1, a novel yeast gene that prevents intrachromosomal excision recombination, shows carboxy-terminal homology to the <i>Saccharomyces cerevisiae</i> TOP1 gene.. <i>Molecular and Cellular Biology</i> , 1990, 10, 1439-1451. | 1.1 | 137 |
| 928 | Courtship in <i>Saccharomyces cerevisiae</i> : an early cell-cell interaction during mating.. <i>Molecular and Cellular Biology</i> , 1990, 10, 2202-2213. | 1.1 | 84 |
| 929 | Expression and function of a human initiator tRNA gene in the yeast <i>Saccharomyces cerevisiae</i> .. <i>Molecular and Cellular Biology</i> , 1990, 10, 4486-4494. | 1.1 | 33 |
| 930 | Thermal energy suppresses mutational defects in DNA unwinding at a yeast replication origin.. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 1990, 87, 2486-2490. | 3.3 | 66 |
| 931 | In vivo expression and mitochondrial targeting of yeast apoiso-1-cytochrome c fusion proteins.. <i>Molecular and Cellular Biology</i> , 1990, 10, 5753-5762. | 1.1 | 18 |
| 932 | Molecular and expression analysis of the negative regulators involved in the transcriptional regulation of acid phosphatase production in <i>Saccharomyces cerevisiae</i> .. <i>Molecular and Cellular Biology</i> , 1990, 10, 5950-5957. | 1.1 | 26 |
| 933 | Stoichiometry of G protein subunits affects the <i>Saccharomyces cerevisiae</i> mating pheromone signal transduction pathway.. <i>Molecular and Cellular Biology</i> , 1990, 10, 510-517. | 1.1 | 157 |
| 934 | Mutations in RAD6, a yeast gene encoding a ubiquitin-conjugating enzyme, stimulate retrotransposition.. <i>Molecular and Cellular Biology</i> , 1990, 10, 1017-1022. | 1.1 | 79 |
| 935 | In vivo analysis of the <i>Saccharomyces cerevisiae</i> HO nuclease recognition site by site-directed mutagenesis.. <i>Molecular and Cellular Biology</i> , 1990, 10, 1174-1179. | 1.1 | 78 |
| 936 | Transcription of a yeast phosphoglucomutase isozyme gene is galactose inducible and glucose repressible.. <i>Molecular and Cellular Biology</i> , 1990, 10, 1415-1422. | 1.1 | 67 |

| # | ARTICLE | IF | CITATIONS |
|-----|--|-----|-----------|
| 937 | The SLP1 gene of <i>Saccharomyces cerevisiae</i> is essential for vacuolar morphogenesis and function.. <i>Molecular and Cellular Biology</i> , 1990, 10, 2214-2223. | 1.1 | 125 |
| 938 | Null alleles of <i>SAC7</i> suppress temperature-sensitive actin mutations in <i>Saccharomyces cerevisiae</i> .. <i>Molecular and Cellular Biology</i> , 1990, 10, 2308-2314. | 1.1 | 53 |
| 939 | Role of vacuolar acidification in protein sorting and zymogen activation: a genetic analysis of the yeast vacuolar proton-translocating ATPase.. <i>Molecular and Cellular Biology</i> , 1990, 10, 3737-3749. | 1.1 | 188 |
| 940 | Phosphorylated forms of <i>GAL4</i> are correlated with ability to activate transcription.. <i>Molecular and Cellular Biology</i> , 1990, 10, 4623-4629. | 1.1 | 70 |
| 941 | The N-terminal TPR region is the functional domain of <i>SSN6</i> , a nuclear phosphoprotein of <i>Saccharomyces cerevisiae</i> .. <i>Molecular and Cellular Biology</i> , 1990, 10, 4744-4756. | 1.1 | 143 |
| 942 | Evidence for a heat shock transcription factor-independent mechanism for heat shock induction of transcription in <i>Saccharomyces cerevisiae</i> .. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 1990, 87, 6550-6554. | 3.3 | 64 |
| 943 | Mutations that increase the activity of a transcriptional activator in yeast and mammalian cells.. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 1990, 87, 2127-2131. | 3.3 | 71 |
| 944 | Mutation of cysteine-88 in the <i>Saccharomyces cerevisiae</i> <i>RAD6</i> protein abolishes its ubiquitin-conjugating activity and its various biological functions.. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 1990, 87, 2695-2699. | 3.3 | 147 |
| 945 | Disruption of genes encoding subunits of yeast vacuolar H(+)-ATPase causes conditional lethality.. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 1990, 87, 3503-3507. | 3.3 | 311 |
| 946 | Adenylate cyclase in <i>Saccharomyces cerevisiae</i> is a peripheral membrane protein.. <i>Molecular and Cellular Biology</i> , 1990, 10, 3873-3883. | 1.1 | 44 |
| 947 | Characterization of yeast <i>Vps33p</i> , a protein required for vacuolar protein sorting and vacuole biogenesis.. <i>Molecular and Cellular Biology</i> , 1990, 10, 4638-4649. | 1.1 | 123 |
| 948 | Human DNA ligase I cDNA: cloning and functional expression in <i>Saccharomyces cerevisiae</i> .. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 1990, 87, 6679-6683. | 3.3 | 145 |
| 949 | The mouse c-rel protein has an N-terminal regulatory domain and a C-terminal transcriptional transactivation domain.. <i>Molecular and Cellular Biology</i> , 1990, 10, 5473-5485. | 1.1 | 172 |
| 950 | The yeast <i>PRP6</i> gene encodes a U4/U6 small nuclear ribonucleoprotein particle (snRNP) protein, and the <i>PRP9</i> gene encodes a protein required for U2 snRNP binding.. <i>Molecular and Cellular Biology</i> , 1990, 10, 6417-6425. | 1.1 | 114 |
| 951 | Mutational mapping of RAS-responsive domains of the <i>Saccharomyces cerevisiae</i> adenylyl cyclase.. <i>Molecular and Cellular Biology</i> , 1990, 10, 2539-2543. | 1.1 | 61 |
| 952 | Coding and noncoding sequences at the 3' end of yeast histone H2B mRNA confer cell cycle regulation.. <i>Molecular and Cellular Biology</i> , 1990, 10, 2687-2694. | 1.1 | 46 |
| 953 | Identification of positive-acting domains in <i>GCN2</i> protein kinase required for translational activation of <i>GCN4</i> expression.. <i>Molecular and Cellular Biology</i> , 1990, 10, 2820-2831. | 1.1 | 123 |
| 954 | Mutational analysis of the consensus sequence of a replication origin from yeast chromosome III.. <i>Molecular and Cellular Biology</i> , 1990, 10, 3917-3925. | 1.1 | 176 |

| # | ARTICLE | IF | CITATIONS |
|-----|---|-----|-----------|
| 955 | Farnesyl cysteine C-terminal methyltransferase activity is dependent upon the STE14 gene product in <i>Saccharomyces cerevisiae</i> .. <i>Molecular and Cellular Biology</i> , 1990, 10, 5071-5076. | 1.1 | 120 |
| 956 | The SNF5 protein of <i>Saccharomyces cerevisiae</i> is a glutamine- and proline-rich transcriptional activator that affects expression of a broad spectrum of genes.. <i>Molecular and Cellular Biology</i> , 1990, 10, 5616-5625. | 1.1 | 155 |
| 957 | The <i>Saccharomyces cerevisiae</i> SIN3 gene, a negative regulator of HO, contains four paired amphipathic helix motifs.. <i>Molecular and Cellular Biology</i> , 1990, 10, 5927-5936. | 1.1 | 166 |
| 958 | Functional analysis of a duplicated linked pair of ribosomal protein genes in <i>Saccharomyces cerevisiae</i> .. <i>Molecular and Cellular Biology</i> , 1990, 10, 6097-6100. | 1.1 | 14 |
| 959 | gcr2, a new mutation affecting glycolytic gene expression in <i>Saccharomyces cerevisiae</i> .. <i>Molecular and Cellular Biology</i> , 1990, 10, 6389-6396. | 1.1 | 76 |
| 960 | HOL1 mutations confer novel ion transport in <i>Saccharomyces cerevisiae</i> .. <i>Molecular and Cellular Biology</i> , 1990, 10, 643-652. | 1.1 | 26 |
| 961 | Self-regulation of 70-kilodalton heat shock proteins in <i>Saccharomyces cerevisiae</i> .. <i>Molecular and Cellular Biology</i> , 1990, 10, 1622-1632. | 1.1 | 132 |
| 962 | MER1, a yeast gene required for chromosome pairing and genetic recombination, is induced in meiosis.. <i>Molecular and Cellular Biology</i> , 1990, 10, 2379-2389. | 1.1 | 149 |
| 963 | Isolation of the gene encoding the <i>Saccharomyces cerevisiae</i> centromere-binding protein CP1.. <i>Molecular and Cellular Biology</i> , 1990, 10, 2458-2467. | 1.1 | 186 |
| 964 | TEC1, a gene involved in the activation of Ty1 and Ty1-mediated gene expression in <i>Saccharomyces cerevisiae</i> : cloning and molecular analysis.. <i>Molecular and Cellular Biology</i> , 1990, 10, 3541-3550. | 1.1 | 109 |
| 965 | Sequences within an upstream activation site in the yeast enolase gene ENO2 modulate repression of ENO2 expression in strains carrying a null mutation in the positive regulatory gene GCR1.. <i>Molecular and Cellular Biology</i> , 1990, 10, 4863-4871. | 1.1 | 30 |
| 966 | A gene family for acidic ribosomal proteins in <i>Schizosaccharomyces pombe</i> : two essential and two nonessential genes.. <i>Molecular and Cellular Biology</i> , 1990, 10, 2341-2348. | 1.1 | 45 |
| 967 | Transcriptional derepression of the <i>Saccharomyces cerevisiae</i> HSP26 gene during heat shock.. <i>Molecular and Cellular Biology</i> , 1990, 10, 6362-6373. | 1.1 | 49 |
| 968 | Transport of the yeast ATP synthase \hat{f} 2-subunit into mitochondria. Effects of amino acid substitutions on targeting. <i>Biochemical Journal</i> , 1990, 266, 227-234. | 1.7 | 9 |
| 969 | Expression of human glutathione S-transferases in <i>Saccharomyces cerevisiae</i> confers resistance to the anticancer drugs adriamycin and chlorambucil. <i>Biochemical Journal</i> , 1990, 268, 309-315. | 1.7 | 130 |
| 970 | Role of NAD-linked glutamate dehydrogenase in nitrogen metabolism in <i>Saccharomyces cerevisiae</i> . <i>Journal of Bacteriology</i> , 1990, 172, 4927-4935. | 1.0 | 144 |
| 971 | Chromosome engineering in <i>Saccharomyces cerevisiae</i> by using a site-specific recombination system of a yeast plasmid. <i>Journal of Bacteriology</i> , 1990, 172, 610-618. | 1.0 | 112 |
| 972 | Structural and functional stability of IncP plasmids during stepwise transmission by trans-kingdom mating: promiscuous conjugation of <i>Escherichia coli</i> and <i>Saccharomyces cerevisiae</i> .. <i>Japanese Journal of Genetics</i> , 1990, 65, 323-334. | 1.0 | 38 |

| # | ARTICLE | IF | CITATIONS |
|-----|---|------|-----------|
| 973 | Yeast suppressor mutations and transfer RNA processing. <i>Methods in Enzymology</i> , 1990, 181, 377-394. | 0.4 | 13 |
| 974 | Genetic methods for study of trans-acting genes involved in processing of precursors to yeast cytoplasmic transfer RNAs. <i>Methods in Enzymology</i> , 1990, 181, 400-421. | 0.4 | 16 |
| 975 | Prolongation of the yeast life span by the v-Ha-RAS oncogene. <i>Molecular Microbiology</i> , 1990, 4, 2081-2086. | 1.2 | 54 |
| 976 | Isolation of a chitin synthase gene (CHS 1) from <i>Candida albicans</i> by expression in <i>Saccharomyces cerevisiae</i> . <i>Molecular Microbiology</i> , 1990, 4, 197-207. | 1.2 | 108 |
| 977 | Identification and characterization of a gene and protein required for glycosylation in the yeast Golgi. <i>Molecular Microbiology</i> , 1990, 4, 1993-2001. | 1.2 | 18 |
| 978 | Plasmid transformation in <i>Bacillus subtilis</i> NB22, an antifungal-antibiotic iturin producer. <i>FEMS Microbiology Letters</i> , 1990, 67, 227-230. | 0.7 | 2 |
| 979 | <i>Kluyveromyces</i> as a Host for Heterologous Gene Expression: Expression and Secretion of Prochymosin. <i>Nature Biotechnology</i> , 1990, 8, 135-139. | 9.4 | 141 |
| 980 | High Efficiency Transformation of Intact Yeast Cells by Electric Field Pulses. <i>Nature Biotechnology</i> , 1990, 8, 223-227. | 9.4 | 119 |
| 981 | Expression in Yeast of Amino-Terminal Peptide Fusions to Hepatitis B Core Antigen and Their Immunological Properties. <i>Nature Biotechnology</i> , 1990, 8, 644-649. | 9.4 | 31 |
| 982 | Homologous activators of ras in fission and budding yeast. <i>Nature</i> , 1990, 344, 355-357. | 13.7 | 158 |
| 983 | Human U2 snRNA can function in pre-mRNA splicing in yeast. <i>Nature</i> , 1990, 345, 270-273. | 13.7 | 48 |
| 984 | Escape of DNA from mitochondria to the nucleus in <i>Saccharomyces cerevisiae</i> . <i>Nature</i> , 1990, 346, 376-379. | 13.7 | 259 |
| 985 | GAP1, the general amino acid permease gene of <i>Saccharomyces cerevisiae</i> . Nucleotide sequence, protein similarity with the other bakers yeast amino acid permeases, and nitrogen catabolite repression. <i>FEBS Journal</i> , 1990, 190, 39-44. | 0.2 | 275 |
| 986 | Phosphorylation of yeast hexokinases. <i>FEBS Journal</i> , 1990, 190, 371-375. | 0.2 | 60 |
| 987 | Isolation of a gene for a regulatory 15-kDa subunit of mitochondrial F1F0-ATPase and construction of mutant yeast lacking the protein. <i>FEBS Journal</i> , 1990, 192, 49-53. | 0.2 | 24 |
| 988 | REPAIR OF UV-DAMAGED INCOMING PLASMID DNA IN <i>Saccharomyces cerevisiae</i> . <i>Photochemistry and Photobiology</i> , 1990, 51, 331-342. | 1.3 | 9 |
| 989 | CHARACTERIZATION OF RAD4 GENE REQUIRED FOR ULTRAVIOLET-INDUCED EXCISION REPAIR OF <i>Saccharomyces cerevisiae</i> PROPAGATED IN <i>Escherichia coli</i> WITHOUT INACTIVATION. <i>Photochemistry and Photobiology</i> , 1990, 52, 395-400. | 1.3 | 4 |
| 990 | Structural and functional analyses of a yeast mitochondrial ribosomal protein homologous to ribosomal protein S15 of <i>Escherichia coli</i> . <i>Nucleic Acids Research</i> , 1990, 18, 6895-6901. | 6.5 | 30 |

| # | ARTICLE | IF | CITATIONS |
|------|--|-----|-----------|
| 991 | A short domain of the plant vacuolar protein phytohemagglutinin targets invertase to the yeast vacuole.. Plant Cell, 1990, 2, 533-546. | 3.1 | 73 |
| 992 | Yeast-derived recombinant human insulin-like growth factor I: Production, purification, and structural characterization. The Protein Journal, 1990, 9, 95-104. | 1.1 | 52 |
| 993 | A system for temperature-controlled expression of a foreign gene with dual mode in <i>Saccharomyces cerevisiae</i> . Journal of Bioscience and Bioengineering, 1990, 69, 322-327. | 0.9 | 12 |
| 994 | Heterogeneity and maintenance of centromere plasmid copy number in <i>Saccharomyces cerevisiae</i> . Chromosoma, 1990, 99, 281-288. | 1.0 | 13 |
| 995 | Detection of specific RNA sequences in yeast by in situ colony hybridization. Yeast, 1990, 6, 31-34. | 0.8 | 1 |
| 996 | The start gene <i>CDC28</i> and the genetic stability of yeast. Yeast, 1990, 6, 231-243. | 0.8 | 34 |
| 997 | The hydrophilic and acidic N-terminus of the integral membrane enzyme phosphatidylserine synthase is required for efficient membrane insertion. Yeast, 1990, 6, 331-343. | 0.8 | 21 |
| 998 | Characterization of the cytochrome c gene from the starch-fermenting yeast <i>Schwanniomyces occidentalis</i> and its expression in bakers's yeast. Yeast, 1990, 6, 429-440. | 0.8 | 25 |
| 999 | The omnipotent suppressor <i>SUP45</i> affects nucleic acid metabolism and mitochondrial structure. Yeast, 1990, 6, 441-450. | 0.8 | 10 |
| 1000 | Ornithine decarboxylase in <i>Saccharomyces cerevisiae</i> : Chromosomal assignment and genetic mapping of the <i>SPE1</i> gene. Yeast, 1990, 6, 455-460. | 0.8 | 13 |
| 1001 | Divergence and conservation of <i>SUP2</i> (<i>SUP35</i>) gene of yeasts <i>Pichia pinus</i> and <i>Saccharomyces cerevisiae</i> . Yeast, 1990, 6, 461-472. | 0.8 | 70 |
| 1002 | Cloning and expression of β -galactosidase gene from <i>Streptococcus thermophilus</i> in <i>Saccharomyces cerevisiae</i> . Biotechnology Letters, 1990, 12, 499-504. | 1.1 | 8 |
| 1003 | Molecular basis for determining the sensitivity of eucaryotes to the antimetabolic drug rhizoxin. Molecular Genetics and Genomics, 1990, 222, 169-175. | 2.4 | 41 |
| 1004 | Transformation of <i>Bacillus subtilis</i> with the treatment by alkali cations. Biotechnology Letters, 1990, 12, 99-104. | 1.1 | 9 |
| 1005 | Genetic mapping and biochemical analysis of mutants in the maltose regulatory gene of the <i>MAL1</i> locus of <i>Saccharomyces cerevisiae</i> . Archives of Microbiology, 1990, 154, 544-9. | 1.0 | 6 |
| 1006 | Development of a genetic transformation system for <i>Histoplasma capsulatum</i> : Complementation of uracil auxotrophy. Molecular Genetics and Genomics, 1990, 221, 358-362. | 2.4 | 51 |
| 1007 | Effects of seven different mutations in the <i>pho1</i> gene on enzymatic activity, glycosylation and secretion of acid phosphatase in <i>Schizosaccharomyces pombe</i> . Molecular Genetics and Genomics, 1990, 221, 403-410. | 2.4 | 11 |
| 1008 | A <i>GCN4</i> protein recognition element is not sufficient for <i>GCN4</i> -dependent regulation of transcription in the <i>ARO7</i> promoter of <i>Saccharomyces cerevisiae</i> . Molecular Genetics and Genomics, 1990, 224, 57-64. | 2.4 | 21 |

| # | ARTICLE | IF | CITATIONS |
|------|--|-----|-----------|
| 1009 | A yeast mutant, PRP20, altered in mRNA metabolism and maintenance of the nuclear structure, is defective in a gene homologous to the human gene RCC1 which is involved in the control of chromosome condensation. <i>Molecular Genetics and Genomics</i> , 1990, 224, 72-80. | 2.4 | 161 |
| 1010 | Molecular analysis of the mitochondrial transcription factor mtf2 of <i>Saccharomyces cerevisiae</i> . <i>Molecular Genetics and Genomics</i> , 1990, 220, 186-190. | 2.4 | 23 |
| 1011 | Ty RNA levels determine the spectrum of retrotransposition events that activate gene expression in <i>Saccharomyces cerevisiae</i> . <i>Molecular Genetics and Genomics</i> , 1990, 220, 213-221. | 2.4 | 98 |
| 1012 | The UGA3 gene regulating the GABA catabolic pathway in <i>Saccharomyces cerevisiae</i> codes for a putative zinc-finger protein acting on RNA amount. <i>Molecular Genetics and Genomics</i> , 1990, 220, 269-276. | 2.4 | 69 |
| 1013 | Yeast telomere length varies in response to changes in the amount of polyC1 " 3A in the cell. <i>Molecular Genetics and Genomics</i> , 1990, 221, 280-282. | 2.4 | 4 |
| 1014 | TUF factor binds to the upstream region of the pyruvate decarboxylase structural gene (PDC1) of <i>Saccharomyces cerevisiae</i> . <i>Molecular Genetics and Genomics</i> , 1990, 223, 449-456. | 2.4 | 39 |
| 1015 | Deletion analysis of the ARG4 promoter of <i>Saccharomyces cerevisiae</i> : A poly(dAdT) stretch involved in gene transcription. <i>Molecular Genetics and Genomics</i> , 1990, 223, 474-480. | 2.4 | 11 |
| 1016 | The yeast mitochondrial leucyl-tRNA synthetase is a splicing factor for the excision of several group I introns. <i>Molecular Genetics and Genomics</i> , 1990, 224, 209-221. | 2.4 | 71 |
| 1017 | Conjugative transfer and autonomous replication of a promiscuous IncQ plasmid in the cyanobacterium <i>Synechocystis</i> PCC 6803. <i>Molecular Genetics and Genomics</i> , 1990, 221, 129-133. | 2.4 | 47 |
| 1018 | Differential regulation of STA genes of <i>Saccharomyces cerevisiae</i> . <i>Molecular Genetics and Genomics</i> , 1990, 222, 87-96. | 2.4 | 8 |
| 1019 | Molecular cloning of the DAC2/FUS3 gene essential for pheromone-induced G1-arrest of the cell cycle in <i>Saccharomyces cerevisiae</i> . <i>Current Genetics</i> , 1990, 18, 395-400. | 0.8 | 20 |
| 1020 | An effective transformation method for <i>Hansenula polymorpha</i> . <i>Current Genetics</i> , 1990, 18, 169-170. | 0.8 | 39 |
| 1021 | Application of the ?-glucuronidase gene fusion system to <i>Saccharomyces cerevisiae</i> . <i>Current Genetics</i> , 1990, 17, 261-264. | 0.8 | 45 |
| 1022 | Biolistic nuclear transformation of <i>Saccharomyces cerevisiae</i> and other fungi. <i>Current Genetics</i> , 1990, 17, 97-103. | 0.8 | 170 |
| 1023 | Evolutionary conservation of transcriptional machinery between yeast and plants as shown by the efficient expression from the CaMV 35S promoter and 35S terminator. <i>Current Genetics</i> , 1990, 17, 473-479. | 0.8 | 34 |
| 1024 | Heterologous expression of human 5-aminolevulinatase in <i>Saccharomyces cerevisiae</i> . <i>Current Genetics</i> , 1990, 17, 1-6. | 0.8 | 4 |
| 1025 | Correct splicing of modified introns of a <i>Rhizopus</i> proteinase gene in <i>Saccharomyces cerevisiae</i> . <i>Molecular Genetics and Genomics</i> , 1990, 223, 11-16. | 2.4 | 6 |
| 1026 | Expression in <i>Saccharomyces cerevisiae</i> of a gene associated with cytoplasmic male sterility from maize: Respiratory dysfunction and uncoupling of yeast mitochondria. <i>Molecular Genetics and Genomics</i> , 1990, 223, 24-32. | 2.4 | 55 |

| # | ARTICLE | IF | CITATIONS |
|------|--|-----|-----------|
| 1027 | Restoration of the yeast LEU2 gene by transcriptionally controlled recombination between tandem repeats. <i>Molecular Genetics and Genomics</i> , 1990, 223, 107-113. | 2.4 | 6 |
| 1028 | Functional analysis of the yeast genome: use of two-dimensional gel electrophoresis to detect genes in randomly cloned DNA sequences. <i>Current Genetics</i> , 1990, 18, 281-286. | 0.8 | 6 |
| 1029 | Functional analysis of the sporulation-specific SPR6 gene of <i>Saccharomyces cerevisiae</i> . <i>Current Genetics</i> , 1990, 18, 293-301. | 0.8 | 15 |
| 1030 | G418-resistance as a dominant marker and reporter for gene expression in <i>Saccharomyces cerevisiae</i> . <i>Current Genetics</i> , 1990, 18, 303-313. | 0.8 | 82 |
| 1031 | A recessive mutant allele of the HNM1 gene of <i>Saccharomyces cerevisiae</i> is responsible for hyper-resistance to nitrogen mustard. <i>Current Genetics</i> , 1990, 18, 187-192. | 0.8 | 4 |
| 1032 | The recombinational hot spot mutation ade6-M26 of <i>Schizosaccharomyces pombe</i> stimulates recombination at sites in a nearby interval. <i>Current Genetics</i> , 1990, 18, 193-197. | 0.8 | 10 |
| 1033 | The structural gene coding for thiamin-repressible acid phosphatase in <i>Schizosaccharomyces pombe</i> . <i>Current Genetics</i> , 1990, 18, 269-272. | 0.8 | 20 |
| 1034 | Fate of highly expressed proteins destined to peroxisomes in <i>Saccharomyces cerevisiae</i> . <i>Current Genetics</i> , 1990, 18, 23-27. | 0.8 | 26 |
| 1035 | Use of the Tn903 neomycin-resistance gene for promoter analysis in the fission yeast <i>Schizosaccharomyces pombe</i> . <i>Current Genetics</i> , 1990, 18, 511-516. | 0.8 | 12 |
| 1036 | Centromeric DNA of <i>Kluyveromyces lactis</i> . <i>Current Genetics</i> , 1990, 18, 517-522. | 0.8 | 66 |
| 1037 | Transformation of commercial baker's yeast strains by electroporation. <i>Biotechnology Letters</i> , 1990, 4, 285-290. | 0.5 | 10 |
| 1038 | Integration of heterologous genes into the chromosome of <i>Saccharomyces cerevisiae</i> using a delta sequence of yeast retrotransposon Ty. <i>Applied Microbiology and Biotechnology</i> , 1990, 33, 302-306. | 1.7 | 94 |
| 1039 | Regulated overproduction and secretion of yeast carboxypeptidase Y. <i>Applied Microbiology and Biotechnology</i> , 1990, 33, 307-312. | 1.7 | 42 |
| 1040 | Fusion of luxA and luxB and its expression in <i>E. coli</i> , <i>S. cerevisiae</i> and <i>D. melanogaster</i> . <i>Luminescence</i> , 1990, 5, 89-97. | 1.3 | 22 |
| 1041 | Yeast myosin heavy chain mutant: Maintenance of the cell type specific budding pattern and the normal deposition of chitin and cell wall components requires an intact myosin heavy chain gene. <i>Cytoskeleton</i> , 1990, 17, 301-308. | 4.4 | 88 |
| 1042 | DNA sequences from <i>Saccharomycopsis fibuligera</i> capable of autonomous replication in <i>Saccharomyces cerevisiae</i> . <i>Journal of Basic Microbiology</i> , 1990, 30, 301-304. | 1.8 | 0 |
| 1043 | The plant G box promoter sequence activates transcription in <i>Saccharomyces cerevisiae</i> and is bound in vitro by a yeast activity similar to GBF, the plant G box binding factor.. <i>EMBO Journal</i> , 1990, 9, 1727-1735. | 3.5 | 46 |
| 1044 | Primary structure, genomic organization and heterologous expression of a glucose transporter from <i>Arabidopsis thaliana</i> .. <i>EMBO Journal</i> , 1990, 9, 3045-3050. | 3.5 | 209 |

| # | ARTICLE | IF | CITATIONS |
|------|--|-----|-----------|
| 1045 | <i>Drosophila cdc2</i> homologs: a functional homolog is coexpressed with a cognate variant.. EMBO Journal, 1990, 9, 3573-3581. | 3.5 | 168 |
| 1046 | <i>Saccharomyces cerevisiae</i> protein phosphatase 2A performs an essential cellular function and is encoded by two genes.. EMBO Journal, 1990, 9, 4339-4346. | 3.5 | 169 |
| 1047 | Peroxisomal protein import is conserved between yeast, plants, insects and mammals.. EMBO Journal, 1990, 9, 85-90. | 3.5 | 272 |
| 1048 | Intermediates of recombination during mating type switching in <i>Saccharomyces cerevisiae</i> .. EMBO Journal, 1990, 9, 663-673. | 3.5 | 373 |
| 1049 | Catabolism of bis(5'-nucleosidyl) tetraphosphates in <i>Saccharomyces cerevisiae</i> . Journal of Bacteriology, 1990, 172, 6892-6899. | 1.0 | 45 |
| 1050 | Characterization of the <i>Bacillus stearothermophilus</i> manganese superoxide dismutase gene and its ability to complement copper/zinc superoxide dismutase deficiency in <i>Saccharomyces cerevisiae</i> . Journal of Bacteriology, 1990, 172, 1539-1546. | 1.0 | 39 |
| 1051 | The CHL 1 (CTF 1) gene product of <i>Saccharomyces cerevisiae</i> is important for chromosome transmission and normal cell cycle progression in G2/M.. EMBO Journal, 1990, 9, 4347-4358. | 3.5 | 135 |
| 1052 | A yeast tRNA precursor containing a pre-mRNA intron is spliced via the pre-mRNA splicing mechanism.. EMBO Journal, 1990, 9, 705-709. | 3.5 | 8 |
| 1053 | Accumulation of pre-tRNA splicing $\hat{\epsilon}^{-2}/3\hat{\epsilon}^{\text{TM}}$ intermediates in a <i>Saccharomyces cerevisiae</i> mutant.. EMBO Journal, 1990, 9, 1245-1252. | 3.5 | 47 |
| 1054 | The <i>rhp6+</i> gene of <i>Schizosaccharomyces pombe</i> : a structural and functional homolog of the RAD6 gene from the distantly related yeast <i>Saccharomyces cerevisiae</i> .. EMBO Journal, 1990, 9, 1423-1430. | 3.5 | 86 |
| 1055 | Effect of transcription of yeast chromatin on DNA topology in vivo.. EMBO Journal, 1990, 9, 1873-1881. | 3.5 | 49 |
| 1056 | Effects of multiple yeast <i>rad3</i> mutant alleles on UV sensitivity, mutability, and mitotic recombination. Journal of Bacteriology, 1990, 172, 6620-6630. | 1.0 | 42 |
| 1057 | Development of an integrative DNA transformation system for the yeast <i>Candida tropicalis</i> . Journal of Bacteriology, 1990, 172, 4571-4577. | 1.0 | 62 |
| 1058 | Three GCN4 responsive elements act synergistically as upstream and as TATA-like elements in the yeast TRP4 promoter.. EMBO Journal, 1990, 9, 2951-2957. | 3.5 | 20 |
| 1059 | Specificity of the mutator effect caused by disruption of the RAD1 excision repair gene of <i>Saccharomyces cerevisiae</i> . Journal of Bacteriology, 1990, 172, 3009-3014. | 1.0 | 52 |
| 1060 | Role of trans-activating proteins in the generation of active chromatin at the PHO5 promoter in <i>S. cerevisiae</i> .. EMBO Journal, 1990, 9, 2523-2528. | 3.5 | 132 |
| 1061 | An essential <i>Saccharomyces cerevisiae</i> single-stranded DNA binding protein is homologous to the large subunit of human RP-A.. EMBO Journal, 1990, 9, 2321-2329. | 3.5 | 170 |
| 1062 | The recognition component of the N-end rule pathway.. EMBO Journal, 1990, 9, 3179-3189. | 3.5 | 352 |

| # | ARTICLE | IF | CITATIONS |
|------|---|-----|-----------|
| 1063 | The adenylate cyclase/protein kinase cascade regulates entry into meiosis in <i>Saccharomyces cerevisiae</i> through the gene <i>IME1</i> .. EMBO Journal, 1990, 9, 3225-3232. | 3.5 | 61 |
| 1064 | Signal sequence for generation of mRNA 3' end in the <i>Saccharomyces cerevisiae</i> <i>GAL7</i> gene.. EMBO Journal, 1990, 9, 3691-3697. | 3.5 | 61 |
| 1065 | In vitro reconstitution of <i>cdc25</i> regulated <i>S. cerevisiae</i> adenylyl cyclase and its kinetic properties.. EMBO Journal, 1990, 9, 641-651. | 3.5 | 49 |
| 1066 | Structural and functional analysis of <i>ypt2</i> , an essential ras-related gene in the fission yeast <i>Schizosaccharomyces pombe</i> encoding a <i>Sec4</i> protein homologue.. EMBO Journal, 1990, 9, 1957-1962. | 3.5 | 72 |
| 1067 | Cloning and characterization of <i>Kluyveromyces lactis</i> <i>SEC14</i> , a gene whose product stimulates Golgi secretory function in <i>Saccharomyces cerevisiae</i> . Journal of Bacteriology, 1990, 172, 4510-4521. | 1.0 | 81 |
| 1068 | Regulation of a yeast <i>HSP70</i> gene by a cAMP responsive transcriptional control element.. EMBO Journal, 1990, 9, 2543-2553. | 3.5 | 110 |
| 1069 | Oncogenic transformation by <i>vrel</i> requires an amino-terminal activation domain.. Molecular and Cellular Biology, 1990, 10, 2840-2847. | 1.1 | 155 |
| 1070 | Structure and molecular analysis of <i>RGR1</i> , a gene required for glucose repression of <i>Saccharomyces cerevisiae</i> .. Molecular and Cellular Biology, 1990, 10, 4130-4138. | 1.1 | 94 |
| 1071 | Curing Industrial <i>Saccharomyces</i> Yeasts of Parasitic 2 $\frac{1}{4}$ m Plasmid. Journal of the American Society of Brewing Chemists, 1990, 48, 107-110. | 0.8 | 1 |
| 1072 | Subthreshold Vicinal Diketone Levels in Lager Brewing Yeast Fermentations by Means of <i>LV5</i> Gene Amplification. Journal of the American Society of Brewing Chemists, 1990, 48, 111-114. | 0.8 | 42 |
| 1073 | Isolation and characterization of a new mutant of <i>Saccharomyces cerevisiae</i> with altered synthesis of 5-aminolevulinic acid. Journal of Bacteriology, 1990, 172, 2855-2861. | 1.0 | 5 |
| 1074 | ADP ribosylation factor is an essential protein in <i>Saccharomyces cerevisiae</i> and is encoded by two genes.. Molecular and Cellular Biology, 1990, 10, 6690-6699. | 1.1 | 265 |
| 1075 | Increased dosage of the <i>MSN1</i> gene restores invertase expression in yeast mutants defective in the <i>SNF1</i> protein kinase. Nucleic Acids Research, 1990, 18, 6959-6964. | 6.5 | 75 |
| 1076 | Two systems of glucose repression of the <i>GAL1</i> promoter in <i>Saccharomyces cerevisiae</i> .. Molecular and Cellular Biology, 1990, 10, 4757-4769. | 1.1 | 195 |
| 1077 | Depletion of U14 small nuclear RNA (<i>snR128</i>) disrupts production of 18S rRNA in <i>Saccharomyces cerevisiae</i> .. Molecular and Cellular Biology, 1990, 10, 1145-1152. | 1.1 | 276 |
| 1078 | Role of arginine 180 and glutamic acid 177 of ricin toxin A chain in enzymatic inactivation of ribosomes.. Molecular and Cellular Biology, 1990, 10, 6257-6263. | 1.1 | 98 |
| 1079 | Yeast clathrin has a distinctive light chain that is important for cell growth.. Journal of Cell Biology, 1990, 111, 1437-1449. | 2.3 | 77 |
| 1080 | Biological activity of the mammalian <i>RAP</i> genes in yeast.. Molecular Biology of the Cell, 1990, 1, 763-769. | 6.5 | 25 |

| # | ARTICLE | IF | CITATIONS |
|------|---|-----|-----------|
| 1081 | Human cDNAs encoding homologs of the small p34Cdc28/Cdc2-associated protein of <i>Saccharomyces cerevisiae</i> and <i>Schizosaccharomyces pombe</i> .. <i>Genes and Development</i> , 1990, 4, 1332-1344. | 2.7 | 168 |
| 1082 | STE11 is a protein kinase required for cell-type-specific transcription and signal transduction in yeast.. <i>Genes and Development</i> , 1990, 4, 1862-1874. | 2.7 | 191 |
| 1083 | Expression of the <i>E.coli ada</i> gene in yeast protects against the toxic and mutagenic effects of N-methyl-N- ϵ -nitro-N-nitrosoguanidine. <i>Nucleic Acids Research</i> , 1990, 18, 331-335. | 6.5 | 17 |
| 1084 | Targeting of a cytosolic protein to the nuclear periphery.. <i>Journal of Cell Biology</i> , 1990, 111, 2829-2837. | 2.3 | 33 |
| 1085 | Sec2 protein contains a coiled-coil domain essential for vesicular transport and a dispensable carboxy terminal domain.. <i>Journal of Cell Biology</i> , 1990, 110, 1897-1909. | 2.3 | 88 |
| 1086 | BIK1, a protein required for microtubule function during mating and mitosis in <i>Saccharomyces cerevisiae</i> , colocalizes with tubulin.. <i>Journal of Cell Biology</i> , 1990, 111, 2573-2586. | 2.3 | 169 |
| 1087 | Yeast KRE genes provide evidence for a pathway of cell wall beta-glucan assembly.. <i>Journal of Cell Biology</i> , 1990, 110, 1833-1843. | 2.3 | 246 |
| 1088 | Use of <i>Saccharomyces cerevisiae</i> expressing beta-galactosidase to screen for antimycotic agents directed against yeast cell wall biosynthesis and possible application to pathogenic fungi. <i>Antimicrobial Agents and Chemotherapy</i> , 1990, 34, 660-662. | 1.4 | 8 |
| 1089 | Molecular analysis of the yeast VPS3 gene and the role of its product in vacuolar protein sorting and vacuolar segregation during the cell cycle.. <i>Journal of Cell Biology</i> , 1990, 111, 877-892. | 2.3 | 112 |
| 1090 | Domains of yeast U4 spliceosomal RNA required for PRP4 protein binding, snRNP-snRNP interactions, and pre-mRNA splicing in vivo.. <i>Genes and Development</i> , 1990, 4, 1185-1196. | 2.7 | 93 |
| 1091 | Subcellular localization of yeast CDC46 varies with the cell cycle.. <i>Genes and Development</i> , 1990, 4, 2252-2263. | 2.7 | 219 |
| 1092 | Site-specific cleavage of a yeast chromosome by oligonucleotide-directed triple-helix formation. <i>Science</i> , 1990, 249, 73-75. | 6.0 | 210 |
| 1093 | Loss of BiP/GRP78 function blocks translocation of secretory proteins in yeast.. <i>Journal of Cell Biology</i> , 1990, 110, 1885-1895. | 2.3 | 449 |
| 1094 | A yeast protein that influences the chromatin structure of UASC and functions as a powerful auxiliary gene activator.. <i>Genes and Development</i> , 1990, 4, 503-514. | 2.7 | 227 |
| 1095 | A yeast H2A-H2B promoter can be regulated by changes in histone gene copy number.. <i>Genes and Development</i> , 1990, 4, 752-763. | 2.7 | 62 |
| 1096 | The DNA unwinding element in a yeast replication origin functions independently of easily unwound sequences present elsewhere on a plasmid. <i>Nucleic Acids Research</i> , 1990, 18, 6601-6605. | 6.5 | 25 |
| 1097 | PRP5: a helicase-like protein required for mRNA splicing in yeast.. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 1990, 87, 4236-4240. | 3.3 | 130 |
| 1098 | Interchangeable RNA polymerase I and II enhancers.. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 1990, 87, 8202-8206. | 3.3 | 19 |

| # | ARTICLE | IF | CITATIONS |
|------|---|-----|-----------|
| 1099 | A DNA replication enhancer in <i>Saccharomyces cerevisiae</i> .. Proceedings of the National Academy of Sciences of the United States of America, 1990, 87, 4665-4669. | 3.3 | 103 |
| 1100 | The SPA2 gene of <i>Saccharomyces cerevisiae</i> is important for pheromone-induced morphogenesis and efficient mating.. Journal of Cell Biology, 1990, 111, 1451-1464. | 2.3 | 151 |
| 1101 | Transcription of a yeast U6 snRNA gene requires a polymerase III promoter element in a novel position.. Genes and Development, 1990, 4, 1345-1356. | 2.7 | 154 |
| 1102 | High-frequency transformation method and library transducing vectors for cloning mammalian cDNAs by trans-complementation of <i>Schizosaccharomyces pombe</i> . Nucleic Acids Research, 1990, 18, 6485-6489. | 6.5 | 471 |
| 1103 | GAL4 protein: purification, association with GAL80 protein, and conserved domain structure.. Molecular and Cellular Biology, 1990, 10, 2916-2923. | 1.1 | 89 |
| 1104 | cAMP-dependent phosphorylation and inactivation of yeast transcription factor ADR1 does not affect DNA binding.. Proceedings of the National Academy of Sciences of the United States of America, 1990, 87, 4098-4102. | 3.3 | 77 |
| 1105 | Yeast structural gene (APN1) for the major apurinic endonuclease: homology to <i>Escherichia coli</i> endonuclease IV.. Proceedings of the National Academy of Sciences of the United States of America, 1990, 87, 4193-4197. | 3.3 | 204 |
| 1106 | Leucine-rich repeats and carboxyl terminus are required for interaction of yeast adenylate cyclase with RAS proteins.. Proceedings of the National Academy of Sciences of the United States of America, 1990, 87, 8711-8715. | 3.3 | 132 |
| 1107 | Generation of deletion derivatives by targeted transformation of human-derived yeast artificial chromosomes.. Proceedings of the National Academy of Sciences of the United States of America, 1990, 87, 1300-1304. | 3.3 | 93 |
| 1108 | The yeast RNA1 gene product necessary for RNA processing is located in the cytosol and apparently excluded from the nucleus.. Journal of Cell Biology, 1990, 111, 309-321. | 2.3 | 217 |
| 1109 | DNA repair in a small yeast plasmid folded into chromatin. Nucleic Acids Research, 1990, 18, 2045-2051. | 6.5 | 35 |
| 1110 | Host genes that influence transposition in yeast: the abundance of a rare tRNA regulates Ty1 transposition frequency.. Proceedings of the National Academy of Sciences of the United States of America, 1990, 87, 8360-8364. | 3.3 | 102 |
| 1111 | Characterization of VPS34, a gene required for vacuolar protein sorting and vacuole segregation in <i>Saccharomyces cerevisiae</i> .. Molecular and Cellular Biology, 1990, 10, 6742-6754. | 1.1 | 450 |
| 1112 | Characterization of TPII gene expression in isogenic wild-type and <i>andgcr1</i> -deletion mutant strains of <i>Saccharomyces cerevisiae</i> . Nucleic Acids Research, 1990, 18, 7099-7107. | 6.5 | 50 |
| 1113 | Association of catalytic and regulatory subunits of cyclic AMP-dependent protein kinase requires a negatively charged side group at a conserved threonine.. Molecular and Cellular Biology, 1990, 10, 1066-1075. | 1.1 | 64 |
| 1114 | Selectivity changes in site-directed mutants of the VDAC ion channel: structural implications. Science, 1990, 247, 1233-1236. | 6.0 | 274 |
| 1115 | Autonomous replication sequences in an extrachromosomal element of a pathogenic <i>Entamoeba histolytica</i> . Nucleic Acids Research, 1990, 18, 5515-5519. | 6.5 | 19 |
| 1116 | RAD10, an excision repair gene of <i>Saccharomyces cerevisiae</i> , is involved in the RAD1 pathway of mitotic recombination.. Molecular and Cellular Biology, 1990, 10, 2485-2491. | 1.1 | 162 |

| # | ARTICLE | IF | CITATIONS |
|------|--|------|-----------|
| 1117 | Site-directed, recombination-mediated mutagenesis of a complex gene locus. <i>Nucleic Acids Research</i> , 1990, 18, 7349-7355. | 6.5 | 28 |
| 1118 | Characterization of TUP1, a mediator of glucose repression in <i>Saccharomyces cerevisiae</i> .. <i>Molecular and Cellular Biology</i> , 1990, 10, 6500-6511. | 1.1 | 235 |
| 1119 | Cloning and characterization of nuclear genes for two mitochondrial ribosomal proteins in <i>Saccharomyces cerevisiae</i> . <i>Nucleic Acids Research</i> , 1990, 18, 1521-1529. | 6.5 | 47 |
| 1120 | Yeast tRNA ^{Trp} genes with anticodons corresponding to UAA and UGA nonsense codons. <i>Nucleic Acids Research</i> , 1990, 18, 4215-4221. | 6.5 | 11 |
| 1121 | Transcript levels of the <i>Saccharomyces cerevisiae</i> DNA repair gene RAD23 increase in response to UV light and in meiosis but remain constant in the mitotic cell cycle. <i>Nucleic Acids Research</i> , 1990, 18, 4737-4742. | 6.5 | 65 |
| 1122 | Expression of yeast 5S RNA is independent of the rDNA enhancer region. <i>Nucleic Acids Research</i> , 1990, 18, 4179-4184. | 6.5 | 14 |
| 1123 | Molecular cloning of the yeast mitochondrial aconitase gene (ACO1) and evidence of a synergistic regulation of expression by glucose plus glutamate.. <i>Molecular and Cellular Biology</i> , 1990, 10, 3551-3561. | 1.1 | 129 |
| 1124 | [23] Manipulating yeast genome using plasmid vectors. <i>Methods in Enzymology</i> , 1990, 185, 280-297. | 0.4 | 74 |
| 1125 | Overproduction of the yeast STE12 protein leads to constitutive transcriptional induction.. <i>Genes and Development</i> , 1990, 4, 492-502. | 2.7 | 130 |
| 1126 | Protein splicing converts the yeast TFP1 gene product to the 69-kD subunit of the vacuolar H(+)-adenosine triphosphatase. <i>Science</i> , 1990, 250, 651-657. | 6.0 | 483 |
| 1127 | BET1, BOS1, and SEC22 are members of a group of interacting yeast genes required for transport from the endoplasmic reticulum to the Golgi complex.. <i>Molecular and Cellular Biology</i> , 1990, 10, 3405-3414. | 1.1 | 190 |
| 1128 | Functional expression of the <i>Chlorella</i> hexose transporter in <i>Schizosaccharomyces pombe</i> .. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 1990, 87, 7949-7952. | 3.3 | 73 |
| 1129 | Mouse Zfx protein is similar to Zfy-2: each contains an acidic activating domain and 13 zinc fingers.. <i>Molecular and Cellular Biology</i> , 1990, 10, 681-688. | 1.1 | 87 |
| 1130 | <i>Saccharomyces cerevisiae</i> homoserine kinase is homologous to prokaryotic homoserine kinases. <i>Gene</i> , 1990, 96, 177-180. | 1.0 | 10 |
| 1131 | Expression of cDNAs encoding barley α -amylase 1 and 2 in yeast and characterization of the secreted proteins. <i>Gene</i> , 1990, 94, 173-179. | 1.0 | 81 |
| 1132 | FUS3 encodes a cdc2+/CDC28-related kinase required for the transition from mitosis into conjugation. <i>Cell</i> , 1990, 60, 649-664. | 13.5 | 481 |
| 1133 | A genetic approach to the study of farnesylation. <i>Methods</i> , 1990, 1, 246-252. | 1.9 | 0 |
| 1134 | Expression and characterization of infectious bursal disease virus polyprotein in yeast. <i>Gene</i> , 1990, 95, 179-186. | 1.0 | 14 |

| # | ARTICLE | IF | CITATIONS |
|------|---|------|-----------|
| 1135 | An improved method for yeast 2 $\frac{1}{4}$ m plasmid curing. <i>Gene</i> , 1990, 88, 241-245. | 1.0 | 8 |
| 1136 | Organelle transformation: Shoot first, ask questions later. <i>Trends in Biochemical Sciences</i> , 1990, 15, 465-468. | 3.7 | 47 |
| 1137 | Nucleotide sequence and characterization of temperature-sensitive pol1 mutants of <i>Saccharomyces cerevisiae</i> . <i>Gene</i> , 1990, 90, 99-104. | 1.0 | 54 |
| 1138 | Yeast centromere binding protein CBF1, of the helix-loop-helix protein family, is required for chromosome stability and methionine prototrophy. <i>Cell</i> , 1990, 61, 437-446. | 13.5 | 360 |
| 1139 | Synthesis of human coagulation factor XIII in yeast. <i>Gene</i> , 1990, 86, 279-283. | 1.0 | 14 |
| 1140 | Synthesis of <i>Drosophila melanogaster</i> alcohol dehydrogenase in yeast. <i>Gene</i> , 1990, 93, 205-212. | 1.0 | 7 |
| 1141 | Functional expression of the <i>Candida albicans</i> β -tubulin gene in <i>Saccharomyces cerevisiae</i> . <i>Gene</i> , 1990, 90, 115-123. | 1.0 | 16 |
| 1142 | A yeast system for stable expression of hepatitis B surface antigen. <i>Journal of Biotechnology</i> , 1990, 14, 411-422. | 1.9 | 5 |
| 1143 | In vivo translocation of the cell wall acid phosphatase across the yeast endoplasmic reticulum membrane: are there multiple signals for the targeting process?. <i>Biochimie</i> , 1990, 72, 103-114. | 1.3 | 6 |
| 1144 | KAR3, a kinesin-related gene required for yeast nuclear fusion. <i>Cell</i> , 1990, 60, 1029-1041. | 13.5 | 520 |
| 1145 | Analysis of wild-type and rad50 mutants of yeast suggests an intimate relationship between meiotic chromosome synapsis and recombination. <i>Cell</i> , 1990, 61, 419-436. | 13.5 | 625 |
| 1146 | YAC cloning: Options and problems. <i>Gene Analysis Techniques</i> , 1990, 7, 94-99. | 1.1 | 11 |
| 1147 | Activity and thermal stability of genetically truncated forms of <i>Aspergillus glucoamylase</i> . <i>Gene</i> , 1990, 91, 131-134. | 1.0 | 53 |
| 1148 | Cloning of the yeast SFL2 gene: its disruption results in pleiotropic phenotypes characteristic for tup1 mutants. <i>Gene</i> , 1990, 89, 93-99. | 1.0 | 61 |
| 1149 | Hydroxylation Reactions by Recombinant Yeast Cells Expressing P450 Monooxygenases. <i>Annals of the New York Academy of Sciences</i> , 1990, 613, 37-43. | 1.8 | 5 |
| 1150 | Genetics and genetic engineering of the industrial yeast <i>Yarrowia lipolytica</i> . , 1990, , 43-73. | | 14 |
| 1151 | <i>Escherichia coli</i> and <i>Saccharomyces cerevisiae</i> acetylornithine aminotransferases: evolutionary relationship with ornithine aminotransferases. <i>Gene</i> , 1990, 90, 69-78. | 1.0 | 63 |
| 1152 | The NUP1 gene encodes an essential component of the yeast nuclear pore complex. <i>Cell</i> , 1990, 61, 965-978. | 13.5 | 230 |

| # | ARTICLE | IF | CITATIONS |
|------|--|------|-----------|
| 1153 | A putative GTP binding protein homologous to interferon-inducible Mx proteins performs an essential function in yeast protein sorting. <i>Cell</i> , 1990, 61, 1063-1074. | 13.5 | 287 |
| 1154 | A pathway for generation and processing of double-strand breaks during meiotic recombination in <i>S. cerevisiae</i> . <i>Cell</i> , 1990, 61, 1089-1101. | 13.5 | 774 |
| 1155 | The yeast heat shock transcription factor contains a transcriptional activation domain whose activity is repressed under nonshock conditions. <i>Cell</i> , 1990, 62, 807-817. | 13.5 | 176 |
| 1156 | Portable encapsidation signal of the L-A double-stranded RNA virus of <i>S. cerevisiae</i> . <i>Cell</i> , 1990, 62, 819-828. | 13.5 | 137 |
| 1157 | Position effect at <i>S. cerevisiae</i> telomeres: Reversible repression of Pol II transcription. <i>Cell</i> , 1990, 63, 751-762. | 13.5 | 1,369 |
| 1158 | The NF1 locus encodes a protein functionally related to mammalian GAP and yeast IRA proteins. <i>Cell</i> , 1990, 63, 851-859. | 13.5 | 808 |
| 1159 | Distinct, essential roles of type 1 and 2A protein phosphatases in the control of the fission yeast cell division cycle. <i>Cell</i> , 1990, 63, 405-415. | 13.5 | 357 |
| 1160 | The fission yeast <i>cut1+</i> gene regulates spindle pole body duplication and has homology to the budding yeast <i>ESP1</i> gene. <i>Cell</i> , 1990, 62, 913-925. | 13.5 | 157 |
| 1161 | Meiotic gene conversion and crossing over: Their relationship to each other and to chromosome synapsis and segregation. <i>Cell</i> , 1990, 62, 927-937. | 13.5 | 209 |
| 1162 | A candidate protein kinase C gene, <i>PKC1</i> , is required for the <i>S. cerevisiae</i> cell cycle. <i>Cell</i> , 1990, 62, 213-224. | 13.5 | 443 |
| 1163 | Ribosomal frameshifting in the yeast retrotransposon <i>Ty</i> : tRNAs induce slippage on a 7 nucleotide minimal site. <i>Cell</i> , 1990, 62, 339-352. | 13.5 | 311 |
| 1164 | A third essential DNA polymerase in <i>S. cerevisiae</i> . <i>Cell</i> , 1990, 62, 1143-1151. | 13.5 | 365 |
| 1165 | <i>GAL11P</i> : A yeast mutation that potentiates the effect of weak <i>GAL4</i> -derived activators. <i>Cell</i> , 1990, 63, 1299-1309. | 13.5 | 169 |
| 1166 | In vivo degradation of a transcriptional regulator: The yeast $\hat{I}\pm 2$ repressor. <i>Cell</i> , 1990, 61, 697-708. | 13.5 | 251 |
| 1167 | Identification of a gene necessary for cell cycle arrest by a negative growth factor of yeast: <i>FAR1</i> is an inhibitor of a G1 cyclin, <i>CLN2</i> . <i>Cell</i> , 1990, 63, 999-1011. | 13.5 | 405 |
| 1168 | Courtship in <i>S. cerevisiae</i> : Both cell types choose mating partners by responding to the strongest pheromone signal. <i>Cell</i> , 1990, 63, 1039-1051. | 13.5 | 207 |
| 1169 | A novel mediator between activator proteins and the RNA polymerase II transcription apparatus. <i>Cell</i> , 1990, 61, 1209-1215. | 13.5 | 442 |
| 1170 | <i>SRV2</i> , a gene required for RAS activation of adenylate cyclase in yeast. <i>Cell</i> , 1990, 61, 329-340. | 13.5 | 257 |

| # | ARTICLE | IF | CITATIONS |
|------|--|------|-----------|
| 1171 | The conserved terminal region of <i>Trichoderma reesei</i> cellulases forms a strong antigenic epitope for polyclonal antibodies. <i>Biochimica Et Biophysica Acta Gene Regulatory Mechanisms</i> , 1990, 1087, 137-141. | 2.4 | 11 |
| 1172 | Characterisation of a repetitive DNA family from <i>Entamoeba histolytica</i> containing <i>Saccharomyces cerevisiae</i> ARS consensus sequences. <i>Gene</i> , 1990, 96, 197-203. | 1.0 | 20 |
| 1173 | Expression of Torpedo nicotinic acetylcholine receptor subunits in yeast is enhanced by use of yeast signal sequences. <i>Gene</i> , 1990, 86, 145-152. | 1.0 | 13 |
| 1174 | Disruption of the gene XRN1, coding for a 5'â€²â†'3'â€² exoribonuclease, restricts yeast cell growth. <i>Gene</i> , 1990, 95, 85-90. | 1.0 | 142 |
| 1175 | Expression of Alcohol-Inducible Rabbit Liver Cytochrome P-450 3a (P-450IIE1) in <i>Saccharomyces cerevisiae</i> with the Copper-Inducible CUP1 Promoter. <i>DNA and Cell Biology</i> , 1990, 9, 111-118. | 0.9 | 26 |
| 1176 | Transfer and expression of heterologous genes in yeasts other than <i>Saccharomyces cerevisiae</i> . , 1990, 43, 75-102. | | 11 |
| 1177 | [34] Î±-Factor leader-directed secretion of heterologous proteins from yeast. <i>Methods in Enzymology</i> , 1990, 185, 408-421. | 0.4 | 53 |
| 1178 | Expression and secretion of rice Î±-amylase by <i>Saccharomyces cerevisiae</i> . <i>Gene</i> , 1990, 94, 209-216. | 1.0 | 57 |
| 1179 | A repeating amino acid motif in CDC23 defines a family of proteins and a new relationship among genes required for mitosis and RNA synthesis. <i>Cell</i> , 1990, 60, 307-317. | 13.5 | 512 |
| 1180 | Snap helix with knob and hole: Essential repeats in <i>S. pombe</i> nuclear protein nuc2+. <i>Cell</i> , 1990, 60, 319-328. | 13.5 | 304 |
| 1181 | Molecular cloning of chromosome I DNA from <i>Saccharomyces cerevisiae</i> : isolation, characterization and regulation of the SPO7 sporulation gene. <i>Gene</i> , 1990, 95, 65-72. | 1.0 | 14 |
| 1182 | An inducible expression vector for both fission and budding yeast. <i>Gene</i> , 1990, 86, 257-261. | 1.0 | 57 |
| 1183 | Expression of human chromosomal proteins HMG-14 and HMG-17 in <i>Saccharomyces cerevisiae</i> . <i>Experimental Cell Research</i> , 1990, 191, 71-75. | 1.2 | 1 |
| 1184 | Isolation and characterization of humanregprotein produced in <i>Saccharomyces cerevisiae</i> . <i>FEBS Letters</i> , 1990, 272, 85-88. | 1.3 | 20 |
| 1185 | Inhibition of oxygen toxicity by targeting Superoxide dismutase to endothelial cell surface. <i>FEBS Letters</i> , 1990, 269, 89-92. | 1.3 | 52 |
| 1186 | Expression of the <i>Drosophila melanogaster</i> metallothionein genes in yeast. <i>FEBS Letters</i> , 1990, 269, 273-276. | 1.3 | 14 |
| 1187 | Expression and pharmacological characterization of the human M1 muscarinic receptor in <i>Saccharomyces cerevisiae</i> . <i>FEBS Letters</i> , 1990, 266, 21-25. | 1.3 | 51 |
| 1188 | Expression of the yeast LEU4 gene is subject to four different modes of control. <i>Archives of Biochemistry and Biophysics</i> , 1990, 276, 294-298. | 1.4 | 18 |

| # | ARTICLE | IF | CITATIONS |
|------|---|-----|-----------|
| 1189 | Expression of human lysosomal β -hexosaminidase in yeast vacuoles. <i>Biochemical and Biophysical Research Communications</i> , 1990, 170, 383-390. | 1.0 | 6 |
| 1190 | Expression and secretion of goat β -lactalbumin as an active protein in <i>Saccharomyces cerevisiae</i> . <i>Biochemical and Biophysical Research Communications</i> , 1990, 173, 741-747. | 1.0 | 5 |
| 1191 | Structure and transcriptional control of the <i>Saccharomyces cerevisiae</i> POX1 gene encoding acylcoenzyme A oxidase. <i>Gene</i> , 1990, 88, 247-252. | 1.0 | 99 |
| 1192 | Xylitol Production by Recombinant <i>Saccharomyces Cerevisiae</i> . <i>Bio/technology</i> , 1991, 9, 1090-1095. | 1.9 | 190 |
| 1193 | Enhanced Secretion of Human Nerve Growth Factor from <i>Saccharomyces cerevisiae</i> using an Advanced λ -Integration System. <i>Bio/technology</i> , 1991, 9, 1382-1385. | 1.9 | 34 |
| 1194 | Primary structure and regulation of a glucoamylase-encoding gene (STA2) in <i>Saccharomyces diastaticus</i> . <i>Gene</i> , 1991, 100, 95-103. | 1.0 | 41 |
| 1195 | Co-expression of a <i>Saccharomyces diastaticus</i> glucoamylase-encoding gene and a <i>Bacillus amyloliquefaciens</i> α -amylase-encoding gene in <i>Saccharomyces cerevisiae</i> . <i>Gene</i> , 1991, 100, 85-93. | 1.0 | 59 |
| 1196 | Glucose-responsive and oleic acid-responsive elements in the gene encoding the peroxisomal trifunctional enzyme of <i>Candida tropicalis</i> . <i>Gene</i> , 1991, 105, 129-134. | 1.0 | 27 |
| 1197 | Mammalian cAMP-dependent protein kinase functionally replaces its homolog in yeast. <i>Gene</i> , 1991, 99, 171-179. | 1.0 | 14 |
| 1198 | Fission yeast promoter-probe vectors based on hygromycin resistance. <i>Gene</i> , 1991, 100, 241-245. | 1.0 | 26 |
| 1199 | Production of hepatitis B virion-like particles in yeast. <i>Gene</i> , 1991, 106, 143-149. | 1.0 | 15 |
| 1200 | A split zinc-finger protein is required for normal yeast growth. <i>Gene</i> , 1991, 107, 101-110. | 1.0 | 34 |
| 1201 | Expression of the <i>Escherichia coli</i> recA gene in the yeast <i>Saccharomyces cerevisiae</i> . <i>Biochimie</i> , 1991, 73, 285-288. | 1.3 | 5 |
| 1202 | Cloning and transcriptional analysis of the ADE6 gene of <i>Saccharomyces cerevisiae</i> . <i>Gene</i> , 1991, 107, 149-154. | 1.0 | 20 |
| 1203 | New shuttle vectors for direct cloning in <i>Saccharomyces cerevisiae</i> . <i>Gene</i> , 1991, 104, 99-102. | 1.0 | 8 |
| 1204 | High level expression of the major core protein VP7 and the non-structural protein NS3 of bluetongue virus in yeast: use of expressed VP7 as a diagnostic, group-reactive antigen in a blocking ELISA. <i>Virus Research</i> , 1991, 18, 165-178. | 1.1 | 32 |
| 1205 | Synthesis and expression of genes encoding tuna, pigeon, and horse cytochromes c in the yeast <i>Saccharomyces cerevisiae</i> . <i>Gene</i> , 1991, 105, 73-81. | 1.0 | 42 |
| 1206 | The cell-cycle-regulated budding yeast gene DBF2, encoding a putative protein kinase, has a homologue that is not under cell-cycle control. <i>Gene</i> , 1991, 104, 63-70. | 1.0 | 75 |

| # | ARTICLE | IF | CITATIONS |
|------|--|-----|-----------|
| 1207 | Expression of the <i>Bacillus subtilis</i> levanase gene in <i>Escherichia coli</i> and <i>Saccharomyces cerevisiae</i> . <i>Journal of Biotechnology</i> , 1991, 18, 243-254. | 1.9 | 12 |
| 1208 | Molecular cloning of the <i>Candida maltosa</i> ADE1 gene. <i>Gene</i> , 1991, 107, 161-164. | 1.0 | 16 |
| 1209 | High level expression of biologically active estrogen receptor in <i>Saccharomyces cerevisiae</i> . <i>Journal of Steroid Biochemistry and Molecular Biology</i> , 1991, 39, 291-297. | 1.2 | 129 |
| 1210 | The pleiotropic UGA35(DURL) regulatory gene of <i>Saccharomyces cerevisiae</i> : cloning, sequence and identity with the DAL81 gene. <i>Gene</i> , 1991, 97, 163-171. | 1.0 | 60 |
| 1211 | Structural and functional conservation between the high-affinity K ⁺ transporters of <i>Saccharomyces uvarum</i> and <i>Saccharomyces cerevisiae</i> . <i>Gene</i> , 1991, 99, 39-46. | 1.0 | 20 |
| 1212 | Protein disulfide isomerase is essential for viability in <i>Saccharomyces cerevisiae</i> . <i>Gene</i> , 1991, 108, 81-89. | 1.0 | 123 |
| 1213 | Cloning, disruption and sequence of the gene encoding yeast C-5 sterol desaturase. <i>Gene</i> , 1991, 102, 39-44. | 1.0 | 157 |
| 1214 | Identification of the DNA binding site for NGFI-B by genetic selection in yeast. <i>Science</i> , 1991, 252, 1296-1300. | 6.0 | 554 |
| 1215 | Breeding of Flocculent Industrial <i>Saccharomyces cerevisiae</i> Strains by Introducing the Flocculation Gene FLO1. <i>Agricultural and Biological Chemistry</i> , 1991, 55, 1547-1552. | 0.3 | 0 |
| 1216 | Identification of nuclear genes which participate to mitochondrial translation in <i>Saccharomyces cerevisiae</i> . <i>Biochimie</i> , 1991, 73, 1525-1532. | 1.3 | 13 |
| 1217 | The highly conserved N-terminal domains of histones H3 and H4 are required for normal cell cycle progression.. <i>Molecular and Cellular Biology</i> , 1991, 11, 4111-4120. | 1.1 | 93 |
| 1218 | Disruption and mutagenesis of the <i>Saccharomyces cerevisiae</i> PDX1 gene encoding the protein X component of the pyruvate dehydrogenase complex. <i>Biochemistry</i> , 1991, 30, 2834-2839. | 1.2 | 79 |
| 1219 | Expression of spinach glycolate oxidase in <i>Saccharomyces cerevisiae</i> : purification and characterization. <i>Biochemistry</i> , 1991, 30, 4612-4619. | 1.2 | 82 |
| 1220 | Intramitochondrial functions regulate nonmitochondrial citrate synthase (CIT2) expression in <i>Saccharomyces cerevisiae</i> .. <i>Molecular and Cellular Biology</i> , 1991, 11, 38-46. | 1.1 | 221 |
| 1221 | Crystallization and preliminary x-ray analysis of the cAMP-dependent protein kinase catalytic subunit from <i>Saccharomyces cerevisiae</i> . <i>Biochemistry</i> , 1991, 30, 10595-10600. | 1.2 | 8 |
| 1222 | Studies on yeast peroxisomal citrate synthase. <i>Archives of Biochemistry and Biophysics</i> , 1991, 286, 132-137. | 1.4 | 18 |
| 1223 | Genetic analysis of yeast Iso-1-cytochrome c structural requirements: Suppression of Gly6 replacements by an Asn52 → Ile replacement. <i>Archives of Biochemistry and Biophysics</i> , 1991, 288, 261-269. | 1.4 | 18 |
| 1224 | Galactose-dependent expression of the recombinant Ca ²⁺ -binding photoprotein aequorin in yeast. <i>Biochemical and Biophysical Research Communications</i> , 1991, 174, 115-122. | 1.0 | 15 |

| # | ARTICLE | IF | CITATIONS |
|------|---|------|-----------|
| 1225 | Proteinase B is, indeed, not required for chitin synthetase 1 function in <i>Saccharomyces cerevisiae</i> . <i>Biochemical and Biophysical Research Communications</i> , 1991, 174, 204-210. | 1.0 | 11 |
| 1226 | The phosphotransferase activity of casein kinase II is required for its physiological function in vivo. <i>Biochemical and Biophysical Research Communications</i> , 1991, 181, 524-528. | 1.0 | 19 |
| 1227 | Molecular cloning and sequencing of genomic DNA encoding yeast vacuolar carboxypeptidase yscS. <i>FEBS Letters</i> , 1991, 283, 27-32. | 1.3 | 18 |
| 1228 | The proteinase yscA-inhibitor, IA3, gene Studies of cytoplasmic proteinase inhibitor deficiency on yeast physiology. <i>FEBS Letters</i> , 1991, 283, 78-84. | 1.3 | 17 |
| 1229 | Cloning and sequencing of the inulinase gene of <i>Kluyveromyces marxianus</i> var. <i>marxianus</i> ATCC 12424. <i>FEBS Letters</i> , 1991, 289, 64-68. | 1.3 | 89 |
| 1230 | Structural and functional analysis of <i>Trichoderma reesei</i> endoglucanase I expressed in yeast <i>Saccharomyces cerevisiae</i> . <i>FEBS Letters</i> , 1991, 291, 45-49. | 1.3 | 9 |
| 1231 | Regions in the promoter of the yeast FBP1 gene implicated in catabolite repression may bind the product of the regulatory gene MIG1. <i>FEBS Letters</i> , 1991, 291, 97-100. | 1.3 | 48 |
| 1232 | Biogenesis of the yeast vacuole (lysosome) Active site mutation in the vacuolar aspartate proteinase yscA blocks maturation of vacuolar proteinases. <i>FEBS Letters</i> , 1991, 293, 62-66. | 1.3 | 20 |
| 1233 | The promoter of <i>Saccharomyces cerevisiae</i> FBA1 gene contains a single positive upstream regulatory element. <i>FEBS Letters</i> , 1991, 293, 97-100. | 1.3 | 18 |
| 1234 | Stable ester conjugate between the <i>Saccharomyces cerevisiae</i> RAD6 protein and ubiquitin has no biological activity. <i>Journal of Molecular Biology</i> , 1991, 221, 745-749. | 2.0 | 57 |
| 1235 | Identification and functional analysis of the nuclear localization signals of ribosomal protein L25 from <i>Saccharomyces cerevisiae</i> . <i>Journal of Molecular Biology</i> , 1991, 221, 225-237. | 2.0 | 66 |
| 1236 | Mutational specificity of DNA precursor pool imbalances in yeast arising from deoxycytidylate deaminase deficiency or treatment with thymidylate. <i>Journal of Molecular Biology</i> , 1991, 220, 933-946. | 2.0 | 82 |
| 1237 | Enhanced survival of yeast expressing an antifreeze gene analogue after freezing. <i>Cryobiology</i> , 1991, 28, 474-482. | 0.3 | 29 |
| 1238 | The expression of human cytochrome P450IA1 in the yeast <i>Saccharomyces cerevisiae</i> . <i>Biochemical Pharmacology</i> , 1991, 42, 753-758. | 2.0 | 25 |
| 1239 | Transformation of <i>Schizophyllum commune</i> : An analysis of specific properties. <i>Experimental Mycology</i> , 1991, 15, 326-335. | 1.8 | 14 |
| 1240 | Centromere-dependent binding of yeast minichromosomes to microtubules in vitro. <i>Cell</i> , 1991, 66, 483-495. | 13.5 | 54 |
| 1241 | Yeast BUD5, encoding a putative GDP-GTP exchange factor, is necessary for bud site selection and interacts with bud formation gene BEM1. <i>Cell</i> , 1991, 65, 1213-1224. | 13.5 | 280 |
| 1242 | Functional cloning of BUD5, a CDC25-related gene from <i>S. cerevisiae</i> that can suppress a dominant-negative RAS2 mutant. <i>Cell</i> , 1991, 65, 1225-1231. | 13.5 | 114 |

| # | ARTICLE | IF | CITATIONS |
|------|--|------|-----------|
| 1243 | An evolutionarily conserved cyclin homolog from <i>Drosophila</i> rescues yeast deficient in G1 cyclins. <i>Cell</i> , 1991, 66, 1207-1216. | 13.5 | 174 |
| 1244 | Meiosis-specific RNA splicing in yeast. <i>Cell</i> , 1991, 66, 1257-1268. | 13.5 | 212 |
| 1245 | Modifiers of position effect are shared between telomeric and silent mating-type loci in <i>S. cerevisiae</i> . <i>Cell</i> , 1991, 66, 1279-1287. | 13.5 | 733 |
| 1246 | A highly conserved domain of TFIIID displays species specificity in vivo. <i>Cell</i> , 1991, 65, 333-340. | 13.5 | 106 |
| 1247 | <i>S. cerevisiae</i> $\hat{\pm}$ pheromone receptors activate a novel signal transduction pathway for mating partner discrimination. <i>Cell</i> , 1991, 67, 389-402. | 13.5 | 137 |
| 1248 | <i>S. pombe</i> gene <i>sds22+</i> essential for a midmitotic transition encodes a leucine-rich repeat protein that positively modulates protein phosphatase-1. <i>Cell</i> , 1991, 64, 149-157. | 13.5 | 206 |
| 1249 | <i>mik1</i> and <i>wee1</i> cooperate in the inhibitory tyrosine phosphorylation of <i>cdc2</i> . <i>Cell</i> , 1991, 64, 1111-1122. | 13.5 | 715 |
| 1250 | <i>SGV1</i> encodes a <i>CDC28/cdc2</i> -related kinase required for a $\hat{\pm}$ subunit-mediated adaptive response to pheromone in <i>S. cerevisiae</i> . <i>Cell</i> , 1991, 65, 785-795. | 13.5 | 88 |
| 1251 | Structural basis for the regulation of splicing of a yeast messenger RNA. <i>Cell</i> , 1991, 65, 797-804. | 13.5 | 168 |
| 1252 | Mutations in yeast U5 snRNA alter the specificity of 5' splice-site cleavage. <i>Cell</i> , 1991, 65, 115-123. | 13.5 | 214 |
| 1253 | Transcriptional activation of <i>CLN1</i> , <i>CLN2</i> , and a putative new G1 cyclin (<i>HCS26</i>) by <i>SWI4</i> , a positive regulator of G1-specific transcription. <i>Cell</i> , 1991, 66, 1015-1026. | 13.5 | 342 |
| 1254 | Pheromone-induced phosphorylation of a G protein $\hat{2}$ subunit in <i>S. cerevisiae</i> is associated with an adaptive response to mating pheromone. <i>Cell</i> , 1991, 64, 703-716. | 13.5 | 111 |
| 1255 | Mutations in the CDP-choline pathway for phospholipid biosynthesis bypass the requirement for an essential phospholipid transfer protein. <i>Cell</i> , 1991, 64, 789-800. | 13.5 | 363 |
| 1256 | The signal recognition particle in <i>S. cerevisiae</i> . <i>Cell</i> , 1991, 67, 131-144. | 13.5 | 317 |
| 1257 | A novel protein kinase homolog essential for protein sorting to the yeast lysosome-like vacuole. <i>Cell</i> , 1991, 64, 425-437. | 13.5 | 206 |
| 1258 | The nuclear coded mitoribosomal proteins <i>YmL27</i> and <i>YmL31</i> are both essential for mitochondrial function in yeast. <i>Biochimie</i> , 1991, 73, 837-844. | 1.3 | 26 |
| 1259 | The overexpression of the 3' terminal region of the <i>CDC25</i> gene of <i>Saccharomyces cerevisiae</i> causes growth inhibition and alteration of purine nucleotides pools. <i>Biochimica Et Biophysica Acta Gene Regulatory Mechanisms</i> , 1991, 1089, 206-212. | 2.4 | 10 |
| 1260 | Posttranscriptional regulation of the expression of <i>MET2</i> gene of <i>Saccharomyces cerevisiae</i> . <i>Biochimica Et Biophysica Acta Gene Regulatory Mechanisms</i> , 1991, 1089, 47-53. | 2.4 | 9 |

| # | ARTICLE | IF | CITATIONS |
|------|--|-----|-----------|
| 1261 | Production of functional IgM Fab fragments by. <i>Journal of Biotechnology</i> , 1991, 20, 291-300. | 1.9 | 12 |
| 1262 | [12] High-efficiency transformation of yeast by electroporation. <i>Methods in Enzymology</i> , 1991, 194, 182-187. | 0.4 | 818 |
| 1263 | [20] In vitro mutagenesis and plasmid shuffling: From cloned gene to mutant yeast. <i>Methods in Enzymology</i> , 1991, 194, 302-318. | 0.4 | 585 |
| 1264 | [24] Transformation of yeast directly with synthetic oligonucleotides. <i>Methods in Enzymology</i> , 1991, 194, 362-369. | 0.4 | 37 |
| 1265 | Sequences of two tandem genes regulated by carbon sources, one being essential for n-alkane assimilation in <i>Candida maltosa</i> . <i>Gene</i> , 1991, 106, 61-69. | 1.0 | 18 |
| 1266 | Recombinant DNA Techniques. , 1991, , 149-167. | | 1 |
| 1267 | [19] Targeting, disruption, replacement, and allele rescue: Integrative DNA transformation in yeast. <i>Methods in Enzymology</i> , 1991, 194, 281-301. | 0.4 | 1,319 |
| 1268 | Extragenic suppressors of a translation initiation defect in the <i>cyc1</i> gene of <i>Saccharomyces cerevisiae</i> . <i>Biochimie</i> , 1991, 73, 1445-1455. | 1.3 | 32 |
| 1269 | Identification of two cell-cycle-controlling <i>cdc2</i> gene homologs in <i>Arabidopsis thaliana</i> . <i>Gene</i> , 1991, 105, 159-165. | 1.0 | 160 |
| 1270 | Differential effect of cysteine-to-serine substitutions in metallothionein on cadmium resistance.. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 1991, 88, 3024-3028. | 3.3 | 43 |
| 1271 | DMSO-enhanced whole cell yeast transformation. <i>Nucleic Acids Research</i> , 1991, 19, 5791-5791. | 6.5 | 517 |
| 1272 | New vectors in fission yeast: application for cloning the <i>his2</i> gene. <i>Gene</i> , 1991, 99, 47-54. | 1.0 | 56 |
| 1273 | Expression in <i>Escherichia coli</i> of the <i>Saccharomyces cerevisiae</i> CCT gene encoding cholinephosphate cytidyltransferase. <i>Journal of Bacteriology</i> , 1991, 173, 2134-2136. | 1.0 | 27 |
| 1274 | Site-directed mutagenesis of the yeast actin gene: a test for actin function in vivo.. <i>EMBO Journal</i> , 1991, 10, 3951-3958. | 3.5 | 47 |
| 1275 | The 3' to 5' exonuclease activity located in the DNA polymerase delta subunit of <i>Saccharomyces cerevisiae</i> is required for accurate replication.. <i>EMBO Journal</i> , 1991, 10, 2165-2170. | 3.5 | 206 |
| 1276 | A novel switch-activating site (SAS1) and its cognate binding factor (SAP1) required for efficient <i>mat1</i> switching in <i>Schizosaccharomyces pombe</i> .. <i>EMBO Journal</i> , 1991, 10, 3025-3032. | 3.5 | 98 |
| 1277 | Mutational analysis of the putative effector domain of the GTP-binding Ypt1 protein in yeast suggests specific regulation by a novel GAP activity.. <i>EMBO Journal</i> , 1991, 10, 785-792. | 3.5 | 83 |
| 1278 | Double-strand-break repair and recombination catalyzed by a nuclear extract of <i>Saccharomyces cerevisiae</i> .. <i>EMBO Journal</i> , 1991, 10, 987-996. | 3.5 | 31 |

| # | ARTICLE | IF | CITATIONS |
|------|---|-----|-----------|
| 1279 | Molecular cloning of the mitochondrial aldehyde dehydrogenase gene of <i>Saccharomyces cerevisiae</i> by genetic complementation. <i>Journal of Bacteriology</i> , 1991, 173, 3199-3208. | 1.0 | 40 |
| 1280 | Transformation system for <i>Amycolatopsis (Nocardia) mediterranei</i> : direct transformation of mycelium with plasmid DNA. <i>Journal of Bacteriology</i> , 1991, 173, 6325-6331. | 1.0 | 43 |
| 1281 | Multiple Ca ²⁺ /calmodulin-dependent protein kinase genes in a unicellular eukaryote.. <i>EMBO Journal</i> , 1991, 10, 1511-1522. | 3.5 | 115 |
| 1282 | Proteinase yscE, the yeast proteasome/multicatalytic-multifunctional proteinase: mutants unravel its function in stress induced proteolysis and uncover its necessity for cell survival.. <i>EMBO Journal</i> , 1991, 10, 555-562. | 3.5 | 383 |
| 1283 | A mutation in the yeast heat-shock factor gene causes temperature-sensitive defects in both mitochondrial protein import and the cell cycle.. <i>Molecular and Cellular Biology</i> , 1991, 11, 2647-2655. | 1.1 | 94 |
| 1284 | Saturation mutagenesis of the UASNTR (GATAA) responsible for nitrogen catabolite repression-sensitive transcriptional activation of the allantoin pathway genes in <i>Saccharomyces cerevisiae</i> . <i>Journal of Bacteriology</i> , 1991, 173, 4977-4982. | 1.0 | 76 |
| 1285 | A genetic and structural analysis of the yeast Vps15 protein kinase: evidence for a direct role of Vps15p in vacuolar protein delivery.. <i>EMBO Journal</i> , 1991, 10, 4049-4060. | 3.5 | 97 |
| 1286 | ABC1, a novel yeast nuclear gene has a dual function in mitochondria: it suppresses a cytochrome b mRNA translation defect and is essential for the electron transfer in the bc 1 complex.. <i>EMBO Journal</i> , 1991, 10, 2023-2031. | 3.5 | 100 |
| 1287 | Identification of regulatory residues of the yeast adenylyl cyclase.. <i>EMBO Journal</i> , 1991, 10, 349-359. | 3.5 | 20 |
| 1288 | Distinct cis-acting signals enhance 3' endpoint formation of CYC1 mRNA in the yeast <i>Saccharomyces cerevisiae</i> .. <i>EMBO Journal</i> , 1991, 10, 563-571. | 3.5 | 121 |
| 1289 | Transcription-induced nucleosome "splitting": an underlying structure for DNase I sensitive chromatin.. <i>EMBO Journal</i> , 1991, 10, 607-615. | 3.5 | 130 |
| 1290 | Genetic and molecular analyses of the SUP201 gene: a tRNA(3Arg) nonsense suppressor of yeast <i>cyr1-2</i> . <i>Journal of Bacteriology</i> , 1991, 173, 2406-2408. | 1.0 | 5 |
| 1291 | Exceptional codon recognition by the glutamine tRNAs in <i>Saccharomyces cerevisiae</i> .. <i>EMBO Journal</i> , 1991, 10, 1481-1491. | 3.5 | 29 |
| 1292 | A specific DNA sequence is required for high frequency of recombination in the <i>ade6</i> gene of fission yeast.. <i>EMBO Journal</i> , 1991, 10, 2157-2163. | 3.5 | 100 |
| 1293 | Cloning of a yeast U1 snRNP 70K protein homologue: functional conservation of an RNA-binding domain between humans and yeast.. <i>EMBO Journal</i> , 1991, 10, 2627-2634. | 3.5 | 62 |
| 1294 | UBA 1: an essential yeast gene encoding ubiquitin-activating enzyme.. <i>EMBO Journal</i> , 1991, 10, 227-236. | 3.5 | 231 |
| 1295 | The small nucleolar RNP protein NOP1 (fibrillarin) is required for pre-rRNA processing in yeast.. <i>EMBO Journal</i> , 1991, 10, 573-583. | 3.5 | 315 |
| 1296 | Negative regulation of transcription of the <i>Saccharomyces cerevisiae</i> catalase T (CTT1) gene by cAMP is mediated by a positive control element.. <i>EMBO Journal</i> , 1991, 10, 585-592. | 3.5 | 100 |

| # | ARTICLE | IF | CITATIONS |
|------|---|-----|-----------|
| 1297 | DAL82, a second gene required for induction of allantoin system gene transcription in <i>Saccharomyces cerevisiae</i> . <i>Journal of Bacteriology</i> , 1991, 173, 255-261. | 1.0 | 33 |
| 1298 | A dominant interfering mutation (CYR3) of the <i>Saccharomyces cerevisiae</i> RAS2 gene. <i>Journal of Bacteriology</i> , 1991, 173, 4533-4536. | 1.0 | 6 |
| 1299 | Identification of the structural gene for glucose-6-phosphate dehydrogenase in yeast. Inactivation leads to a nutritional requirement for organic sulfur.. <i>EMBO Journal</i> , 1991, 10, 547-553. | 3.5 | 118 |
| 1300 | Isolation from <i>Candida albicans</i> of a functional homolog of the <i>Saccharomyces cerevisiae</i> KRE1 gene, which is involved in cell wall beta-glucan synthesis. <i>Journal of Bacteriology</i> , 1991, 173, 6859-6864. | 1.0 | 59 |
| 1301 | [18] Characterization of human P450IIC isozymes by using yeast expression system. <i>Methods in Enzymology</i> , 1991, 206, 183-190. | 0.4 | 5 |
| 1302 | Null mutants of <i>Saccharomyces cerevisiae</i> Cu,Zn superoxide dismutase: characterization and spontaneous mutation rates. <i>Journal of Bacteriology</i> , 1991, 173, 5918-5920. | 1.0 | 194 |
| 1303 | Cloning and analysis of the nuclear gene for YmL33, a protein of the large subunit of the mitochondrial ribosome in <i>Saccharomyces cerevisiae</i> . <i>Journal of Bacteriology</i> , 1991, 173, 4013-4020. | 1.0 | 24 |
| 1304 | Suppression of ribosomal reinitiation at upstream open reading frames in amino acid-starved cells forms the basis for GCN4 translational control.. <i>Molecular and Cellular Biology</i> , 1991, 11, 486-496. | 1.1 | 251 |
| 1305 | Mutations in a conserved region of RNA polymerase II influence the accuracy of mRNA start site selection.. <i>Molecular and Cellular Biology</i> , 1991, 11, 5781-5791. | 1.1 | 58 |
| 1306 | RPD3 encodes a second factor required to achieve maximum positive and negative transcriptional states in <i>Saccharomyces cerevisiae</i> .. <i>Molecular and Cellular Biology</i> , 1991, 11, 6317-6327. | 1.1 | 300 |
| 1307 | The role of RAP1 in the regulation of the MAT alpha locus.. <i>Molecular and Cellular Biology</i> , 1991, 11, 1069-1079. | 1.1 | 70 |
| 1308 | Genetic evidence that different functional domains of the PET54 gene product facilitate expression of the mitochondrial genes COX1 and COX3 in <i>Saccharomyces cerevisiae</i> .. <i>Molecular and Cellular Biology</i> , 1991, 11, 2399-2405. | 1.1 | 32 |
| 1309 | Molecular and genetic analysis of the gene encoding the <i>Saccharomyces cerevisiae</i> strand exchange protein Sep1.. <i>Molecular and Cellular Biology</i> , 1991, 11, 2593-2608. | 1.1 | 116 |
| 1310 | The HXT1 gene product of <i>Saccharomyces cerevisiae</i> is a new member of the family of hexose transporters.. <i>Molecular and Cellular Biology</i> , 1991, 11, 3804-3813. | 1.1 | 111 |
| 1311 | A -1 ribosomal frameshift in a double-stranded RNA virus of yeast forms a gag-pol fusion protein.. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 1991, 88, 174-178. | 3.3 | 354 |
| 1312 | Complementation of a yeast cell cycle mutant by an alfalfa cDNA encoding a protein kinase homologous to p34cdc2.. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 1991, 88, 1636-1640. | 3.3 | 202 |
| 1313 | CDC55, a <i>Saccharomyces cerevisiae</i> gene involved in cellular morphogenesis: identification, characterization, and homology to the B subunit of mammalian type 2A protein phosphatase.. <i>Molecular and Cellular Biology</i> , 1991, 11, 5767-5780. | 1.1 | 292 |
| 1314 | The fission yeast <i>dis3+</i> gene encodes a 110-kDa essential protein implicated in mitotic control.. <i>Molecular and Cellular Biology</i> , 1991, 11, 5839-5847. | 1.1 | 82 |

| # | ARTICLE | IF | CITATIONS |
|------|---|-----|-----------|
| 1315 | Expression of the DAL80 gene, whose product is homologous to the GATA factors and is a negative regulator of multiple nitrogen catabolic genes in <i>Saccharomyces cerevisiae</i> , is sensitive to nitrogen catabolite repression.. <i>Molecular and Cellular Biology</i> , 1991, 11, 6205-6215. | 1.1 | 160 |
| 1316 | p68 RNA helicase: identification of a nucleolar form and cloning of related genes containing a conserved intron in yeasts.. <i>Molecular and Cellular Biology</i> , 1991, 11, 1326-1333. | 1.1 | 129 |
| 1317 | Identification of DNA regions required for mitotic and meiotic functions within the centromere of <i>Schizosaccharomyces pombe</i> chromosome I.. <i>Molecular and Cellular Biology</i> , 1991, 11, 2206-2215. | 1.1 | 81 |
| 1318 | Integrative selection of human chromosome-specific yeast artificial chromosomes.. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 1991, 88, 7788-7791. | 3.3 | 10 |
| 1319 | Yeast beta-glucan synthesis: KRE6 encodes a predicted type II membrane protein required for glucan synthesis in vivo and for glucan synthase activity in vitro.. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 1991, 88, 11295-11299. | 3.3 | 143 |
| 1320 | Altered response to growth rate changes in <i>Kluyveromyces lactis</i> versus <i>Saccharomyces cerevisiae</i> demonstrated by heterologous expression of ribosomal protein 59 (CRY1). <i>Nucleic Acids Research</i> , 1991, 19, 4701-4707. | 6.5 | 20 |
| 1321 | The yeast putative transcriptional repressor RGM1 is a proline-rich zinc finger protein. <i>Nucleic Acids Research</i> , 1991, 19, 4873-4877. | 6.5 | 27 |
| 1322 | Yeast Flocculation Mediated by Overexpression of the Chitin Synthase 2 Gene in <i>Saccharomyces cerevisiae</i> .. <i>Agricultural and Biological Chemistry</i> , 1991, 55, 2901-2903. | 0.3 | 8 |
| 1323 | The glucoamylase cDNA from <i>Aspergillus oryzae</i> : Its cloning, nucleotide sequence, and expression in <i>Saccharomyces cerevisiae</i> .. <i>Agricultural and Biological Chemistry</i> , 1991, 55, 941-949. | 0.3 | 36 |
| 1324 | Genetic Engineering of a Sake Yeast Producing No Urea. <i>Journal of the Brewing Society of Japan</i> , 1991, 86, 99-107. | 0.1 | 1 |
| 1325 | Production of a site specifically cleavable P-glycoprotein-BETA-galactosidase fusion protein.. <i>Agricultural and Biological Chemistry</i> , 1991, 55, 1075-1080. | 0.3 | 7 |
| 1326 | The yeast rad18 mutator specifically increases G.C→T.A transversions without reducing correction of G-A or C-T mismatches to G.C pairs.. <i>Molecular and Cellular Biology</i> , 1991, 11, 218-225. | 1.1 | 48 |
| 1327 | Heteroduplex formation and mismatch repair of the "stuck" mutation during mating-type switching in <i>Saccharomyces cerevisiae</i> .. <i>Molecular and Cellular Biology</i> , 1991, 11, 5372-5380. | 1.1 | 91 |
| 1328 | Sequence and expression of GLN3, a positive nitrogen regulatory gene of <i>Saccharomyces cerevisiae</i> encoding a protein with a putative zinc finger DNA-binding domain.. <i>Molecular and Cellular Biology</i> , 1991, 11, 6216-6228. | 1.1 | 213 |
| 1329 | Role of the complex upstream region of the GDH2 gene in nitrogen regulation of the NAD-linked glutamate dehydrogenase in <i>Saccharomyces cerevisiae</i> .. <i>Molecular and Cellular Biology</i> , 1991, 11, 6229-6247. | 1.1 | 63 |
| 1330 | Specific cis-acting sequence for PHO8 expression interacts with PHO4 protein, a positive regulatory factor, in <i>Saccharomyces cerevisiae</i> .. <i>Molecular and Cellular Biology</i> , 1991, 11, 785-794. | 1.1 | 31 |
| 1331 | Yeast CBP1 mRNA 3' end formation is regulated during the induction of mitochondrial function.. <i>Molecular and Cellular Biology</i> , 1991, 11, 813-821. | 1.1 | 30 |
| 1332 | byr2, a <i>Schizosaccharomyces pombe</i> gene encoding a protein kinase capable of partial suppression of the ras1 mutant phenotype.. <i>Molecular and Cellular Biology</i> , 1991, 11, 3554-3563. | 1.1 | 207 |

| # | ARTICLE | IF | CITATIONS |
|------|---|-----|-----------|
| 1333 | Assessment of the transcriptional activation potential of the HMG chromosomal proteins.. Molecular and Cellular Biology, 1991, 11, 4483-4489. | 1.1 | 52 |
| 1334 | Differential induction of 3-deoxy-D-arabino-heptulosonate 7-phosphate synthase genes in <i>Arabidopsis thaliana</i> by wounding and pathogenic attack.. Proceedings of the National Academy of Sciences of the United States of America, 1991, 88, 8821-8825. | 3.3 | 159 |
| 1335 | Mutational analysis of the functional domains of yeast K1 killer toxin.. Molecular and Cellular Biology, 1991, 11, 175-181. | 1.1 | 41 |
| 1336 | Isolation, nucleotide sequence analysis, and disruption of the MDH2 gene from <i>Saccharomyces cerevisiae</i> : evidence for three isozymes of yeast malate dehydrogenase.. Molecular and Cellular Biology, 1991, 11, 370-380. | 1.1 | 110 |
| 1337 | In vivo analysis of the <i>Saccharomyces cerevisiae</i> centromere CDEIII sequence: requirements for mitotic chromosome segregation.. Molecular and Cellular Biology, 1991, 11, 5212-5221. | 1.1 | 48 |
| 1338 | A putative zinc finger protein, <i>Saccharomyces cerevisiae</i> Vps18p, affects late Golgi functions required for vacuolar protein sorting and efficient alpha-factor prohormone maturation.. Molecular and Cellular Biology, 1991, 11, 5813-5824. | 1.1 | 89 |
| 1339 | RPD1 (SIN3/UME4) is required for maximal activation and repression of diverse yeast genes.. Molecular and Cellular Biology, 1991, 11, 6306-6316. | 1.1 | 181 |
| 1340 | Membrane insertion of uracil permease, a polytopic yeast plasma membrane protein.. Molecular and Cellular Biology, 1991, 11, 1114-1124. | 1.1 | 105 |
| 1341 | The yeast SLY gene products, suppressors of defects in the essential GTP-binding Ypt1 protein, may act in endoplasmic reticulum-to-Golgi transport.. Molecular and Cellular Biology, 1991, 11, 2980-2993. | 1.1 | 180 |
| 1342 | Translation initiation factor 5A and its hypusine modification are essential for cell viability in the yeast <i>Saccharomyces cerevisiae</i> .. Molecular and Cellular Biology, 1991, 11, 3105-3114. | 1.1 | 353 |
| 1343 | The <i>Saccharomyces cerevisiae</i> SRK1 gene, a suppressor of <i>bcy1</i> and <i>ins1</i> , may be involved in protein phosphatase function.. Molecular and Cellular Biology, 1991, 11, 3369-3373. | 1.1 | 81 |
| 1344 | DPB2, the gene encoding DNA polymerase II subunit B, is required for chromosome replication in <i>Saccharomyces cerevisiae</i> .. Proceedings of the National Academy of Sciences of the United States of America, 1991, 88, 4601-4605. | 3.3 | 103 |
| 1345 | A wild-type DNA ligase I gene is expressed in Bloom's syndrome cells.. Proceedings of the National Academy of Sciences of the United States of America, 1991, 88, 7615-7619. | 3.3 | 45 |
| 1346 | In vitro assembly of apophytochrome and apophytochrome deletion mutants expressed in yeast with phycocyanobilin.. Proceedings of the National Academy of Sciences of the United States of America, 1991, 88, 10392-10396. | 3.3 | 59 |
| 1347 | Mutants of <i>Escherichia coli</i> initiator tRNA that suppress amber codons in <i>Saccharomyces cerevisiae</i> and are aminoacylated with tyrosine by yeast extracts.. Proceedings of the National Academy of Sciences of the United States of America, 1991, 88, 11378-11382. | 3.3 | 50 |
| 1348 | Mutational analysis of yeast vacuolar H(+)-ATPase.. Proceedings of the National Academy of Sciences of the United States of America, 1991, 88, 1938-1942. | 3.3 | 156 |
| 1349 | Chapter 20 Mutations That Affect Nuclear Organization in Yeast. Methods in Cell Biology, 1991, 35, 525-541. | 0.5 | 1 |
| 1350 | Different classes of polyadenylation sites in the yeast <i>Saccharomyces cerevisiae</i> .. Molecular and Cellular Biology, 1991, 11, 3060-3069. | 1.1 | 111 |

| # | ARTICLE | IF | CITATIONS |
|------|--|-----|-----------|
| 1351 | A poly(dA.dT) tract is a component of the recombination initiation site at the ARG4 locus in <i>Saccharomyces cerevisiae</i> .. <i>Molecular and Cellular Biology</i> , 1991, 11, 322-328. | 1.1 | 125 |
| 1352 | Quantitation of alpha-factor internalization and response during the <i>Saccharomyces cerevisiae</i> cell cycle.. <i>Molecular and Cellular Biology</i> , 1991, 11, 5251-5258. | 1.1 | 32 |
| 1353 | Mutations in SPT16/CDC68 suppress cis- and trans-acting mutations that affect promoter function in <i>Saccharomyces cerevisiae</i> .. <i>Molecular and Cellular Biology</i> , 1991, 11, 5710-5717. | 1.1 | 172 |
| 1354 | A functional 125-kDa core polypeptide of fission yeast DNA topoisomerase II.. <i>Molecular and Cellular Biology</i> , 1991, 11, 6093-6102. | 1.1 | 74 |
| 1355 | Association of RAP1 binding sites with stringent control of ribosomal protein gene transcription in <i>Saccharomyces cerevisiae</i> .. <i>Molecular and Cellular Biology</i> , 1991, 11, 2723-2735. | 1.1 | 177 |
| 1356 | GCD2, a translational repressor of the GCN4 gene, has a general function in the initiation of protein synthesis in <i>Saccharomyces cerevisiae</i> .. <i>Molecular and Cellular Biology</i> , 1991, 11, 3203-3216. | 1.1 | 175 |
| 1357 | In situ distinction between steroid receptor binding and transactivation at a target gene.. <i>Molecular and Cellular Biology</i> , 1991, 11, 4350-4355. | 1.1 | 89 |
| 1358 | Physiological and genetic analysis of the carbon regulation of the NAD-dependent glutamate dehydrogenase of <i>Saccharomyces cerevisiae</i> .. <i>Molecular and Cellular Biology</i> , 1991, 11, 4455-4465. | 1.1 | 46 |
| 1359 | GRR1 of <i>Saccharomyces cerevisiae</i> is required for glucose repression and encodes a protein with leucine-rich repeats.. <i>Molecular and Cellular Biology</i> , 1991, 11, 5101-5112. | 1.1 | 177 |
| 1360 | AAR2, a gene for splicing pre-mRNA of the MATa1 cistron in cell type control of <i>Saccharomyces cerevisiae</i> .. <i>Molecular and Cellular Biology</i> , 1991, 11, 5693-5700. | 1.1 | 23 |
| 1361 | The <i>Saccharomyces cerevisiae</i> ADR1 gene is a positive regulator of transcription of genes encoding peroxisomal proteins.. <i>Molecular and Cellular Biology</i> , 1991, 11, 699-704. | 1.1 | 158 |
| 1362 | The yeast homolog to mouse Tcp-1 affects microtubule-mediated processes.. <i>Molecular and Cellular Biology</i> , 1991, 11, 2629-2640. | 1.1 | 172 |
| 1363 | The CDC25 protein of <i>Saccharomyces cerevisiae</i> promotes exchange of guanine nucleotides bound to ras.. <i>Molecular and Cellular Biology</i> , 1991, 11, 2641-2646. | 1.1 | 215 |
| 1364 | SPT5, an essential gene important for normal transcription in <i>Saccharomyces cerevisiae</i> , encodes an acidic nuclear protein with a carboxy-terminal repeat.. <i>Molecular and Cellular Biology</i> , 1991, 11, 3009-3019. | 1.1 | 171 |
| 1365 | Complex formation by positive and negative translational regulators of GCN4.. <i>Molecular and Cellular Biology</i> , 1991, 11, 3217-3228. | 1.1 | 131 |
| 1366 | In vivo characterization of the <i>Saccharomyces cerevisiae</i> centromere DNA element I, a binding site for the helix-loop-helix protein CPF1.. <i>Molecular and Cellular Biology</i> , 1991, 11, 3545-3553. | 1.1 | 49 |
| 1367 | Complex transcriptional regulation of the <i>Saccharomyces cerevisiae</i> CYB2 gene encoding cytochrome b2: CYP1(HAP1) activator binds to the CYB2 upstream activation site UAS1-B2.. <i>Molecular and Cellular Biology</i> , 1991, 11, 3762-3772. | 1.1 | 65 |
| 1368 | Viability of clathrin heavy-chain-deficient <i>Saccharomyces cerevisiae</i> is compromised by mutations at numerous loci: implications for the suppression hypothesis.. <i>Molecular and Cellular Biology</i> , 1991, 11, 3868-3878. | 1.1 | 33 |

| # | ARTICLE | IF | CITATIONS |
|------|---|-----|-----------|
| 1369 | Cloning and characterization of a gene which determines osmotic stability in <i>Saccharomyces cerevisiae</i> .. <i>Molecular and Cellular Biology</i> , 1991, 11, 4235-4243. | 1.1 | 31 |
| 1370 | Upstream induction sequence, the cis-acting element required for response to the allantoin pathway inducer and enhancement of operation of the nitrogen-regulated upstream activation sequence in <i>Saccharomyces cerevisiae</i> . <i>Journal of Bacteriology</i> , 1991, 173, 7186-7195. | 1.0 | 42 |
| 1371 | Transformation system for an asporogenous methylotrophic yeast, <i>Candida boidinii</i> : cloning of the orotidine-5'-phosphate decarboxylase gene (URA3), isolation of uracil auxotrophic mutants, and use of the mutants for integrative transformation. <i>Journal of Bacteriology</i> , 1991, 173, 7458-7463. | 1.0 | 76 |
| 1372 | [51] Functional expression of mammalian adenosine cyclic monophosphate-dependent protein kinase in <i>saccharomyces cerevisiae</i> . <i>Methods in Enzymology</i> , 1991, 200, 605-627. | 0.4 | 3 |
| 1373 | Identification and structure of four yeast genes (SLY) that are able to suppress the functional loss of YPT1, a member of the RAS superfamily.. <i>Molecular and Cellular Biology</i> , 1991, 11, 872-885. | 1.1 | 324 |
| 1374 | CLONING OF A YEAST GENE WHICH CAUSES PHENOLIC OFF-FLAVOURS IN BEER. <i>Journal of the Institute of Brewing</i> , 1991, 97, 353-357. | 0.8 | 32 |
| 1375 | The CDC20 gene product of <i>Saccharomyces cerevisiae</i> , a beta-transducin homolog, is required for a subset of microtubule-dependent cellular processes.. <i>Molecular and Cellular Biology</i> , 1991, 11, 5592-5602. | 1.1 | 105 |
| 1376 | Characteristics of <i>Nicotiana tabacum</i> nitrate reductase protein produced in <i>Saccharomyces cerevisiae</i> . <i>Biochemical Journal</i> , 1991, 278, 393-397. | 1.7 | 26 |
| 1377 | Expression and Stability of Plasmid with Ethionine Rosistance Gene as a Marker in <i>Saccharomyces cerevisiae</i> .. <i>Kagaku Kogaku Ronbunshu</i> , 1991, 17, 674-679. | 0.1 | 0 |
| 1378 | Protection against cadmium toxicity in yeast by alcohol dehydrogenase. <i>Journal of Inorganic Biochemistry</i> , 1991, 44, 155-161. | 1.5 | 15 |
| 1379 | Purification and crystallization of yeast hexokinase isoenzymes. <i>Journal of Chromatography A</i> , 1991, 587, 85-92. | 1.8 | 15 |
| 1380 | A mutated ARO4 gene for feedback-resistant DAHP synthase which causes both o-fluoro-dl-phenylalanine resistance and ?-phenethyl-alcohol overproduction in <i>Saccharomyces cerevisiae</i> . <i>Current Genetics</i> , 1991, 20, 453-456. | 0.8 | 31 |
| 1381 | Isolation of xylose reductase gene of <i>Pichia stipitis</i> and its expression in <i>Saccharomyces cerevisiae</i> . <i>Applied Biochemistry and Biotechnology</i> , 1991, 28-29, 327-340. | 1.4 | 67 |
| 1382 | Repair of exogenous (plasmid) DNA damaged by photoaddition of 8-methoxypsoralen in the yeast <i>Saccharomyces cerevisiae</i> . <i>Mutation Research - Fundamental and Molecular Mechanisms of Mutagenesis</i> , 1991, 251, 123-131. | 0.4 | 15 |
| 1383 | The genetic activity of N6-hydroxyadenine and 2-amino-N6-hydroxyadenine in <i>Escherichia coli</i> , <i>Salmonella typhimurium</i> and <i>Saccharomyces cerevisiae</i> . <i>Mutation Research - Environmental Mutagenesis and Related Subjects Including Methodology</i> , 1991, 253, 33-46. | 0.4 | 49 |
| 1384 | Optimization of the expression of a foreign gene from galactose-inducible promoter in a recombinant yeast. <i>Journal of Bioscience and Bioengineering</i> , 1991, 72, 187-192. | 0.9 | 6 |
| 1385 | Nucleotide sequence and characterization of a Gene conferring resistance to ethionine in yeast <i>Saccharomyces cerevisiae</i> . <i>Journal of Bioscience and Bioengineering</i> , 1991, 71, 211-215. | 0.9 | 43 |
| 1386 | Autonomously replicating fragments of mitochondrial DNA from <i>Penicillium urticae</i> in <i>Saccharomyces cerevisiae</i> . <i>Journal of Bioscience and Bioengineering</i> , 1991, 71, 275-277. | 0.9 | 3 |

| # | ARTICLE | IF | CITATIONS |
|------|---|------|-----------|
| 1387 | Expression of the xylan-degrading genes of <i>Bacillus pumilus</i> IPO in <i>Saccharomyces cerevisiae</i> . <i>Journal of Bioscience and Bioengineering</i> , 1991, 71, 303-308. | 0.9 | 1 |
| 1388 | Characterization of a nuclear localization sequence in the polyomavirus capsid protein VP1. <i>Virology</i> , 1991, 185, 513-518. | 1.1 | 54 |
| 1389 | Cryopreservation of competent <i>Schizosaccharomyces pombe</i> protoplasts. <i>Trends in Genetics</i> , 1991, 7, 40. | 2.9 | 9 |
| 1390 | Competent <i>Saccharomyces cerevisiae</i> cells can be frozen and used for transformation with high frequency. <i>Trends in Genetics</i> , 1991, 7, 242. | 2.9 | 7 |
| 1391 | PRP16 is an RNA-dependent ATPase that interacts transiently with the spliceosome. <i>Nature</i> , 1991, 349, 494-499. | 13.7 | 353 |
| 1392 | A homologue of the bacterial heat-shock gene DnaJ that alters protein sorting in yeast. <i>Nature</i> , 1991, 349, 627-630. | 13.7 | 121 |
| 1393 | A suppressor of a yeast splicing mutation (<i>prp8-l</i>) encodes a putative ATP-dependent RNA helicase. <i>Nature</i> , 1991, 349, 715-717. | 13.7 | 130 |
| 1394 | Regulated import and degradation of a cytosolic protein in the yeast vacuole. <i>Nature</i> , 1991, 350, 313-318. | 13.7 | 167 |
| 1395 | A suppressor of yeast <i>spp81/ded1</i> mutations encodes a very similar putative ATP-dependent RNA helicase. <i>Molecular Microbiology</i> , 1991, 5, 805-812. | 1.2 | 57 |
| 1396 | A new system for amplifying 2 μ m plasmid copy number in <i>Saccharomyces cerevisiae</i> . <i>Molecular Microbiology</i> , 1991, 5, 1539-1548. | 1.2 | 15 |
| 1397 | Thirty-three nucleotides of 5' flanking sequence including the 'TATA' box are necessary and sufficient for efficient U2 snRNA transcription in <i>Schizosaccharomyces pombe</i> . <i>Molecular Microbiology</i> , 1991, 5, 1621-1625. | 1.2 | 9 |
| 1398 | Deletion of the COX7 gene in <i>Saccharomyces cerevisiae</i> reveals a role for cytochrome c oxidase subunit VII in assembly of remaining subunits. <i>Molecular Microbiology</i> , 1991, 5, 1769-1777. | 1.2 | 36 |
| 1399 | Molecular cloning, characterization and analysis of the regulation of the <i>ARO2</i> gene, encoding chorismate synthase, of <i>Saccharomyces cerevisiae</i> . <i>Molecular Microbiology</i> , 1991, 5, 2143-2153. | 1.2 | 21 |
| 1400 | Secretion of oligomeric Val ⁸ -human calcitonin by <i>Saccharomyces cerevisiae</i> . <i>FEMS Microbiology Letters</i> , 1991, 83, 23-28. | 0.7 | 5 |
| 1401 | Transformation of the yeast <i>Saccharomyces kluyveri</i> by <i>Saccharomyces cerevisiae</i> -based plasmids. <i>FEMS Microbiology Letters</i> , 1991, 82, 149-152. | 0.7 | 13 |
| 1402 | The proteinase yscB inhibitor (PBI2) gene of yeast and studies on the function of its protein product. <i>FEBS Journal</i> , 1991, 197, 1-7. | 0.2 | 26 |
| 1403 | Carboxypeptidase yscS: gene structure and function of the vacuolar enzyme. <i>FEBS Journal</i> , 1991, 197, 399-405. | 0.2 | 37 |
| 1404 | Production and purification of recombinant human interleukin-6 secreted by the yeast <i>Saccharomyces cerevisiae</i> . <i>FEBS Journal</i> , 1991, 198, 217-222. | 0.2 | 38 |

| # | ARTICLE | IF | CITATIONS |
|------|---|-----|-----------|
| 1405 | Characterisation of the specific p-nitrophenylphosphatase gene and protein of <i>Schizosaccharomyces pombe</i> . <i>FEBS Journal</i> , 1991, 198, 493-497. | 0.2 | 12 |
| 1406 | Purification, characterisation and mutagenesis of highly expressed recombinant yeast pyruvate kinase. <i>FEBS Journal</i> , 1991, 198, 513-519. | 0.2 | 17 |
| 1407 | Glucose repression in <i>Saccharomyces cerevisiae</i> is directly associated with hexose phosphorylation by hexokinases PI and PII. <i>FEBS Journal</i> , 1991, 199, 511-518. | 0.2 | 196 |
| 1408 | Characterization of Hex2 protein, a negative regulatory element necessary for glucose repression in yeast. <i>FEBS Journal</i> , 1991, 200, 311-319. | 0.2 | 68 |
| 1409 | Identification and characterization of a <i>Saccharomyces cerevisiae</i> gene (PAR1) conferring resistance to iron chelators. <i>FEBS Journal</i> , 1991, 200, 487-493. | 0.2 | 75 |
| 1410 | Monoclonal antibodies against core and cellulose-binding domains of <i>Trichoderma reesei</i> cellobiohydrolases I and II and endoglucanase I. <i>FEBS Journal</i> , 1991, 200, 643-649. | 0.2 | 55 |
| 1411 | Molecular cloning of soluble aminopeptidases from <i>Saccharomyces cerevisiae</i> . Sequence analysis of aminopeptidase yscII, a putative zinc-metallopeptidase. <i>FEBS Journal</i> , 1991, 202, 993-1002. | 0.2 | 38 |
| 1412 | Interference with myosin subfragment-1 binding by site-directed mutagenesis of actin. <i>FEBS Journal</i> , 1991, 200, 35-41. | 0.2 | 54 |
| 1413 | PURPLE FISSION YEAST: OVEREXPRESSION and PROCESSING OF THE PIGMENT BACTERIORHODOPSIN IN <i>Schizosaccharomyces pombe</i> . <i>Photochemistry and Photobiology</i> , 1991, 54, 1009-1016. | 1.3 | 17 |
| 1414 | Endoglucanase a gene fusion vectors for monitoring protein secretion and glycosylation in yeast. <i>Analytical Biochemistry</i> , 1991, 197, 290-295. | 1.1 | 11 |
| 1415 | Rapid determination of plasmid-carrying yeast cells by using an imaging sensor system. <i>Biotechnology and Bioengineering</i> , 1991, 38, 1331-1336. | 1.7 | 2 |
| 1416 | High-level secretion of biologically active aprotinin from the yeast <i>Pichia pastoris</i> . <i>Journal of Industrial Microbiology</i> , 1991, 7, 197-201. | 0.9 | 74 |
| 1417 | Cu(II)-Binding properties of a cytochrome c with a synthetic metal-binding site: His-X3-His in an α -helix. <i>Proteins: Structure, Function and Bioinformatics</i> , 1991, 10, 156-161. | 1.5 | 84 |
| 1418 | RCS1, a gene involved in controlling cell size in <i>Saccharomyces cerevisiae</i> . <i>Yeast</i> , 1991, 7, 1-14. | 0.8 | 18 |
| 1419 | Yeast mutants with increased bacterial transposon Tn5 excision. <i>Yeast</i> , 1991, 7, 37-50. | 0.8 | 26 |
| 1420 | Mapping of the ARS-like activity and transcription initiation sites in the non-canonical yeast mitochondrial ori 6 region. <i>Yeast</i> , 1991, 7, 51-60. | 0.8 | 11 |
| 1421 | Sequence and genetic analysis of NHP2: A moderately abundant high mobility group-like nuclear protein with an essential function in <i>Saccharomyces cerevisiae</i> . <i>Yeast</i> , 1991, 7, 79-90. | 0.8 | 31 |
| 1422 | Influence of the codon following the initiation codon on the expression of the lacZ gene in <i>Saccharomyces cerevisiae</i> . <i>Yeast</i> , 1991, 7, 157-165. | 0.8 | 10 |

| # | ARTICLE | IF | CITATIONS |
|------|---|-----|-----------|
| 1423 | Applications of high efficiency lithium acetate transformation of intact yeast cells using single-stranded nucleic acids as carrier. <i>Yeast</i> , 1991, 7, 253-263. | 0.8 | 424 |
| 1424 | CDC15, an essential cell cycle gene in <i>Saccharomyces cerevisiae</i> , encodes a protein kinased domain. <i>Yeast</i> , 1991, 7, 265-273. | 0.8 | 115 |
| 1425 | Inositol biosynthesis: <i>Candida albicans</i> and <i>Saccharomyces cerevisiae</i> genes share common regulation. <i>Yeast</i> , 1991, 7, 325-336. | 0.8 | 14 |
| 1426 | Optimization of <i>Bacillus</i> α -amylase production by <i>Saccharomyces cerevisiae</i> . <i>Yeast</i> , 1991, 7, 337-346. | 0.8 | 32 |
| 1427 | DNA insertions in the "silent"™ regions of the 2 $\frac{1}{4}$ m plasmid of <i>Saccharomyces cerevisiae</i> influence plasmid stability. <i>Yeast</i> , 1991, 7, 347-356. | 0.8 | 7 |
| 1428 | A new glucose-repressible gene identified from the analysis of chromatin structure in deletion mutants of yeast SUC2 locus. <i>Yeast</i> , 1991, 7, 379-389. | 0.8 | 52 |
| 1429 | Intracellular expression of <i>Kluyveromyces lactis</i> toxin β subunit mimics treatment with exogenous toxin and distinguishes two classes of toxin-resistant mutant. <i>Yeast</i> , 1991, 7, 617-625. | 0.8 | 75 |
| 1430 | The sequence of 8.8 kb of yeast chromosome III cloned in lambda PM3270 contains an unusual long ORF (YCR601). <i>Yeast</i> , 1991, 7, 631-641. | 0.8 | 4 |
| 1431 | Cloning and sequencing of the yeast <i>Saccharomyces cerevisiae</i> SEC1 gene localized on chromosome IV. <i>Yeast</i> , 1991, 7, 643-650. | 0.8 | 70 |
| 1432 | Functional analysis of a conserved amino-terminal region of HSP70 by site-directed mutagenesis. <i>Yeast</i> , 1991, 7, 699-716. | 0.8 | 9 |
| 1433 | Yeast mutant affected for viability upon nutrient starvation: Characterization and cloning of the RVS161 gene. <i>Yeast</i> , 1991, 7, 727-743. | 0.8 | 122 |
| 1434 | Four major transcriptional responses in the methionine/threonine biosynthetic pathway of <i>Saccharomyces cerevisiae</i> . <i>Yeast</i> , 1991, 7, 781-803. | 0.8 | 107 |
| 1435 | Electroporation-stimulated recombination in yeast. <i>Yeast</i> , 1991, 7, 823-831. | 0.8 | 20 |
| 1436 | TDH2 is linked to MET3 on chromosome X of <i>Saccharomyces cerevisiae</i> . <i>Yeast</i> , 1991, 7, 873-880. | 0.8 | 38 |
| 1437 | Molecular cloning of the β -glutamylcysteine synthetase gene of <i>Saccharomyces cerevisiae</i> . <i>Yeast</i> , 1991, 7, 953-961. | 0.8 | 124 |
| 1438 | Cloning of the HIS3 gene of <i>Yarrowia lipolytica</i> . <i>Antonie Van Leeuwenhoek</i> , 1991, 60, 95-99. | 0.7 | 3 |
| 1439 | High level expression of isocitrate lyase gene of n-alkane-utilizing yeast <i>Candida tropicalis</i> in <i>Saccharomyces cerevisiae</i> . <i>Archives of Microbiology</i> , 1991, 156, 439-443. | 1.0 | 9 |
| 1440 | Analysis of a <i>Candida albicans</i> gene that encodes a novel mechanism for resistance to benomyl and methotrexate. <i>Molecular Genetics and Genomics</i> , 1991, 227, 318-329. | 2.4 | 208 |

| # | ARTICLE | IF | CITATIONS |
|------|---|-----|-----------|
| 1441 | The <i>Saccharomyces cerevisiae</i> genes (CMP1 and CMP2) encoding calmodulin-binding proteins homologous to the catalytic subunit of mammalian protein phosphatase 2B. <i>Molecular Genetics and Genomics</i> , 1991, 227, 52-59. | 2.4 | 182 |
| 1442 | Expression in yeast of a cDNA copy of the K2 killer toxin gene. <i>Molecular Genetics and Genomics</i> , 1991, 227, 127-136. | 2.4 | 63 |
| 1443 | APMR2 tandem repeat with a modified C-terminus is located downstream from the KRS1 gene encoding lysyl-tRNA synthetase in <i>Saccharomyces cerevisiae</i> . <i>Molecular Genetics and Genomics</i> , 1991, 227, 149-54. | 2.4 | 20 |
| 1444 | Mitochondrial protein import: isolation and characterization of the <i>Saccharomyces cerevisiae</i> MFT1 gene. <i>Molecular Genetics and Genomics</i> , 1991, 225, 483-491. | 2.4 | 34 |
| 1445 | The KNS1 gene of <i>Saccharomyces cerevisiae</i> encodes a nonessential protein kinase homologue that is distantly related to members of the CDC28/cdc2 gene family. <i>Molecular Genetics and Genomics</i> , 1991, 229, 1-9. | 2.4 | 38 |
| 1446 | The CCR1 (SNF1) and SCH9 protein kinases act independently of cAMP-dependent protein kinase and the transcriptional activator ADR1 in controlling yeast ADH2 expression. <i>Molecular Genetics and Genomics</i> , 1991, 229, 395-399. | 2.4 | 49 |
| 1447 | Cloning of the two essential yeast genes, PRP6 and PRP9, and their rapid mapping, disruption and partial sequencing using a linker insertion strategy. <i>Molecular Genetics and Genomics</i> , 1991, 225, 199-202. | 2.4 | 16 |
| 1448 | Mutations in a <i>Saccharomyces cerevisiae</i> host showing increased holding stability of the heterologous plasmid pSRI. <i>Molecular Genetics and Genomics</i> , 1991, 225, 257-265. | 2.4 | 26 |
| 1449 | DNA sequences in chromosomes 11 and VII code for pyruvate carboxylase isoenzymes in <i>Saccharomyces cerevisiae</i> : analysis of pyruvate carboxylase-deficient strains. <i>Molecular Genetics and Genomics</i> , 1991, 229, 307-315. | 2.4 | 90 |
| 1450 | The synthesis of the two S-adenosyl-methionine synthetases is differently regulated in <i>Saccharomyces cerevisiae</i> . <i>Molecular Genetics and Genomics</i> , 1991, 226-226, 224-232. | 2.4 | 75 |
| 1451 | Correlation of biochemical blocks and genetic lesions in leucine auxotrophic strains of the imperfect yeast <i>Candida maltosa</i> . <i>Molecular Genetics and Genomics</i> , 1991, 227, 361-368. | 2.4 | 9 |
| 1452 | The <i>Escherichia coli</i> recA gene increases resistance of the yeast <i>Saccharomyces cerevisiae</i> to ionizing and ultraviolet radiation. <i>Molecular Genetics and Genomics</i> , 1991, 227, 473-480. | 2.4 | 13 |
| 1453 | Mitochondrial inner membrane protease 1 of <i>Saccharomyces cerevisiae</i> shows sequence similarity to the <i>Escherichia coli</i> leader peptidase. <i>Molecular Genetics and Genomics</i> , 1991, 228, 167-176. | 2.4 | 118 |
| 1454 | CYP1 (HAP1) is a determinant effector of alternative expression of heme-dependent transcription in yeast. <i>Molecular Genetics and Genomics</i> , 1991, 228, 300-306. | 2.4 | 40 |
| 1455 | TFS1: A suppressor of cdc25 mutations in <i>Saccharomyces cerevisiae</i> . <i>Molecular Genetics and Genomics</i> , 1991, 230, 241-250. | 2.4 | 52 |
| 1456 | The <i>Saccharomyces cerevisiae</i> SPT14 gene is essential for normal expression of the yeast transposon, Ty, as well as for expression of the HIS4 gene and several genes in the mating pathway. <i>Molecular Genetics and Genomics</i> , 1991, 230, 310-320. | 2.4 | 33 |
| 1457 | A multi-component upstream activation sequence of the <i>Saccharomyces cerevisiae</i> glyceraldehyde-3-phosphate dehydrogenase gene promoter. <i>Molecular Genetics and Genomics</i> , 1991, 231, 22-32. | 2.4 | 74 |
| 1458 | An improved lithium acetate method for yeast transformation. <i>Biotechnology Letters</i> , 1991, 5, 13-18. | 0.5 | 10 |

| # | ARTICLE | IF | CITATIONS |
|------|---|-----|-----------|
| 1459 | Heterologous gene expression in continuous cultures of budding yeast. <i>Applied Microbiology and Biotechnology</i> , 1991, 34, 632-636. | 1.7 | 17 |
| 1460 | Hyper-resistance to nitrogen mustard in <i>Saccharomyces cerevisiae</i> is caused by defective choline transport. <i>Current Genetics</i> , 1991, 19, 423-427. | 0.8 | 14 |
| 1461 | The SNQ3 gene of <i>Saccharomyces cerevisiae</i> confers hyper-resistance to several functionally unrelated chemicals. <i>Current Genetics</i> , 1991, 19, 429-433. | 0.8 | 79 |
| 1462 | Ethanol improves the transformation efficiency of intact yeast cells. <i>Current Genetics</i> , 1991, 20, 1-3. | 0.8 | 20 |
| 1463 | Ubiquitin gene expression: response to environmental changes. <i>Current Genetics</i> , 1991, 20, 17-23. | 0.8 | 35 |
| 1464 | The TSM1 gene of <i>Saccharomyces cerevisiae</i> overlaps the MAT locus. <i>Current Genetics</i> , 1991, 20, 25-31. | 0.8 | 21 |
| 1465 | Molecular cloning and characterization of a <i>Candida tsukubaensis</i> β -glucosidase gene in the yeast <i>Saccharomyces cerevisiae</i> . <i>Current Genetics</i> , 1991, 20, 45-52. | 0.8 | 11 |
| 1466 | Sequence and localization of the gene encoding yeast phosphoglycerate mutase. <i>Current Genetics</i> , 1991, 20, 167-171. | 0.8 | 21 |
| 1467 | Complex interaction of yeast nuclear proteins with the enhancer/promoter region of SV40. <i>Current Genetics</i> , 1991, 20, 359-363. | 0.8 | 3 |
| 1468 | A novel method for in situ screening of yeast colonies with the β -glucuronidase reporter gene. <i>Current Genetics</i> , 1991, 20, 437-439. | 0.8 | 12 |
| 1469 | The CDC8 gene product is required for transformation with episomal and integrative plasmids in <i>Saccharomyces cerevisiae</i> . <i>Current Genetics</i> , 1991, 20, 265-267. | 0.8 | 1 |
| 1470 | Mapping of four ras superfamily genes by physical and genetic means in <i>Schizosaccharomyces pombe</i> . <i>Current Genetics</i> , 1991, 20, 277-281. | 0.8 | 1 |
| 1471 | Cloning, sequencing and expression of the <i>Schwanniomyces occidentalis</i> NADP-dependent glutamate dehydrogenase gene. <i>Current Genetics</i> , 1991, 20, 219-224. | 0.8 | 22 |
| 1472 | Isolation and genetic study of p-fluoro-dl-phenylalanine-resistant mutants overproducing β -phenethyl-alcohol in <i>Saccharomyces cerevisiae</i> . <i>Current Genetics</i> , 1991, 20, 449-452. | 0.8 | 27 |
| 1473 | Replication analysis of plasmid DNAs injected into <i>Drosophila</i> embryos. <i>Chromosoma</i> , 1991, 100, 267-277. | 1.0 | 5 |
| 1474 | Interactions between chromosomal omnipotent suppressors and extrachromosomal effectors in <i>Saccharomyces cerevisiae</i> . <i>Current Genetics</i> , 1991, 19, 243-248. | 0.8 | 12 |
| 1475 | Analysis of centromere function in <i>Saccharomyces cerevisiae</i> using synthetic centromere mutants. <i>Chromosoma</i> , 1991, 101, 189-197. | 1.0 | 40 |
| 1476 | A gene tightly linked to CEN6 is important for growth of <i>Saccharomyces cerevisiae</i> . <i>Current Genetics</i> , 1991, 19, 1-8. | 0.8 | 29 |

| # | ARTICLE | IF | CITATIONS |
|------|---|-----|-----------|
| 1477 | Yeast cell cycle protein CDC48p shows full-length homology to the mammalian protein VCP and is a member of a protein family involved in secretion, peroxisome formation, and gene expression.. Journal of Cell Biology, 1991, 114, 443-453. | 2.3 | 301 |
| 1478 | Rapid colony transformation of <i>Saccharomyces cerevisiae</i> . Nucleic Acids Research, 1991, 19, 1945-1945. | 6.5 | 13 |
| 1479 | The use of a synthetic tRNA gene as a novel approach to study <i>in vivo</i> transcription and chromatin structure in yeast. Nucleic Acids Research, 1991, 19, 3849-3855. | 6.5 | 14 |
| 1480 | Analysis of sequences in the <i>INO1</i> promoter that are involved in its regulation by phospholipid precursors. Nucleic Acids Research, 1991, 19, 1687-1693. | 6.5 | 122 |
| 1481 | Lack of a requirement for strict rotational alignment among transcription factor binding sites in yeast. Nucleic Acids Research, 1991, 19, 3099-3103. | 6.5 | 4 |
| 1482 | The <i>BOS1</i> gene encodes an essential 27-kD putative membrane protein that is required for vesicular transport from the ER to the Golgi complex in yeast.. Journal of Cell Biology, 1991, 113, 55-64. | 2.3 | 109 |
| 1483 | The <i>PHO84</i> gene of <i>Saccharomyces cerevisiae</i> encodes an inorganic phosphate transporter.. Molecular and Cellular Biology, 1991, 11, 3229-3238. | 1.1 | 343 |
| 1484 | Genetic fine-structural analysis of the <i>Saccharomyces cerevisiae</i> alpha-pheromone receptor.. Molecular Biology of the Cell, 1991, 2, 439-452. | 6.5 | 25 |
| 1485 | The yeast <i>MCK1</i> gene encodes a protein kinase homolog that activates early meiotic gene expression.. Genes and Development, 1991, 5, 533-548. | 2.7 | 146 |
| 1486 | Pheromone-dependent phosphorylation of the yeast <i>STE12</i> protein correlates with transcriptional activation.. Genes and Development, 1991, 5, 741-750. | 2.7 | 149 |
| 1487 | Suppressors of a U4 snRNA mutation define a novel U6 snRNP protein with RNA-binding motifs.. Genes and Development, 1991, 5, 773-785. | 2.7 | 153 |
| 1488 | A suppressor of a centromere DNA mutation encodes a putative protein kinase (<i>MCK1</i>).. Genes and Development, 1991, 5, 549-560. | 2.7 | 115 |
| 1489 | Different legumin protein domains act as vacuolar targeting signals.. Plant Cell, 1991, 3, 695-708. | 3.1 | 109 |
| 1490 | Improvement of bacterial α -glucanase thermostability by glycosylation. Journal of General Microbiology, 1991, 137, 579-585. | 2.3 | 64 |
| 1491 | Fission yeast genes that confer resistance to staurosporine encode an AP-1-like transcription factor and a protein kinase related to the mammalian ERK1/MAP2 and budding yeast FUS3 and KSS1 kinases.. Genes and Development, 1991, 5, 60-73. | 2.7 | 361 |
| 1492 | Direct introduction and transient expression of capped and non-capped RNA in <i>Saccharomyces cerevisiae</i> . Nucleic Acids Research, 1991, 19, 4949-4953. | 6.5 | 41 |
| 1493 | <i>Sec15</i> protein, an essential component of the exocytotic apparatus, is associated with the plasma membrane and with a soluble 19.5S particle.. Journal of Cell Biology, 1991, 112, 1117-1131. | 2.3 | 91 |
| 1494 | The <i>Arabidopsis</i> functional homolog of the <i>p34cdc2</i> protein kinase.. Plant Cell, 1991, 3, 531-540. | 3.1 | 245 |

| # | ARTICLE | IF | CITATIONS |
|------|--|-----|-----------|
| 1495 | Isolation and characterization of a gene encoding a G-protein alpha subunit from <i>Schizosaccharomyces pombe</i> : involvement in mating and sporulation pathways.. Proceedings of the National Academy of Sciences of the United States of America, 1991, 88, 5877-5881. | 3.3 | 162 |
| 1496 | Protein folding causes an arrest of preprotein translocation into mitochondria in vivo.. Journal of Cell Biology, 1991, 115, 1601-1609. | 2.3 | 117 |
| 1497 | A role for unsaturated fatty acids in mitochondrial movement and inheritance.. Journal of Cell Biology, 1991, 115, 1249-1257. | 2.3 | 102 |
| 1498 | The NSR1 gene encodes a protein that specifically binds nuclear localization sequences and has two RNA recognition motifs.. Journal of Cell Biology, 1991, 113, 1-12. | 2.3 | 189 |
| 1499 | A cytoskeleton-related gene, <i>uso1</i> , is required for intracellular protein transport in <i>Saccharomyces cerevisiae</i> .. Journal of Cell Biology, 1991, 113, 245-260. | 2.3 | 167 |
| 1500 | Yeast ribosomal proteins: XII. YS11 of <i>Saccharomyces cerevisiae</i> is a homologue to <i>E.coli</i> S4 according to the gene analysis. Nucleic Acids Research, 1991, 19, 2603-2608. | 6.5 | 22 |
| 1501 | In vivo mapping of a sequence required for interference with the yeast killer virus.. Proceedings of the National Academy of Sciences of the United States of America, 1991, 88, 1271-1275. | 3.3 | 25 |
| 1502 | Mcm2 and Mcm3, two proteins important for ARS activity, are related in structure and function.. Genes and Development, 1991, 5, 944-957. | 2.7 | 184 |
| 1503 | CAL1, a gene required for activity of chitin synthase 3 in <i>Saccharomyces cerevisiae</i> .. Journal of Cell Biology, 1991, 114, 101-109. | 2.3 | 174 |
| 1504 | [17] Preparation of clone libraries in yeast artificial-chromosome vectors. Methods in Enzymology, 1991, 194, 251-270. | 0.4 | 80 |
| 1505 | [22] Shuttle mutagenesis: Bacterial transposons for genetic manipulations in yeast. Methods in Enzymology, 1991, 194, 329-342. | 0.4 | 36 |
| 1506 | [26] Vectors for constitutive and inducible gene expression in yeast. Methods in Enzymology, 1991, 194, 389-398. | 0.4 | 209 |
| 1507 | [34] Production of proteins by secretion from yeast. Methods in Enzymology, 1991, 194, 491-507. | 0.4 | 29 |
| 1508 | MOD5 translation initiation sites determine N6-isopentenyladenosine modification of mitochondrial and cytoplasmic tRNA.. Molecular and Cellular Biology, 1991, 11, 2382-2390. | 1.1 | 94 |
| 1509 | Determinants for glycopospholipid anchoring of the <i>Saccharomyces cerevisiae</i> GAS1 protein to the plasma membrane.. Molecular and Cellular Biology, 1991, 11, 27-37. | 1.1 | 229 |
| 1510 | RAP1 protein activates and silences transcription of mating-type genes in yeast.. Genes and Development, 1991, 5, 616-628. | 2.7 | 178 |
| 1511 | The CYC8 and TUP1 proteins involved in glucose repression in <i>Saccharomyces cerevisiae</i> are associated in a protein complex.. Molecular and Cellular Biology, 1991, 11, 3307-3316. | 1.1 | 240 |
| 1512 | Cloning DPB3, the gene encoding the third subunit of DNA polymerase II of <i>Saccharomyces cerevisiae</i> . Nucleic Acids Research, 1991, 19, 4867-4872. | 6.5 | 98 |

| # | ARTICLE | IF | CITATIONS |
|------|---|-----|-----------|
| 1513 | ramutations which increase artificial chromosome stability in <i>Saccharomyces cerevisiae</i> identify transcription and recombination proteins. <i>Nucleic Acids Research</i> , 1991, 19, 1385-1391. | 6.5 | 81 |
| 1514 | [21] Recovery of plasmids from yeast into <i>Escherichia coli</i> : Shuttle vectors. <i>Methods in Enzymology</i> , 1991, 194, 319-329. | 0.4 | 147 |
| 1515 | TRK1 and TRK2 encode structurally related K ⁺ transporters in <i>Saccharomyces cerevisiae</i> .. <i>Molecular and Cellular Biology</i> , 1991, 11, 4266-4273. | 1.1 | 277 |
| 1516 | Synthesis of Wild Type and Mutant Human Hemoglobins in <i>Saccharomyces cerevisiae</i> . <i>Nature Biotechnology</i> , 1991, 9, 57-61. | 9.4 | 99 |
| 1517 | Heritable Damage to Yeast Caused by Transformation. <i>Nature Biotechnology</i> , 1991, 9, 179-182. | 9.4 | 20 |
| 1518 | Alanine scanning site-directed mutagenesis of the zinc fingers of transcription factor ADR1: residues that contact DNA and that transactivate.. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 1991, 88, 9188-9192. | 3.3 | 78 |
| 1519 | mRNA leader length and initiation codon context determine alternative AUG selection for the yeast gene MOD5.. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 1991, 88, 9789-9793. | 3.3 | 127 |
| 1520 | Expression of three mammalian cDNAs that interfere with RAS function in <i>Saccharomyces cerevisiae</i> .. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 1991, 88, 2913-2917. | 3.3 | 109 |
| 1521 | A fission-yeast gene encoding a protein with features of protein-tyrosine-phosphatases.. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 1991, 88, 3455-3459. | 3.3 | 49 |
| 1522 | The function of chitin synthases 2 and 3 in the <i>Saccharomyces cerevisiae</i> cell cycle.. <i>Journal of Cell Biology</i> , 1991, 114, 111-123. | 2.3 | 436 |
| 1523 | Synthesis of large rRNAs by RNA polymerase II in mutants of <i>Saccharomyces cerevisiae</i> defective in RNA polymerase I.. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 1991, 88, 3962-3966. | 3.3 | 176 |
| 1524 | Mutations in the HML E silencer of <i>Saccharomyces cerevisiae</i> yield metastable inheritance of transcriptional repression.. <i>Genes and Development</i> , 1991, 5, 605-615. | 2.7 | 112 |
| 1525 | Antiestrogen can establish nonproductive receptor complexes and alter chromatin structure at target enhancers.. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 1991, 88, 3125-3129. | 3.3 | 89 |
| 1526 | Characterization of YDJ1: a yeast homologue of the bacterial dnaJ protein.. <i>Journal of Cell Biology</i> , 1991, 114, 609-621. | 2.3 | 270 |
| 1527 | N-myristoylation is required for function of the pheromone-responsive G alpha protein of yeast: conditional activation of the pheromone response by a temperature-sensitive N-myristoyl transferase.. <i>Genes and Development</i> , 1991, 5, 1969-1981. | 2.7 | 84 |
| 1528 | Monitoring of intracellular calcium in <i>Saccharomyces cerevisiae</i> with an apoaequorin cDNA expression system.. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 1991, 88, 6878-6882. | 3.3 | 117 |
| 1529 | Glucose repression of transcription of the <i>Schizosaccharomyces pombe</i> fbp1 gene occurs by a cAMP signaling pathway.. <i>Genes and Development</i> , 1991, 5, 561-571. | 2.7 | 156 |
| 1530 | Unexpected point mutations activate cryptic 3' splice sites by perturbing a natural secondary structure within a yeast intron.. <i>Genes and Development</i> , 1991, 5, 1252-1263. | 2.7 | 79 |

| # | ARTICLE | IF | CITATIONS |
|------|--|-----|-----------|
| 1531 | The Large Surface Protein of Hepatitis B Virus Is Retained in the Yeast Endoplasmic Reticulum and Provokes Its Unique Enlargement. <i>DNA and Cell Biology</i> , 1991, 10, 191-200. | 0.9 | 29 |
| 1532 | DNA binding of CPF1 is required for optimal centromere function but not for maintaining methionine prototrophy in yeast. <i>Nucleic Acids Research</i> , 1991, 19, 2961-2969. | 6.5 | 64 |
| 1533 | Sequence of the clathrin heavy chain from <i>Saccharomyces cerevisiae</i> and requirement of the COOH terminus for clathrin function.. <i>Journal of Cell Biology</i> , 1991, 112, 65-80. | 2.3 | 66 |
| 1534 | Spermidine or spermine is essential for the aerobic growth of <i>Saccharomyces cerevisiae</i> .. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 1991, 88, 5872-5876. | 3.3 | 100 |
| 1535 | Expression of RNase P RNA in <i>Saccharomyces cerevisiae</i> is controlled by an unusual RNA polymerase III promoter.. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 1991, 88, 6986-6990. | 3.3 | 63 |
| 1536 | Stable Multicopy Vectors for High-Level Secretion of Recombinant Human Serum Albumin by <i>Kluyveromyces</i> Yeasts. <i>Bio/technology</i> , 1991, 9, 968-975. | 1.9 | 142 |
| 1537 | Myristic acid auxotrophy caused by mutation of <i>S. cerevisiae</i> myristoyl-CoA:protein N-myristoyltransferase.. <i>Journal of Cell Biology</i> , 1991, 113, 1313-1330. | 2.3 | 68 |
| 1538 | The yeast SRM1 protein and human RCC1 protein share analogous functions.. <i>Molecular Biology of the Cell</i> , 1991, 2, 781-792. | 6.5 | 47 |
| 1539 | sar1, a gene from <i>Schizosaccharomyces pombe</i> encoding a protein that regulates ras1.. <i>Molecular Biology of the Cell</i> , 1991, 2, 453-465. | 6.5 | 83 |
| 1540 | Direct selection for sequences encoding proteases of known specificity.. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 1991, 88, 5159-5162. | 3.3 | 35 |
| 1541 | Generation of a nested series of interstitial deletions in yeast artificial chromosomes carrying human DNA.. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 1991, 88, 5744-5748. | 3.3 | 19 |
| 1542 | Analysis of the interactions of functional domains of a nuclear origin of replication from <i>Saccharomyces cerevisiae</i> . <i>Nucleic Acids Research</i> , 1991, 19, 6255-6262. | 6.5 | 74 |
| 1543 | Characterization of RPR1, an essential gene encoding the RNA component of <i>Saccharomyces cerevisiae</i> nuclear RNase P.. <i>Molecular and Cellular Biology</i> , 1991, 11, 721-730. | 1.1 | 94 |
| 1544 | The URE2 gene product of <i>Saccharomyces cerevisiae</i> plays an important role in the cellular response to the nitrogen source and has homology to glutathione s-transferases.. <i>Molecular and Cellular Biology</i> , 1991, 11, 822-832. | 1.1 | 272 |
| 1545 | The Tissue-Specific Mammalian Transcription Factor, Pit-1, Activates Transcription in <i>Saccharomyces cerevisiae</i> . <i>Molecular Endocrinology</i> , 1991, 5, 1239-1245. | 3.7 | 14 |
| 1546 | Roles of SWI1, SWI2, and SWI3 proteins for transcriptional enhancement by steroid receptors. <i>Science</i> , 1992, 258, 1598-1604. | 6.0 | 483 |
| 1547 | Functional complementation of yeast ste6 by a mammalian multidrug resistance mdr gene. <i>Science</i> , 1992, 256, 232-234. | 6.0 | 246 |
| 1548 | Biochemical and genetic characterization of a yeast TFIIID mutant that alters transcription in vivo and DNA binding in vitro.. <i>Molecular and Cellular Biology</i> , 1992, 12, 2372-2382. | 1.1 | 46 |

| # | ARTICLE | IF | CITATIONS |
|------|---|-----|-----------|
| 1549 | Parallel pathways of gene regulation: homologous regulators SWI5 and ACE2 differentially control transcription of HO and chitinase.. <i>Genes and Development</i> , 1992, 6, 93-104. | 2.7 | 223 |
| 1550 | Ligand-dependent and -independent function of the transactivation regions of the human estrogen receptor in yeast.. <i>Molecular Endocrinology</i> , 1992, 6, 1043-1050. | 3.7 | 90 |
| 1551 | RelB, a new Rel family transcription activator that can interact with p50-NF-kappa B.. <i>Molecular and Cellular Biology</i> , 1992, 12, 674-684. | 1.1 | 390 |
| 1552 | Yeast ribosomal proteins: XIII.Saccharomyces cerevisiaeYL8A gene, interrupted with two introns, encodes a homolog of mammalian L7. <i>Nucleic Acids Research</i> , 1992, 20, 1011-1016. | 6.5 | 29 |
| 1553 | Dominant negative selection of heterologous genes: isolation of Candida albicans genes that interfere with Saccharomyces cerevisiae mating factor-induced cell cycle arrest.. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 1992, 89, 9410-9414. | 3.3 | 126 |
| 1554 | A replication map of a 61-kb circular derivative of Saccharomyces cerevisiae chromosome III.. <i>Molecular Biology of the Cell</i> , 1992, 3, 999-1013. | 0.9 | 74 |
| 1555 | Molecular cloning of RAD16, a gene involved in differential repair inSaccharomyces cerevisiae. <i>Nucleic Acids Research</i> , 1992, 20, 3925-3931. | 6.5 | 112 |
| 1556 | Differentially regulated malate synthase genes participate in carbon and nitrogen metabolism ofS.cerevisiae. <i>Nucleic Acids Research</i> , 1992, 20, 5677-5686. | 6.5 | 98 |
| 1557 | SPO12andSIT4suppress mutations inDBF2, which encodes a cell cycle protein kinase that is periodically expressed. <i>Nucleic Acids Research</i> , 1992, 20, 5617-5623. | 6.5 | 45 |
| 1558 | TheSchizosaccharomyces pombe rhp3+gene required for DNA repair and cell viability is functionally interchangeable with theRAD3gene ofSaccharomyces cerevisiae. <i>Nucleic Acids Research</i> , 1992, 20, 2327-2334. | 6.5 | 28 |
| 1559 | Determination of lysine residues affinity labeled in the active site of yeast RNA polymerase II(B) by mutagenesis. <i>Nucleic Acids Research</i> , 1992, 20, 4721-4725. | 6.5 | 35 |
| 1560 | The yeast actin intron contains a cryptic promoter that can be switched on by preventing transcriptional interference. <i>Nucleic Acids Research</i> , 1992, 20, 4733-4739. | 6.5 | 32 |
| 1561 | Assignment of a yeast protein necessary for mitochondrial transcription initiation. <i>Nucleic Acids Research</i> , 1992, 20, 1053-1059. | 6.5 | 79 |
| 1562 | Mutations in yeast calmodulin cause defects in spindle pole body functions and nuclear integrity.. <i>Journal of Cell Biology</i> , 1992, 119, 1625-1639. | 2.3 | 95 |
| 1563 | Functional dissection of the phosphorylated termini of fission yeast DNA topoisomerase II.. <i>Journal of Cell Biology</i> , 1992, 119, 1023-1036. | 2.3 | 95 |
| 1564 | Regulation of the yeast DNA replication genes through the Mlu I cell cycle box is dependent on SWI6.. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 1992, 89, 9479-9483. | 3.3 | 53 |
| 1565 | Fission yeast pap1-dependent transcription is negatively regulated by an essential nuclear protein, crm1.. <i>Molecular and Cellular Biology</i> , 1992, 12, 5474-5484. | 1.1 | 109 |
| 1566 | Requirement for ESP1 in the nuclear division of Saccharomyces cerevisiae.. <i>Molecular Biology of the Cell</i> , 1992, 3, 1443-1454. | 0.9 | 107 |

| # | ARTICLE | IF | CITATIONS |
|------|---|-----|-----------|
| 1567 | Characterization and mutagenesis of the gene encoding the A49 subunit of RNA polymerase A in <i>Saccharomyces cerevisiae</i> .. Proceedings of the National Academy of Sciences of the United States of America, 1992, 89, 9302-9305. | 3.3 | 58 |
| 1568 | Regulation of fitness in yeast overexpressing glycolytic enzymes: responses to heat shock and nitrogen starvation. <i>Genetical Research</i> , 1992, 59, 167-177. | 0.3 | 5 |
| 1569 | Fission yeast <i>sts1+</i> gene encodes a protein similar to the chicken lamin B receptor and is implicated in pleiotropic drug-sensitivity, divalent cation-sensitivity, and osmoregulation.. <i>Molecular Biology of the Cell</i> , 1992, 3, 263-273. | 0.9 | 31 |
| 1570 | Genetic and biochemical analysis of the adenyl cyclase-associated protein, <i>cap</i> , in <i>Schizosaccharomyces pombe</i> .. <i>Molecular Biology of the Cell</i> , 1992, 3, 167-180. | 0.9 | 114 |
| 1571 | Direct induction of G1-specific transcripts following reactivation of the Cdc28 kinase in the absence of de novo protein synthesis.. <i>Genes and Development</i> , 1992, 6, 557-567. | 2.7 | 50 |
| 1572 | Role of astral microtubules and actin in spindle orientation and migration in the budding yeast, <i>Saccharomyces cerevisiae</i> .. <i>Journal of Cell Biology</i> , 1992, 119, 583-593. | 2.3 | 307 |
| 1573 | Cis and trans regulatory elements required for regulation of the <i>CHO1</i> gene of <i>Saccharomyces cerevisiae</i> . <i>Nucleic Acids Research</i> , 1992, 20, 1411-1418. | 6.5 | 63 |
| 1574 | Characterization of TPM1 disrupted yeast cells indicates an involvement of tropomyosin in directed vesicular transport.. <i>Journal of Cell Biology</i> , 1992, 118, 285-299. | 2.3 | 161 |
| 1575 | Human and human-yeast chimeric U6 snRNA genes identify structural elements required for expression in yeast. <i>Nucleic Acids Research</i> , 1992, 20, 479-485. | 6.5 | 19 |
| 1576 | Separate domains of KAR1 mediate distinct functions in mitosis and nuclear fusion. <i>Journal of Cell Biology</i> , 1992, 117, 1277-1287. | 2.3 | 99 |
| 1577 | Signal transduction in <i>Saccharomyces cerevisiae</i> requires tyrosine and threonine phosphorylation of FUS3 and KSS1.. <i>Genes and Development</i> , 1992, 6, 1280-1292. | 2.7 | 287 |
| 1578 | <i>Saccharomyces telomeres</i> assume a non-nucleosomal chromatin structure.. <i>Genes and Development</i> , 1992, 6, 197-210. | 2.7 | 267 |
| 1579 | Two genes in <i>Saccharomyces cerevisiae</i> encode a membrane-bound form of casein kinase-1.. <i>Molecular Biology of the Cell</i> , 1992, 3, 275-286. | 0.9 | 100 |
| 1580 | Signal recognition particle receptor is important for cell growth and protein secretion in <i>Saccharomyces cerevisiae</i> .. <i>Molecular Biology of the Cell</i> , 1992, 3, 895-911. | 0.9 | 109 |
| 1581 | The VPS1 protein, a homolog of dynamin required for vacuolar protein sorting in <i>Saccharomyces cerevisiae</i> , is a GTPase with two functionally separable domains.. <i>Journal of Cell Biology</i> , 1992, 119, 773-786. | 2.3 | 204 |
| 1582 | Cloning and Characterization of an <i>Arabidopsis thaliana</i> Topoisomerase I Gene. <i>Plant Physiology</i> , 1992, 99, 1493-1501. | 2.3 | 40 |
| 1583 | The <i>Saccharomyces cerevisiae</i> MGT1 DNA repair methyltransferase gene: its promoter and entire coding sequence, regulation and in vivo biological functions. <i>Nucleic Acids Research</i> , 1992, 20, 3599-3606. | 6.5 | 44 |
| 1584 | The yeast UME6 gene product is required for transcriptional repression mediated by the CAR1 URS1 repressor binding site. <i>Nucleic Acids Research</i> , 1992, 20, 1909-1915. | 6.5 | 94 |

| # | ARTICLE | IF | CITATIONS |
|------|--|-----|-----------|
| 1585 | Cyclin-B homologs in <i>Saccharomyces cerevisiae</i> function in S phase and in G2.. <i>Genes and Development</i> , 1992, 6, 2021-2034. | 2.7 | 274 |
| 1586 | Kinesin-related proteins required for assembly of the mitotic spindle.. <i>Journal of Cell Biology</i> , 1992, 118, 95-108. | 2.3 | 347 |
| 1587 | Meiotic recombination and segregation of human-derived artificial chromosomes in <i>Saccharomyces cerevisiae</i> .. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 1992, 89, 5296-5300. | 3.3 | 65 |
| 1588 | Regulatory gene <i>INO4</i> of yeast phospholipid biosynthesis is positively autoregulated and functions as a trans-activator of fatty acid synthase genes <i>FAS1</i> and <i>FAS2</i> from <i>Saccharomyces cerevisiae</i> . <i>Nucleic Acids Research</i> , 1992, 20, 5955-5961. | 6.5 | 51 |
| 1589 | Alternative pathways for the sorting of soluble vacuolar proteins in yeast: a <i>vps35</i> null mutant missorts and secretes only a subset of vacuolar hydrolases.. <i>Molecular Biology of the Cell</i> , 1992, 3, 415-427. | 0.9 | 99 |
| 1590 | Mutation of a tyrosine localization signal in the cytosolic tail of yeast <i>Kex2</i> protease disrupts Golgi retention and results in default transport to the vacuole.. <i>Molecular Biology of the Cell</i> , 1992, 3, 1353-1371. | 0.9 | 224 |
| 1591 | Characterization of cDNA encoding the mouse DNA topoisomerase II that can complement the budding yeast <i>top2</i> mutation. <i>Nucleic Acids Research</i> , 1992, 20, 5297-5303. | 6.5 | 37 |
| 1592 | Identification of Fungal Genes Involved in Plant Pathogenesis and Host Range. <i>Plant Gene Research</i> , 1992, , 129-151. | 0.4 | 1 |
| 1593 | Cloning of the α -Amylase cDNA of <i>Aspergillus shirousamii</i> and Its Expression in <i>Saccharomyces cerevisiae</i> . <i>Bioscience, Biotechnology and Biochemistry</i> , 1992, 56, 174-179. | 0.6 | 24 |
| 1594 | The C terminus of the NF- κ B p50 precursor and an κ B isoform contain transcription activation domains. <i>Nucleic Acids Research</i> , 1992, 20, 2453-2458. | 6.5 | 50 |
| 1595 | Mutants in the <i>S. cerevisiae</i> <i>PKC1</i> gene display a cell cycle-specific osmotic stability defect.. <i>Journal of Cell Biology</i> , 1992, 116, 1221-1229. | 2.3 | 401 |
| 1596 | Selective and immediate effects of clathrin heavy chain mutations on Golgi membrane protein retention in <i>Saccharomyces cerevisiae</i> .. <i>Journal of Cell Biology</i> , 1992, 118, 531-540. | 2.3 | 151 |
| 1597 | <i>SED5</i> encodes a 39-kD integral membrane protein required for vesicular transport between the ER and the Golgi complex.. <i>Journal of Cell Biology</i> , 1992, 119, 513-521. | 2.3 | 305 |
| 1598 | <i>NOP3</i> is an essential yeast protein which is required for pre-rRNA processing.. <i>Journal of Cell Biology</i> , 1992, 119, 737-747. | 2.3 | 112 |
| 1599 | The yeast Ca(2+)-ATPase homologue, <i>PMR1</i> , is required for normal Golgi function and localizes in a novel Golgi-like distribution.. <i>Molecular Biology of the Cell</i> , 1992, 3, 633-654. | 0.9 | 431 |
| 1600 | A gene encoding a protein with seven zinc finger domains acts on the sexual differentiation pathways of <i>Schizosaccharomyces pombe</i> .. <i>Molecular Biology of the Cell</i> , 1992, 3, 721-734. | 0.9 | 46 |
| 1601 | The protein sequence and some intron positions are conserved between the switching genes <i>wi10</i> of <i>Schizosaccharomyces pombe</i> and the human excision repair gene <i>ERCC1</i> . <i>Nucleic Acids Research</i> , 1992, 20, 6347-6353. | 6.5 | 59 |
| 1602 | The <i>wi4+</i> gene of <i>Schizosaccharomyces pombe</i> encodes a homologue of mismatch repair enzymes. <i>Nucleic Acids Research</i> , 1992, 20, 2271-2278. | 6.5 | 57 |

| # | ARTICLE | IF | CITATIONS |
|------|--|-----|-----------|
| 1603 | SPT3 interacts with TFIID to allow normal transcription in <i>Saccharomyces cerevisiae</i> .. <i>Genes and Development</i> , 1992, 6, 1319-1331. | 2.7 | 217 |
| 1604 | Small sequence insertions within the branch point region dictate alternative sites of lariat formation in a yeast intron. <i>Nucleic Acids Research</i> , 1992, 20, 6649-6655. | 6.5 | 6 |
| 1605 | Chitinase and chitin synthase 1: counterbalancing activities in cell separation of <i>Saccharomyces cerevisiae</i> . <i>Journal of General Microbiology</i> , 1992, 138, 97-102. | 2.3 | 147 |
| 1606 | A new family of yeast nuclear pore complex proteins.. <i>Journal of Cell Biology</i> , 1992, 119, 705-723. | 2.3 | 256 |
| 1607 | Functional analysis of the transcriptional activator encoded by the maize B gene: evidence for a direct functional interaction between two classes of regulatory proteins.. <i>Genes and Development</i> , 1992, 6, 864-875. | 2.7 | 378 |
| 1608 | Sec8p and Sec15p are components of a plasma membrane-associated 19.5S particle that may function downstream of Sec4p to control exocytosis.. <i>Journal of Cell Biology</i> , 1992, 118, 1041-1056. | 2.3 | 116 |
| 1609 | A mutant nuclear protein with similarity to RNA binding proteins interferes with nuclear import in yeast.. <i>Molecular Biology of the Cell</i> , 1992, 3, 875-893. | 0.9 | 133 |
| 1610 | A genetic system for studying the activity of a proteolytic enzyme.. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 1992, 89, 4159-4162. | 3.3 | 26 |
| 1611 | Molecular analysis of POP2 gene, a gene required for glucose-derepression of gene expression in <i>Saccharomyces cerevisiae</i> . <i>Nucleic Acids Research</i> , 1992, 20, 6227-6233. | 6.5 | 67 |
| 1612 | Nuclear and mitochondrial inheritance in yeast depends on novel cytoplasmic structures defined by the MDM1 protein.. <i>Journal of Cell Biology</i> , 1992, 118, 385-395. | 2.3 | 99 |
| 1613 | Membrane protein sorting in the yeast secretory pathway: evidence that the vacuole may be the default compartment.. <i>Journal of Cell Biology</i> , 1992, 119, 69-83. | 2.3 | 181 |
| 1614 | Mitochondrial DNA maintenance in yeast requires a protein containing a region related to the GTP-binding domain of dynamin.. <i>Genes and Development</i> , 1992, 6, 380-389. | 2.7 | 237 |
| 1615 | Control of Gene Expression from the SUC2 Promoter of <i>Saccharomyces cerevisiae</i> using Two Carbon Sources.. <i>Kagaku Kogaku Ronbunshu</i> , 1992, 18, 693-700. | 0.1 | 3 |
| 1616 | Breeding and Properties of Sake Yeasts. <i>Journal of the Brewing Society of Japan</i> , 1992, 87, 801-813. | 0.1 | 2 |
| 1617 | [49] Yeast artificial chromosome modification and manipulation. <i>Methods in Enzymology</i> , 1992, 216, 584-603. | 0.4 | 18 |
| 1618 | [33] Purification of Sec4 Protein from <i>Saccharomyces Cerevisiae</i> and <i>Escherichia coli</i> . <i>Methods in Enzymology</i> , 1992, 219, 352-362. | 0.4 | 7 |
| 1619 | A transcriptionally active tRNA gene interferes with nucleosome positioning in vivo.. <i>Molecular and Cellular Biology</i> , 1992, 12, 4015-4025. | 1.1 | 87 |
| 1620 | Multiple mechanisms mediate glucose repression of the yeast GAL1 gene.. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 1992, 89, 5922-5926. | 3.3 | 66 |

| # | ARTICLE | IF | CITATIONS |
|------|--|-----|-----------|
| 1621 | A putative ATP-dependent RNA helicase involved in <i>Saccharomyces cerevisiae</i> ribosome assembly.. Proceedings of the National Academy of Sciences of the United States of America, 1992, 89, 11131-11135. | 3.3 | 103 |
| 1622 | Ectopic recombination between Ty elements in <i>Saccharomyces cerevisiae</i> is not induced by DNA damage.. Molecular and Cellular Biology, 1992, 12, 4441-4448. | 1.1 | 37 |
| 1623 | Involvement of the SIN4 global transcriptional regulator in the chromatin structure of <i>Saccharomyces cerevisiae</i> .. Molecular and Cellular Biology, 1992, 12, 4503-4514. | 1.1 | 170 |
| 1624 | The yeast EUG1 gene encodes an endoplasmic reticulum protein that is functionally related to protein disulfide isomerase.. Molecular and Cellular Biology, 1992, 12, 4601-4611. | 1.1 | 151 |
| 1625 | A REB1-binding site is required for GCN4-independent ILV1 basal level transcription and can be functionally replaced by an ABF1-binding site.. Molecular and Cellular Biology, 1992, 12, 5516-5526. | 1.1 | 48 |
| 1626 | The EGD1 product, a yeast homolog of human BTF3, may be involved in GAL4 DNA binding.. Molecular and Cellular Biology, 1992, 12, 5683-5689. | 1.1 | 50 |
| 1627 | RSR1, a ras-like gene homologous to Krev-1 (smg21A/rap1A): role in the development of cell polarity and interactions with the Ras pathway in <i>Saccharomyces cerevisiae</i> .. Molecular and Cellular Biology, 1992, 12, 758-766. | 1.1 | 97 |
| 1628 | Dissection of a carboxy-terminal region of the yeast regulatory protein RAP1 with effects on both transcriptional activation and silencing.. Molecular and Cellular Biology, 1992, 12, 1209-1217. | 1.1 | 83 |
| 1629 | Two alternative pathways of double-strand break repair that are kinetically separable and independently modulated.. Molecular and Cellular Biology, 1992, 12, 1292-1303. | 1.1 | 364 |
| 1630 | Hydrolysis of GTP by Sec4 protein plays an important role in vesicular transport and is stimulated by a GTPase-activating protein in <i>Saccharomyces cerevisiae</i> .. Molecular and Cellular Biology, 1992, 12, 2017-2028. | 1.1 | 132 |
| 1631 | STP1, a gene involved in pre-tRNA processing, encodes a nuclear protein containing zinc finger motifs.. Molecular and Cellular Biology, 1992, 12, 2633-2643. | 1.1 | 22 |
| 1632 | Dominant mutations in a gene encoding a putative protein kinase (BCK1) bypass the requirement for a <i>Saccharomyces cerevisiae</i> protein kinase C homolog.. Molecular and Cellular Biology, 1992, 12, 172-182. | 1.1 | 361 |
| 1633 | RPC53 encodes a subunit of <i>Saccharomyces cerevisiae</i> RNA polymerase C (III) whose inactivation leads to a predominantly G1 arrest.. Molecular and Cellular Biology, 1992, 12, 4314-4326. | 1.1 | 49 |
| 1634 | RPC82 encodes the highly conserved, third-largest subunit of RNA polymerase C (III) from <i>Saccharomyces cerevisiae</i> .. Molecular and Cellular Biology, 1992, 12, 4433-4440. | 1.1 | 29 |
| 1635 | Truncated protein phosphatase GLC7 restores translational activation of GCN4 expression in yeast mutants defective for the eIF-2 alpha kinase GCN2.. Molecular and Cellular Biology, 1992, 12, 5700-5710. | 1.1 | 106 |
| 1636 | Connections between RNA splicing and DNA intron mobility in yeast mitochondria: RNA maturase and DNA endonuclease switching experiments.. Molecular and Cellular Biology, 1992, 12, 696-705. | 1.1 | 42 |
| 1637 | IME4, a gene that mediates MAT and nutritional control of meiosis in <i>Saccharomyces cerevisiae</i> .. Molecular and Cellular Biology, 1992, 12, 1078-1086. | 1.1 | 133 |
| 1638 | ADR1c mutations enhance the ability of ADR1 to activate transcription by a mechanism that is independent of effects on cyclic AMP-dependent protein kinase phosphorylation of Ser-230.. Molecular and Cellular Biology, 1992, 12, 1507-1514. | 1.1 | 47 |

| # | ARTICLE | IF | CITATIONS |
|------|--|-----|-----------|
| 1639 | A presumptive helicase (MOT1 gene product) affects gene expression and is required for viability in the yeast <i>Saccharomyces cerevisiae</i> .. <i>Molecular and Cellular Biology</i> , 1992, 12, 1879-1892. | 1.1 | 179 |
| 1640 | Posttranslational control of Ty1 retrotransposition occurs at the level of protein processing.. <i>Molecular and Cellular Biology</i> , 1992, 12, 2813-2825. | 1.1 | 86 |
| 1641 | Role of GCR2 in transcriptional activation of yeast glycolytic genes.. <i>Molecular and Cellular Biology</i> , 1992, 12, 3834-3842. | 1.1 | 82 |
| 1642 | Yeast casein kinase I homologues: an essential gene pair.. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 1992, 89, 28-32. | 3.3 | 146 |
| 1643 | Functional expression of human <i>mdr1</i> in the yeast <i>Saccharomyces cerevisiae</i> .. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 1992, 89, 2302-2306. | 3.3 | 97 |
| 1644 | Gene products that promote mRNA turnover in <i>Saccharomyces cerevisiae</i> .. <i>Molecular and Cellular Biology</i> , 1992, 12, 2165-2177. | 1.1 | 308 |
| 1645 | Regulation of fitness in yeast overexpressing glycolytic enzymes: parameters of growth and viability. <i>Genetical Research</i> , 1992, 59, 35-48. | 0.3 | 15 |
| 1646 | dOct2, a <i>Drosophila</i> Oct transcription factor that functions in yeast.. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 1992, 89, 7080-7084. | 3.3 | 24 |
| 1647 | Cloning of a human galactokinase gene (GK2) on chromosome 15 by complementation in yeast.. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 1992, 89, 10887-10891. | 3.3 | 31 |
| 1648 | The <i>Saccharomyces cerevisiae</i> ARG4 initiator of meiotic gene conversion and its associated double-strand DNA breaks can be inhibited by transcriptional interference.. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 1992, 89, 12068-12072. | 3.3 | 70 |
| 1649 | A yeast protein phosphatase related to the vaccinia virus VH1 phosphatase is induced by nitrogen starvation.. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 1992, 89, 12175-12179. | 3.3 | 96 |
| 1650 | MAS5, a yeast homolog of DnaJ involved in mitochondrial protein import.. <i>Molecular and Cellular Biology</i> , 1992, 12, 283-291. | 1.1 | 146 |
| 1651 | The 70-kilodalton adenylyl cyclase-associated protein is not essential for interaction of <i>Saccharomyces cerevisiae</i> adenylyl cyclase with RAS proteins.. <i>Molecular and Cellular Biology</i> , 1992, 12, 4937-4945. | 1.1 | 34 |
| 1652 | Molecular and genetic analysis of the yeast early meiotic recombination genes REC102 and REC107/MER2.. <i>Molecular and Cellular Biology</i> , 1992, 12, 1248-1256. | 1.1 | 56 |
| 1653 | Regulation of yeast COX6 by the general transcription factor ABF1 and separate HAP2- and heme-responsive elements.. <i>Molecular and Cellular Biology</i> , 1992, 12, 2302-2314. | 1.1 | 44 |
| 1654 | PMI40, an intron-containing gene required for early steps in yeast mannosylation.. <i>Molecular and Cellular Biology</i> , 1992, 12, 2924-2930. | 1.1 | 65 |
| 1655 | <i>Saccharomyces cerevisiae</i> RAD5-encoded DNA repair protein contains DNA helicase and zinc-binding sequence motifs and affects the stability of simple repetitive sequences in the genome.. <i>Molecular and Cellular Biology</i> , 1992, 12, 3807-3818. | 1.1 | 213 |
| 1656 | NAM9 nuclear suppressor of mitochondrial ochre mutations in <i>Saccharomyces cerevisiae</i> codes for a protein homologous to S4 ribosomal proteins from chloroplasts, bacteria, and eucaryotes.. <i>Molecular and Cellular Biology</i> , 1992, 12, 402-412. | 1.1 | 34 |

| # | ARTICLE | IF | CITATIONS |
|------|--|-----|-----------|
| 1657 | Characterization of the DNA target site for the yeast ARGR regulatory complex, a sequence able to mediate repression or induction by arginine.. <i>Molecular and Cellular Biology</i> , 1992, 12, 68-81. | 1.1 | 62 |
| 1658 | Yeast RHO3 and RHO4 ras superfamily genes are necessary for bud growth, and their defect is suppressed by a high dose of bud formation genes CDC42 and BEM1.. <i>Molecular and Cellular Biology</i> , 1992, 12, 5690-5699. | 1.1 | 166 |
| 1659 | MEI4, a meiosis-specific yeast gene required for chromosome synapsis.. <i>Molecular and Cellular Biology</i> , 1992, 12, 1340-1351. | 1.1 | 65 |
| 1660 | Human p53 and CDC2Hs genes combine to inhibit the proliferation of <i>Saccharomyces cerevisiae</i> .. <i>Molecular and Cellular Biology</i> , 1992, 12, 1357-1365. | 1.1 | 126 |
| 1661 | MET4, a leucine zipper protein, and centromere-binding factor 1 are both required for transcriptional activation of sulfur metabolism in <i>Saccharomyces cerevisiae</i> .. <i>Molecular and Cellular Biology</i> , 1992, 12, 1719-1727. | 1.1 | 165 |
| 1662 | Fused protein domains inhibit DNA binding by LexA.. <i>Molecular and Cellular Biology</i> , 1992, 12, 3006-3014. | 1.1 | 185 |
| 1663 | COT1, a gene involved in cobalt accumulation in <i>Saccharomyces cerevisiae</i> .. <i>Molecular and Cellular Biology</i> , 1992, 12, 3678-3688. | 1.1 | 197 |
| 1664 | A delay in the <i>Saccharomyces cerevisiae</i> cell cycle that is induced by a dicentric chromosome and dependent upon mitotic checkpoints.. <i>Molecular and Cellular Biology</i> , 1992, 12, 3857-3864. | 1.1 | 49 |
| 1665 | PRP38 encodes a yeast protein required for pre-mRNA splicing and maintenance of stable U6 small nuclear RNA levels.. <i>Molecular and Cellular Biology</i> , 1992, 12, 3939-3947. | 1.1 | 79 |
| 1666 | Replication forks pause at yeast centromeres.. <i>Molecular and Cellular Biology</i> , 1992, 12, 4056-4066. | 1.1 | 134 |
| 1667 | Genetic interaction between transcription elongation factor TFIIIS and RNA polymerase II.. <i>Molecular and Cellular Biology</i> , 1992, 12, 4142-4152. | 1.1 | 133 |
| 1668 | Identification of pre-mRNA polyadenylation sites in <i>Saccharomyces cerevisiae</i> .. <i>Molecular and Cellular Biology</i> , 1992, 12, 4215-4229. | 1.1 | 61 |
| 1669 | Chromosomal context dependence of a eukaryotic recombinational hot spot.. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 1992, 89, 227-231. | 3.3 | 75 |
| 1670 | Discrimination between related DNA sites by a single amino acid residue of Myc-related basic-helix-loop-helix proteins.. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 1992, 89, 599-602. | 3.3 | 179 |
| 1671 | Coordinated regulation and inositol-mediated and fatty acid-mediated repression of fatty acid synthase genes in <i>Saccharomyces cerevisiae</i> .. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 1992, 89, 10232-10236. | 3.3 | 73 |
| 1672 | Preferential repair of cyclobutane pyrimidine dimers in the transcribed strand of a gene in yeast chromosomes and plasmids is dependent on transcription.. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 1992, 89, 10696-10700. | 3.3 | 173 |
| 1673 | RNA-dependent RNA polymerase consensus sequence of the L-A double-stranded RNA virus: definition of essential domains.. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 1992, 89, 2185-2189. | 3.3 | 64 |
| 1674 | A dominant activating mutation in the effector region of RAS abolishes IRA2 sensitivity.. <i>Molecular and Cellular Biology</i> , 1992, 12, 631-637. | 1.1 | 20 |

| # | ARTICLE | IF | CITATIONS |
|------|---|-----|-----------|
| 1675 | Identification of a human cDNA encoding a protein that is structurally and functionally related to the yeast adenylyl cyclase-associated CAP proteins.. <i>Molecular and Cellular Biology</i> , 1992, 12, 5033-5040. | 1.1 | 74 |
| 1676 | CTF4 (CHL15) mutants exhibit defective DNA metabolism in the yeast <i>Saccharomyces cerevisiae</i> .. <i>Molecular and Cellular Biology</i> , 1992, 12, 5736-5747. | 1.1 | 88 |
| 1677 | Characterization of double-strand break-induced recombination: homology requirements and single-stranded DNA formation.. <i>Molecular and Cellular Biology</i> , 1992, 12, 563-575. | 1.1 | 387 |
| 1678 | Role of multifunctional autonomously replicating sequence binding factor 1 in the initiation of DNA replication and transcriptional control in <i>Saccharomyces cerevisiae</i> .. <i>Molecular and Cellular Biology</i> , 1992, 12, 1064-1077. | 1.1 | 84 |
| 1679 | Mutational analysis of conserved positions potentially important for initiator tRNA function in <i>Saccharomyces cerevisiae</i> .. <i>Molecular and Cellular Biology</i> , 1992, 12, 1432-1442. | 1.1 | 44 |
| 1680 | Different sequence elements are required for function of the cauliflower mosaic virus polyadenylation site in <i>Saccharomyces cerevisiae</i> compared with in plants.. <i>Molecular and Cellular Biology</i> , 1992, 12, 2322-2330. | 1.1 | 40 |
| 1681 | Overproduction of the GAL1 or GAL3 protein causes galactose-independent activation of the GAL4 protein: evidence for a new model of induction for the yeast GAL/MEL regulon.. <i>Molecular and Cellular Biology</i> , 1992, 12, 2701-2707. | 1.1 | 109 |
| 1682 | In vivo production of a stable single-stranded cDNA in <i>Saccharomyces cerevisiae</i> by means of a bacterial retron.. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 1992, 89, 5735-5739. | 3.3 | 20 |
| 1683 | The osmotic integrity of the yeast cell requires a functional PKC1 gene product.. <i>Molecular and Cellular Biology</i> , 1992, 12, 4896-4905. | 1.1 | 251 |
| 1684 | Semidominant suppressors of Srs2 helicase mutations of <i>Saccharomyces cerevisiae</i> map in the RAD51 gene, whose sequence predicts a protein with similarities to procaryotic RecA proteins.. <i>Molecular and Cellular Biology</i> , 1992, 12, 3224-3234. | 1.1 | 354 |
| 1685 | DPH5, a methyltransferase gene required for diphthamide biosynthesis in <i>Saccharomyces cerevisiae</i> .. <i>Molecular and Cellular Biology</i> , 1992, 12, 4026-4037. | 1.1 | 69 |
| 1686 | Identification of a mammalian gene structurally and functionally related to the CDC25 gene of <i>Saccharomyces cerevisiae</i> .. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 1992, 89, 7100-7104. | 3.3 | 84 |
| 1687 | Control of mRNA turnover as a mechanism of glucose repression in <i>Saccharomyces cerevisiae</i> .. <i>Molecular and Cellular Biology</i> , 1992, 12, 2941-2948. | 1.1 | 68 |
| 1688 | The ARS consensus sequence is required for chromosomal origin function in <i>Saccharomyces cerevisiae</i> .. <i>Molecular and Cellular Biology</i> , 1992, 12, 4305-4313. | 1.1 | 117 |
| 1689 | SEN1, a positive effector of tRNA-splicing endonuclease in <i>Saccharomyces cerevisiae</i> .. <i>Molecular and Cellular Biology</i> , 1992, 12, 2154-2164. | 1.1 | 80 |
| 1690 | Nitrogen catabolite repression of arginase (CAR1) expression in <i>Saccharomyces cerevisiae</i> is derived from regulated inducer exclusion. <i>Journal of Bacteriology</i> , 1992, 174, 48-55. | 1.0 | 19 |
| 1691 | Expression of the thermostable beta-galactosidase gene from the archaeobacterium <i>Sulfolobus solfataricus</i> in <i>Saccharomyces cerevisiae</i> and characterization of a new inducible promoter for heterologous expression. <i>Journal of Bacteriology</i> , 1992, 174, 873-882. | 1.0 | 63 |
| 1692 | Lysine biosynthesis in selected pathogenic fungi: characterization of lysine auxotrophs and the cloned LYS1 gene of <i>Candida albicans</i> . <i>Journal of Bacteriology</i> , 1992, 174, 7379-7384. | 1.0 | 53 |

| # | ARTICLE | IF | CITATIONS |
|------|---|------|-----------|
| 1693 | [29] Probing molecular structure and structural changes of voltage-gated channel by expressing mutant channels in yeast and reconstituting them into planar membranes. <i>Methods in Enzymology</i> , 1992, 207, 432-444. | 0.4 | 4 |
| 1694 | Cloning and characterization of the CYS3 (CY11) gene of <i>Saccharomyces cerevisiae</i> . <i>Journal of Bacteriology</i> , 1992, 174, 3339-3347. | 1.0 | 44 |
| 1695 | Automatic elimination of unnecessary bacterial sequences from yeast vectors. <i>Gene</i> , 1992, 121, 161-165. | 1.0 | 2 |
| 1696 | Efficient synthesis of secreted murine interleukin-2 by <i>Saccharomyces cerevisiae</i> : influence of 3' untranslated regions and codon usage. <i>Gene</i> , 1992, 111, 207-213. | 1.0 | 16 |
| 1697 | Secretion of a <i>Cryptococcus albidus</i> xylanase in <i>Saccharomyces cerevisiae</i> . <i>Gene</i> , 1992, 116, 109-113. | 1.0 | 30 |
| 1698 | Cloning and sequence analysis of a <i>Candida maltosa</i> gene which confers resistance to cycloheximide. <i>Gene</i> , 1992, 116, 105-108. | 1.0 | 49 |
| 1699 | Strategies for the Genetic Manipulation of <i>Saccharomyces cerevisiae</i> . <i>Critical Reviews in Biotechnology</i> , 1992, 12, 157-188. | 5.1 | 16 |
| 1700 | Expression and secretion in yeast of active insect defensin, an inducible antibacterial peptide from the fleshfly <i>Phormia terranova</i> . <i>Invertebrate Reproduction and Development</i> , 1992, 21, 15-24. | 0.3 | 25 |
| 1701 | Protocol for High-Efficiency Yeast Transformation. , 1992, , 501-505. | | 1 |
| 1702 | Unipolar cell divisions in the yeast <i>S. cerevisiae</i> lead to filamentous growth: Regulation by starvation and RAS. <i>Cell</i> , 1992, 68, 1077-1090. | 13.5 | 1,202 |
| 1703 | A new essential gene of <i>Saccharomyces cerevisiae</i> , a defect in it may result in instability of nucleus. <i>Experimental Cell Research</i> , 1992, 200, 48-57. | 1.2 | 23 |
| 1704 | High-level production of a peroxisomal enzyme: <i>Aspergillus flavus</i> uricase accumulates intracellularly and is active in <i>Saccharomyces cerevisiae</i> . <i>Gene</i> , 1992, 122, 139-145. | 1.0 | 41 |
| 1705 | The phenotype of a dihydrofolate reductase mutant of <i>Saccharomyces cerevisiae</i> . <i>Gene</i> , 1992, 121, 167-171. | 1.0 | 45 |
| 1706 | A homolog of the proteasome-related RING10 gene is essential for yeast cell growth. <i>Gene</i> , 1992, 122, 203-206. | 1.0 | 15 |
| 1707 | A novel, highly regulated, rapidly inducible system for the expression of chicken progesterone receptor, cPRA, in <i>Saccharomyces cerevisiae</i> . <i>Gene</i> , 1992, 114, 51-58. | 1.0 | 20 |
| 1708 | Cloning and regulation of <i>Schizosaccharomyces pombe</i> thi2, a gene involved in thiamine biosynthesis. <i>Gene</i> , 1992, 117, 141-143. | 1.0 | 29 |
| 1709 | Expression of recombinant cytochromes c from various species in <i>Saccharomyces cerevisiae</i> : post-translational modifications. <i>Protein Expression and Purification</i> , 1992, 3, 441-452. | 0.6 | 22 |
| 1710 | The cell division cycle gene CDC60 encodes cytosolic leucyl-tRNA synthetase in <i>Saccharomyces cerevisiae</i> . <i>Gene</i> , 1992, 120, 43-49. | 1.0 | 28 |

| # | ARTICLE | IF | CITATIONS |
|------|--|------|-----------|
| 1711 | Purification and characterization of wheat $\hat{\iota}$ -gliadin synthesized in the yeast, <i>Saccharomyces cerevisiae</i> . <i>Gene</i> , 1992, 116, 119-127. | 1.0 | 18 |
| 1712 | Molecular analysis of the yeast Ty4 element: homology with Ty1, copia, and plant retrotransposons. <i>Gene</i> , 1992, 122, 119-128. | 1.0 | 25 |
| 1713 | Cloning and analysis of a <i>Candida maltosa</i> gene which confers resistance to formaldehyde in <i>Saccharomyces cerevisiae</i> . <i>Gene</i> , 1992, 122, 207-211. | 1.0 | 63 |
| 1714 | Gene transfer strategies in plants. <i>Biotechnology Advances</i> , 1992, 10, 393-412. | 6.0 | 16 |
| 1715 | A coding region segment is necessary, but not sufficient for rapid decay of the HIS3 mRNA in yeast. <i>Gene</i> , 1992, 114, 35-41. | 1.0 | 33 |
| 1716 | Production of cytochrome P450 reductase yeast-rat hybrid proteins in <i>Saccharomyces cerevisiae</i> . <i>Gene</i> , 1992, 110, 33-39. | 1.0 | 18 |
| 1717 | Structure and Function of the Yeast Plasma-Membrane H ⁺ -ATPase. <i>Annals of the New York Academy of Sciences</i> , 1992, 671, 195-203. | 1.8 | 29 |
| 1718 | The yeast SUA7 gene encodes a homolog of human transcription factor TFIIB and is required for normal start site selection in vivo. <i>Cell</i> , 1992, 68, 977-988. | 13.5 | 216 |
| 1719 | Ssn6-Tup1 is a general repressor of transcription in yeast. <i>Cell</i> , 1992, 68, 709-719. | 13.5 | 726 |
| 1720 | Characterization of SAP-1, a protein recruited by serum response factor to the c-fos serum response element. <i>Cell</i> , 1992, 68, 597-612. | 13.5 | 682 |
| 1721 | The mechanism of RU486 antagonism is dependent on the conformation of the carboxy-terminal tail of the human progesterone receptor. <i>Cell</i> , 1992, 69, 703-713. | 13.5 | 388 |
| 1722 | Translation initiation requires the PAB-dependent poly(A) ribonuclease in yeast. <i>Cell</i> , 1992, 70, 961-973. | 13.5 | 214 |
| 1723 | The translation machinery and 70 kd heat shock protein cooperate in protein synthesis. <i>Cell</i> , 1992, 71, 97-105. | 13.5 | 512 |
| 1724 | The arrest of replication forks in the rDNA of yeast occurs independently of transcription. <i>Cell</i> , 1992, 71, 267-276. | 13.5 | 266 |
| 1725 | Autoregulation of the yeast lysyl-tRNA synthetase gene GCD5/KRS1 by translational and transcriptional control mechanisms. <i>Cell</i> , 1992, 70, 647-657. | 13.5 | 63 |
| 1726 | DMC1: A meiosis-specific yeast homolog of <i>E. coli</i> recA required for recombination, synaptonemal complex formation, and cell cycle progression. <i>Cell</i> , 1992, 69, 439-456. | 13.5 | 1,201 |
| 1727 | Rad51 protein involved in repair and recombination in <i>S. cerevisiae</i> is a RecA-like protein. <i>Cell</i> , 1992, 69, 457-470. | 13.5 | 1,249 |
| 1728 | A position effect on the time of replication origin activation in yeast. <i>Cell</i> , 1992, 68, 333-339. | 13.5 | 268 |

| # | ARTICLE | IF | CITATIONS |
|------|---|------|-----------|
| 1729 | SHR3: A novel component of the secretory pathway specifically required for localization of amino acid permeases in yeast. <i>Cell</i> , 1992, 71, 463-478. | 13.5 | 210 |
| 1730 | Nucleosome disruption at the yeast PHO5 promoter upon PHO5 induction occurs in the absence of DNA replication. <i>Cell</i> , 1992, 71, 853-864. | 13.5 | 169 |
| 1731 | A new signal peptide useful for secretion of heterologous proteins from yeast and its application for synthesis of hirudin. <i>Gene</i> , 1992, 110, 25-31. | 1.0 | 37 |
| 1732 | Multifunctional yeast high-copy-number shuttle vectors. <i>Gene</i> , 1992, 110, 119-122. | 1.0 | 1,658 |
| 1733 | High-frequency binding of IgE to the Der p allergen expressed in yeast*3. <i>Journal of Allergy and Clinical Immunology</i> , 1992, 89, 95-102. | 1.5 | 54 |
| 1734 | YDJ1p facilitates polypeptide translocation across different intracellular membranes by a conserved mechanism. <i>Cell</i> , 1992, 71, 1143-1155. | 13.5 | 271 |
| 1735 | Mutagenesis of the yeast plasma membrane H(+)-ATPase. A novel expression system. <i>Biophysical Journal</i> , 1992, 62, 228-237. | 0.2 | 20 |
| 1736 | The C-terminus of the succinate dehydrogenase IP peptide of <i>Saccharomyces cerevisiae</i> is significant for assembly of complex II. <i>Biochemistry</i> , 1992, 31, 8442-8448. | 1.2 | 18 |
| 1737 | Production of a soluble and secreted antigenic fragment of HBsAg in yeast. <i>Biochemical and Biophysical Research Communications</i> , 1992, 184, 50-59. | 1.0 | 2 |
| 1738 | Mammalian and viral DNA sequences which interfere with the maintenance of a centromeric vector in yeast. <i>Biochemical and Biophysical Research Communications</i> , 1992, 187, 737-743. | 1.0 | 2 |
| 1739 | Molecular cloning of acetyl coenzyme a: Deacetylcephalosporin C o-acetyltransferase cDNA from <i>Acremonium chrysogenum</i> : Sequence and expression of catalytic activity in yeast. <i>Biochemical and Biophysical Research Communications</i> , 1992, 182, 995-1001. | 1.0 | 27 |
| 1740 | Expression and function of heterologous forms of malate dehydrogenase in yeast. <i>Archives of Biochemistry and Biophysics</i> , 1992, 293, 93-102. | 1.4 | 11 |
| 1741 | Expression of human ferredoxin in <i>Saccharomyces cerevisiae</i> : Mitochondrial import of the protein and assembly of the [2Fe-2S] center. <i>Archives of Biochemistry and Biophysics</i> , 1992, 294, 603-608. | 1.4 | 13 |
| 1742 | Evolution of organellar proton-ATPases. <i>Biochimica Et Biophysica Acta - Bioenergetics</i> , 1992, 1100, 109-124. | 0.5 | 180 |
| 1743 | Increased sterigmatocystin-induced mutation frequency in <i>Saccharomyces cerevisiae</i> expressing cytochrome P450 CYP2B1. <i>Biochemical Pharmacology</i> , 1992, 43, 374-376. | 2.0 | 6 |
| 1744 | Molecular cloning of an essential yeast gene encoding a proteasomal subunit. <i>FEBS Letters</i> , 1992, 299, 39-43. | 1.3 | 27 |
| 1745 | Production of a unique multi-lamella structure in the nuclei of yeast expressing <i>Drosophila copia</i> gagprecursor. <i>FEBS Letters</i> , 1992, 302, 5-7. | 1.3 | 8 |
| 1746 | Cloning and characteristics of a positive regulatory gene, TH12(PHO6), of thiamin biosynthesis in <i>Saccharomyces cerevisiae</i> . <i>FEBS Letters</i> , 1992, 297, 155-158. | 1.3 | 48 |

| # | ARTICLE | IF | CITATIONS |
|------|--|-----|-----------|
| 1747 | Cloning of a human choline kinase cDNA by complementation of the yeastckimutation. FEBS Letters, 1992, 304, 229-232. | 1.3 | 40 |
| 1748 | Yeast ADP/ATP carrier (AAC) proteins exhibit similar enzymatic properties but their deletion produces different phenotypes. FEBS Letters, 1992, 304, 277-280. | 1.3 | 37 |
| 1749 | Upstream activation element of thePH03gene encoding for thiamine-repressible acid phosphatase inSaccharomyces cerevisiae. FEBS Letters, 1992, 305, 244-248. | 1.3 | 9 |
| 1750 | Protein engineering ofDrosophilaalcohol dehydrogenase The hydroxyl group of Tyr152is involved in the active site of the enzyme. FEBS Letters, 1992, 308, 235-239. | 1.3 | 49 |
| 1751 | Regulatory regions in the yeastFBP1andPCK1genes. FEBS Letters, 1992, 311, 110-114. | 1.3 | 57 |
| 1752 | Identification of cryptic lox sites in the yeast genome by selection for Cre-mediated chromosome translocations that confer multiple drug resistance. Journal of Molecular Biology, 1992, 223, 911-928. | 2.0 | 78 |
| 1753 | NAM7 nuclear gene encodes a novel member of a family of helicases with a Zn-ligand motif and is involved in mitochondrial functions in Saccharomyces cerevisiae. Journal of Molecular Biology, 1992, 224, 575-587. | 2.0 | 85 |
| 1754 | GCR3 encodes an acidic protein that is required for expression of glycolytic genes in Saccharomyces cerevisiae. Journal of Bacteriology, 1992, 174, 5526-5532. | 1.0 | 45 |
| 1755 | SRN1, a yeast gene involved in RNA processing, is identical to HEX2/REG1, a negative regulator in glucose repression.. Molecular and Cellular Biology, 1992, 12, 2673-2680. | 1.1 | 54 |
| 1756 | Proline biosynthesis in Saccharomyces cerevisiae: molecular analysis of the PRO1 gene, which encodes gamma-glutamyl kinase. Journal of Bacteriology, 1992, 174, 4148-4156. | 1.0 | 23 |
| 1757 | Identification and functional analysis of two U3 binding sites on yeast pre-ribosomal RNA.. EMBO Journal, 1992, 11, 1531-1542. | 3.5 | 234 |
| 1758 | A 22 bp cis-acting element is necessary and sufficient for the induction of the yeast KAR2 (BiP) gene by unfolded proteins.. EMBO Journal, 1992, 11, 2583-2593. | 3.5 | 334 |
| 1759 | Dominant-negative mutants of a yeast G-protein beta subunit identify two functional regions involved in pheromone signalling.. EMBO Journal, 1992, 11, 4805-4813. | 3.5 | 83 |
| 1760 | U14 function in Saccharomyces cerevisiae can be provided by large deletion variants of yeast U14 and hybrid mouse-yeast U14 RNAs.. EMBO Journal, 1992, 11, 683-689. | 3.5 | 35 |
| 1761 | The Saccharomyces cerevisiae NPS1 gene, a novel CDC gene which encodes a 160 kDa nuclear protein involved in G2 phase control.. EMBO Journal, 1992, 11, 4017-4026. | 3.5 | 70 |
| 1762 | The protein kinase homologue Ste20p is required to link the yeast pheromone response G-protein beta gamma subunits to downstream signalling components.. EMBO Journal, 1992, 11, 4815-4824. | 3.5 | 391 |
| 1763 | Directed mutagenesis in an asporogenous methylotrophic yeast: cloning, sequencing, and one-step gene disruption of the 3-isopropylmalate dehydrogenase gene (LEU2) of Candida boidinii to derive doubly auxotrophic marker strains. Journal of Bacteriology, 1992, 174, 5988-5993. | 1.0 | 49 |
| 1764 | Species specific protein-DNA interactions may determine the chromatin units of genes in S.cerevisiae and in S.pombe.. EMBO Journal, 1992, 11, 1177-1185. | 3.5 | 26 |

| # | ARTICLE | IF | CITATIONS |
|------|--|-----|-----------|
| 1765 | OCH1 encodes a novel membrane bound mannosyltransferase: outer chain elongation of asparagine-linked oligosaccharides.. EMBO Journal, 1992, 11, 2511-2519. | 3.5 | 248 |
| 1766 | Two distinct yeast transcriptional activators require the function of the GCN5 protein to promote normal levels of transcription.. EMBO Journal, 1992, 11, 4145-4152. | 3.5 | 272 |
| 1767 | GAC1 may encode a regulatory subunit for protein phosphatase type 1 in <i>Saccharomyces cerevisiae</i> .. EMBO Journal, 1992, 11, 87-96. | 3.5 | 145 |
| 1768 | Coordinate genetic control of yeast fatty acid synthase genes FAS1 and FAS2 by an upstream activation site common to genes involved in membrane lipid biosynthesis.. EMBO Journal, 1992, 11, 107-114. | 3.5 | 116 |
| 1769 | A role for clathrin in the sorting of vacuolar proteins in the Golgi complex of yeast.. EMBO Journal, 1992, 11, 2811-2818. | 3.5 | 140 |
| 1770 | A new subclass of nucleoporins that functionally interact with nuclear pore protein NSP1.. EMBO Journal, 1992, 11, 5051-5061. | 3.5 | 162 |
| 1771 | High-affinity ouabain binding by yeast cells expressing Na ⁺ , K ⁽⁺⁾ -ATPase alpha subunits and the gastric H ⁺ , K ⁽⁺⁾ -ATPase beta subunit.. Proceedings of the National Academy of Sciences of the United States of America, 1992, 89, 2834-2838. | 3.3 | 95 |
| 1772 | A novel and conserved salt-induced protein is an important determinant of salt tolerance in yeast.. EMBO Journal, 1992, 11, 3157-3164. | 3.5 | 177 |
| 1773 | Proline biosynthesis in <i>Saccharomyces cerevisiae</i> : analysis of the PRO3 gene, which encodes delta 1-pyrroline-5-carboxylate reductase. Journal of Bacteriology, 1992, 174, 3782-3788. | 1.0 | 49 |
| 1774 | Sequence-specific initiator elements focus initiation of transcription to distinct sites in the yeast TRP4 promoter.. EMBO Journal, 1992, 11, 4583-4590. | 3.5 | 16 |
| 1775 | Allelism of IMP1 and GAL2 genes of <i>Saccharomyces cerevisiae</i> . Journal of Bacteriology, 1992, 174, 3411-3415. | 1.0 | 17 |
| 1776 | Artificial nucleosome positioning sequences tested in yeast minichromosomes: a strong rotational setting is not sufficient to position nucleosomes in vivo.. EMBO Journal, 1992, 11, 1187-1193. | 3.5 | 36 |
| 1777 | Cloning by functional complementation of a mouse cDNA encoding a homologue of CDC25, a <i>Saccharomyces cerevisiae</i> RAS activator.. EMBO Journal, 1992, 11, 2151-2157. | 3.5 | 222 |
| 1778 | A zinc finger protein from <i>Candida albicans</i> is involved in sucrose utilization. Journal of Bacteriology, 1992, 174, 222-232. | 1.0 | 41 |
| 1779 | A positive regulatory gene, THI3, is required for thiamine metabolism in <i>Saccharomyces cerevisiae</i> . Journal of Bacteriology, 1992, 174, 4701-4706. | 1.0 | 39 |
| 1780 | Mutations activating the yeast eIF-2 alpha kinase GCN2: isolation of alleles altering the domain related to histidyl-tRNA synthetases.. Molecular and Cellular Biology, 1992, 12, 5801-5815. | 1.1 | 126 |
| 1781 | Drastic alteration of cycloheximide sensitivity by substitution of one amino acid in the L41 ribosomal protein of yeasts. Journal of Bacteriology, 1992, 174, 254-262. | 1.0 | 75 |
| 1782 | Ubiquitin as a degradation signal.. EMBO Journal, 1992, 11, 497-505. | 3.5 | 234 |

| # | ARTICLE | IF | CITATIONS |
|------|---|-----|-----------|
| 1783 | Expression of high levels of human tissue plasminogen activator in yeast under the control of an inducible GAL promoter. <i>Applied Microbiology and Biotechnology</i> , 1992, 37, 604-8. | 1.7 | 21 |
| 1784 | Studies on plasmid stability, cell metabolism and superoxide dismutase production by P _{gk} ? strains of <i>Saccharomyces cerevisiae</i> . <i>Applied Microbiology and Biotechnology</i> , 1992, 37, 615-20. | 1.7 | 8 |
| 1785 | Molecular cloning and analysis of <i>Schizosaccharomyces pombe</i> rad9, a gene involved in DNA repair and mutagenesis. <i>Molecular Genetics and Genomics</i> , 1992, 232, 367-376. | 2.4 | 46 |
| 1786 | Isolation and phenotypic analysis of conditional-lethal, linker-insertion mutations in the gene encoding the largest subunit of RNA polymerase II in <i>Saccharomyces cerevisiae</i> . <i>Molecular Genetics and Genomics</i> , 1992, 232, 408-414. | 2.4 | 37 |
| 1787 | Evolution of the alcohol dehydrogenase (ADH) genes in yeast: characterization of a fourth ADH in <i>Kluyveromyces lactis</i> . <i>Molecular Genetics and Genomics</i> , 1992, 232, 479-488. | 2.4 | 46 |
| 1788 | The role of promoter elements of a ribosomal protein gene in <i>Saccharomyces cerevisiae</i> under various physiological conditions. <i>Molecular Genetics and Genomics</i> , 1992, 234, 22-32. | 2.4 | 2 |
| 1789 | NPK1, a nonessential protein kinase gene in <i>Saccharomyces cerevisiae</i> with similarity to <i>Aspergillus nidulans</i> nimA. <i>Molecular Genetics and Genomics</i> , 1992, 234, 164-167. | 2.4 | 32 |
| 1790 | The cauliflower mosaic virus 35S promoter is regulated by cAMP in <i>Saccharomyces cerevisiae</i> . <i>Molecular Genetics and Genomics</i> , 1992, 235, 365-372. | 2.4 | 15 |
| 1791 | The DAC2/FUS3 protein kinase is not essential for transcriptional activation of the mating pheromone response pathway in <i>Saccharomyces cerevisiae</i> . <i>Molecular Genetics and Genomics</i> , 1992, 235, 450-452. | 2.4 | 11 |
| 1792 | The regulatory gene nit-2 of <i>Neurospora crassa</i> complements a nnu mutant of <i>Gibberella zeae</i> (<i>Fusarium graminearum</i>). <i>Molecular Genetics and Genomics</i> , 1992, 235, 458-462. | 2.4 | 8 |
| 1793 | Downstream activating sequence within the coding region of a yeast gene: specific binding in vitro of RAP1 protein. <i>Molecular Genetics and Genomics</i> , 1992, 236, 65-75. | 2.4 | 25 |
| 1794 | Positive regulation of the LPD1 gene of <i>Saccharomyces cerevisiae</i> by the HAP2/HAP3/HAP4 activation system. <i>Molecular Genetics and Genomics</i> , 1992, 231, 296-303. | 2.4 | 38 |
| 1795 | Inhibition of cell proliferation in <i>Saccharomyces cerevisiae</i> by expression of human NAD ⁺ ADP-ribosyltransferase requires the DNA binding domain (zinc fingers). <i>Molecular Genetics and Genomics</i> , 1992, 232, 231-239. | 2.4 | 39 |
| 1796 | ore2, a mutation affecting proline biosynthesis in the yeast <i>Saccharomyces cerevisiae</i> , leads to a cdc phenotype. <i>Molecular Genetics and Genomics</i> , 1992, 234, 193-200. | 2.4 | 2 |
| 1797 | Characterization of a staurosporine- and temperature-sensitive mutant, stt1, of <i>Saccharomyces cerevisiae</i> : STT1 is allelic to PKC1. <i>Molecular Genetics and Genomics</i> , 1992, 231, 337-344. | 2.4 | 86 |
| 1798 | Translational readthrough at nonsense mutations in the HSF1 gene of <i>Saccharomyces cerevisiae</i> . <i>Molecular Genetics and Genomics</i> , 1992, 234, 369-378. | 2.4 | 23 |
| 1799 | Molecular cloning of the imidazoleglycerolphosphate dehydratase gene of <i>Trichoderma harzianum</i> by genetic complementation in <i>Saccharomyces cerevisiae</i> using a direct expression vector. <i>Molecular Genetics and Genomics</i> , 1992, 234, 481-488. | 2.4 | 16 |
| 1800 | Dual function of a new nuclear gene for oxidative phosphorylation and vegetative growth in yeast. <i>Molecular Genetics and Genomics</i> , 1992, 232, 58-64. | 2.4 | 110 |

| # | ARTICLE | IF | CITATIONS |
|------|---|-----|-----------|
| 1801 | A novel mutation occurring in the PHO80 gene suppresses the PHO4 c mutations of <i>Saccharomyces cerevisiae</i> . <i>Current Genetics</i> , 1992, 21, 95-99. | 0.8 | 20 |
| 1802 | Physical and biochemical characterization of the cloned LYS5 gene required for γ -aminoadipate reductase activity in the lysine biosynthetic pathway of <i>Saccharomyces cerevisiae</i> . <i>Current Genetics</i> , 1992, 21, 13-16. | 0.8 | 6 |
| 1803 | One-step transformation of yeast in stationary phase. <i>Current Genetics</i> , 1992, 21, 83-84. | 0.8 | 673 |
| 1804 | The <i>Candida albicans</i> PMM1 gene encoding phosphomannomutase complements a <i>Saccharomyces cerevisiae</i> sec 53-6 mutation. <i>Current Genetics</i> , 1992, 22, 501-503. | 0.8 | 38 |
| 1805 | Positive cis-acting regulatory sequences mediate proper control of POL1 transcription in <i>Saccharomyces cerevisiae</i> . <i>Current Genetics</i> , 1992, 21, 183-189. | 0.8 | 14 |
| 1806 | Transformation of the thermophilic fungus <i>Humicola grisea</i> var. <i>thermoidea</i> and overproduction of <i>Humicola glucoamylase</i> . <i>Current Genetics</i> , 1992, 21, 225-229. | 0.8 | 20 |
| 1807 | The gene for cadmium metallothionein from a cadmium-resistant yeast appears to be identical to CUP 1 in a copper-resistant strain. <i>Current Genetics</i> , 1992, 21, 275-280. | 0.8 | 16 |
| 1808 | Cloning and mapping of the CYS4 gene of <i>Saccharomyces cerevisiae</i> . <i>Current Genetics</i> , 1992, 21, 285-289. | 0.8 | 16 |
| 1809 | Genetic analysis of serine biosynthesis and glucose repression in yeast. <i>Current Genetics</i> , 1992, 21, 295-300. | 0.8 | 49 |
| 1810 | The UGA43 negative regulatory gene of <i>Saccharomyces cerevisiae</i> contains both a GATA-1 type zinc finger and a putative leucine zipper. <i>Current Genetics</i> , 1992, 21, 301-307. | 0.8 | 61 |
| 1811 | Isolation and characterization of additional genes influencing resistance to various mutagens in the yeast <i>Saccharomyces cerevisiae</i> . <i>Current Genetics</i> , 1992, 21, 319-324. | 0.8 | 28 |
| 1812 | The CCS1 gene from <i>Saccharomyces cerevisiae</i> which is involved in mitochondrial functions is identified as IRA2 an attenuator of RAS1 and RAS2 gene products. <i>Current Genetics</i> , 1992, 21, 325-329. | 0.8 | 18 |
| 1813 | Complementation of the <i>Saccharomyces cerevisiae</i> srb1-1 mutation: an autoselection system for stable plasmid maintenance. <i>Current Genetics</i> , 1992, 21, 339-344. | 0.8 | 20 |
| 1814 | Construction of a Tetracycline-inducible promoter in <i>Schizosaccharomyces pombe</i> . <i>Current Genetics</i> , 1992, 21, 345-349. | 0.8 | 52 |
| 1815 | Mutational analysis of a variant of ARS1 from <i>Saccharomyces cerevisiae</i> . <i>Current Genetics</i> , 1992, 22, 175-180. | 0.8 | 5 |
| 1816 | The UASMAL is a bidirectional promoter element required for the expression of both the MAL61 and MAL62 genes of the <i>Saccharomyces</i> MAL6 locus. <i>Current Genetics</i> , 1992, 22, 181-189. | 0.8 | 60 |
| 1817 | In vitro mutagenesis of the mitochondrial leucyl-tRNA synthetase of <i>S. cerevisiae</i> reveals residues critical for its in vivo activities. <i>Current Genetics</i> , 1992, 22, 69-74. | 0.8 | 12 |
| 1818 | Complementation of <i>Saccharomyces cerevisiae</i> acid phosphatase mutation by a genomic sequence from the yeast <i>Yarrowia lipolytica</i> identifies a new phosphatase. <i>Current Genetics</i> , 1992, 22, 345-355. | 0.8 | 26 |

| # | ARTICLE | IF | CITATIONS |
|------|---|-----|-----------|
| 1819 | Identification of UAS elements and binding proteins necessary for derepression of <i>Saccharomyces cerevisiae</i> fructose-1,6-bisphosphatase. <i>Current Genetics</i> , 1992, 22, 363-370. | 0.8 | 40 |
| 1820 | Cloning, sequence analysis and yeast expression of the <i>egl1</i> gene from <i>Trichoderma longibrachiatum</i> . <i>Applied Microbiology and Biotechnology</i> , 1992, 38, 370-5. | 1.7 | 37 |
| 1821 | The activity of a model heterologous protein in <i>pep4-3</i> mutants of <i>Saccharomyces cerevisiae</i> . <i>Applied Microbiology and Biotechnology</i> , 1992, 36, 754-8. | 1.7 | 4 |
| 1822 | Enhanced expression of heterologous proteins by the use of a superinducible vector in budding yeast. <i>Applied Microbiology and Biotechnology</i> , 1992, 36, 655-8. | 1.7 | 15 |
| 1823 | Expression and secretion of pea-seed lipoxygenase isoenzymes in <i>Saccharomyces cerevisiae</i> . <i>Applied Microbiology and Biotechnology</i> , 1992, 37, 342-351. | 1.7 | 16 |
| 1824 | The complete sequence of a 10.8kb fragment to the right of the chromosome III centromere of <i>Saccharomyces cerevisiae</i> . <i>Yeast</i> , 1992, 8, 61-70. | 0.8 | 15 |
| 1825 | A rapid method for localized mutagenesis of yeast genes. <i>Yeast</i> , 1992, 8, 79-82. | 0.8 | 481 |
| 1826 | <i>IMP2</i> , a nuclear gene controlling the mitochondrial dependence of galactose, maltose and raffinose utilization in <i>Saccharomyces cerevisiae</i> . <i>Yeast</i> , 1992, 8, 83-93. | 0.8 | 26 |
| 1827 | Identification of a novel secreted glycoprotein of the yeast <i>Saccharomyces cerevisiae</i> stimulated by heat shock. <i>Yeast</i> , 1992, 8, 157-169. | 0.8 | 22 |
| 1828 | Molecular cloning of <i>CIF1</i> , a yeast gene necessary for growth on glucose. <i>Yeast</i> , 1992, 8, 183-192. | 0.8 | 114 |
| 1829 | The peroxisomal import signal of amine oxidase from the yeast <i>Hansenula polymorpha</i> is not universal. <i>Yeast</i> , 1992, 8, 243-252. | 0.8 | 18 |
| 1830 | Molecular cloning and analysis of autonomous replicating sequence of <i>Candida maltosa</i> . <i>Yeast</i> , 1992, 8, 253-259. | 0.8 | 18 |
| 1831 | Genetic and molecular analysis of <i>DNA43</i> and <i>DNA52</i> : Two new cell-cycle genes in <i>Saccharomyces cerevisiae</i> . <i>Yeast</i> , 1992, 8, 273-289. | 0.8 | 67 |
| 1832 | Isolation and sequence analysis of the gene encoding translation elongation factor 3 from <i>Candida albicans</i> . <i>Yeast</i> , 1992, 8, 337-352. | 0.8 | 47 |
| 1833 | N-linked glycosylation of proteinase B precursors of the yeast <i>Saccharomyces cerevisiae</i> is not require for proper targeting or processing of the enzyme. <i>Yeast</i> , 1992, 8, 353-359. | 0.8 | 3 |
| 1834 | Genetic analysis of maintenance and expression of L and M double-stranded RNAs from yeast killer virus K28. <i>Yeast</i> , 1992, 8, 373-384. | 0.8 | 23 |
| 1835 | Molecular analysis of yeast chromosome II between <i>CMD1</i> and <i>LYS2</i> : The excision repair gene <i>RAD16</i> located in this region belongs to a novel group of double-finger proteins. <i>Yeast</i> , 1992, 8, 397-408. | 0.8 | 26 |
| 1836 | Foreign gene expression in yeast: a review. <i>Yeast</i> , 1992, 8, 423-488. | 0.8 | 948 |

| # | ARTICLE | IF | CITATIONS |
|------|---|------|-----------|
| 1837 | Dosage-dependent translational suppression in yeast <i>Saccharomyces cerevisiae</i> . <i>Yeast</i> , 1992, 8, 489-499. | 0.8 | 63 |
| 1838 | Factors affecting homologous overexpression of the <i>Saccharomyces cerevisiae</i> Lanosterol 14 α -demethylase gene. <i>Yeast</i> , 1992, 8, 519-533. | 0.8 | 8 |
| 1839 | Isolation of new temperature-sensitive mutants of <i>Saccharomyces cerevisiae</i> deficient in mannose outer chain elongation. <i>Yeast</i> , 1992, 8, 535-547. | 0.8 | 79 |
| 1840 | SEC6 encodes an 85 kDa soluble protein required for exocytosis in yeast. <i>Yeast</i> , 1992, 8, 549-558. | 0.8 | 38 |
| 1841 | Mutational analysis of <i>Schizosaccharomyces pombe</i> U4 snRNA by plasmid exchange. <i>Yeast</i> , 1992, 8, 647-653. | 0.8 | 7 |
| 1842 | The telomere-associated MAL3 locus of <i>Saccharomyces</i> is a tandem array of repeated genes. <i>Yeast</i> , 1992, 8, 655-665. | 0.8 | 54 |
| 1843 | Efficient selection of phleomycin-resistant <i>Saccharomyces cerevisiae</i> transformants. <i>Yeast</i> , 1992, 8, 667-668. | 0.8 | 22 |
| 1844 | Efficient electropulse transformation of intact <i>Candida maltosa</i> cells by different homologous vector plasmids. <i>Yeast</i> , 1992, 8, 691-697. | 0.8 | 24 |
| 1845 | Disruption and mapping of <i>IDI1</i> , the gene for isopentenyl diphosphate isomerase in <i>Saccharomyces cerevisiae</i> . <i>Yeast</i> , 1992, 8, 743-748. | 0.8 | 35 |
| 1846 | II. Yeast sequencing reports. An 11.4 kb DNA segment on the left arm of yeast chromosome II carries the carboxypeptidase Y sorting gene <i>PEP1</i> , as well as <i>ACH1</i> , <i>FUS3</i> and a putative ars. <i>Yeast</i> , 1992, 8, 769-776. | 0.8 | 25 |
| 1847 | Structural and putative regulatory sequences of <i>Kluyveromyces</i> ribosomal protein genes. <i>Yeast</i> , 1992, 8, 903-922. | 0.8 | 24 |
| 1848 | Isolation and characterization of cDNA clones encoding <i>cdc2</i> homologues from <i>Oryza sativa</i> : a functional homologue and cognate variants. <i>Molecular Genetics and Genomics</i> , 1992, 233, 10-16. | 2.4 | 72 |
| 1849 | The <i>NAM8</i> gene in <i>Saccharomyces cerevisiae</i> encodes a protein with putative RNA binding motifs and acts as a suppressor of mitochondrial splicing deficiencies when overexpressed. <i>Molecular Genetics and Genomics</i> , 1992, 233, 136-144. | 2.4 | 20 |
| 1850 | Development of high cell density cultures of engineered <i>Saccharomyces cerevisiae</i> cells able to grow on lactose. <i>Biotechnology Letters</i> , 1992, 14, 1085-1088. | 1.1 | 17 |
| 1851 | Hyperglycosylation does not alter the properties of A bacterial cellulase secreted in yeast. <i>Biotechnology Letters</i> , 1992, 14, 409-414. | 1.1 | 2 |
| 1852 | Expression of glucoamylase gene using <i>SUC2</i> promoter in <i>Saccharomyces cerevisiae</i> . <i>Biotechnology Letters</i> , 1992, 14, 747-752. | 1.1 | 13 |
| 1853 | Growth and copper resistance of recombinant <i>Saccharomyces cerevisiae</i> containing a metallothionein gene. <i>Biotechnology Letters</i> , 1992, 14, 7-10. | 1.1 | 3 |
| 1854 | SEC65 gene product is a subunit of the yeast signal recognition particle required for its integrity. <i>Nature</i> , 1992, 356, 532-533. | 13.7 | 74 |

| # | ARTICLE | IF | CITATIONS |
|------|--|-----|-----------|
| 1855 | Functional expression of the rat brown adipose tissue uncoupling protein in <i>Saccharomyces cerevisiae</i> . <i>Molecular Microbiology</i> , 1992, 6, 363-370. | 1.2 | 37 |
| 1856 | Use of a cis-acting mutation to study the role of FLP-mediated recombination in the maintenance of native yeast 2 μ m plasmids. <i>Molecular Microbiology</i> , 1992, 6, 3101-3107. | 1.2 | 6 |
| 1857 | A leptomyacin B resistance gene of <i>Schizosaccharomyces pombe</i> encodes a protein similar to the mammalian P-glycoproteins. <i>Molecular Microbiology</i> , 1992, 6, 761-769. | 1.2 | 80 |
| 1858 | Fast high-performance liquid chromatographic purification of <i>Saccharomyces cerevisiae</i> phosphoenolpyruvate carboxykinase. <i>Journal of Chromatography A</i> , 1992, 625, 47-54. | 1.8 | 12 |
| 1859 | Expression of Japanese quail ovalbumin in <i>Saccharomyces cerevisiae</i> . <i>Folia Microbiologica</i> , 1992, 37, 273-278. | 1.1 | 1 |
| 1860 | Construction of a new <i>Escherichia coli</i> "Saccharomyces cerevisiae" shuttle plasmid cloning vector allowing positive selection for cloned fragments. <i>Folia Microbiologica</i> , 1992, 37, 193-198. | 1.1 | 4 |
| 1861 | Isolation and nucleotide sequence of <i>Hansenula anomala</i> URA3 gene encoding orotidine-5 β -phosphate decarboxylase. <i>Journal of Bioscience and Bioengineering</i> , 1992, 74, 352-357. | 0.9 | 10 |
| 1862 | Stabilization of a recombinant plasmid in yeast. <i>Journal of Bioscience and Bioengineering</i> , 1992, 74, 139-144. | 0.9 | 7 |
| 1863 | Cloning and characterization of SKT5, a <i>Saccharomyces cerevisiae</i> gene that affects protoplast regeneration and resistance to killer toxin of <i>Kluyveromyces lactis</i> . <i>Journal of Bioscience and Bioengineering</i> , 1992, 74, 199-208. | 0.9 | 18 |
| 1864 | The orotidine-5 β -phosphate decarboxylase gene (URA3) of a methylotrophic yeast, <i>Candida boidinii</i> : Nucleotide sequence and its expression in <i>Escherichia coli</i> . <i>Journal of Bioscience and Bioengineering</i> , 1992, 73, 255-260. | 0.9 | 10 |
| 1865 | Partial restoration of sporulation defect in sake yeasts, kyokai no. 7 and no. 9, by increased dosage of the IME1 gene. <i>Journal of Bioscience and Bioengineering</i> , 1992, 73, 265-270. | 0.9 | 41 |
| 1866 | Molecular breeding of a sake yeast with a mutated ARO4 gene which causes both resistance to o-fluoro-dl-phenylalanine and increased production of l ² -phenethyl alcohol. <i>Journal of Bioscience and Bioengineering</i> , 1992, 73, 366-369. | 0.9 | 19 |
| 1867 | QRI8, a novel ubiquitin-conjugating enzyme in <i>Saccharomyces cerevisiae</i> . <i>Biochimica Et Biophysica Acta Gene Regulatory Mechanisms</i> , 1992, 1132, 211-213. | 2.4 | 9 |
| 1868 | Characterization of recombinant human and rat pancreatic phospholipases A2 secreted from <i>Saccharomyces cerevisiae</i> : Difference in proteolytic processing. <i>Biochimica Et Biophysica Acta Gene Regulatory Mechanisms</i> , 1992, 1171, 1-10. | 2.4 | 5 |
| 1869 | Molecular cloning and expression of <i>Bacillus subtilis</i> bglS gene in <i>Saccharomyces cerevisiae</i> . <i>Current Microbiology</i> , 1992, 25, 279-282. | 1.0 | 3 |
| 1870 | Differential proteolytic sensitivity of yeast fatty acid synthetase subunits alpha and beta contributing to a balanced ratio of both fatty acid synthetase components. <i>FEBS Journal</i> , 1992, 203, 607-614. | 0.2 | 38 |
| 1871 | The ICL1 gene from <i>Saccharomyces cerevisiae</i> . <i>FEBS Journal</i> , 1992, 204, 983-990. | 0.2 | 92 |
| 1872 | The C-terminal region of subunit 4 (subunit b) is essential for assembly of the F0 portion of yeast mitochondrial ATP synthase. <i>FEBS Journal</i> , 1992, 205, 163-172. | 0.2 | 28 |

| # | ARTICLE | IF | CITATIONS |
|------|---|-----|-----------|
| 1873 | Uroporphyrinogen decarboxylase in <i>Saccharomyces cerevisiae</i> . HEM12 gene sequence and evidence for two conserved glycines essential for enzymatic activity. <i>FEBS Journal</i> , 1992, 205, 1011-1016. | 0.2 | 27 |
| 1874 | YmL9, a nucleus-encoded mitochondrial ribosomal protein of yeast, is homologous to L3 ribosomal proteins from all natural kingdoms and photosynthetic organelles. <i>FEBS Journal</i> , 1992, 206, 373-380. | 0.2 | 29 |
| 1875 | The polyanion-binding domain of cytoplasmic Lys-tRNA synthetase from <i>Saccharomyces cerevisiae</i> is not essential for cell viability. <i>FEBS Journal</i> , 1992, 207, 1-11. | 0.2 | 15 |
| 1876 | Autoactivation of proteinase A initiates activation of yeast vacuolar zymogens. <i>FEBS Journal</i> , 1992, 207, 277-283. | 0.2 | 68 |
| 1877 | Characterization of yeast-expressed beta-actins, site-specifically mutated at the tumor-related residue Gly245. <i>FEBS Journal</i> , 1992, 207, 315-320. | 0.2 | 17 |
| 1878 | Development of a mutagenesis, expression and purification system for yeast phosphoglycerate mutase. Investigation of the role of active-site His181. <i>FEBS Journal</i> , 1992, 207, 709-714. | 0.2 | 54 |
| 1879 | Expression and secretion of wheat germ agglutinin by <i>Saccharomyces cerevisiae</i> . <i>FEBS Journal</i> , 1992, 210, 989-997. | 0.2 | 20 |
| 1880 | Lactose/whey utilization and ethanol production by transformed <i>Saccharomyces cerevisiae</i> cells. <i>Biotechnology and Bioengineering</i> , 1992, 39, 799-805. | 1.7 | 61 |
| 1881 | Minimal extent of homology required for completion of meiotic recombination in <i>Saccharomyces cerevisiae</i> . <i>Genesis</i> , 1992, 13, 498-514. | 3.1 | 23 |
| 1882 | In vitro mutagenesis of the yeast <i>SUP4</i> gene to identify all substitutions that can be detected in vivo with the <i>SUP4</i> system. <i>Environmental and Molecular Mutagenesis</i> , 1992, 19, 282-287. | 0.9 | 24 |
| 1883 | Enhancement of cloned gene product synthesis via autoselection in recombinant <i>Saccharomyces cerevisiae</i> . <i>Biotechnology and Bioengineering</i> , 1993, 41, 801-810. | 1.7 | 21 |
| 1884 | Bioconversion of lactose/whey to fructose diphosphate with recombinant <i>saccharomyces cerevisiae</i> cells. <i>Biotechnology and Bioengineering</i> , 1993, 42, 398-400. | 1.7 | 25 |
| 1885 | Phorbol ester activation of functional rat protein kinase C β -1 causes phenotype in yeast. <i>Journal of Cellular Biochemistry</i> , 1993, 52, 320-329. | 1.2 | 20 |
| 1886 | The murine immune response to the male-specific antigen mouse testicular cytochrome c. <i>European Journal of Immunology</i> , 1993, 23, 1992-1998. | 1.6 | 0 |
| 1887 | Creation of an active estrogen-responsive element by a single base change in the flanking sequence of a cellular oncogene: A possible mechanism for hormonal carcinogenesis?. <i>Molecular Carcinogenesis</i> , 1993, 7, 76-82. | 1.3 | 10 |
| 1888 | Enhanced recovery and purification of <i>Aspergillus glucoamylase</i> from <i>Saccharomyces cerevisiae</i> by the addition of poly(aspartic acid) tails. <i>Enzyme and Microbial Technology</i> , 1993, 15, 593-600. | 1.6 | 22 |
| 1889 | Genetic transformation of industrial yeasts using an amino acid analog resistance gene as a directly selectable marker. <i>Enzyme and Microbial Technology</i> , 1993, 15, 874-876. | 1.6 | 13 |
| 1890 | Influence of culture conditions on the production of heterologous interleukin 1 β by <i>Kluyveromyces lactis</i> . <i>Biotechnology Letters</i> , 1993, 7, 609-614. | 0.5 | 8 |

| # | ARTICLE | IF | CITATIONS |
|------|---|-----|-----------|
| 1891 | Biologically active human and mouse nerve growth factors secreted by the yeast <i>Saccharomyces cerevisiae</i> . <i>Applied Microbiology and Biotechnology</i> , 1993, 38, 624-30. | 1.7 | 12 |
| 1892 | Co-expression of an <i>Erwinia chrysanthemi</i> pectate lyase-encoding gene (<i>pelE</i>) and an <i>E. carotovora</i> polygalacturonase-encoding gene (<i>peh1</i>) in <i>Saccharomyces cerevisiae</i> . <i>Applied Microbiology and Biotechnology</i> , 1993, 39, 181-8. | 1.7 | 29 |
| 1893 | Increased activity of a model heterologous protein in <i>Saccharomyces cerevisiae</i> strains with reduced vacuolar proteinases. <i>Applied Microbiology and Biotechnology</i> , 1993, 39, 211. | 1.7 | 4 |
| 1894 | A nucleotide substitution in one of the β -tubulin genes of <i>Trichoderma viride</i> confers resistance to the antimitotic drug methyl benzimidazole-2-yl-carbamate. <i>Molecular Genetics and Genomics</i> , 1993, 240, 73-80. | 2.4 | 57 |
| 1895 | Potentially rapid walking in cellular regulatory networks using the gene-gene interference method in yeast. <i>Molecular Genetics and Genomics</i> , 1993, 240, 245-257. | 2.4 | 18 |
| 1896 | A gene, <i>SMP2</i> , involved in plasmid maintenance and respiration in <i>Saccharomyces cerevisiae</i> encodes a highly charged protein. <i>Molecular Genetics and Genomics</i> , 1993, 236-236, 283-288. | 2.4 | 58 |
| 1897 | The consensus sequence of <i>Kluyveromyces lactis</i> centromeres shows homology to functional centromeric DNA from <i>Saccharomyces cerevisiae</i> . <i>Molecular Genetics and Genomics</i> , 1993, 236-236, 355-362. | 2.4 | 29 |
| 1898 | Differential expression of two genes encoding isoforms of the ATPase involved in sodium efflux in <i>Saccharomyces cerevisiae</i> . <i>Molecular Genetics and Genomics</i> , 1993, 236-236, 363-368. | 2.4 | 198 |
| 1899 | A pair of putative protein kinase genes (<i>YPK1</i> and <i>YPK2</i>) is required for cell growth in <i>Saccharomyces cerevisiae</i> . <i>Molecular Genetics and Genomics</i> , 1993, 236-236, 443-447. | 2.4 | 66 |
| 1900 | Molecular structure and genetic regulation of <i>SFA</i> , a gene responsible for resistance to formaldehyde in <i>Saccharomyces cerevisiae</i> , and characterization of its protein product. <i>Molecular Genetics and Genomics</i> , 1993, 237, 351-358. | 2.4 | 91 |
| 1901 | A new yeast gene with a myosin-like heptad repeat structure. <i>Molecular Genetics and Genomics</i> , 1993, 237, 359-369. | 2.4 | 41 |
| 1902 | Molecular and genetic characterization of <i>SPT4</i> , a gene important for transcription initiation in <i>Saccharomyces cerevisiae</i> . <i>Molecular Genetics and Genomics</i> , 1993, 237, 449-459. | 2.4 | 60 |
| 1903 | Characterization of the <i>MKS1</i> gene, a new negative regulator of the Ras-cyclic AMP pathway in <i>Saccharomyces cerevisiae</i> . <i>Molecular Genetics and Genomics</i> , 1993, 238-238, 6-16. | 2.4 | 56 |
| 1904 | A specific host factor binds at a cis-acting transcriptionally silent locus required for stability control of yeast plasmid <i>pSR1</i> . <i>Molecular Genetics and Genomics</i> , 1993, 238-238, 120-128. | 2.4 | 1 |
| 1905 | Activity of the yeast MAP kinase homologue <i>Slr2</i> is critically required for cell integrity at 37°C. <i>Molecular Genetics and Genomics</i> , 1993, 241-241, 177-184. | 2.4 | 126 |
| 1906 | Transcription of the yeast mitochondrial genome requires cyclic AMP. <i>Molecular Genetics and Genomics</i> , 1993, 241-241, 213-224. | 2.4 | 19 |
| 1907 | <i>MSI3</i> , a multicopy suppressor of mutants hyperactivated in the RAS-CAMP pathway, encodes a novel HSP70 protein of <i>Saccharomyces cerevisiae</i> . <i>Molecular Genetics and Genomics</i> , 1993, 240, 323-332. | 2.4 | 45 |
| 1908 | The centromere and promoter factor 1, <i>CPF1</i> , of <i>Saccharomyces cerevisiae</i> modulates gene activity through a family of factors including <i>SPT21</i> , <i>RPD1</i> (<i>SIN3</i>), <i>RPD3</i> and <i>CCR4</i> . <i>Molecular Genetics and Genomics</i> , 1993, 240, 374-386. | 2.4 | 48 |

| # | ARTICLE | IF | CITATIONS |
|------|---|-----|-----------|
| 1909 | A general suppressor of RNA polymerase I, II and III mutations in <i>Saccharomyces cerevisiae</i> . <i>Molecular Genetics and Genomics</i> , 1993, 239, 169-176. | 2.4 | 94 |
| 1910 | Cloning and expression of the UGA4 gene coding for the inducible GABA-specific transport protein of <i>Saccharomyces cerevisiae</i> . <i>Molecular Genetics and Genomics</i> , 1993, 237-237, 17-25. | 2.4 | 104 |
| 1911 | A point mutation in the core subunit gene of yeast mitochondrial RNA polymerase is suppressed by a high level of specificity factor MTF1. <i>Molecular Genetics and Genomics</i> , 1993, 237-237, 49-57. | 2.4 | 17 |
| 1912 | Cloning of <i>Saccharomyces cerevisiae</i> STE5 as a suppressor of a Ste20 protein kinase mutant: structural and functional similarity of Ste5 to Far1. <i>Molecular Genetics and Genomics</i> , 1993, 241-241, 241-254. | 2.4 | 64 |
| 1913 | Functional expression of the transcriptional activator Opaque-2 of <i>Zea mays</i> in transformed yeast. <i>Molecular Genetics and Genomics</i> , 1993, 241-241, 319-326. | 2.4 | 38 |
| 1914 | An impaired RNA polymerase II activity in <i>Saccharomyces cerevisiae</i> causes cell-cycle inhibition at START. <i>Molecular Genetics and Genomics</i> , 1993, 241-241, 327-334. | 2.4 | 14 |
| 1915 | Mating of the fission yeast occurs independently of pmd1 + gene product, a structural homologue of budding yeast STE6 and mammalian P-glycoproteins. <i>Archives of Microbiology</i> , 1993, 160, 162-165. | 1.0 | 12 |
| 1916 | Isolation of the DLD gene of <i>Saccharomyces cerevisiae</i> encoding the mitochondrial enzyme D-lactate ferricytochrome c oxidoreductase. <i>Molecular Genetics and Genomics</i> , 1993, 238, 315-324. | 2.4 | 70 |
| 1917 | Expression of the <i>Saccharomyces cerevisiae</i> RAD50 gene during meiosis: steady-state transcript levels rise and fall while steady-state protein levels remain constant. <i>Molecular Genetics and Genomics</i> , 1993, 238, 390-400. | 2.4 | 19 |
| 1918 | A soybean coproporphyrinogen oxidase gene is highly expressed in root nodules. <i>Plant Molecular Biology</i> , 1993, 23, 35-43. | 2.0 | 62 |
| 1919 | Secretion and localization of invertase and inulinase in recombinant <i>Saccharomyces cerevisiae</i> . <i>Biotechnology Letters</i> , 1993, 15, 1049-1054. | 1.1 | 20 |
| 1920 | Mutations in the yeast fructose 1,6-biphosphatase structural gene affect expression of a fructose 1,6-biphosphatase-endoglucanase A hybrid protein. <i>Current Genetics</i> , 1993, 23, 370-372. | 0.8 | 1 |
| 1921 | A high copy number of yeast γ -glutamylcysteine synthetase suppresses a nuclear mutation affecting mitochondrial translation. <i>Current Genetics</i> , 1993, 23, 408-413. | 0.8 | 33 |
| 1922 | Donation of information to the unbroken chromosome in double-strand break repair. <i>Current Genetics</i> , 1993, 23, 414-422. | 0.8 | 16 |
| 1923 | A sequence-specific endonuclease, Endo.SceI, can efficiently induce gene conversion in yeast mitochondria lacking a major exonuclease. <i>Current Genetics</i> , 1993, 23, 537-541. | 0.8 | 10 |
| 1924 | Asd-homothallism of <i>Saccharomyces cerevisiae</i> : identification of asd1-1 as an allele of sir4 and detection of γ -specific suppressors of it. <i>Current Genetics</i> , 1993, 24, 1-6. | 0.8 | 3 |
| 1925 | Expression enhancement of the Tn5 neomycin-resistance gene by removal of upstream ATG sequences and its use for probing heterologous upstream activating sequences in yeast. <i>Current Genetics</i> , 1993, 24, 12-20. | 0.8 | 5 |
| 1926 | Expression of the <i>Klebsiella pneumoniae</i> pullulanase-encoding gene in <i>Saccharomyces cerevisiae</i> . <i>Current Genetics</i> , 1993, 24, 32-37. | 0.8 | 15 |

| # | ARTICLE | IF | CITATIONS |
|------|---|-----|-----------|
| 1927 | Normal mitochondrial structure and genome maintenance in yeast requires the dynamin-like product of the MGM1 gene. <i>Current Genetics</i> , 1993, 24, 141-148. | 0.8 | 135 |
| 1928 | High efficiency transformation of <i>Kluyveromyces marxianus</i> by a replicative plasmid. <i>Current Genetics</i> , 1993, 24, 181-183. | 0.8 | 28 |
| 1929 | Molecular cloning of the PEL1 gene of <i>Saccharomyces cerevisiae</i> that is essential for the viability of petite mutants. <i>Current Genetics</i> , 1993, 24, 307-312. | 0.8 | 53 |
| 1930 | Messenger RNA 3'-end formation of a DNA fragment from the human c-myc 3'-end region in <i>Saccharomyces cerevisiae</i> . <i>Current Genetics</i> , 1993, 23, 201-204. | 0.8 | 7 |
| 1931 | Cloning of the C-URA3 gene and construction of a triple auxotroph (his5, ade1, ura3) as a useful host for the genetic engineering of <i>Candida maltosa</i> . <i>Current Genetics</i> , 1993, 23, 205-210. | 0.8 | 31 |
| 1932 | Physical mapping of the MEL gene family in <i>Saccharomyces cerevisiae</i> . <i>Current Genetics</i> , 1993, 24, 461-464. | 0.8 | 26 |
| 1933 | Isolation of a mutant allele that deregulates the threonine biosynthesis in <i>Saccharomyces cerevisiae</i> . <i>Current Genetics</i> , 1993, 24, 465-471. | 0.8 | 24 |
| 1934 | Regulation of the ADE2 gene from <i>Saccharomyces cerevisiae</i> . <i>Current Genetics</i> , 1993, 24, 472-480. | 0.8 | 19 |
| 1935 | The <i>ogd1</i> and <i>kgd1</i> mutants lacking 2-oxoglutarate dehydrogenase activity in yeast are allelic and can be differentiated by the cloned amber suppressor. <i>Current Genetics</i> , 1993, 24, 377-381. | 0.8 | 8 |
| 1936 | Use of reporter genes for the isolation and characterisation of different classes of sporulation mutants in the yeast <i>Saccharomyces cerevisiae</i> . <i>Current Genetics</i> , 1993, 24, 451-454. | 0.8 | 3 |
| 1937 | Parameters affecting lithium acetate-mediated transformation of <i>Saccharomyces cerevisiae</i> and development of a rapid and simplified procedure. <i>Current Genetics</i> , 1993, 24, 455-459. | 0.8 | 136 |
| 1938 | Titration of replication activity by increasing ARS dosage in yeast plasmids. <i>Current Genetics</i> , 1993, 23, 141-147. | 0.8 | 3 |
| 1939 | Physical localization of the flocculation gene FLO1 on chromosome I of <i>Saccharomyces cerevisiae</i> . <i>Yeast</i> , 1993, 9, 1-10. | 0.8 | 70 |
| 1940 | Identification and genetic mapping of CHL genes controlling mitotic chromosome transmission in yeast. <i>Yeast</i> , 1993, 9, 11-19. | 0.8 | 58 |
| 1941 | The SCH9 protein kinase mRNA contains a long 5' leader with a small open reading frame. <i>Yeast</i> , 1993, 9, 21-32. | 0.8 | 12 |
| 1942 | A yeast antiviral protein, SKI8, shares a repeated amino acid sequence pattern with β -subunits of G proteins and several other proteins. <i>Yeast</i> , 1993, 9, 43-51. | 0.8 | 36 |
| 1943 | Human catalase is imported and assembled in peroxisomes of <i>Saccharomyces cerevisiae</i> . <i>Yeast</i> , 1993, 9, 59-69. | 0.8 | 4 |
| 1944 | The VPH2 gene encodes a 25 kDa protein required for activity of the yeast vacuolar H ⁺ -ATPase. <i>Yeast</i> , 1993, 9, 175-184. | 0.8 | 33 |

| # | ARTICLE | IF | CITATIONS |
|------|---|-----|-----------|
| 1945 | Reduced efficiency in the glycosylation of the first sequon of <i>Saccharomyces cerevisiae</i> exoglucanase leads to the synthesis and secretion of a new glycoform of the molecule. <i>Yeast</i> , 1993, 9, 221-234. | 0.8 | 14 |
| 1946 | YMC1, a yeast gene encoding a new putative mitochondrial carrier protein. <i>Yeast</i> , 1993, 9, 301-305. | 0.8 | 20 |
| 1947 | Secretion of mouse α -amylase from fission yeast <i>Schizosaccharomyces pombe</i> : Presence of chymostatin-sensitive protease activity in the culture medium. <i>Yeast</i> , 1993, 9, 379-387. | 0.8 | 33 |
| 1948 | Genetics of heat-curability of killer virus of yeast. <i>Yeast</i> , 1993, 9, 411-418. | 0.8 | 13 |
| 1949 | Three yeast genes, PIR1, PIR2 and PIR3, containing internal tandem repeats, are related to each other, and PIR1 and PIR2 are required for tolerance to heat shock. <i>Yeast</i> , 1993, 9, 481-494. | 0.8 | 110 |
| 1950 | Increased transformation levels in intact cells of <i>Saccharomyces cerevisiae</i> aculeacin A-resistant mutants. <i>Yeast</i> , 1993, 9, 523-526. | 0.8 | 6 |
| 1951 | Genes required for derepression of an extracellular glucoamylase gene, STA2, in the yeast <i>Saccharomyces</i> . <i>Yeast</i> , 1993, 9, 533-541. | 0.8 | 26 |
| 1952 | Alteration of cell population structure due to cell lysis in <i>Saccharomyces cerevisiae</i> cells overexpressing the GAL4 gene. <i>Yeast</i> , 1993, 9, 575-582. | 0.8 | 27 |
| 1953 | A new promoter-probe vector for <i>Saccharomyces cerevisiae</i> using fungal glucoamylase cDNA as the reporter gene. <i>Yeast</i> , 1993, 9, 599-605. | 0.8 | 7 |
| 1954 | Molecular analysis of HEM6 (HEM12) in <i>Saccharomyces cerevisiae</i> , the gene for uroporphyrinogen decarboxylase. <i>Yeast</i> , 1993, 9, 613-623. | 0.8 | 10 |
| 1955 | Molecular characterization of the SEC1 gene of <i>Saccharomyces cerevisiae</i> : Subcellular distribution of a protein required for yeast protein secretion. <i>Yeast</i> , 1993, 9, 703-713. | 0.8 | 9 |
| 1956 | Vectors for the inducible overexpression of glutathione S-transferase fusion proteins in yeast. <i>Yeast</i> , 1993, 9, 715-722. | 0.8 | 289 |
| 1957 | Vector YFRp1 allows transformant selection in <i>Saccharomyces cerevisiae</i> via resistance to formaldehyde. <i>Yeast</i> , 1993, 9, 783-785. | 0.8 | 8 |
| 1958 | Cloning and characterization of the SEC18 gene from <i>Candida albicans</i> . <i>Yeast</i> , 1993, 9, 875-887. | 0.8 | 25 |
| 1959 | Comparison of the biochemical and biological functions of tyrosine phosphatases from fission yeast, budding yeast and animal cells. <i>Yeast</i> , 1993, 9, 1039-1052. | 0.8 | 10 |
| 1960 | KTR2: A new member of the KRE2 mannosyltransferase gene family. <i>Yeast</i> , 1993, 9, 1057-1063. | 0.8 | 28 |
| 1961 | Identification of two divergently transcribed genes centromere-proximal to the ARG4 locus on chromosome VIII of <i>Saccharomyces cerevisiae</i> . <i>Yeast</i> , 1993, 9, 1111-1120. | 0.8 | 9 |
| 1962 | GLT2, a gene for mitochondrial glycerol 3-phosphate dehydrogenase of <i>Saccharomyces cerevisiae</i> . <i>Yeast</i> , 1993, 9, 1121-1130. | 0.8 | 136 |

| # | ARTICLE | IF | CITATIONS |
|------|--|------|-----------|
| 1963 | Transformation in the methylotrophic yeast <i>Pichia methanolica</i> utilizing homologous ADE1 and heterologous <i>Saccharomyces cerevisiae</i> ADE2 and LEU2 genes as genetic markers. <i>Yeast</i> , 1993, 9, 1189-1197. | 0.8 | 17 |
| 1964 | The 5-aminoimidazole ribonucleotide-carboxylase structural gene of the methylotrophic yeast <i>Pichia methanolica</i> : Cloning, sequencing and homology analysis. <i>Yeast</i> , 1993, 9, 1251-1258. | 0.8 | 14 |
| 1965 | The sequence of a 17.5 kb DNA fragment on the left arm of yeast chromosome XI identifies the protein kinase gene ELM1, the DNA primase gene PRI2, a new gene encoding a putative histone and seven new open reading frames. <i>Yeast</i> , 1993, 9, 1379-1384. | 0.8 | 11 |
| 1966 | Preparation and characterization of the E168Q site-directed mutant of yeast enolase 1. <i>Proteins: Structure, Function and Bioinformatics</i> , 1993, 17, 426-434. | 1.5 | 21 |
| 1967 | Characterization of two fluorescent tryptophans in recombinant human granulocyte-colony stimulating factor: Comparison of native sequence protein and tryptophan-deficient mutants. <i>The Protein Journal</i> , 1993, 12, 229-236. | 1.1 | 21 |
| 1968 | Cloning and expression of cytochrome P450 genes controlling flower colour. <i>Nature</i> , 1993, 366, 276-279. | 13.7 | 367 |
| 1969 | Suppression of a dominant G protein β -subunit mutation in yeast by G γ protein expression. <i>Molecular Microbiology</i> , 1993, 9, 813-821. | 1.2 | 6 |
| 1970 | Conversion of Starch to Ethanol in a Recombinant <i>Saccharomyces cerevisiae</i> Strain Expressing Rice α -Amylase from a Novel <i>Pichia pastoris</i> Alcohol Oxidase Promoter. <i>Nature Biotechnology</i> , 1993, 11, 606-610. | 9.4 | 12 |
| 1971 | Introduction and expression of the 400 kilobase precursor amyloid protein gene in transgenic mice. <i>Nature Genetics</i> , 1993, 5, 22-30. | 9.4 | 331 |
| 1972 | A mammalian guanine-nucleotide-releasing protein enhances function of yeast secretory protein Sec4. <i>Nature</i> , 1993, 361, 464-467. | 13.7 | 118 |
| 1973 | Interactions of three domains distinguishing the Ras-related GTP-binding proteins Ypt1 and Sec4. <i>Nature</i> , 1993, 362, 560-563. | 13.7 | 188 |
| 1974 | A gene encoding sn-glycerol 3-phosphate dehydrogenase (NAD ⁺) complements an osmosensitive mutant of <i>Saccharomyces cerevisiae</i> . <i>Molecular Microbiology</i> , 1993, 10, 1101-1111. | 1.2 | 186 |
| 1975 | The general amino acid control regulates MET4, which encodes a methionine-pathway-specific transcriptional activator of <i>Saccharomyces cerevisiae</i> . <i>Molecular Microbiology</i> , 1993, 7, 215-228. | 1.2 | 38 |
| 1976 | AUA1, a gene involved in ammonia regulation of amino acid transport in <i>Saccharomyces cerevisiae</i> . <i>Molecular Microbiology</i> , 1993, 8, 167-178. | 1.2 | 14 |
| 1977 | Identification of a new nuclear gene (CEM1) encoding a protein homologous to a β -keto-acyl synthase which is essential for mitochondrial respiration in <i>Saccharomyces cerevisiae</i> . <i>Molecular Microbiology</i> , 1993, 9, 545-555. | 1.2 | 64 |
| 1978 | The <i>fdp1</i> and <i>cif1</i> mutations are caused by different single nucleotide changes in the yeast <i>CIF1</i> gene. <i>FEMS Microbiology Letters</i> , 1993, 107, 251-253. | 0.7 | 7 |
| 1979 | Expression of <i>Aspergillus oryzae</i> α -amylase gene in <i>Saccharomyces cerevisiae</i> . <i>FEMS Microbiology Letters</i> , 1993, 112, 119-124. | 0.7 | 20 |
| 1980 | Purification and characterization of recombinant human beta1-4 galactosyltransferase expressed in <i>Saccharomyces cerevisiae</i> . <i>FEBS Journal</i> , 1993, 212, 113-120. | 0.2 | 37 |

| # | ARTICLE | IF | CITATIONS |
|------|---|-----|-----------|
| 1981 | Fructose-1,6-bisphosphatase of the yeast <i>Kluyveromyces lactis</i> . <i>FEBS Journal</i> , 1993, 212, 193-199. | 0.2 | 15 |
| 1982 | A <i>Candida albicans</i> homolog of CDC25 is functional in <i>Saccharomyces cerevisiae</i> . <i>FEBS Journal</i> , 1993, 213, 195-204. | 0.2 | 14 |
| 1983 | Transcriptional control of AAC3 gene encoding mitochondrial ADP/ATP translocator in <i>Saccharomyces cerevisiae</i> by oxygen, heme and ROX1 factor. <i>FEBS Journal</i> , 1993, 213, 547-553. | 0.2 | 70 |
| 1984 | Natural cycloheximide resistance in yeast. The role of ribosomal protein L41. <i>FEBS Journal</i> , 1993, 213, 841-848. | 0.2 | 35 |
| 1985 | In vitro formation of a photoreversible adduct of phycocyanobilin and tobacco apophytochrome B. <i>FEBS Journal</i> , 1993, 215, 587-594. | 0.2 | 70 |
| 1986 | Both isoforms of protein phosphatase Z are essential for the maintenance of cell size and integrity in <i>Saccharomyces cerevisiae</i> in response to osmotic stress. <i>FEBS Journal</i> , 1993, 216, 269-279. | 0.2 | 58 |
| 1987 | Structure and metabolic control of the <i>Yarrowia lipolytica</i> peroxisomal 3-oxoacyl-CoA-thiolase gene. <i>FEBS Journal</i> , 1993, 216, 607-613. | 0.2 | 36 |
| 1988 | A note on the primary structure and expression of an <i>Erwinia carotovora</i> polygalacturonase-encoding gene (<i>peh1</i>) in <i>Escherichia coli</i> and <i>Saccharomyces cerevisiae</i> . <i>Journal of Applied Bacteriology</i> , 1993, 75, 149-158. | 1.1 | 11 |
| 1989 | Pleiotropic effects of heterozygosity at the mating-type locus of the yeast <i>Saccharomyces cerevisiae</i> on repair, recombination and transformation. <i>Mutation Research - Fundamental and Molecular Mechanisms of Mutagenesis</i> , 1993, 290, 239-247. | 0.4 | 11 |
| 1990 | Use of the PDR4 gene as a dominant selective marker in combination with cerulenin for prototrophic strains in <i>Saccharomyces cerevisiae</i> . <i>Journal of Bioscience and Bioengineering</i> , 1993, 76, 60-63. | 0.9 | 15 |
| 1991 | Cloning and phenotypic characterization of a gene enhancing resistance against oxidative stress in <i>saccharomyces cerevisiae</i> . <i>Journal of Bioscience and Bioengineering</i> , 1993, 75, 327-331. | 0.9 | 17 |
| 1992 | Cloning and nucleotide sequence of the urea amidolyase gene from <i>Candida utilis</i> . <i>Journal of Bioscience and Bioengineering</i> , 1993, 75, 245-253. | 0.9 | 8 |
| 1993 | Construction of xylose-assimilating <i>Saccharomyces cerevisiae</i> . <i>Journal of Bioscience and Bioengineering</i> , 1993, 75, 83-88. | 0.9 | 154 |
| 1994 | Molecular cloning of guinea pig CYP1A1: Complete primary structure and fast mobility of expressed protein on electrophoresis. <i>Biochimica Et Biophysica Acta Gene Regulatory Mechanisms</i> , 1993, 1216, 237-244. | 2.4 | 8 |
| 1995 | Rat liver flavin-containing monooxygenase (FMO): cDNA cloning and expression in yeast. <i>Biochimica Et Biophysica Acta Gene Regulatory Mechanisms</i> , 1993, 1173, 165-171. | 2.4 | 44 |
| 1996 | Changes in ion fluxes and the energy demand during spore development in <i>Phytophthora infestans</i> zoospores. <i>Folia Microbiologica</i> , 1993, 38, 193-200. | 1.1 | 5 |
| 1997 | Substitution of Gly-224 residue to ile in yeast alcohol dehydrogenase and enzyme reaction mechanism. <i>Archives of Pharmacal Research</i> , 1993, 16, 231-236. | 2.7 | 1 |
| 1998 | Identification of a protein from <i>Saccharomyces cerevisiae</i> with E2F-like DNA-binding and transactivating properties. <i>FEBS Letters</i> , 1993, 321, 153-158. | 1.3 | 7 |

| # | ARTICLE | IF | CITATIONS |
|------|--|-----|-----------|
| 1999 | Cloning, sequence and expression of the gene encoding the malolactic enzyme from <i>Lactococcus lactis</i> . <i>FEBS Letters</i> , 1993, 332, 74-80. | 1.3 | 61 |
| 2000 | Site-directed mutagenesis of yeast phosphoglycerate kinase. <i>FEBS Letters</i> , 1993, 320, 193-197. | 1.3 | 6 |
| 2001 | The <i>pep2</i> gene of <i>Saccharomyces cerevisiae</i> encodes a phosphodiesterase which protects the cell from extracellular cAMP. <i>FEBS Letters</i> , 1993, 325, 191-195. | 1.3 | 57 |
| 2002 | Regulation of Cu,Zn- and Mn-superoxide dismutase transcription in <i>Saccharomyces cerevisiae</i> . <i>FEBS Letters</i> , 1993, 315, 197-200. | 1.3 | 44 |
| 2003 | Evolutionary relationships between yeast and bacterial homoserine dehydrogenases. <i>FEBS Letters</i> , 1993, 323, 289-293. | 1.3 | 19 |
| 2004 | Centromeres of the fission yeast <i>Schizosaccharomyces pombe</i> are highly variable genetic loci. <i>Molecular and Cellular Biology</i> , 1993, 13, 4578-4587. | 1.1 | 94 |
| 2005 | Gene SNQ2 of <i>Saccharomyces cerevisiae</i> , which confers resistance to 4-nitroquinoline-N-oxide and other chemicals, encodes a 169 kDa protein homologous to ATP-dependent permeases. <i>Molecular Genetics and Genomics</i> , 1993, 236-236, 214-218. | 2.4 | 192 |
| 2006 | Biolistic transformation of <i>Trichoderma harzianum</i> and <i>Gliocladium virens</i> using plasmid and genomic DNA. <i>Current Genetics</i> , 1993, 24, 349-356. | 0.8 | 90 |
| 2007 | A <i>Saccharomyces cerevisiae</i> upstream activating sequence mediates induction of peroxisome proliferation by fatty acids. <i>Gene</i> , 1993, 132, 49-55. | 1.0 | 82 |
| 2008 | Isolation and nucleotide sequences of the genes encoding killer toxins from <i>Hansenula mrakii</i> and <i>H. saturnus</i> . <i>Gene</i> , 1993, 137, 265-270. | 1.0 | 51 |
| 2009 | Effect of a <i>pmr1</i> disruption and different signal sequences on the intracellular processing and secretion of <i>Cyamopsis tetragonoloba</i> β -galactosidase by <i>Saccharomyces cerevisiae</i> . <i>Gene</i> , 1993, 125, 115-123. | 1.0 | 49 |
| 2010 | Two functional soybean genes encoding p34cdc2 protein kinases are regulated by different plant developmental pathways. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 1993, 90, 943-947. | 3.3 | 103 |
| 2011 | Cloning, sequencing, and heterologous expression of a cellulase-encoding cDNA (<i>cbhl</i>) from <i>Penicillium janthinellum</i> . <i>Gene</i> , 1993, 124, 57-65. | 1.0 | 35 |
| 2012 | A random mutagenesis procedure: application to the <i>POL3</i> gene of <i>Saccharomyces cerevisiae</i> . <i>Gene</i> , 1993, 127, 139-144. | 1.0 | 12 |
| 2013 | Bcl-2 inhibition of neural death: decreased generation of reactive oxygen species. <i>Science</i> , 1993, 262, 1274-1277. | 6.0 | 1,670 |
| 2014 | A protein containing conserved RNA-recognition motifs is associated with ribosomal subunits in <i>Saccharomyces cerevisiae</i> . <i>Nucleic Acids Research</i> , 1993, 21, 3211-3216. | 6.5 | 11 |
| 2015 | Kinetic analysis of NAD ⁺ -isocitrate dehydrogenase with altered isocitrate binding sites: Contribution of IDH1 and IDH2 subunits to regulation and catalysis. <i>Biochemistry</i> , 1993, 32, 9323-9328. | 1.2 | 72 |
| 2016 | Chicken progesterone receptor expressed in <i>Saccharomyces cerevisiae</i> is correctly phosphorylated at all four Ser-Pro phosphorylation sites. <i>Biochemistry</i> , 1993, 32, 9563-9569. | 1.2 | 25 |

| # | ARTICLE | IF | CITATIONS |
|------|---|------|-----------|
| 2017 | Destabilizing effects of replacing a surface lysine of cytochrome c with aromatic amino acids: implications for the denatured state. <i>Biochemistry</i> , 1993, 32, 183-190. | 1.2 | 114 |
| 2018 | Multiple tandem integrations of transforming DNA sequences in yeast chromosomes suggest a mechanism for integrative transformation by homologous recombination. <i>Gene</i> , 1993, 134, 41-50. | 1.0 | 39 |
| 2019 | Isolation and characterization of SSE1 and SSE2, new members of the yeast HSP70 multigene family. <i>Gene</i> , 1993, 132, 57-66. | 1.0 | 115 |
| 2020 | Reconstitution of protein kinase C alpha function by the protein kinase C beta-I carboxy terminus. <i>Molecular and Cellular Endocrinology</i> , 1993, 98, 9-16. | 1.6 | 5 |
| 2021 | Cloning of the dihydroxyacid dehydratase-encoding gene (ILV3) from <i>Saccharomyces cerevisiae</i> . <i>Gene</i> , 1993, 137, 179-185. | 1.0 | 29 |
| 2022 | Isolation, overexpression and disruption of a <i>Saccharomyces cerevisiae</i> YNK gene encoding nucleoside diphosphate kinase. <i>Gene</i> , 1993, 129, 141-146. | 1.0 | 57 |
| 2023 | Identification and genetic analysis of <i>Schizosaccharomyces pombe</i> cDNAs that suppress deletion of IRA1 in <i>Saccharomyces cerevisiae</i> . <i>Gene</i> , 1993, 129, 147-152. | 1.0 | 4 |
| 2024 | <i>Saccharomyces</i> telomeres acquire single-strand TG1â€³ tails late in S phase. <i>Cell</i> , 1993, 72, 51-60. | 13.5 | 392 |
| 2025 | Genetic evidence that an activation domain of GAL4 does not require acidity and may form a β sheet. <i>Cell</i> , 1993, 72, 575-585. | 13.5 | 160 |
| 2026 | Sct1 functions in partnership with Cdc10 in a transcription complex that activates cell cycle START and inhibits differentiation. <i>Cell</i> , 1993, 72, 607-619. | 13.5 | 126 |
| 2027 | ZIP1 is a synaptonemal complex protein required for meiotic chromosome synapsis. <i>Cell</i> , 1993, 72, 365-378. | 13.5 | 625 |
| 2028 | Temperature-sensitive mutations demonstrate roles for yeast fibrillar in pre-rRNA processing, pre-rRNA methylation, and ribosome assembly. <i>Cell</i> , 1993, 72, 443-457. | 13.5 | 482 |
| 2029 | Target of rapamycin in yeast, TOR2, is an essential phosphatidylinositol kinase homolog required for G1 progression. <i>Cell</i> , 1993, 73, 585-596. | 13.5 | 819 |
| 2030 | Identification of revertants for the cystic fibrosis Δ F508 mutation using STE6-CFTR chimeras in yeast. <i>Cell</i> , 1993, 73, 335-346. | 13.5 | 183 |
| 2031 | Mutations in U1 snRNA bypass the requirement for a cell type-specific RNA splicing factor. <i>Cell</i> , 1993, 73, 407-415. | 13.5 | 56 |
| 2032 | FAR1 links the signal transduction pathway to the cell cycle machinery in yeast. <i>Cell</i> , 1993, 73, 747-760. | 13.5 | 359 |
| 2033 | Identification of essential components of the <i>S. cerevisiae</i> kinetochore. <i>Cell</i> , 1993, 73, 761-774. | 13.5 | 215 |
| 2034 | Δ splice site recognition in <i>S. cerevisiae</i> does not require base pairing with U1 snRNA. <i>Cell</i> , 1993, 73, 803-812. | 13.5 | 50 |

| # | ARTICLE | IF | CITATIONS |
|------|--|------|-----------|
| 2035 | A novel programmed frameshift expresses the POL3 gene of retrotransposon Ty3 of yeast: Frameshifting without tRNA slippage. <i>Cell</i> , 1993, 74, 93-103. | 13.5 | 161 |
| 2036 | A mechanism to enhance mRNA splicing fidelity: The RNA-dependent ATPase Prp16 governs usage of a discard pathway for aberrant lariat intermediates. <i>Cell</i> , 1993, 73, 1377-1391. | 13.5 | 180 |
| 2037 | Fission yeast cut5+, required for S phase onset and M phase restraint, is identical to the radiation-damage repair gene rad4+. <i>Cell</i> , 1993, 74, 383-393. | 13.5 | 241 |
| 2038 | The essential function of yeast protein disulfide isomerase does not reside in its isomerase activity. <i>Cell</i> , 1993, 74, 899-908. | 13.5 | 221 |
| 2039 | A transmembrane protein with a cdc2+CDC28-related kinase activity is required for signaling from the ER to the nucleus. <i>Cell</i> , 1993, 74, 743-756. | 13.5 | 759 |
| 2040 | Splice site choice and splicing efficiency are positively influenced by pre-mRNA intramolecular base pairing in yeast. <i>Cell</i> , 1993, 72, 893-901. | 13.5 | 100 |
| 2041 | RNA-dependent replication, transcription, and persistence of brome mosaic virus RNA replicons in <i>S. cerevisiae</i> . <i>Cell</i> , 1993, 72, 961-970. | 13.5 | 222 |
| 2042 | Transcriptional induction of genes encoding endoplasmic reticulum resident proteins requires a transmembrane protein kinase. <i>Cell</i> , 1993, 73, 1197-1206. | 13.5 | 1,101 |
| 2043 | Mxi1, a protein that specifically interacts with Max to bind Myc-Max recognition sites. <i>Cell</i> , 1993, 72, 223-232. | 13.5 | 792 |
| 2044 | Specific dephosphorylation of phosphopeptides by the yeast alkaline phosphatase encoded by PHO8 gene. <i>Biochimica Et Biophysica Acta - Molecular Cell Research</i> , 1993, 1177, 221-228. | 1.9 | 12 |
| 2045 | Phenotypic effects in <i>Saccharomyces cerevisiae</i> after regulated expression of Î²-(1,3)-glucanase from <i>Nicotiana plumbaginifolia</i> . <i>Journal of Biotechnology</i> , 1993, 27, 295-305. | 1.9 | 5 |
| 2046 | The putative phosphoinositide-specific phospholipase C gene, PLC1, of the yeast <i>Saccharomyces cerevisiae</i> is important for cell growth.. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 1993, 90, 1804-1808. | 3.3 | 164 |
| 2047 | In vivo evidence for endogenous DNA alkylation damage as a source of spontaneous mutation in eukaryotic cells.. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 1993, 90, 2117-2121. | 3.3 | 162 |
| 2048 | Membrane protein retention in the yeast Golgi apparatus: dipeptidyl aminopeptidase A is retained by a cytoplasmic signal containing aromatic residues.. <i>Journal of Cell Biology</i> , 1993, 121, 1197-1209. | 2.3 | 144 |
| 2049 | Multiple regions of NSR1 are sufficient for accumulation of a fusion protein within the nucleolus.. <i>Journal of Cell Biology</i> , 1993, 123, 1081-1091. | 2.3 | 63 |
| 2050 | Oxygen toxicity in a polyamine-depleted spe2 delta mutant of <i>Saccharomyces cerevisiae</i> .. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 1993, 90, 4693-4697. | 3.3 | 57 |
| 2051 | mRNA destabilization triggered by premature translational termination depends on at least three cis-acting sequence elements and one trans-acting factor.. <i>Genes and Development</i> , 1993, 7, 1737-1754. | 2.7 | 275 |
| 2052 | An integrated approach for identifying and mapping human genes.. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 1993, 90, 4364-4368. | 3.3 | 12 |

| # | ARTICLE | IF | CITATIONS |
|------|---|-----|-----------|
| 2053 | Futile cycles in <i>Saccharomyces cerevisiae</i> strains expressing the gluconeogenic enzymes during growth on glucose.. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 1993, 90, 1290-1294. | 3.3 | 65 |
| 2054 | NUP2, a novel yeast nucleoporin, has functional overlap with other proteins of the nuclear pore complex.. <i>Molecular Biology of the Cell</i> , 1993, 4, 209-222. | 0.9 | 120 |
| 2055 | Four yeast spliceosomal proteins (PRP5, PRP9, PRP11, and PRP21) interact to promote U2 snRNP binding to pre-mRNA.. <i>Genes and Development</i> , 1993, 7, 1909-1925. | 2.7 | 134 |
| 2056 | FUS3 phosphorylates multiple components of the mating signal transduction cascade: evidence for STE12 and FAR1.. <i>Molecular Biology of the Cell</i> , 1993, 4, 495-510. | 0.9 | 268 |
| 2057 | GAL4- Δ and GAL4- Δ activate transcription by different mechanisms. <i>Nucleic Acids Research</i> , 1993, 21, 2157-2163. | 6.5 | 15 |
| 2058 | Concerted action of the transcriptional activators REB1, RAP1, and GCR1 in the high-level expression of the glycolytic gene TPI.. <i>Molecular and Cellular Biology</i> , 1993, 13, 543-550. | 1.1 | 95 |
| 2059 | The effect of posttranslational modifications on the interaction of Ras2 with adenylyl cyclase. <i>Science</i> , 1993, 259, 683-686. | 6.0 | 147 |
| 2060 | Nucleosome disruption by transcription factor binding in yeast. <i>Science</i> , 1993, 262, 1563-1566. | 6.0 | 84 |
| 2061 | lxr1, a yeast protein that binds to platinated DNA and confers sensitivity to cisplatin. <i>Science</i> , 1993, 261, 603-605. | 6.0 | 188 |
| 2062 | The promoter region of the yeast KAR2 (BiP) gene contains a regulatory domain that responds to the presence of unfolded proteins in the endoplasmic reticulum.. <i>Molecular and Cellular Biology</i> , 1993, 13, 877-890. | 1.1 | 326 |
| 2063 | Gene RRN4 in <i>Saccharomyces cerevisiae</i> encodes the A12.2 subunit of RNA polymerase I and is essential only at high temperatures.. <i>Molecular and Cellular Biology</i> , 1993, 13, 114-122. | 1.1 | 112 |
| 2064 | [35] Transformation of intact yeast cells by electroporation. <i>Methods in Enzymology</i> , 1993, 217, 478-483. | 0.4 | 12 |
| 2065 | Mutations in a protein tyrosine phosphatase gene (PTP2) and a protein serine/threonine phosphatase gene (PTC1) cause a synthetic growth defect in <i>Saccharomyces cerevisiae</i> .. <i>Molecular and Cellular Biology</i> , 1993, 13, 5408-5417. | 1.1 | 166 |
| 2066 | Involvement of U6 snRNA in 5' splice site selection. <i>Science</i> , 1993, 262, 2035-2039. | 6.0 | 264 |
| 2067 | Mutation Analysis of <i>Saccharomyces cerevisiae</i> CDC6 Promoter: Defining Its UAS Domain and Cell Cycle Regulating Element. <i>DNA and Cell Biology</i> , 1993, 12, 363-370. | 0.9 | 15 |
| 2068 | GAL4 disrupts a repressing nucleosome during activation of GAL1 transcription in vivo.. <i>Genes and Development</i> , 1993, 7, 857-869. | 2.7 | 109 |
| 2069 | Two independent peroxisomal targeting signals in catalase A of <i>Saccharomyces cerevisiae</i> .. <i>Journal of Cell Biology</i> , 1993, 120, 665-673. | 2.3 | 178 |
| 2070 | Identification and functional analysis of a novel yeast small nucleolar RNA. <i>Nucleic Acids Research</i> , 1993, 21, 5386-5390. | 6.5 | 14 |

| # | ARTICLE | IF | CITATIONS |
|------|---|-----|-----------|
| 2071 | The extremely conserved amino terminus of RAD6 ubiquitin-conjugating enzyme is essential for amino-end rule-dependent protein degradation.. <i>Genes and Development</i> , 1993, 7, 250-261. | 2.7 | 81 |
| 2072 | The alpha and beta subunits of nematode actin capping protein function in yeast.. <i>Molecular Biology of the Cell</i> , 1993, 4, 907-917. | 0.9 | 30 |
| 2073 | Suppression of a sec63 mutation identifies a novel component of the yeast endoplasmic reticulum translocation apparatus.. <i>Molecular Biology of the Cell</i> , 1993, 4, 919-930. | 0.9 | 46 |
| 2074 | Patterns of mitochondrial sorting in yeast zygotes.. <i>Molecular Biology of the Cell</i> , 1993, 4, 21-36. | 0.9 | 76 |
| 2075 | Suppression of yeast geranylgeranyl transferase I defect by alternative prenylation of two target GTPases, Rho1p and Cdc42p.. <i>Molecular Biology of the Cell</i> , 1993, 4, 1017-1025. | 0.9 | 74 |
| 2076 | Functional homology of protein kinases required for sexual differentiation in <i>Schizosaccharomyces pombe</i> and <i>Saccharomyces cerevisiae</i> suggests a conserved signal transduction module in eukaryotic organisms.. <i>Molecular Biology of the Cell</i> , 1993, 4, 107-120. | 0.9 | 164 |
| 2077 | Recombinant human sickle hemoglobin expressed in yeast.. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 1993, 90, 918-922. | 3.3 | 47 |
| 2078 | Genetic interactions between KAR2 and SEC63, encoding eukaryotic homologues of DnaK and DnaJ in the endoplasmic reticulum.. <i>Molecular Biology of the Cell</i> , 1993, 4, 1145-1159. | 0.9 | 137 |
| 2079 | Centromere function on minichromosomes isolated from budding yeast.. <i>Molecular Biology of the Cell</i> , 1993, 4, 859-870. | 0.9 | 18 |
| 2080 | Negative regulation of mitosis by the fission yeast protein phosphatase ppa2.. <i>Genes and Development</i> , 1993, 7, 1059-1071. | 2.7 | 131 |
| 2081 | SAC1p is an integral membrane protein that influences the cellular requirement for phospholipid transfer protein function and inositol in yeast. <i>Journal of Cell Biology</i> , 1993, 122, 79-94. | 2.3 | 171 |
| 2082 | Dominant missense mutations in a novel yeast protein related to mammalian phosphatidylinositol 3-kinase and VPS34 abrogate rapamycin cytotoxicity.. <i>Molecular and Cellular Biology</i> , 1993, 13, 6012-6023. | 1.1 | 259 |
| 2083 | Selective protein degradation in the yeast exocytic pathway.. <i>Molecular Biology of the Cell</i> , 1993, 4, 729-736. | 0.9 | 37 |
| 2084 | Synthetic-lethal interactions identify two novel genes, SLA1 and SLA2, that control membrane cytoskeleton assembly in <i>Saccharomyces cerevisiae</i> . <i>Journal of Cell Biology</i> , 1993, 122, 635-644. | 2.3 | 290 |
| 2085 | Fission yeast genes <i>nda1+</i> and <i>nda4+</i> , mutations of which lead to S-phase block, chromatin alteration and Ca ²⁺ suppression, are members of the CDC46/MCM2 family.. <i>Molecular Biology of the Cell</i> , 1993, 4, 1003-1015. | 0.9 | 88 |
| 2086 | The a-factor transporter (STE6 gene product) and cell polarity in the yeast <i>Saccharomyces cerevisiae</i> .. <i>Journal of Cell Biology</i> , 1993, 120, 1203-1215. | 2.3 | 104 |
| 2087 | Molecular analysis of the PHO81 gene of <i>Saccharomyces cerevisiae</i> . <i>Nucleic Acids Research</i> , 1993, 21, 1975-1982. | 6.5 | 37 |
| 2088 | The carboxy-terminal tail of the homeo domain protein alpha 2 is required for function with a second homeo domain protein.. <i>Genes and Development</i> , 1993, 7, 1862-1870. | 2.7 | 68 |

| # | ARTICLE | IF | CITATIONS |
|------|--|-----|-----------|
| 2089 | Roles of multiple glucose transporters in <i>Saccharomyces cerevisiae</i> .. <i>Molecular and Cellular Biology</i> , 1993, 13, 638-648. | 1.1 | 146 |
| 2090 | A mitotic role for a novel fission yeast protein kinase <i>dsk1</i> with cell cycle stage dependent phosphorylation and localization.. <i>Molecular Biology of the Cell</i> , 1993, 4, 247-260. | 0.9 | 87 |
| 2091 | Isolation and characterization of a gene (<i>CBF2</i>) specifying a protein component of the budding yeast kinetochore.. <i>Journal of Cell Biology</i> , 1993, 121, 513-519. | 2.3 | 127 |
| 2092 | The hexokinase gene is required for transcriptional regulation of the glucose transporter gene <i>RAG1</i> in <i>Kluyveromyces lactis</i> .. <i>Molecular and Cellular Biology</i> , 1993, 13, 3882-3889. | 1.1 | 108 |
| 2093 | Clathrin facilitates the internalization of seven transmembrane segment receptors for mating pheromones in yeast.. <i>Journal of Cell Biology</i> , 1993, 123, 1707-1716. | 2.3 | 148 |
| 2094 | Purification and Characterization of the Biologically Active Human Truncated Macrophage Colony-Stimulating Factor Expressed in <i>Saccharomyces cerevisiae</i> . <i>Biological Chemistry Hoppe-Seyler</i> , 1993, 374, 903-908. | 1.4 | 1 |
| 2095 | Influence of the three nucleotides upstream of the initiation codon on expression of the <i>Escherichia coli lacZ</i> gene in <i>Saccharomyces Cerevisiae</i> . <i>Nucleic Acids Research</i> , 1993, 21, 4268-4271. | 6.5 | 19 |
| 2096 | Architecture of a yeast U6 RNA gene promoter.. <i>Molecular and Cellular Biology</i> , 1993, 13, 3015-3026. | 1.1 | 95 |
| 2097 | Biogenesis of the Yeast Vacuole (Lysosome). Signal Sequence Deletion of the Vacuolar Aspartic Proteinase <i>yscA</i> Does not Block Maturation of Vacuolar Proteinases. <i>Biological Chemistry Hoppe-Seyler</i> , 1993, 374, 1109-1116. | 1.4 | 5 |
| 2098 | Structural requirement of C-terminal region of chicken lysozyme signal peptide. <i>Animal Biotechnology</i> , 1993, 4, 113-120. | 0.7 | 2 |
| 2099 | The fission yeast <i>rad22</i> gene, having a function in matingtype switching and repair of DNA damages, encodes a protein homolog to <i>Rad52</i> of <i>Saccharomyces cerevisiae</i> . <i>Nucleic Acids Research</i> , 1993, 21, 5940-5944. | 6.5 | 142 |
| 2100 | The biochemical defects of <i>prp4-1</i> and <i>prp6-1</i> yeast splicing mutants reveal that the <i>PRP6</i> protein is required for the accumulation of the [U4/U6.U5] tri-snRNP. <i>Nucleic Acids Research</i> , 1993, 21, 1555-1562. | 6.5 | 50 |
| 2101 | A temperature-sensitive <i>NUP116</i> null mutant forms a nuclear envelope seal over the yeast nuclear pore complex thereby blocking nucleocytoplasmic traffic.. <i>Journal of Cell Biology</i> , 1993, 123, 275-284. | 2.3 | 203 |
| 2102 | Yeast actin with a mutation in the "hydrophobic plug" between subdomains 3 and 4 (L266D) displays a cold-sensitive polymerization defect.. <i>Journal of Cell Biology</i> , 1993, 123, 1185-1195. | 2.3 | 104 |
| 2103 | Regulatory circuit for responses of nitrogen catabolic gene expression to the <i>GLN3</i> and <i>DAL80</i> proteins and nitrogen catabolite repression in <i>Saccharomyces cerevisiae</i> . <i>Journal of Bacteriology</i> , 1993, 175, 64-73. | 1.0 | 144 |
| 2104 | [21] Expression and screening in yeast of genes mutagenized in vitro. <i>Methods in Enzymology</i> , 1993, 217, 301-312. | 0.4 | 0 |
| 2105 | [46] Experimental determination of rates of concerted evolution. <i>Methods in Enzymology</i> , 1993, 224, 631-646. | 0.4 | 19 |
| 2106 | Cofilin is an essential component of the yeast cortical cytoskeleton.. <i>Journal of Cell Biology</i> , 1993, 120, 421-435. | 2.3 | 245 |

| # | ARTICLE | IF | CITATIONS |
|------|---|-----|-----------|
| 2107 | Effect of heterologous expression of acyl-CoA-binding protein on acyl-CoA level and composition in yeast. <i>Biochemical Journal</i> , 1993, 290, 369-374. | 1.7 | 95 |
| 2108 | Expression of rat liver ketohexokinase in yeast results in fructose intolerance. <i>Biochemical Journal</i> , 1993, 291, 179-186. | 1.7 | 18 |
| 2109 | Demonstration by electrospray mass spectrometry that the peptidyl dipeptidase activity of cathepsin B is capable of rat cathepsin B C-terminal processing. <i>Biochemical Journal</i> , 1993, 294, 923-927. | 1.7 | 26 |
| 2110 | Cysteine biosynthesis in <i>Saccharomyces cerevisiae</i> occurs through the transsulfuration pathway which has been built up by enzyme recruitment. <i>Journal of Bacteriology</i> , 1993, 175, 5366-5374. | 1.0 | 68 |
| 2111 | Physiological analysis of mutants indicates involvement of the <i>Saccharomyces cerevisiae</i> GPI-anchored protein gp115 in morphogenesis and cell separation. <i>Journal of Bacteriology</i> , 1993, 175, 1879-1885. | 1.0 | 116 |
| 2112 | Yeast ribosomal protein L1 is required for the stability of newly synthesized 5S rRNA and the assembly of 60S ribosomal subunits.. <i>Molecular and Cellular Biology</i> , 1993, 13, 2835-2845. | 1.1 | 143 |
| 2113 | HSP78 encodes a yeast mitochondrial heat shock protein in the Clp family of ATP-dependent proteases.. <i>Molecular and Cellular Biology</i> , 1993, 13, 6304-6313. | 1.1 | 112 |
| 2114 | Interactions among the subunits of the G protein involved in <i>Saccharomyces cerevisiae</i> mating.. <i>Molecular and Cellular Biology</i> , 1993, 13, 1-8. | 1.1 | 76 |
| 2115 | A conserved alternative splice in the von Recklinghausen neurofibromatosis (NF1) gene produces two neurofibromin isoforms, both of which have GTPase-activating protein activity.. <i>Molecular and Cellular Biology</i> , 1993, 13, 487-495. | 1.1 | 157 |
| 2116 | Short artificial hairpins sequester splicing signals and inhibit yeast pre-mRNA splicing.. <i>Molecular and Cellular Biology</i> , 1993, 13, 6841-6848. | 1.1 | 73 |
| 2117 | DRS1 to DRS7, novel genes required for ribosome assembly and function in <i>Saccharomyces cerevisiae</i> .. <i>Molecular and Cellular Biology</i> , 1993, 13, 7901-7912. | 1.1 | 105 |
| 2118 | Antibody mimicking the action of RAS proteins on yeast adenylyl cyclase: implication for RAS-effector interaction.. <i>Molecular and Cellular Biology</i> , 1993, 13, 769-774. | 1.1 | 12 |
| 2119 | Pheromone-induced signal transduction in <i>Saccharomyces cerevisiae</i> requires the sequential function of three protein kinases.. <i>Molecular and Cellular Biology</i> , 1993, 13, 2069-2080. | 1.1 | 152 |
| 2120 | A complex regulatory element from the yeast gene ENO2 modulates GCR1-dependent transcriptional activation.. <i>Molecular and Cellular Biology</i> , 1993, 13, 2623-2633. | 1.1 | 35 |
| 2121 | GCN1, a translational activator of GCN4 in <i>Saccharomyces cerevisiae</i> , is required for phosphorylation of eukaryotic translation initiation factor 2 by protein kinase GCN2.. <i>Molecular and Cellular Biology</i> , 1993, 13, 3541-3556. | 1.1 | 117 |
| 2122 | Evidence that the SKI antiviral system of <i>Saccharomyces cerevisiae</i> acts by blocking expression of viral mRNA.. <i>Molecular and Cellular Biology</i> , 1993, 13, 4331-4341. | 1.1 | 118 |
| 2123 | Promoter elements determining weak expression of the GAL4 regulatory gene of <i>Saccharomyces cerevisiae</i> .. <i>Molecular and Cellular Biology</i> , 1993, 13, 4999-5009. | 1.1 | 42 |
| 2124 | A protein complex of translational regulators of GCN4 mRNA is the guanine nucleotide-exchange factor for translation initiation factor 2 in yeast.. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 1993, 90, 5350-5354. | 3.3 | 88 |

| # | ARTICLE | IF | CITATIONS |
|------|---|-----|-----------|
| 2125 | Mutations that alter the third cytoplasmic loop of the a-factor receptor lead to a constitutive and hypersensitive phenotype.. Proceedings of the National Academy of Sciences of the United States of America, 1993, 90, 9921-9925. | 3.3 | 61 |
| 2126 | The Arabidopsis endoplasmic reticulum retention receptor functions in yeast. Proceedings of the National Academy of Sciences of the United States of America, 1993, 90, 11433-11437. | 3.3 | 109 |
| 2127 | Essential role of the HMG domain in the function of yeast mitochondrial histone HM: functional complementation of HM by the nuclear nonhistone protein NHP6A.. Proceedings of the National Academy of Sciences of the United States of America, 1993, 90, 5598-5602. | 3.3 | 44 |
| 2128 | Cytoplasmic dynein is required for normal nuclear segregation in yeast. Proceedings of the National Academy of Sciences of the United States of America, 1993, 90, 11172-11176. | 3.3 | 375 |
| 2129 | Function of yeast cytoplasmic C1-tetrahydrofolate synthase.. Proceedings of the National Academy of Sciences of the United States of America, 1993, 90, 2636-2640. | 3.3 | 10 |
| 2130 | Yeast calmodulin and a conserved nuclear protein participate in the in vivo binding of a matrix association region.. Proceedings of the National Academy of Sciences of the United States of America, 1993, 90, 5623-5627. | 3.3 | 40 |
| 2131 | Characterization of HIR1 and HIR2, two genes required for regulation of histone gene transcription in Saccharomyces cerevisiae.. Molecular and Cellular Biology, 1993, 13, 28-38. | 1.1 | 137 |
| 2132 | The effect on chromosome stability of deleting replication origins.. Molecular and Cellular Biology, 1993, 13, 391-398. | 1.1 | 87 |
| 2133 | AFR1 acts in conjunction with the alpha-factor receptor to promote morphogenesis and adaptation.. Molecular and Cellular Biology, 1993, 13, 6876-6888. | 1.1 | 43 |
| 2134 | A common element involved in transcriptional regulation of two DNA alkylation repair genes (MAG) Tj ETQq1 1 0.784314 rgBT /Overl | 1.1 | 60 |
| 2135 | GSP1 and GSP2, genetic suppressors of the prp20-1 mutant in Saccharomyces cerevisiae: GTP-binding proteins involved in the maintenance of nuclear organization.. Molecular and Cellular Biology, 1993, 13, 2152-2161. | 1.1 | 155 |
| 2136 | A U5 small nuclear ribonucleoprotein particle protein involved only in the second step of pre-mRNA splicing in Saccharomyces cerevisiae.. Molecular and Cellular Biology, 1993, 13, 2959-2970. | 1.1 | 65 |
| 2137 | Location and characterization of autonomously replicating sequences from chromosome VI of Saccharomyces cerevisiae.. Molecular and Cellular Biology, 1993, 13, 5043-5056. | 1.1 | 107 |
| 2138 | Alteration of a yeast SH3 protein leads to conditional viability with defects in cytoskeletal and budding patterns.. Molecular and Cellular Biology, 1993, 13, 5070-5084. | 1.1 | 174 |
| 2139 | On the directional specificity of ribosome frameshifting at a "hungry" codon.. Proceedings of the National Academy of Sciences of the United States of America, 1993, 90, 5469-5473. | 3.3 | 60 |
| 2140 | Mutations in the alpha subunit of eukaryotic translation initiation factor 2 (eIF-2 alpha) that overcome the inhibitory effect of eIF-2 alpha phosphorylation on translation initiation.. Proceedings of the National Academy of Sciences of the United States of America, 1993, 90, 7215-7219. | 3.3 | 61 |
| 2141 | A multicopy suppressor gene of the Saccharomyces cerevisiae G1 cell cycle mutant gene dbf4 encodes a protein kinase and is identified as CDC5.. Molecular and Cellular Biology, 1993, 13, 4445-4457. | 1.1 | 261 |
| 2142 | SPT13 (GAL11) of Saccharomyces cerevisiae negatively regulates activity of the MCM1 transcription factor in Ty1 elements.. Molecular and Cellular Biology, 1993, 13, 63-71. | 1.1 | 43 |

| # | ARTICLE | IF | CITATIONS |
|------|--|-----|-----------|
| 2143 | p62cdc23 of <i>Saccharomyces cerevisiae</i> : a nuclear tetratricopeptide repeat protein with two mutable domains.. <i>Molecular and Cellular Biology</i> , 1993, 13, 1212-1221. | 1.1 | 69 |
| 2144 | Transcriptional repression in <i>Saccharomyces cerevisiae</i> by a SIN3-LexA fusion protein.. <i>Molecular and Cellular Biology</i> , 1993, 13, 1805-1814. | 1.1 | 117 |
| 2145 | The COT2 gene is required for glucose-dependent divalent cation transport in <i>Saccharomyces cerevisiae</i> .. <i>Molecular and Cellular Biology</i> , 1993, 13, 2041-2049. | 1.1 | 44 |
| 2146 | Interspersion of an unusual GCN4 activation site with a complex transcriptional repression site in Ty2 elements of <i>Saccharomyces cerevisiae</i> .. <i>Molecular and Cellular Biology</i> , 1993, 13, 2091-2103. | 1.1 | 18 |
| 2147 | MKK1 and MKK2, which encode <i>Saccharomyces cerevisiae</i> mitogen-activated protein kinase-kinase homologs, function in the pathway mediated by protein kinase C.. <i>Molecular and Cellular Biology</i> , 1993, 13, 3076-3083. | 1.1 | 327 |
| 2148 | Analysis of the function of the 70-kilodalton cyclase-associated protein (CAP) by using mutants of yeast adenylyl cyclase defective in CAP binding.. <i>Molecular and Cellular Biology</i> , 1993, 13, 4087-4097. | 1.1 | 37 |
| 2149 | Guanine nucleotide exchange factor for eukaryotic translation initiation factor 2 in <i>Saccharomyces cerevisiae</i> : interactions between the essential subunits GCD2, GCD6, and GCD7 and the regulatory subunit GCN3.. <i>Molecular and Cellular Biology</i> , 1993, 13, 4618-4631. | 1.1 | 56 |
| 2150 | Yeast phenotype classifies mammalian protein kinase C cDNA mutants.. <i>Molecular and Cellular Biology</i> , 1993, 13, 4728-4735. | 1.1 | 17 |
| 2151 | An essential yeast protein, CBF5p, binds in vitro to centromeres and microtubules.. <i>Molecular and Cellular Biology</i> , 1993, 13, 4884-4893. | 1.1 | 167 |
| 2152 | Translation of the yeast transcriptional activator GCN4 is stimulated by purine limitation: implications for activation of the protein kinase GCN2.. <i>Molecular and Cellular Biology</i> , 1993, 13, 5099-5111. | 1.1 | 123 |
| 2153 | Cloning of the STE5 gene of <i>Saccharomyces cerevisiae</i> as a suppressor of the mating defect of cdc25 temperature-sensitive mutants.. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 1993, 90, 5474-5478. | 3.3 | 30 |
| 2154 | A genetic method for defining DNA-binding domains: application to the nuclear receptor NGFI-B.. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 1993, 90, 9186-9190. | 3.3 | 30 |
| 2155 | Three downstream sites repress transcription of a Ty2 retrotransposon in <i>Saccharomyces cerevisiae</i> .. <i>Molecular and Cellular Biology</i> , 1993, 13, 2081-2090. | 1.1 | 28 |
| 2156 | Origin activation and formation of single-strand TG1-3 tails occur sequentially in late S phase on a yeast linear plasmid.. <i>Molecular and Cellular Biology</i> , 1993, 13, 4057-4065. | 1.1 | 91 |
| 2157 | Genetic evidence for in vivo cross-specificity of the CaaX-box protein prenyltransferases farnesyltransferase and geranylgeranyltransferase-I in <i>Saccharomyces cerevisiae</i> .. <i>Molecular and Cellular Biology</i> , 1993, 13, 4260-4275. | 1.1 | 124 |
| 2158 | The <i>Saccharomyces cerevisiae</i> PRP21 gene product is an integral component of the prespliceosome.. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 1993, 90, 6771-6775. | 3.3 | 43 |
| 2159 | Heat-shock protein hsp90 governs the activity of pp60v-src kinase.. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 1993, 90, 7074-7078. | 3.3 | 422 |
| 2160 | The yeast KRE9 gene encodes an O glycoprotein involved in cell surface beta-glucan assembly.. <i>Molecular and Cellular Biology</i> , 1993, 13, 6346-6356. | 1.1 | 91 |

| # | ARTICLE | IF | CITATIONS |
|------|---|-----|-----------|
| 2161 | The <i>Saccharomyces cerevisiae</i> Cdc68 transcription activator is antagonized by San1, a protein implicated in transcriptional silencing.. <i>Molecular and Cellular Biology</i> , 1993, 13, 7553-7565. | 1.1 | 39 |
| 2162 | The rRNA enhancer regulates rRNA transcription in <i>Saccharomyces cerevisiae</i> .. <i>Molecular and Cellular Biology</i> , 1993, 13, 1283-1289. | 1.1 | 37 |
| 2163 | Differential function and expression of <i>Saccharomyces cerevisiae</i> B-type cyclins in mitosis and meiosis.. <i>Molecular and Cellular Biology</i> , 1993, 13, 2113-2125. | 1.1 | 172 |
| 2164 | Functional domains of the yeast STE12 protein, a pheromone-responsive transcriptional activator.. <i>Molecular and Cellular Biology</i> , 1993, 13, 3765-3772. | 1.1 | 53 |
| 2165 | An essential yeast gene with homology to the exonuclease-encoding XRN1/KEM1 gene also encodes a protein with exoribonuclease activity.. <i>Molecular and Cellular Biology</i> , 1993, 13, 341-350. | 1.1 | 115 |
| 2166 | GCD11, a negative regulator of GCN4 expression, encodes the gamma subunit of eIF-2 in <i>Saccharomyces cerevisiae</i> .. <i>Molecular and Cellular Biology</i> , 1993, 13, 506-520. | 1.1 | 95 |
| 2167 | PUB1: a major yeast poly(A)+ RNA-binding protein.. <i>Molecular and Cellular Biology</i> , 1993, 13, 6114-6123. | 1.1 | 43 |
| 2168 | DNA topoisomerase I controls the kinetics of promoter activation and DNA topology in <i>Saccharomyces cerevisiae</i> .. <i>Molecular and Cellular Biology</i> , 1993, 13, 6702-6710. | 1.1 | 13 |
| 2169 | The SH2 domain is required for stable phosphorylation of p56lck at tyrosine 505, the negative regulatory site.. <i>Molecular and Cellular Biology</i> , 1993, 13, 7112-7121. | 1.1 | 51 |
| 2170 | A new transcriptional-activation motif restricted to a class of helix-loop-helix proteins is functionally conserved in both yeast and mammalian cells.. <i>Molecular and Cellular Biology</i> , 1993, 13, 792-800. | 1.1 | 134 |
| 2171 | Conservation of transcriptional activation functions of the NF-kappa B p50 and p65 subunits in mammalian cells and <i>Saccharomyces cerevisiae</i> .. <i>Molecular and Cellular Biology</i> , 1993, 13, 1666-1674. | 1.1 | 82 |
| 2172 | A 5'-3' exonuclease from <i>Saccharomyces cerevisiae</i> is required for in vitro recombination between linear DNA molecules with overlapping homology.. <i>Molecular and Cellular Biology</i> , 1993, 13, 3125-3134. | 1.1 | 41 |
| 2173 | Differential transcriptional activation by v-myb and c-myb in animal cells and <i>Saccharomyces cerevisiae</i> .. <i>Molecular and Cellular Biology</i> , 1993, 13, 4423-4431. | 1.1 | 36 |
| 2174 | NPK1, a tobacco gene that encodes a protein with a domain homologous to yeast BCK1, STE11, and Byr2 protein kinases.. <i>Molecular and Cellular Biology</i> , 1993, 13, 4745-4752. | 1.1 | 141 |
| 2175 | Expression of p60v-src in <i>Saccharomyces cerevisiae</i> results in elevation of p34CDC28 kinase activity and release of the dependence of DNA replication on mitosis.. <i>Molecular and Cellular Biology</i> , 1993, 13, 5112-5121. | 1.1 | 21 |
| 2176 | A yeast mitogen-activated protein kinase homolog (Mpk1p) mediates signalling by protein kinase C.. <i>Molecular and Cellular Biology</i> , 1993, 13, 3067-3075. | 1.1 | 385 |
| 2177 | Regulation of the urea active transporter gene (DUR3) in <i>Saccharomyces cerevisiae</i> . <i>Journal of Bacteriology</i> , 1993, 175, 4688-4698. | 1.0 | 81 |
| 2178 | A <i>Saccharomyces cerevisiae</i> UAS element controlled by protein kinase A activates transcription in response to a variety of stress conditions.. <i>EMBO Journal</i> , 1993, 12, 1997-2003. | 3.5 | 421 |

| # | ARTICLE | IF | CITATIONS |
|------|---|-----|-----------|
| 2179 | Analysis of feedback-resistant anthranilate synthases from <i>Saccharomyces cerevisiae</i> . <i>Journal of Bacteriology</i> , 1993, 175, 1061-1068. | 1.0 | 42 |
| 2180 | Suppressors of clathrin deficiency: overexpression of ubiquitin rescues lethal strains of clathrin-deficient <i>Saccharomyces cerevisiae</i> .. <i>Molecular and Cellular Biology</i> , 1993, 13, 521-532. | 1.1 | 58 |
| 2181 | Properties of <i>Saccharomyces cerevisiae</i> wee1 and its differential regulation of p34CDC28 in response to G1 and G2 cyclins.. <i>EMBO Journal</i> , 1993, 12, 3417-3426. | 3.5 | 311 |
| 2182 | A new pair of B-type cyclins from <i>Saccharomyces cerevisiae</i> that function early in the cell cycle.. <i>EMBO Journal</i> , 1993, 12, 3437-3447. | 3.5 | 87 |
| 2183 | A membrane-associated complex containing the Vps15 protein kinase and the Vps34 PI 3-kinase is essential for protein sorting to the yeast lysosome-like vacuole.. <i>EMBO Journal</i> , 1993, 12, 2195-2204. | 3.5 | 311 |
| 2184 | GCR1, a transcriptional activator in <i>Saccharomyces cerevisiae</i> , complexes with RAP1 and can function without its DNA binding domain.. <i>EMBO Journal</i> , 1993, 12, 2431-2437. | 3.5 | 88 |
| 2185 | Vps1p, a member of the dynamin GTPase family, is necessary for Golgi membrane protein retention in <i>Saccharomyces cerevisiae</i> .. <i>EMBO Journal</i> , 1993, 12, 3049-3059. | 3.5 | 122 |
| 2186 | Chromatin transitions during activation and repression of galactose-regulated genes in yeast.. <i>EMBO Journal</i> , 1993, 12, 4603-4613. | 3.5 | 98 |
| 2187 | Characterization of NGG1, a novel yeast gene required for glucose repression of GAL4p-regulated transcription.. <i>EMBO Journal</i> , 1993, 12, 5255-5265. | 3.5 | 51 |
| 2188 | Guanosine diphosphatase is required for protein and sphingolipid glycosylation in the Golgi lumen of <i>Saccharomyces cerevisiae</i> . <i>Journal of Cell Biology</i> , 1993, 122, 307-323. | 2.3 | 175 |
| 2189 | Identification and purification of a protein that binds DNA cooperatively with the yeast SWI5 protein.. <i>Molecular and Cellular Biology</i> , 1993, 13, 5524-5537. | 1.1 | 50 |
| 2190 | Far1 and Fus3 link the mating pheromone signal transduction pathway to three G1-phase Cdc28 kinase complexes.. <i>Molecular and Cellular Biology</i> , 1993, 13, 5659-5669. | 1.1 | 177 |
| 2191 | High-frequency transformation of a methylotrophic yeast, <i>Candida boidinii</i> , with autonomously replicating plasmids which are also functional in <i>Saccharomyces cerevisiae</i> . <i>Journal of Bacteriology</i> , 1993, 175, 3556-3562. | 1.0 | 35 |
| 2192 | SPL1-1, a <i>Saccharomyces cerevisiae</i> mutation affecting tRNA splicing. <i>Journal of Bacteriology</i> , 1993, 175, 1433-1442. | 1.0 | 53 |
| 2193 | The control in cis of the position and the amount of the ARG4 meiotic double-strand break of <i>Saccharomyces cerevisiae</i> .. <i>EMBO Journal</i> , 1993, 12, 1459-1466. | 3.5 | 70 |
| 2194 | The yeast SSS1 gene is essential for secretory protein translocation and encodes a conserved protein of the endoplasmic reticulum.. <i>EMBO Journal</i> , 1993, 12, 4083-4093. | 3.5 | 91 |
| 2195 | Phospholipid transfer activity is relevant to but not sufficient for the essential function of the yeast SEC14 gene product.. <i>EMBO Journal</i> , 1993, 12, 4775-4784. | 3.5 | 107 |
| 2196 | A growth rate-limiting process in the last growth phase of the yeast life cycle involves RPB4, a subunit of RNA polymerase II. <i>Journal of Bacteriology</i> , 1993, 175, 6358-6363. | 1.0 | 45 |

| # | ARTICLE | IF | CITATIONS |
|------|---|-----|-----------|
| 2197 | Evidence for an interaction between the CYP1(HAP1) activator and a cellular factor during heme-dependent transcriptional regulation in the yeast <i>Saccharomyces cerevisiae</i> .. EMBO Journal, 1993, 12, 1209-1218. | 3.5 | 55 |
| 2198 | A new yeast translation initiation factor suppresses a mutation in the eIF-4A RNA helicase.. EMBO Journal, 1993, 12, 4005-4011. | 3.5 | 86 |
| 2199 | <i>Drosophila</i> protein phosphatase V functionally complements a SIT4 mutant in <i>Saccharomyces cerevisiae</i> and its amino-terminal region can confer this complementation to a heterologous phosphatase catalytic domain.. EMBO Journal, 1993, 12, 4833-4842. | 3.5 | 44 |
| 2200 | The <i>Schizosaccharomyces pombe</i> <i>cwg2+</i> gene codes for the beta subunit of a geranylgeranyltransferase type I required for beta-glucan synthesis.. EMBO Journal, 1993, 12, 5245-5254. | 3.5 | 67 |
| 2201 | Cloning and bacterial expression of the CYS3 gene encoding cystathionine gamma-lyase of <i>Saccharomyces cerevisiae</i> and the physicochemical and enzymatic properties of the protein. Journal of Bacteriology, 1993, 175, 4800-4808. | 1.0 | 52 |
| 2202 | Isolation of a <i>Candida albicans</i> DNA sequence conferring adhesion and aggregation on <i>Saccharomyces cerevisiae</i> . Journal of Bacteriology, 1993, 175, 5683-5689. | 1.0 | 42 |
| 2203 | A DNA unwinding element and an ARS consensus comprise a replication origin within a yeast chromosome.. EMBO Journal, 1993, 12, 4521-4531. | 3.5 | 97 |
| 2204 | A pair of functionally redundant yeast genes (PPZ1 and PPZ2) encoding type 1-related protein phosphatases function within the PKC1-mediated pathway.. Molecular and Cellular Biology, 1993, 13, 5843-5853. | 1.1 | 141 |
| 2205 | SPK1 is an essential S-phase-specific gene of <i>Saccharomyces cerevisiae</i> that encodes a nuclear serine/threonine/tyrosine kinase.. Molecular and Cellular Biology, 1993, 13, 5829-5842. | 1.1 | 101 |
| 2206 | A <i>cdc</i> -like autolytic <i>Saccharomyces cerevisiae</i> mutant altered in budding site selection is complemented by SPO12, a sporulation gene. Journal of Bacteriology, 1993, 175, 6562-6570. | 1.0 | 16 |
| 2207 | The frequency and accuracy of replication past a thymine-thymine cyclobutane dimer are very different in <i>Saccharomyces cerevisiae</i> and <i>Escherichia coli</i> . Journal of Bacteriology, 1993, 175, 2607-2612. | 1.0 | 67 |
| 2208 | Inactivation of YME1, a member of the ftsH-SEC18-PAS1-CDC48 family of putative ATPase-encoding genes, causes increased escape of DNA from mitochondria in <i>Saccharomyces cerevisiae</i> .. Molecular and Cellular Biology, 1993, 13, 5418-5426. | 1.1 | 201 |
| 2209 | Isolation and characterization of <i>S. cerevisiae</i> mutants defective in somatostatin expression: cloning and functional role of a yeast gene encoding an aspartyl protease in precursor processing at monobasic cleavage sites.. EMBO Journal, 1993, 12, 285-294. | 3.5 | 52 |
| 2210 | Two novel protein kinase C-related genes of fission yeast are essential for cell viability and implicated in cell shape control.. EMBO Journal, 1993, 12, 1987-1995. | 3.5 | 125 |
| 2211 | A <i>Saccharomyces cerevisiae</i> homologue of mammalian translation initiation factor 4B contributes to RNA helicase activity.. EMBO Journal, 1993, 12, 3997-4003. | 3.5 | 116 |
| 2212 | Novel members of the mitogen-activated protein kinase activator family in <i>Xenopus laevis</i> .. Molecular and Cellular Biology, 1993, 13, 5738-5748. | 1.1 | 66 |
| 2213 | SSG1, a gene encoding a sporulation-specific 1,3-beta-glucanase in <i>Saccharomyces cerevisiae</i> . Journal of Bacteriology, 1993, 175, 3823-3837. | 1.0 | 50 |
| 2214 | Salt tolerance and methionine biosynthesis in <i>Saccharomyces cerevisiae</i> involve a putative phosphatase gene.. EMBO Journal, 1993, 12, 3105-3110. | 3.5 | 131 |

| # | ARTICLE | IF | CITATIONS |
|------|---|-----|-----------|
| 2215 | Functional cooperation of mitochondrial protein import receptors in yeast.. EMBO Journal, 1993, 12, 4115-4123. | 3.5 | 202 |
| 2216 | A U3 snoRNP protein with homology to splicing factor PRP4 and G beta domains is required for ribosomal RNA processing.. EMBO Journal, 1993, 12, 2549-2558. | 3.5 | 111 |
| 2217 | Transcriptional regulator Leu3 of <i>Saccharomyces cerevisiae</i> : separation of activator and repressor functions.. Molecular and Cellular Biology, 1993, 13, 5702-5709. | 1.1 | 35 |
| 2218 | MOT2 encodes a negative regulator of gene expression that affects basal expression of pheromone-responsive genes in <i>Saccharomyces cerevisiae</i> .. Molecular and Cellular Biology, 1994, 14, 3139-3149. | 1.1 | 30 |
| 2219 | The HOG pathway controls osmotic regulation of transcription via the stress response element (STRE) of the <i>Saccharomyces cerevisiae</i> CTT1 gene.. EMBO Journal, 1994, 13, 4382-4389. | 3.5 | 443 |
| 2220 | Wort Aldehyde Reduction Potential in Free and Immobilized Yeast Systems. Journal of the American Society of Brewing Chemists, 1994, 52, 100-106. | 0.8 | 27 |
| 2221 | The Dbf2 and Dbf20 protein kinases of budding yeast are activated after the metaphase to anaphase cell cycle transition.. EMBO Journal, 1994, 13, 1103-1113. | 3.5 | 120 |
| 2222 | Molecular cloning of chromosome I DNA from <i>Saccharomyces cerevisiae</i> : analysis of the genes in the FUN38-MAK16-SPO7 region. Journal of Bacteriology, 1994, 176, 1872-1880. | 1.0 | 43 |
| 2223 | Fission yeast cut3 and cut14, members of a ubiquitous protein family, are required for chromosome condensation and segregation in mitosis.. EMBO Journal, 1994, 13, 4938-4952. | 3.5 | 300 |
| 2224 | MSG5, a novel protein phosphatase promotes adaptation to pheromone response in <i>S. cerevisiae</i> .. EMBO Journal, 1994, 13, 61-70. | 3.5 | 216 |
| 2225 | Functional differences between mammalian transcription activation domains at the yeast GAL1 promoter.. EMBO Journal, 1994, 13, 641-645. | 3.5 | 66 |
| 2226 | A dominant negative mutation in the conserved RNA helicase motif $\hat{\text{S}}\text{AT}^{\text{TM}}$ causes splicing factor PRP2 to stall in spliceosomes.. EMBO Journal, 1994, 13, 879-887. | 3.5 | 86 |
| 2227 | VPS21 encodes a rab5-like GTP binding protein that is required for the sorting of yeast vacuolar proteins.. EMBO Journal, 1994, 13, 1297-1309. | 3.5 | 177 |
| 2228 | GDI1 encodes a GDP dissociation inhibitor that plays an essential role in the yeast secretory pathway.. EMBO Journal, 1994, 13, 1718-1728. | 3.5 | 180 |
| 2229 | A mitochondrial homolog of bacterial GrpE interacts with mitochondrial hsp70 and is essential for viability.. EMBO Journal, 1994, 13, 1998-2006. | 3.5 | 162 |
| 2230 | A functional interaction between Rev and yeast pre-mRNA is related to splicing complex formation.. EMBO Journal, 1994, 13, 4096-4104. | 3.5 | 71 |
| 2231 | A zinc finger protein, essential for chromosome segregation, constitutes a putative DNA binding subunit of the <i>Saccharomyces cerevisiae</i> kinetochore complex, Cbf3.. EMBO Journal, 1994, 13, 5203-5211. | 3.5 | 75 |
| 2232 | Fission yeast cut5 links nuclear chromatin and M phase regulator in the replication checkpoint control.. EMBO Journal, 1994, 13, 5319-5329. | 3.5 | 128 |

| # | ARTICLE | IF | CITATIONS |
|------|---|-----|-----------|
| 2233 | tRNA genes as transcriptional repressor elements.. <i>Molecular and Cellular Biology</i> , 1994, 14, 1266-1277. | 1.1 | 119 |
| 2234 | The <i>Saccharomyces cerevisiae</i> APS1 gene encodes a homolog of the small subunit of the mammalian clathrin AP-1 complex: evidence for functional interaction with clathrin at the Golgi complex.. <i>EMBO Journal</i> , 1994, 13, 1706-1717. | 3.5 | 70 |
| 2235 | Characterization of two protein kinases from <i>Schizosaccharomyces pombe</i> involved in the regulation of DNA repair.. <i>EMBO Journal</i> , 1994, 13, 2777-2788. | 3.5 | 96 |
| 2236 | The yeast NOP4 gene product is an essential nucleolar protein required for pre-rRNA processing and accumulation of 60S ribosomal subunits.. <i>EMBO Journal</i> , 1994, 13, 3127-3135. | 3.5 | 89 |
| 2237 | Molecular characterization of SIG1, a <i>Saccharomyces cerevisiae</i> gene involved in negative regulation of G-protein-mediated signal transduction.. <i>EMBO Journal</i> , 1994, 13, 3050-3064. | 3.5 | 14 |
| 2238 | The UGA4 UASNTR site required for GLN3-dependent transcriptional activation also mediates DAL80-responsive regulation and DAL80 protein binding in <i>Saccharomyces cerevisiae</i> . <i>Journal of Bacteriology</i> , 1994, 176, 4718-4725. | 1.0 | 59 |
| 2239 | The URE2 protein regulates nitrogen catabolic gene expression through the GATAA-containing UASNTR element in <i>Saccharomyces cerevisiae</i> . <i>Journal of Bacteriology</i> , 1994, 176, 7476-7483. | 1.0 | 54 |
| 2240 | Complete DNA sequence of yeast chromosome II.. <i>EMBO Journal</i> , 1994, 13, 5795-5809. | 3.5 | 216 |
| 2241 | Cloning and expression of the MEP1 gene encoding an ammonium transporter in <i>Saccharomyces cerevisiae</i> .. <i>EMBO Journal</i> , 1994, 13, 3456-3463. | 3.5 | 261 |
| 2242 | NUP145 encodes a novel yeast glycine-leucine-phenylalanine-glycine (GLFG) nucleoporin required for nuclear envelope structure.. <i>Journal of Cell Biology</i> , 1994, 125, 955-969. | 2.3 | 135 |
| 2243 | Trehalose-6-P synthase is dispensable for growth on glucose but not for spore germination in <i>Schizosaccharomyces pombe</i> . <i>Journal of Bacteriology</i> , 1994, 176, 3895-3902. | 1.0 | 82 |
| 2244 | Phosphorylation of dis2 protein phosphatase at the C-terminal cdc2 consensus and its potential role in cell cycle regulation.. <i>EMBO Journal</i> , 1994, 13, 5310-5318. | 3.5 | 110 |
| 2245 | ACT3: a putative centractin homologue in <i>S. cerevisiae</i> is required for proper orientation of the mitotic spindle.. <i>Journal of Cell Biology</i> , 1994, 127, 129-138. | 2.3 | 111 |
| 2246 | The presence of an active S-adenosylmethionine decarboxylase gene increases the growth defect observed in <i>Saccharomyces cerevisiae</i> mutants unable to synthesize putrescine, spermidine, and spermine. <i>Journal of Bacteriology</i> , 1994, 176, 6407-6409. | 1.0 | 20 |
| 2247 | A nucleosome precludes binding of the transcription factor Pho4 in vivo to a critical target site in the PHO5 promoter.. <i>EMBO Journal</i> , 1994, 13, 4848-4855. | 3.5 | 115 |
| 2248 | Cloning, nucleotide sequence, and regulation of <i>Schizosaccharomyces pombe</i> thi4, a thiamine biosynthetic gene. <i>Journal of Bacteriology</i> , 1994, 176, 6631-6635. | 1.0 | 37 |
| 2249 | The mutation DGT1-1 decreases glucose transport and alleviates carbon catabolite repression in <i>Saccharomyces cerevisiae</i> . <i>Journal of Bacteriology</i> , 1994, 176, 7423-7429. | 1.0 | 61 |
| 2250 | Identification of cis- and trans-acting elements involved in the expression of cold shock-inducible TIP1 gene of yeast <i>Saccharomyces cerevisiae</i> . <i>Nucleic Acids Research</i> , 1994, 22, 560-568. | 6.5 | 13 |

| # | ARTICLE | IF | CITATIONS |
|------|---|--------|-----------|
| 2251 | A screen for yeast mutants with defects in the dolichol-mediated pathway for N-glycosylation.. Proceedings of the National Academy of Sciences of the United States of America, 1994, 91, 1485-1489. | 3.3 | 27 |
| 2252 | Growth site localization of Rho1 small GTP-binding protein and its involvement in bud formation in <i>Saccharomyces cerevisiae</i> .. Journal of Cell Biology, 1994, 125, 1077-1093. | 2.3 | 251 |
| 2253 | Mapping actin surfaces required for functional interactions in vivo.. Journal of Cell Biology, 1994, 126, 423-432. | 2.3 | 63 |
| 2254 | Signal-mediated retrieval of a membrane protein from the Golgi to the ER in yeast.. Journal of Cell Biology, 1994, 127, 653-665. | 2.3 | 256 |
| 2255 | Glucose-transport-deficient mutants of <i>Schizosaccharomyces pombe</i> : phenotype, genetics and use for genetic complementation. Microbiology (United Kingdom), 1994, 140, 2617-2623. | 0.7 | 41 |
| 2256 | A <i>Candida albicans</i> cyclic nucleotide phosphodiesterase: cloning and expression in <i>Saccharomyces cerevisiae</i> and biochemical characterization of the recombinant enzyme. Microbiology (United Kingdom), 1994, 140, 2617-2623. | 0.7843 | 147 |
| 2257 | Feedback regulation of 3-hydroxy-3-methylglutaryl coenzyme A reductase in <i>Saccharomyces cerevisiae</i> .. Molecular Biology of the Cell, 1994, 5, 655-665. | 0.9 | 71 |
| 2258 | Identification and analysis of the <i>Saccharomyces cerevisiae</i> SYR1 gene reveals that ergosterol is involved in the action of syringomycin. Microbiology (United Kingdom), 1994, 140, 353-359. | 0.7 | 53 |
| 2259 | A role for autophosphorylation revealed by activated alleles of FUS3, the yeast MAP kinase homolog.. Molecular Biology of the Cell, 1994, 5, 297-312. | 0.9 | 74 |
| 2260 | Mutational analysis of an essential binding site for the U3 snoRNA in the 5' external transcribed spacer of yeast pre-rRNA. Nucleic Acids Research, 1994, 22, 4057-4065. | 6.5 | 61 |
| 2261 | SRD1, a <i>S. cerevisiae</i> gene affecting pre-rRNA processing contains a C2/C2 zinc finger motif. Nucleic Acids Research, 1994, 22, 1265-1271. | 6.5 | 20 |
| 2262 | Two <i>Saccharomyces cerevisiae</i> genes which control sensitivity to G1 arrest induced by <i>Kluyveromyces lactis</i> toxin.. Molecular and Cellular Biology, 1994, 14, 6306-6316. | 1.1 | 88 |
| 2263 | Introduction of YACs into intact yeast cells by a procedure which shows low levels of recombination and co-transformation. Nucleic Acids Research, 1994, 22, 5011-5015. | 6.5 | 14 |
| 2264 | His-154 is involved in the linkage of the <i>Saccharomyces cerevisiae</i> L-A double-stranded RNA virus Gag protein to the cap structure of mRNAs and is essential for M1 satellite virus expression.. Molecular and Cellular Biology, 1994, 14, 2664-2674. | 1.1 | 56 |
| 2265 | Internal tracts of telomeric DNA act as silencers in <i>Saccharomyces cerevisiae</i> .. Genes and Development, 1994, 8, 1411-1422. | 2.7 | 83 |
| 2266 | The essential yeast Tcp1 protein affects actin and microtubules.. Molecular Biology of the Cell, 1994, 5, 1065-1080. | 0.9 | 99 |
| 2267 | Dual requirement in yeast DNA mismatch repair for MLH1 and PMS1, two homologs of the bacterial mutL gene.. Molecular and Cellular Biology, 1994, 14, 407-415. | 1.1 | 229 |
| 2268 | Fine-resolution mapping of spontaneous and double-strand break-induced gene conversion tracts in <i>Saccharomyces cerevisiae</i> reveals reversible mitotic conversion polarity.. Molecular and Cellular Biology, 1994, 14, 3863-3875. | 1.1 | 100 |

| # | ARTICLE | IF | CITATIONS |
|------|---|-----|-----------|
| 2269 | Overcoming telomeric silencing: a trans-activator competes to establish gene expression in a cell cycle-dependent way.. <i>Genes and Development</i> , 1994, 8, 1133-1146. | 2.7 | 271 |
| 2270 | New Developments in Fungal Virology. <i>Advances in Virus Research</i> , 1994, 43, 303-388. | 0.9 | 80 |
| 2271 | Identification and characterization of a novel yeast gene: the YGP1 gene product is a highly glycosylated secreted protein that is synthesized in response to nutrient limitation.. <i>Molecular and Cellular Biology</i> , 1994, 14, 2740-2754. | 1.1 | 65 |
| 2272 | Histone H1 expressed in <i>Saccharomyces cerevisiae</i> binds to chromatin and affects survival, growth, transcription, and plasmid stability but does not change nucleosomal spacing.. <i>Molecular and Cellular Biology</i> , 1994, 14, 2822-2835. | 1.1 | 27 |
| 2273 | Nuclear congression and membrane fusion: two distinct events in the yeast karyogamy pathway.. <i>Journal of Cell Biology</i> , 1994, 126, 911-923. | 2.3 | 111 |
| 2274 | A novel FK506- and rapamycin-binding protein (FPR3 gene product) in the yeast <i>Saccharomyces cerevisiae</i> is a proline rotamase localized to the nucleolus.. <i>Journal of Cell Biology</i> , 1994, 127, 623-639. | 2.3 | 78 |
| 2275 | Bypassing anaphase by fission yeast cut9 mutation: requirement of cut9+ to initiate anaphase.. <i>Journal of Cell Biology</i> , 1994, 127, 1655-1670. | 2.3 | 103 |
| 2276 | Metabolic instability and constitutive endocytosis of STE6, the a-factor transporter of <i>Saccharomyces cerevisiae</i> .. <i>Molecular Biology of the Cell</i> , 1994, 5, 1185-1198. | 0.9 | 102 |
| 2277 | Delineation of a small region within the major transactivation domain of the human glucocorticoid receptor that mediates transactivation of gene expression.. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 1994, 91, 1619-1623. | 3.3 | 92 |
| 2278 | An internal region of the peroxisomal membrane protein PMP47 is essential for sorting to peroxisomes. <i>Journal of Cell Biology</i> , 1994, 124, 915-925. | 2.3 | 84 |
| 2279 | The MAP kinase Fus3 associates with and phosphorylates the upstream signaling component Ste5.. <i>Genes and Development</i> , 1994, 8, 313-327. | 2.7 | 126 |
| 2280 | Yeast NPI46 encodes a novel prolyl cis-trans isomerase that is located in the nucleolus.. <i>Journal of Cell Biology</i> , 1994, 126, 853-862. | 2.3 | 66 |
| 2281 | Synthesis of yeast cell wall glucan and evidence for glucan metabolism in a <i>Saccharomyces cerevisiae</i> whole cell system. <i>Microbiology (United Kingdom)</i> , 1994, 140, 2229-2237. | 0.7 | 10 |
| 2282 | A nuclear protein with sequence similarity to proteins implicated in human acute leukemias is important for cellular morphogenesis and actin cytoskeletal function in <i>Saccharomyces cerevisiae</i> .. <i>Molecular Biology of the Cell</i> , 1994, 5, 617-632. | 0.9 | 49 |
| 2283 | Subcellular locations of MOD5 proteins: mapping of sequences sufficient for targeting to mitochondria and demonstration that mitochondrial and nuclear isoforms commingle in the cytosol.. <i>Molecular and Cellular Biology</i> , 1994, 14, 2298-2306. | 1.1 | 73 |
| 2284 | T7 RNA polymerase-dependent expression of COXII in yeast mitochondria.. <i>Molecular and Cellular Biology</i> , 1994, 14, 4643-4652. | 1.1 | 41 |
| 2285 | Multiple genes encode the translation elongation factor EF-1 β in <i>Saccharomyces cerevisiae</i> . <i>Nucleic Acids Research</i> , 1994, 22, 2703-2707. | 6.5 | 37 |
| 2286 | The budding yeast U5 snRNP Prp8 is a highly conserved protein which links RNA splicing with cell cycle progression. <i>Nucleic Acids Research</i> , 1994, 22, 5555-5564. | 6.5 | 51 |

| # | ARTICLE | IF | CITATIONS |
|------|---|-----|-----------|
| 2287 | Isolation and characterization of a <i>Saccharomyces cerevisiae</i> peptide transport gene.. <i>Molecular and Cellular Biology</i> , 1994, 14, 104-115. | 1.1 | 136 |
| 2288 | An essential gene, <i>ESR1</i> , is required for mitotic growth, DNA repair and meiotic recombination <i>Saccharomyces cerevisiae</i> . <i>Nucleic Acids Research</i> , 1994, 22, 3104-3112. | 6.5 | 223 |
| 2289 | Analysis of <i>FAS3/ACC</i> regulatory region of <i>Saccharomyces cerevisiae</i> : identification of a functional <i>UASINO</i> and sequences responsible for fatty acid mediated repression. <i>Nucleic Acids Research</i> , 1994, 22, 412-418. | 6.5 | 36 |
| 2290 | Collaboration of G1 cyclins in the functional inactivation of the retinoblastoma protein.. <i>Genes and Development</i> , 1994, 8, 1759-1771. | 2.7 | 225 |
| 2291 | Transcription through the yeast origin of replication <i>ARS1</i> ends at the <i>ABFI</i> binding site and affects extrachromosomal maintenance of minichromosomes. <i>Nucleic Acids Research</i> , 1994, 22, 3904-3910. | 6.5 | 31 |
| 2292 | Expression of the <i>E.coli ada</i> gene in <i>S.cerevisiae</i> provides cellular resistance to <i>N</i> -methyl- <i>A</i> '-nitro- <i>A</i> '-nitrosoguanidine in <i>rad6</i> but not in <i>rad52</i> mutants. <i>Nucleic Acids Research</i> , 1994, 22, 5717-5722. | 6.5 | 8 |
| 2293 | Interaction between the first and last nucleotides of pre-mRNA introns is a determinant of 3' splice site selection in <i>S.cerevisiae</i> . <i>Nucleic Acids Research</i> , 1994, 22, 1981-1987. | 6.5 | 44 |
| 2294 | Genetic and physical interactions between <i>Srp1p</i> and nuclear pore complex proteins <i>Nup1p</i> and <i>Nup2p</i> .. <i>Journal of Cell Biology</i> , 1994, 126, 619-630. | 2.3 | 135 |
| 2295 | The <i>END3</i> gene encodes a protein that is required for the internalization step of endocytosis and for actin cytoskeleton organization in yeast.. <i>Molecular Biology of the Cell</i> , 1994, 5, 1023-1037. | 0.9 | 265 |
| 2296 | The <i>Aspergillus nidulans abaA</i> gene encodes a transcriptional activator that acts as a genetic switch to control development.. <i>Molecular and Cellular Biology</i> , 1994, 14, 2503-2515. | 1.1 | 246 |
| 2297 | <i>POM152</i> is an integral protein of the pore membrane domain of the yeast nuclear envelope.. <i>Journal of Cell Biology</i> , 1994, 125, 31-42. | 2.3 | 139 |
| 2298 | <i>P40SDB25</i> , a putative CDK inhibitor, has a role in the M/G1 transition in <i>Saccharomyces cerevisiae</i> .. <i>Genes and Development</i> , 1994, 8, 1640-1653. | 2.7 | 141 |
| 2299 | Induction of pseudohyphal growth by overexpression of <i>PHD1</i> , a <i>Saccharomyces cerevisiae</i> gene related to transcriptional regulators of fungal development.. <i>Molecular and Cellular Biology</i> , 1994, 14, 2100-2112. | 1.1 | 290 |
| 2300 | Endocytosis is required for the growth of vacuolar H(+)-ATPase-defective yeast: identification of six new <i>END</i> genes.. <i>Journal of Cell Biology</i> , 1994, 127, 373-386. | 2.3 | 260 |
| 2301 | <i>GPD1</i> , which encodes glycerol-3-phosphate dehydrogenase, is essential for growth under osmotic stress in <i>Saccharomyces cerevisiae</i> , and its expression is regulated by the high-osmolarity glycerol response pathway.. <i>Molecular and Cellular Biology</i> , 1994, 14, 4135-4144. | 1.1 | 641 |
| 2302 | The <i>SAD1/RAD53</i> protein kinase controls multiple checkpoints and DNA damage-induced transcription in yeast.. <i>Genes and Development</i> , 1994, 8, 2401-2415. | 2.7 | 460 |
| 2303 | <i>nup1</i> mutants exhibit pleiotropic defects in nuclear pore complex function.. <i>Journal of Cell Biology</i> , 1994, 127, 319-332. | 2.3 | 87 |
| 2304 | Mutation of the gene encoding protein kinase C 1 stimulates mitotic recombination in <i>Saccharomyces cerevisiae</i> .. <i>Molecular and Cellular Biology</i> , 1994, 14, 6039-6045. | 1.1 | 84 |

| # | ARTICLE | IF | CITATIONS |
|------|--|-----|-----------|
| 2305 | Requirements for intercistronic distance and level of eukaryotic initiation factor 2 activity in reinitiation on GCN4 mRNA vary with the downstream cistron.. <i>Molecular and Cellular Biology</i> , 1994, 14, 2616-2628. | 1.1 | 95 |
| 2306 | The centromeric K-type repeat and the central core are together sufficient to establish a functional <i>Schizosaccharomyces pombe</i> centromere.. <i>Molecular Biology of the Cell</i> , 1994, 5, 747-761. | 0.9 | 154 |
| 2307 | Localization of the Kar3 kinesin heavy chain-related protein requires the Cik1 interacting protein. <i>Journal of Cell Biology</i> , 1994, 124, 507-519. | 2.3 | 115 |
| 2308 | <i>Candida albicans</i> gene encoding resistance to benomyl and methotrexate is a multidrug resistance gene. <i>Antimicrobial Agents and Chemotherapy</i> , 1994, 38, 648-652. | 1.4 | 158 |
| 2309 | Regulation of splicing at an intermediate step in the formation of the spliceosome.. <i>Genes and Development</i> , 1994, 8, 211-220. | 2.7 | 89 |
| 2310 | The JNM1 gene in the yeast <i>Saccharomyces cerevisiae</i> is required for nuclear migration and spindle orientation during the mitotic cell cycle.. <i>Journal of Cell Biology</i> , 1994, 125, 143-158. | 2.3 | 97 |
| 2311 | Involvement of SRE element of Ty1 transposon in TEC1-dependent transcriptional activation. <i>Nucleic Acids Research</i> , 1994, 22, 999-1005. | 6.5 | 60 |
| 2312 | Mutational analysis of the PRP4 protein of <i>Saccharomyces cerevisiae</i> suggests domain structure and snRNP interactions. <i>Nucleic Acids Research</i> , 1994, 22, 1724-1734. | 6.5 | 27 |
| 2313 | The WD repeats of Tup1 interact with the homeo domain protein alpha 2.. <i>Genes and Development</i> , 1994, 8, 2857-2867. | 2.7 | 210 |
| 2314 | Large-scale analysis of gene expression, protein localization, and gene disruption in <i>Saccharomyces cerevisiae</i> .. <i>Genes and Development</i> , 1994, 8, 1087-1105. | 2.7 | 500 |
| 2315 | TOR1 and TOR2 are structurally and functionally similar but not identical phosphatidylinositol kinase homologues in yeast.. <i>Molecular Biology of the Cell</i> , 1994, 5, 105-118. | 0.9 | 365 |
| 2316 | A Small GTP-Binding Protein from <i>Arabidopsis thaliana</i> Functionally Complements the Yeast YPT6 Null Mutant. <i>Plant Physiology</i> , 1994, 104, 591-596. | 2.3 | 94 |
| 2317 | Regulation of mitochondrial morphology and inheritance by Mdm10p, a protein of the mitochondrial outer membrane.. <i>Journal of Cell Biology</i> , 1994, 126, 1361-1373. | 2.3 | 270 |
| 2318 | Fission yeast minichromosome loss mutants mis cause lethal aneuploidy and replication abnormality.. <i>Molecular Biology of the Cell</i> , 1994, 5, 1145-1158. | 0.9 | 211 |
| 2319 | Initiation preference at a yeast origin of replication.. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 1994, 91, 3418-3422. | 3.3 | 67 |
| 2320 | Regulated degradation of HMG-CoA reductase, an integral membrane protein of the endoplasmic reticulum, in yeast.. <i>Journal of Cell Biology</i> , 1994, 125, 299-312. | 2.3 | 212 |
| 2321 | Identification of cut8+ and cek1+, a novel protein kinase gene, which complement a fission yeast mutation that blocks anaphase.. <i>Molecular and Cellular Biology</i> , 1994, 14, 6361-6371. | 1.1 | 39 |
| 2322 | The yeast TEM1 gene, which encodes a GTP-binding protein, is involved in termination of M phase.. <i>Molecular and Cellular Biology</i> , 1994, 14, 7476-7482. | 1.1 | 182 |

| # | ARTICLE | IF | CITATIONS |
|------|---|-----|-----------|
| 2323 | Mutant isolation of mouse DNA topoisomerase II \pm in yeast. <i>Nucleic Acids Research</i> , 1994, 22, 4229-4233. | 6.5 | 11 |
| 2324 | Are yeast chitin synthases regulated at the transcriptional or the posttranslational level?. <i>Molecular and Cellular Biology</i> , 1994, 14, 7685-7694. | 1.1 | 104 |
| 2325 | The interaction between the first and last intron nucleotides in the second step of pre-mRNA splicing is independent of other conserved intron nucleotides. <i>Nucleic Acids Research</i> , 1994, 22, 5190-5195. | 6.5 | 23 |
| 2326 | Mutational analysis of an essential binding site for the U3 snoRNA in the 5' external transcribed spacer of yeast pre-rRNA. <i>Nucleic Acids Research</i> , 1994, 22, 5139-5147. | 6.5 | 91 |
| 2327 | Methylglyoxal and Lipid Hydroperoxide as Endogenous Cytotoxic Molecular Species: Detoxification and Regulation of Gene Expression in Yeasts. <i>Biotechnology and Genetic Engineering Reviews</i> , 1994, 12, 467-508. | 2.4 | 1 |
| 2328 | Genetic interactions between SIN3 mutations and the <i>Saccharomyces cerevisiae</i> transcriptional activators encoded by MCM1, STE12, and SWI1. <i>Molecular Genetics and Genomics</i> , 1994, 245, 675-685. | 2.4 | 20 |
| 2329 | Expression of human poly(ADP-ribose) polymerase in <i>Saccharomyces cerevisiae</i> . <i>Molecular Genetics and Genomics</i> , 1994, 245, 686-693. | 2.4 | 15 |
| 2330 | SLK1, a yeast homolog of MAP kinase activators, has a RAS/cAMP-independent role in nutrient sensing. <i>Molecular Genetics and Genomics</i> , 1994, 243, 286-296. | 2.4 | 52 |
| 2331 | New in-vivo cloning methods by homologous recombination in yeast. <i>Current Genetics</i> , 1994, 25, 180-183. | 0.8 | 17 |
| 2332 | MRG1-1, a dominant allele that confers methomyl resistance in yeast expressing the cytoplasmic male sterility T-urf13 gene from maize. <i>Current Genetics</i> , 1994, 26, 477-485. | 0.8 | 6 |
| 2333 | The byp1-3 allele of the <i>Saccharomyces cerevisiae</i> GGS1/TPS1 gene and its multi-copy suppressor tRNAGLN (CAG): Ggs1/Tps1 protein levels restraining growth on fermentable sugars and trehalose accumulation. <i>Current Genetics</i> , 1994, 26, 295-301. | 0.8 | 20 |
| 2334 | UV hypersensitivity of yeast linear plasmids. <i>Current Genetics</i> , 1994, 26, 369-373. | 0.8 | 23 |
| 2335 | ERV1 is involved in the cell-division cycle and the maintenance of mitochondrial genomes in <i>Saccharomyces cerevisiae</i> . <i>Current Genetics</i> , 1994, 26, 15-20. | 0.8 | 69 |
| 2336 | <i>Saccharomyces cerevisiae</i> ribosomal protein L37 is encoded by duplicate genes that are differentially expressed. <i>Current Genetics</i> , 1994, 25, 480-487. | 0.8 | 15 |
| 2337 | Cloning and analysis of a FLO5 flocculation gene from <i>S. cerevisiae</i> . <i>Current Genetics</i> , 1994, 25, 196-201. | 0.8 | 45 |
| 2338 | Chimeric evolution of the 2-m genome in <i>Saccharomyces cerevisiae</i> . <i>Journal of Molecular Evolution</i> , 1994, 38, 363-368. | 0.8 | 3 |
| 2339 | Shared control of maltose induction and catabolite repression of the MAL structural genes in <i>Saccharomyces</i> . <i>Molecular Genetics and Genomics</i> , 1994, 243, 622-630. | 2.4 | 24 |
| 2340 | RPK1, an essential yeast protein kinase involved in the regulation of the onset of mitosis, shows homology to mammalian dual-specificity kinases. <i>Molecular Genetics and Genomics</i> , 1994, 243, 641-653. | 2.4 | 31 |

| # | ARTICLE | IF | CITATIONS |
|------|---|-----|-----------|
| 2341 | A new yeast gene, HTR1, required for growth at high temperature, is needed for recovery from mating pheromone-induced G1 arrest. <i>Molecular Genetics and Genomics</i> , 1994, 245, 107-116. | 2.4 | 40 |
| 2342 | Cerulenin-resistant mutants of <i>Saccharomyces cerevisiae</i> with an altered fatty acid synthase gene. <i>Molecular Genetics and Genomics</i> , 1994, 244, 90-96. | 2.4 | 80 |
| 2343 | Increases in cell size at START caused by hyperactivation of the cAMP pathway in <i>Saccharomyces cerevisiae</i> . <i>Molecular Genetics and Genomics</i> , 1994, 243, 158-165. | 2.4 | 20 |
| 2344 | Stress-induced transcriptional activation mediated by YAP1 and YAP2 genes that encode the Jun family of transcriptional activators in <i>Saccharomyces cerevisiae</i> . <i>Molecular Genetics and Genomics</i> , 1994, 242, 250-256. | 2.4 | 93 |
| 2345 | A putative catabolite-repressed cell wall protein from the mycoparasitic fungus <i>Trichoderma harzianum</i> . <i>Molecular Genetics and Genomics</i> , 1994, 242, 461-466. | 2.4 | 33 |
| 2346 | Carbon catabolite repression in <i>Kluyveromyces lactis</i> : isolation and characterization of the KINLD gene encoding the mitochondrial enzyme D-lactate ferricytochrome c oxidoreductase. <i>Molecular Genetics and Genomics</i> , 1994, 244, 622-629. | 2.4 | 39 |
| 2347 | Increase in incidence of chromosome instability and non-conservative recombination between repeats in <i>Saccharomyces cerevisiae</i> hpr1 ⁺ strains. <i>Molecular Genetics and Genomics</i> , 1994, 245, 224-236. | 2.4 | 40 |
| 2348 | Genetic interactions among genes involved in the STT4-PKC1 pathway of <i>Saccharomyces cerevisiae</i> . <i>Molecular Genetics and Genomics</i> , 1994, 242, 631-640. | 2.4 | 92 |
| 2349 | Homologs of the yeast neck filament associated genes: isolation and sequence analysis of <i>Candida albicans</i> CDC3 and CDC10. <i>Molecular Genetics and Genomics</i> , 1994, 242, 689-698. | 2.4 | 36 |
| 2350 | Functional analysis of the zinc cluster domain of the CYP1 (HAP1) complex regulator in heme-sufficient and heme-deficient yeast cells. <i>Molecular Genetics and Genomics</i> , 1994, 242, 699-707. | 2.4 | 23 |
| 2351 | Isolation and characterization of phosphoprotein phosphatase 1 from alfalfa. <i>Molecular Genetics and Genomics</i> , 1994, 244, 176-182. | 2.4 | 12 |
| 2352 | The archaeobacterial membrane protein bacterio-opsin is expressed and N-terminally processed in the yeast <i>Saccharomyces cerevisiae</i> . <i>Molecular Genetics and Genomics</i> , 1994, 244, 183-188. | 2.4 | 11 |
| 2353 | The positive and negative cis-acting elements for methanol regulation in the <i>Pichia pastoris</i> AOX2 gene. <i>Molecular Genetics and Genomics</i> , 1994, 243, 489-499. | 2.4 | 53 |
| 2354 | MBR1 and MBR3, two related yeast genes that can suppress the growth defect of hap2, hap3 and hap4 mutants. <i>Molecular Genetics and Genomics</i> , 1994, 243, 575-583. | 2.4 | 22 |
| 2355 | SCM2, a tryptophan permease in <i>Saccharomyces cerevisiae</i> , is important for cell growth. <i>Molecular Genetics and Genomics</i> , 1994, 244, 260-268. | 2.4 | 26 |
| 2356 | Interactions between gene products involved in divalent cation transport in <i>Saccharomyces cerevisiae</i> . <i>Molecular Genetics and Genomics</i> , 1994, 244, 303-311. | 2.4 | 64 |
| 2357 | Differential accumulation of S-adenosylmethionine synthetase transcripts in response to salt stress. <i>Plant Molecular Biology</i> , 1994, 25, 217-227. | 2.0 | 175 |
| 2358 | The plant transcription factor TGA1 stimulates expression of the CaMV 35S promoter in <i>Saccharomyces cerevisiae</i> . <i>Plant Molecular Biology</i> , 1994, 25, 323-328. | 2.0 | 16 |

| # | ARTICLE | IF | CITATIONS |
|------|---|-----|-----------|
| 2359 | Expression and secretion of human lipocortin-1 by promoter and signal sequence of STA1 from <i>Saccharomyces diastaticus</i> . <i>Biotechnology Letters</i> , 1994, 16, 897-902. | 1.1 | 7 |
| 2360 | Molecular characterization of the SPT23 gene: A dosage-dependent suppressor of ty-induced promoter mutations from <i>Saccharomyces cerevisiae</i> . <i>Yeast</i> , 1994, 10, 81-92. | 0.8 | 24 |
| 2361 | Selectable marker replacement in <i>Saccharomyces cerevisiae</i> . <i>Yeast</i> , 1994, 10, 141-149. | 0.8 | 15 |
| 2362 | Sequence comparison of the ARG4 chromosomal regions from the two related yeasts, <i>Saccharomyces cerevisiae</i> and <i>Saccharomyces douglasii</i> . <i>Yeast</i> , 1994, 10, 309-317. | 0.8 | 21 |
| 2363 | A new family of yeast genes implicated in ergosterol synthesis is related to the human oxysterol binding protein. <i>Yeast</i> , 1994, 10, 341-353. | 0.8 | 93 |
| 2364 | XIII. Yeast sequencing reports. Cloning and sequencing of the NES24 gene of <i>Saccharomyces cerevisiae</i> . <i>Yeast</i> , 1994, 10, 371-376. | 0.8 | 2 |
| 2365 | Isolation of a CDC25 family gene, MSI2/LTE1, as a multicopy suppressor of <i>ira1</i> . <i>Yeast</i> , 1994, 10, 451-461. | 0.8 | 102 |
| 2366 | A search for an essential function of the replication origin ARS1 in the life cycle of <i>Saccharomyces cerevisiae</i> . <i>Yeast</i> , 1994, 10, 491-496. | 0.8 | 3 |
| 2367 | Use of β -lactamase as a secreted reporter of promoter function in yeast. <i>Yeast</i> , 1994, 10, 497-508. | 0.8 | 27 |
| 2368 | <i>Saccharomyces cerevisiae</i> mata mutant cells defective in pointed projection formation in response to β -factor at high concentrations. <i>Yeast</i> , 1994, 10, 579-594. | 0.8 | 14 |
| 2369 | V. Yeast mapping reports. The MAG1 3-methyladenine DNA glycosylase gene is closely linked to the SPT15 TATA-binding TFIIID gene on chromosome V-R in <i>Saccharomyces cerevisiae</i> . <i>Yeast</i> , 1994, 10, 687-691. | 0.8 | 4 |
| 2370 | A new nuclear suppressor system for a mitochondrial RNA polymerase mutant identifies an unusual zinc-finger protein and a polyglutamine domain protein in <i>Saccharomyces cerevisiae</i> . <i>Yeast</i> , 1994, 10, 719-731. | 0.8 | 29 |
| 2371 | Yeast β -glucanases can be used as efficient and readily detectable reporter genes in <i>Saccharomyces cerevisiae</i> . <i>Yeast</i> , 1994, 10, 747-756. | 0.8 | 26 |
| 2372 | Phosphoribosylpyrophosphate synthetase (PRS): A new gene family in <i>Saccharomyces cerevisiae</i> . <i>Yeast</i> , 1994, 10, 1031-1044. | 0.8 | 42 |
| 2373 | Activation of trehalase during growth induction by nitrogen sources in the yeast <i>Saccharomyces cerevisiae</i> depends on the free catalytic subunits of camp-dependent protein kinase, but not on functional ras proteins. <i>Yeast</i> , 1994, 10, 1049-1064. | 0.8 | 59 |
| 2374 | Predominant localization of non-specific lipid-transfer protein of the yeast <i>Candida tropicalis</i> in the matrix of peroxisomes. <i>Yeast</i> , 1994, 10, 1065-1074. | 0.8 | 16 |
| 2375 | Analysis of β -glucans and chitin in a <i>Saccharomyces cerevisiae</i> cell wall mutant using high-performance liquid chromatography. <i>Yeast</i> , 1994, 10, 1083-1092. | 0.8 | 31 |
| 2376 | VII. Yeast sequencing reports. Cloning and characterization of a sulphite-resistance gene of <i>Saccharomyces cerevisiae</i> . <i>Yeast</i> , 1994, 10, 1101-1110. | 0.8 | 23 |

| # | ARTICLE | IF | CITATIONS |
|------|---|-----|-----------|
| 2377 | II. Yeast sequencing reports. The nucleotide sequence of TTP1, a gene encoding a predicted type II membrane protein. <i>Yeast</i> , 1994, 10, 1111-1115. | 0.8 | 5 |
| 2378 | Yeast sequencing reports. Cloning and DNA sequence of a <i>Kluyveromyces lactis</i> ERD1 homologue. <i>Yeast</i> , 1994, 10, 1117-1124. | 0.8 | 4 |
| 2379 | Identification of a set of yeast genes coding for a novel family of putative ATPases with high similarity to constituents of the 26S protease complex. <i>Yeast</i> , 1994, 10, 1141-1155. | 0.8 | 127 |
| 2380 | Molecular characterization of a gene that confers 2-deoxyglucose resistance in yeast. <i>Yeast</i> , 1994, 10, 1195-1202. | 0.8 | 29 |
| 2381 | IV. Yeast sequencing reports. Nucleotide sequence of the SAC2 gene of <i>Saccharomyces cerevisiae</i> . <i>Yeast</i> , 1994, 10, 1211-1216. | 0.8 | 6 |
| 2382 | A family of vectors that facilitate transposon and insertional mutagenesis of cloned genes in yeast. <i>Yeast</i> , 1994, 10, 1267-1272. | 0.8 | 35 |
| 2383 | A genetic screen to isolate genes regulated by the yeast CCAAT-box binding protein Hap2p. <i>Yeast</i> , 1994, 10, 1273-1283. | 0.8 | 55 |
| 2384 | Physiological approach to heterologous human serum albumin production by <i>Kluyveromyces lactis</i> in chemostat culture. <i>Yeast</i> , 1994, 10, 1297-1303. | 0.8 | 22 |
| 2385 | IX. Yeast sequencing reports. Cloning and sequence of REV7, a gene whose function is required for DNA damage-induced mutagenesis in <i>Saccharomyces cerevisiae</i> . <i>Yeast</i> , 1994, 10, 1503-1509. | 0.8 | 31 |
| 2386 | New heterologous modules for classical or PCR-based gene disruptions in <i>Saccharomyces cerevisiae</i> . <i>Yeast</i> , 1994, 10, 1793-1808. | 0.8 | 2,547 |
| 2387 | In <i>Saccharomyces cerevisiae</i> overexpression of hybrid Virus-Like-Particles correlates with altered cell volume distributions. <i>Biotechnology Letters</i> , 1994, 16, 1131-1134. | 1.1 | 1 |
| 2388 | Basidiomycetous rDNA functionally replaces its homolog genes in yeast. <i>Current Genetics</i> , 1994, 25, 30-33. | 0.8 | 3 |
| 2389 | PDR3, a new yeast regulatory gene, is homologous to PDR1 and controls the multidrug resistance phenomenon. <i>Molecular Genetics and Genomics</i> , 1994, 244, 501-511. | 2.4 | 190 |
| 2390 | The vacuolar compartment is required for sulfur amino acid homeostasis in <i>Saccharomyces cerevisiae</i> . <i>Molecular Genetics and Genomics</i> , 1994, 244, 519-529. | 2.4 | 25 |
| 2391 | Multifactorial experimental design for optimizing transformation: Electroporation of <i>Streptococcus thermophilus</i> . <i>Biotechnology and Bioengineering</i> , 1994, 43, 490-496. | 1.7 | 29 |
| 2392 | Regulation of the expression of the H ⁺ -ATPase genes PMA1 and PMA2 during growth and effects of octanoic acid in <i>Saccharomyces cerevisiae</i> . <i>Biochimica Et Biophysica Acta Gene Regulatory Mechanisms</i> , 1994, 1217, 65-73. | 2.4 | 32 |
| 2393 | Primary structure of aspergillopepsin I deduced from nucleotide sequence of the gene and aspartic acid-76 is an essential active site of the enzyme for trypsinogen activation. <i>BBA - Proteins and Proteomics</i> , 1994, 1204, 257-264. | 2.1 | 36 |
| 2394 | Comparison of heme environment at the putative distal region of P-450s utilizing their external and internal nitrogenous ligand bound forms. <i>BBA - Proteins and Proteomics</i> , 1994, 1207, 49-57. | 2.1 | 9 |

| # | ARTICLE | IF | CITATIONS |
|------|--|-----|-----------|
| 2395 | Control of secretory production of human lysozyme from <i>Saccharomyces cerevisiae</i> by incubation temperature and phosphate concentration. <i>Journal of Bioscience and Bioengineering</i> , 1994, 77, 376-381. | 0.9 | 2 |
| 2396 | Overexpression and secretion of cellulolytic enzymes by $\hat{\Gamma}$ -sequence-mediated multicopy integration of heterologous DNA sequences into the chromosomes of <i>Saccharomyces cerevisiae</i> . <i>Journal of Bioscience and Bioengineering</i> , 1994, 77, 468-473. | 0.9 | 20 |
| 2397 | Induction of yeast sporulation by lysine-related compounds and glutathione in nutrition-rich conditions. <i>Journal of Bioscience and Bioengineering</i> , 1994, 77, 568-571. | 0.9 | 5 |
| 2398 | A method for direct selection of mating-competent clones from mating-incompetent industrial strains of <i>Saccharomyces cerevisiae</i> . <i>Journal of Bioscience and Bioengineering</i> , 1994, 78, 6-11. | 0.9 | 12 |
| 2399 | Cloning and nucleotide sequence of a gene conferring ability to grow at a low temperature on <i>Saccharomyces cerevisiae</i> tryptophan auxotrophs. <i>Journal of Bioscience and Bioengineering</i> , 1994, 77, 1-9. | 0.9 | 12 |
| 2400 | Expression of the carboxymethylcellulase gene, CMC1, from <i>Cryptococcus flavus</i> in <i>Saccharomyces cerevisiae</i> . <i>Journal of Bioscience and Bioengineering</i> , 1994, 77, 235-238. | 0.9 | 5 |
| 2401 | Introduction of a feed back resistant $\hat{\Gamma}$ -isopropylmalate synthase gene of <i>Saccharomyces cerevisiae</i> into sake yeast. <i>Journal of Bioscience and Bioengineering</i> , 1994, 77, 119-124. | 0.9 | 5 |
| 2402 | A DNA construct useful for specific chromosome loss in <i>Saccharomyces cerevisiae</i> . <i>Journal of Bioscience and Bioengineering</i> , 1994, 77, 125-130. | 0.9 | 4 |
| 2403 | PDC2, a yeast gene essential for synthesis of pyruvate decarboxylase, encodes a novel transcription factor. <i>Journal of Genetics</i> , 1994, 73, 17-32. | 0.4 | 14 |
| 2404 | Direct selection shuttle plasmid vector, pPW264, used for cloning the $\hat{\Gamma}$ -amylase gene of <i>Schwanniomyces occidentalis</i> . <i>Folia Microbiologica</i> , 1994, 39, 255-260. | 1.1 | 2 |
| 2405 | Industrial production of heterologous proteins by fed-batch cultures of the yeast <i>Saccharomyces cerevisiae</i> . <i>FEMS Microbiology Reviews</i> , 1994, 15, 369-410. | 3.9 | 75 |
| 2406 | <i>Saccharomyces cerevisiae</i> mutants sensitive to the antimalarial and antiarrhythmic drug, quinidine. <i>FEMS Microbiology Letters</i> , 1994, 119, 221-227. | 0.7 | 6 |
| 2407 | Investigation of the role of sterol $\hat{\Gamma}$ 7-isomerase in the sensitivity of <i>Saccharomyces cerevisiae</i> to fenpropimorph. <i>FEMS Microbiology Letters</i> , 1994, 122, 223-226. | 0.7 | 16 |
| 2408 | Mixed Lactic Acid-Alcoholic Fermentation by <i>Saccharomyces cerevisiae</i> Expressing the <i>Lactobacillus casei</i> L(+)-LDH. <i>Nature Biotechnology</i> , 1994, 12, 173-177. | 9.4 | 113 |
| 2409 | A Reb1p-binding site is required for efficient activation of the yeast RAP1 gene, but multiple binding sites for Rap1p are not essential. <i>Molecular Microbiology</i> , 1994, 12, 931-940. | 1.2 | 20 |
| 2410 | The Levels of Yeast Gluconeogenic mRNAs Respond to Environmental Factors. <i>FEBS Journal</i> , 1994, 224, 473-481. | 0.2 | 51 |
| 2411 | Isolation and expression of a gene specifying a cdc2-like protein kinase from the human malaria parasite <i>Plasmodium falciparum</i> . <i>FEBS Journal</i> , 1994, 220, 693-701. | 0.2 | 91 |
| 2412 | Human beta1,4 galactosyltransferase and alpha2,6 sialyltransferase expressed in <i>Saccharomyces cerevisiae</i> are retained as active enzymes in the endoplasmic reticulum. <i>FEBS Journal</i> , 1994, 220, 809-817. | 0.2 | 24 |

| # | ARTICLE | IF | CITATIONS |
|------|--|------|-----------|
| 2413 | The NAM1 protein (NAM1p), which is selectively required for cox1, cytb and atp6 transcript processing/stabilisation, is located in the yeast mitochondrial matrix. FEBS Journal, 1994, 222, 27-32. | 0.2 | 35 |
| 2414 | ATP synthase of yeast mitochondria. Isolation of the F1 delta subunit, sequence and disruption of the structural gene. FEBS Journal, 1994, 222, 851-859. | 0.2 | 39 |
| 2415 | Catabolite inactivation of heterologous fructose-1,6-bisphosphatases and fructose-1,6-bisphosphatase-beta-galactosidase fusion proteins in <i>Saccharomyces cerevisiae</i> . FEBS Journal, 1994, 222, 879-884. | 0.2 | 17 |
| 2416 | Production of active, insect-specific scorpion neurotoxin in yeast. FEBS Journal, 1994, 223, 637-645. | 0.2 | 31 |
| 2417 | Proteinase yscD (oligopeptidase yscD). Structure, function and relationship of the yeast enzyme with mammalian thimet oligopeptidase (metalloendopeptidase, EP 24.15). FEBS Journal, 1994, 219, 627-639. | 0.2 | 51 |
| 2418 | The Folding of the Bifunctional TRP3 Protein in Yeast is Influenced by a Translational Pause which Lies in a Region of Structural Divergence with <i>Escherichia coli</i> Indoleglycerol-Phosphate Synthase. FEBS Journal, 1994, 226, 657-664. | 0.2 | 33 |
| 2419 | Evidence That CYP2C19 is the Major (S)-Mephenytoin 4'-Hydroxylase in Humans. <i>Biochemistry</i> , 1994, 33, 1743-1752. | 1.2 | 333 |
| 2420 | Crossover interference is abolished in the absence of a synaptonemal complex protein. <i>Cell</i> , 1994, 79, 283-292. | 13.5 | 298 |
| 2421 | Nup145p is required for nuclear export of mRNA and binds homopolymeric RNA in vitro via a novel conserved motif. <i>Cell</i> , 1994, 78, 275-289. | 13.5 | 139 |
| 2422 | Cleavage of prosomatostatins by the yeast Yap3 and Kex2 endoprotease. <i>Biochimie</i> , 1994, 76, 226-233. | 1.3 | 26 |
| 2423 | Mutation of a fungal aspartic proteinase, <i>Mucor pusillus</i> rennin, to decrease thermostability for use as a milk coagulant. <i>Journal of Biotechnology</i> , 1994, 32, 17-28. | 1.9 | 21 |
| 2424 | Human interferon- β , expressed in <i>Saccharomyces cerevisiae</i> , is predominantly directed to the vacuoles Influence of modified co-expression of secretion factors and chaperones. <i>Journal of Biotechnology</i> , 1994, 32, 179-189. | 1.9 | 5 |
| 2425 | Production of recombinant hirudin by high cell density fed-batch cultivations of a <i>Saccharomyces cerevisiae</i> strain: physiological considerations during the bioprocess design. <i>Journal of Biotechnology</i> , 1994, 32, 249-259. | 1.9 | 24 |
| 2426 | Introduction of DNA into Yeast Cells. <i>Current Protocols in Molecular Biology</i> , 1994, 27, Unit13.7. | 2.9 | 48 |
| 2427 | Expression of the outer capsid proteins VP2 and VP5 of bluetongue virus in <i>Saccharomyces cerevisiae</i> . <i>Virus Research</i> , 1994, 33, 11-25. | 1.1 | 8 |
| 2428 | Analysis of <i>Saccharomyces cerevisiae</i> proteins induced by peroxide and superoxide stress. <i>Microbiology (United Kingdom)</i> , 1994, 140, 3277-3283. | 0.7 | 119 |
| 2429 | Construction of squalene-accumulating <i>Saccharomyces cerevisiae</i> mutants by gene disruption through homologous recombination. <i>Applied Microbiology and Biotechnology</i> , 1994, 42, 353-357. | 1.7 | 14 |
| 2430 | Characterization of the catalase of the n-alkane-utilizing yeast <i>Candida tropicalis</i> functionally expressed in <i>Saccharomyces cerevisiae</i> . <i>Applied Microbiology and Biotechnology</i> , 1994, 40, 682-686. | 1.7 | 7 |

| # | ARTICLE | IF | CITATIONS |
|------|---|-----|-----------|
| 2431 | Saccharomyces cerevisiae cells lacking the homologous pairing protein p175 SEP1 arrest at pachytene during meiotic prophase. Chromosoma, 1994, 103, 129-141. | 1.0 | 39 |
| 2432 | Subunit requirements for expression of functional sodium pumps in yeast cells. Biochimica Et Biophysica Acta - Biomembranes, 1994, 1193, 226-234. | 1.4 | 43 |
| 2433 | Constitutive expression of the human D2S-dopamine receptor in the unicellular yeast Saccharomyces cerevisiae. Biochimica Et Biophysica Acta - Biomembranes, 1994, 1193, 255-262. | 1.4 | 27 |
| 2434 | Expression of the human D2Sdopamine receptor in the yeasts Saccharomyces cerevisiae and Schizosaccharomyces pombe: A comparative study. FEBS Letters, 1994, 344, 41-46. | 1.3 | 56 |
| 2435 | Involvement of cell wall β -glucan in the action of HM-1 killer toxin. FEBS Letters, 1994, 348, 27-32. | 1.3 | 49 |
| 2436 | The human ubiquitin-conjugating enzyme UbcH1 is involved in the repair of UV-damaged, alkylated and cross-linked DNA. FEBS Letters, 1994, 350, 1-4. | 1.3 | 13 |
| 2437 | A 70-kDa heat shock cognate protein suppresses the defects caused by a proteasome mutation in Saccharomyces cerevisiae. FEBS Letters, 1994, 351, 263-266. | 1.3 | 26 |
| 2438 | Yta10p, a member of a novel ATPase family in yeast, is essential for mitochondrial function. FEBS Letters, 1994, 353, 197-200. | 1.3 | 71 |
| 2439 | Polymannosylation to asparagine-19 in hen egg white lysozyme in yeast. FEBS Letters, 1994, 355, 76-80. | 1.3 | 13 |
| 2440 | Anatomy of the stimulative sequences flanking the ARS consensus sequence of chromosome VI in Saccharomyces cerevisiae. Gene, 1994, 150, 213-220. | 1.0 | 12 |
| 2441 | Isolation and functional analysis of a Kluyveromyces lactis RAP1 homologue. Gene, 1994, 150, 35-41. | 1.0 | 50 |
| 2442 | Anti-estrogen activity in the yeast transcription system: Estrogen receptor mediated agonist response. Steroids, 1994, 59, 572-578. | 0.8 | 50 |
| 2443 | Survival of genetically engineered and wild-type strains of the yeast Saccharomyces cerevisiae under simulated environmental conditions: a contribution on risk assessment. Journal of Applied Bacteriology, 1994, 77, 689-693. | 1.1 | 12 |
| 2444 | Heterologous Expression of the Human D2S Dopamine Receptor in Protease-Deficient Saccharomyces cerevisiae Strains. FEBS Journal, 1994, 226, 697-705. | 0.2 | 44 |
| 2445 | The Schizosaccharomyces pombe pka1 gene, encoding a homolog of cAMP-dependent protein kinase. Gene, 1994, 151, 215-220. | 1.0 | 17 |
| 2446 | A new system for the release of heterologous proteins from yeast based on mutant strains deficient in cell integrity. Journal of Biotechnology, 1994, 38, 81-88. | 1.9 | 17 |
| 2447 | Isolation of a Family of cDNA Clones from Ricinus communis L. with Close Homology to the Hexose Carriers. Journal of Plant Physiology, 1994, 143, 178-183. | 1.6 | 45 |
| 2448 | The MCS1/SSD1/SRK1/SSL1 gene is involved in stable maintenance of the chromosome in yeast. Gene, 1994, 143, 135-138. | 1.0 | 50 |

| # | ARTICLE | IF | CITATIONS |
|------|--|-----|-----------|
| 2449 | Genetic and biochemical evidence for yeast GCN2 protein kinase polymerization. <i>Gene</i> , 1994, 143, 21-27. | 1.0 | 19 |
| 2450 | The mating-type region of <i>Schizosaccharomyces pombe</i> contains an essential gene encoding a protein homologous to human modulators of HIV transactivation. <i>Gene</i> , 1994, 145, 205-210. | 1.0 | 15 |
| 2451 | Cloning of a rat cDNA encoding dihydroxypolyprenylbenzoate methyltransferase by functional complementation of a <i>Saccharomyces cerevisiae</i> mutant deficient in ubiquinone biosynthesis. <i>Gene</i> , 1994, 138, 213-217. | 1.0 | 39 |
| 2452 | The <i>E. coli recA</i> gene can restore the defect in mutagenesis of the <i>pso4-1</i> mutant of <i>S. cerevisiae</i> . <i>Mutation Research DNA Repair</i> , 1994, 314, 209-220. | 3.8 | 13 |
| 2453 | Thiamine-repressible genes in <i>Schizosaccharomyces pombe</i> are regulated by a Cys6 zinc-finger motif-containing protein. <i>Gene</i> , 1994, 147, 141-144. | 1.0 | 21 |
| 2454 | Expression of the non-structural protein NS1 of bluetongue virus in bacteria and yeast: identification of two antigenic sites at the amino terminus. <i>Virus Research</i> , 1994, 31, 291-303. | 1.1 | 11 |
| 2455 | Sensitivity to nitrogen mustard in <i>Saccharomyces cerevisiae</i> is independently determined by regulated choline permease and DNA repair. <i>Mutation Research DNA Repair</i> , 1994, 315, 139-145. | 3.8 | 5 |
| 2456 | The isolation and characterization of the gene (<i>dfr1</i>) encoding dihydrofolate reductase (DHFR) in <i>Schizosaccharomyces pombe</i> . <i>Gene</i> , 1994, 147, 131-135. | 1.0 | 10 |
| 2457 | Expression in yeast of an <i>Escherichia coli</i> gene encoding a phospholipid biosynthetic enzyme. <i>Gene</i> , 1994, 147, 111-114. | 1.0 | 1 |
| 2458 | The <i>STL1</i> gene of <i>Saccharomyces cerevisiae</i> is predicted to encode a sugar transporter-like protein. <i>Gene</i> , 1994, 146, 215-219. | 1.0 | 29 |
| 2459 | GSH1, which encodes gamma-glutamylcysteine synthetase, is a target gene for <i>yAP-1</i> transcriptional regulation.. <i>Molecular and Cellular Biology</i> , 1994, 14, 5832-5839. | 1.1 | 291 |
| 2460 | Copper Amine Oxidase: Heterologous Expression, Purification, and Characterization of An Active Enzyme in <i>Saccharomyces cerevisiae</i> . <i>Biochemistry</i> , 1994, 33, 7647-7653. | 1.2 | 113 |
| 2461 | Mutation in Hydrophobic Domain of ATP Synthase Subunit 4 (Subunit b) of Yeast Mitochondria Disturbs Coupling between Proton Translocation and Catalysis. <i>Biochemistry</i> , 1994, 33, 9684-9691. | 1.2 | 18 |
| 2462 | Isolation, Characterization, and Disruption of the Yeast Gene Encoding Cytosolic NADP-Specific Isocitrate Dehydrogenase. <i>Biochemistry</i> , 1994, 33, 9661-9667. | 1.2 | 59 |
| 2463 | Metabolic studies on <i>Saccharomyces cerevisiae</i> containing fused citrate synthase/malate dehydrogenase. <i>Biochemistry</i> , 1994, 33, 11684-11691. | 1.2 | 21 |
| 2464 | Potentiation of Progesterone Receptor-Mediated Transcription by the Immunosuppressant FK506. <i>Biochemistry</i> , 1994, 33, 10666-10671. | 1.2 | 58 |
| 2465 | Two alternative pathways of transcription initiation in the yeast negative regulatory gene <i>GAL80</i> .. <i>Molecular and Cellular Biology</i> , 1994, 14, 6819-6828. | 1.1 | 26 |
| 2466 | Substitution of asparagine residues in <i>Aspergillus awamori</i> glucoamylase by site-directed mutagenesis to eliminate N-glycosylation and inactivation by deamidation. <i>Biochemical Journal</i> , 1994, 301, 275-281. | 1.7 | 66 |

| # | ARTICLE | IF | CITATIONS |
|------|--|-----|-----------|
| 2467 | Studies on the effect of an heterologous fatty acid-binding protein on acyl-CoA oxidase induction in <i>Saccharomyces cerevisiae</i> . <i>Biochemical Journal</i> , 1994, 301, 615-620. | 1.7 | 9 |
| 2468 | Selective elongation of the oligosaccharide attached to the second potential glycosylation site of yeast exoglucanase: effects on the activity and properties of the enzyme. <i>Biochemical Journal</i> , 1994, 304, 917-922. | 1.7 | 10 |
| 2469 | The UAS of the Yeast GAPDH Promoter Consists of Multiple General Functional Elements Including RAP1 and GRF2 Binding Sites.. <i>Journal of Veterinary Medical Science</i> , 1994, 56, 235-244. | 0.3 | 17 |
| 2470 | [25] Purification and characterization of recombinant human sickle hemoglobin expressed in yeast. <i>Methods in Enzymology</i> , 1994, 231, 390-403. | 0.4 | 3 |
| 2471 | [5] Activation of cholera toxin by ADP-ribosylation factors: 20-kDa guanine nucleotide-binding proteins. <i>Methods in Enzymology</i> , 1994, 237, 44-63. | 0.4 | 6 |
| 2472 | SPT10 and SPT21 are required for transcription of particular histone genes in <i>Saccharomyces cerevisiae</i> .. <i>Molecular and Cellular Biology</i> , 1994, 14, 5223-5228. | 1.1 | 83 |
| 2473 | Saturation mutagenesis of a polyadenylation signal reveals a hexanucleotide element essential for mRNA 3' end formation in <i>Saccharomyces cerevisiae</i> .. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 1994, 91, 257-261. | 3.3 | 72 |
| 2474 | Functional expression of P-glycoprotein in <i>Saccharomyces cerevisiae</i> confers cellular resistance to the immunosuppressive and antifungal agent FK520.. <i>Molecular and Cellular Biology</i> , 1994, 14, 277-286. | 1.1 | 70 |
| 2475 | Transcriptional control of the yeast PDR5 gene by the PDR3 gene product.. <i>Molecular and Cellular Biology</i> , 1994, 14, 4653-4661. | 1.1 | 202 |
| 2476 | The <i>sua8</i> suppressors of <i>Saccharomyces cerevisiae</i> encode replacements of conserved residues within the largest subunit of RNA polymerase II and affect transcription start site selection similarly to <i>sua7</i> (TFIIB) mutations.. <i>Molecular and Cellular Biology</i> , 1994, 14, 226-237. | 1.1 | 87 |
| 2477 | Dissection of the ADR1 protein reveals multiple, functionally redundant activation domains interspersed with inhibitory regions: evidence for a repressor binding to the ADR1c region.. <i>Molecular and Cellular Biology</i> , 1994, 14, 629-640. | 1.1 | 50 |
| 2478 | Special peptidyl-tRNA molecules can promote translational frameshifting without slippage.. <i>Molecular and Cellular Biology</i> , 1994, 14, 8107-8116. | 1.1 | 62 |
| 2479 | Spermidine deficiency increases +1 ribosomal frameshifting efficiency and inhibits Ty1 retrotransposition in <i>Saccharomyces cerevisiae</i> .. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 1994, 91, 172-176. | 3.3 | 73 |
| 2480 | Negative regulation of mitosis in fission yeast by catalytically inactive <i>pyp1</i> and <i>pyp2</i> mutants.. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 1994, 91, 10084-10088. | 3.3 | 20 |
| 2481 | Cloning of the <i>Saccharomyces cerevisiae</i> gene whose overexpression overcomes the effects of HM-1 killer toxin, which inhibits beta-glucan synthesis. <i>Journal of Bacteriology</i> , 1994, 176, 1488-1499. | 1.0 | 47 |
| 2482 | Ntk: a Csk-related protein-tyrosine kinase expressed in brain and T lymphocytes.. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 1994, 91, 4975-4979. | 3.3 | 69 |
| 2483 | Colocalization of vertebrate lamin B and lamin B receptor (LBR) in nuclear envelopes and in LBR-induced membrane stacks of the yeast <i>Saccharomyces cerevisiae</i> .. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 1994, 91, 10124-10128. | 3.3 | 59 |
| 2484 | The GLC7 type 1 protein phosphatase is required for glucose repression in <i>Saccharomyces cerevisiae</i> .. <i>Molecular and Cellular Biology</i> , 1994, 14, 6789-6796. | 1.1 | 113 |

| # | ARTICLE | IF | CITATIONS |
|------|---|-----|-----------|
| 2485 | The yeast MOT2 gene encodes a putative zinc finger protein that serves as a global negative regulator affecting expression of several categories of genes, including mating-pheromone-responsive genes.. Molecular and Cellular Biology, 1994, 14, 3150-3157. | 1.1 | 32 |
| 2486 | Mutations in the GCD7 subunit of yeast guanine nucleotide exchange factor eIF-2B overcome the inhibitory effects of phosphorylated eIF-2 on translation initiation.. Molecular and Cellular Biology, 1994, 14, 3208-3222. | 1.1 | 67 |
| 2487 | Type 1 protein phosphatase acts in opposition to IpL1 protein kinase in regulating yeast chromosome segregation.. Molecular and Cellular Biology, 1994, 14, 4731-4740. | 1.1 | 259 |
| 2488 | Low levels of exogenous histone H1 in yeast cause cell death.. Proceedings of the National Academy of Sciences of the United States of America, 1994, 91, 11567-11570. | 3.3 | 25 |
| 2489 | CDC44: a putative nucleotide-binding protein required for cell cycle progression that has homology to subunits of replication factor C.. Molecular and Cellular Biology, 1994, 14, 255-267. | 1.1 | 85 |
| 2490 | Differential effects of Cdc68 on cell cycle-regulated promoters in <i>Saccharomyces cerevisiae</i> .. Molecular and Cellular Biology, 1994, 14, 7455-7465. | 1.1 | 45 |
| 2491 | Identification and analysis of a <i>Saccharomyces cerevisiae</i> copper homeostasis gene encoding a homeodomain protein.. Molecular and Cellular Biology, 1994, 14, 7792-7804. | 1.1 | 31 |
| 2492 | G1 cyclin degradation: the PEST motif of yeast Cln2 is necessary, but not sufficient, for rapid protein turnover.. Molecular and Cellular Biology, 1994, 14, 7953-7966. | 1.1 | 105 |
| 2493 | Interactions among three proteins that specifically activate translation of the mitochondrial COX3 mRNA in <i>Saccharomyces cerevisiae</i> .. Molecular and Cellular Biology, 1994, 14, 1045-1053. | 1.1 | 74 |
| 2494 | Mutations in XRS2 and RAD50 delay but do not prevent mating-type switching in <i>Saccharomyces cerevisiae</i> .. Molecular and Cellular Biology, 1994, 14, 3414-3425. | 1.1 | 222 |
| 2495 | Two types of RAS mutants that dominantly interfere with activators of RAS.. Molecular and Cellular Biology, 1994, 14, 3707-3718. | 1.1 | 52 |
| 2496 | Domain B of ARS307 contains two functional elements and contributes to chromosomal replication origin function.. Molecular and Cellular Biology, 1994, 14, 7652-7659. | 1.1 | 109 |
| 2497 | Genetic identification of residues involved in association of alpha and beta G-protein subunits.. Molecular and Cellular Biology, 1994, 14, 3223-3229. | 1.1 | 51 |
| 2498 | The yeast and mammalian Ras pathways control transcription of heat shock genes independently of heat shock transcription factor.. Molecular and Cellular Biology, 1994, 14, 4929-4937. | 1.1 | 74 |
| 2499 | Concerted action of RAS and G proteins in the sexual response pathways of <i>Schizosaccharomyces pombe</i> .. Molecular and Cellular Biology, 1994, 14, 50-58. | 1.1 | 81 |
| 2500 | Mutational activation of the STE5 gene product bypasses the requirement for G protein beta and gamma subunits in the yeast pheromone response pathway.. Molecular and Cellular Biology, 1994, 14, 1054-1065. | 1.1 | 74 |
| 2501 | The third cytoplasmic loop of a yeast G-protein-coupled receptor controls pathway activation, ligand discrimination, and receptor internalization.. Molecular and Cellular Biology, 1994, 14, 3339-3349. | 1.1 | 76 |
| 2502 | TPD1 of <i>Saccharomyces cerevisiae</i> encodes a protein phosphatase 2C-like activity implicated in tRNA splicing and cell separation.. Molecular and Cellular Biology, 1994, 14, 3634-3645. | 1.1 | 64 |

| # | ARTICLE | IF | CITATIONS |
|------|--|-----|-----------|
| 2503 | Induction of recombination between homologous and diverged DNAs by double-strand gaps and breaks and role of mismatch repair.. <i>Molecular and Cellular Biology</i> , 1994, 14, 4802-4814. | 1.1 | 65 |
| 2504 | A <i>Drosophila</i> gene promoter is subject to glucose repression in yeast cells.. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 1994, 91, 11109-11112. | 3.3 | 19 |
| 2505 | The mutant type 1 protein phosphatase encoded by <i>glc7-1</i> from <i>Saccharomyces cerevisiae</i> fails to interact productively with the GAC1-encoded regulatory subunit.. <i>Molecular and Cellular Biology</i> , 1994, 14, 896-905. | 1.1 | 106 |
| 2506 | Biochemical and genetic analysis of dominant-negative mutations affecting a yeast G-protein gamma subunit.. <i>Molecular and Cellular Biology</i> , 1994, 14, 4571-4578. | 1.1 | 32 |
| 2507 | Effect of sequence context at stop codons on efficiency of reinitiation in GCN4 translational control.. <i>Molecular and Cellular Biology</i> , 1994, 14, 606-618. | 1.1 | 112 |
| 2508 | MDS1, a dosage suppressor of an <i>mck1</i> mutant, encodes a putative yeast homolog of glycogen synthase kinase 3.. <i>Molecular and Cellular Biology</i> , 1994, 14, 831-839. | 1.1 | 82 |
| 2509 | Two FK506 resistance-conferring genes in <i>Saccharomyces cerevisiae</i> , TAT1 and TAT2, encode amino acid permeases mediating tyrosine and tryptophan uptake.. <i>Molecular and Cellular Biology</i> , 1994, 14, 6597-6606. | 1.1 | 161 |
| 2510 | Protein phosphatase 2C, encoded by <i>ptc1+</i> , is important in the heat shock response of <i>Schizosaccharomyces pombe</i> .. <i>Molecular and Cellular Biology</i> , 1994, 14, 3742-3751. | 1.1 | 71 |
| 2511 | Myocyte nuclear factor, a novel winged-helix transcription factor under both developmental and neural regulation in striated myocytes.. <i>Molecular and Cellular Biology</i> , 1994, 14, 4596-4605. | 1.1 | 94 |
| 2512 | Flexibility and interchangeability of polyadenylation signals in <i>Saccharomyces cerevisiae</i> .. <i>Molecular and Cellular Biology</i> , 1994, 14, 4633-4642. | 1.1 | 29 |
| 2513 | Transcriptional activation by CTF proteins is mediated by a bipartite low-proline domain.. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 1994, 91, 3901-3905. | 3.3 | 44 |
| 2514 | The Essential Helicase Gene <i><i>RAD3</i></i> Suppresses Short-Sequence Recombination in <i><i>Saccharomyces cerevisiae</i></i> . <i>Molecular and Cellular Biology</i> , 1995, 15, 3998-4008. | 1.1 | 42 |
| 2515 | Positive Autoregulation of the Yeast Transcription Factor Pdr3p, Which is Involved in Control of Drug Resistance. <i>Molecular and Cellular Biology</i> , 1995, 15, 4043-4051. | 1.1 | 137 |
| 2516 | The <i>Saccharomyces cerevisiae</i> phosphatidylinositol-transfer protein effects a ligand-dependent inhibition of choline-phosphate cytidyltransferase activity.. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 1995, 92, 112-116. | 3.3 | 148 |
| 2517 | Disruption of the nucleoporin gene NUP133 results in clustering of nuclear pore complexes.. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 1995, 92, 1187-1191. | 3.3 | 84 |
| 2518 | Single-Stranded DNA Arising at Telomeres in <i><i>cdc13</i></i> Mutants May Constitute a Specific Signal for the <i><i>RAD9</i></i> Checkpoint. <i>Molecular and Cellular Biology</i> , 1995, 15, 6128-6138. | 1.1 | 619 |
| 2519 | REF2 Encodes an RNA-Binding Protein Directly Involved in Yeast mRNA 3' End Formation. <i>Molecular and Cellular Biology</i> , 1995, 15, 1689-1697. | 1.1 | 39 |
| 2520 | Roles of <i><i>URE2</i></i> and <i><i>GLN3</i></i> in the Proline Utilization Pathway in <i><i>Saccharomyces cerevisiae</i></i> . <i>Molecular and Cellular Biology</i> , 1995, 15, 2321-2330. | 1.1 | 64 |

| # | ARTICLE | IF | CITATIONS |
|------|---|-----|-----------|
| 2521 | GAL4 Interacts with TATA-Binding Protein and Coactivators. <i>Molecular and Cellular Biology</i> , 1995, 15, 2839-2848. | 1.1 | 159 |
| 2522 | Bdf1, a Yeast Chromosomal Protein Required for Sporulation. <i>Molecular and Cellular Biology</i> , 1995, 15, 3685-3696. | 1.1 | 104 |
| 2523 | The <i>Saccharomyces cerevisiae</i> Gene <i>SDS22</i> Encodes a Potential Regulator of the Mitotic Function of Yeast Type 1 Protein Phosphatase. <i>Molecular and Cellular Biology</i> , 1995, 15, 3777-3785. | 1.1 | 95 |
| 2524 | DNA Ligase I Mediates Essential Functions in Mammalian Cells. <i>Molecular and Cellular Biology</i> , 1995, 15, 4303-4308. | 1.1 | 83 |
| 2525 | Identification of an additional gene required for eukaryotic nonsense mRNA turnover.. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 1995, 92, 10354-10358. | 3.3 | 109 |
| 2526 | Replication protein A binds to regulatory elements in yeast DNA repair and DNA metabolism genes.. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 1995, 92, 4907-4911. | 3.3 | 67 |
| 2527 | Expression of a plant viral polycistronic mRNA in yeast, <i>Saccharomyces cerevisiae</i> , mediated by a plant virus translational transactivator.. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 1995, 92, 8911-8915. | 3.3 | 16 |
| 2528 | <i>Escherichia coli</i> iron superoxide dismutase targeted to the mitochondria of yeast cells protects the cells against oxidative stress.. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 1995, 92, 4219-4223. | 3.3 | 27 |
| 2529 | Identification of essential nucleotides in an upstream repressing sequence of <i>Saccharomyces cerevisiae</i> by selection for increased expression of <i>TRK2</i> .. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 1995, 92, 2370-2374. | 3.3 | 25 |
| 2530 | Binding of a Cell-Type-Specific RNA Splicing Factor to Its Target Regulatory Sequence. <i>Molecular and Cellular Biology</i> , 1995, 15, 1953-1960. | 1.1 | 51 |
| 2531 | Multiple Genes, Including a Member of the AAA Family, Are Essential for Degradation of Unassembled Subunit 2 of Cytochrome <i>c</i> Oxidase in Yeast Mitochondria. <i>Molecular and Cellular Biology</i> , 1995, 15, 4441-4452. | 1.1 | 140 |
| 2532 | Isolation of yeast artificial chromosomes free of endogenous yeast chromosomes: construction of alternate hosts with defined karyotypic alterations.. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 1995, 92, 11706-11710. | 3.3 | 28 |
| 2533 | Transcription in the Yeast rRNA Gene Locus: Distribution of the Active Gene Copies and Chromatin Structure of Their Flanking Regulatory Sequences. <i>Molecular and Cellular Biology</i> , 1995, 15, 5294-5303. | 1.1 | 130 |
| 2534 | Correlation of Two-Hybrid Affinity Data with In Vitro Measurements. <i>Molecular and Cellular Biology</i> , 1995, 15, 5820-5829. | 1.1 | 537 |
| 2535 | The Amino-Terminal Domain of Yeast U1-70K Is Necessary and Sufficient for Function. <i>Molecular and Cellular Biology</i> , 1995, 15, 6341-6350. | 1.1 | 26 |
| 2536 | Feedback Inhibition of the Yeast Ribosomal Protein Gene <i>CRY2</i> Is Mediated by the Nucleotide Sequence and Secondary Structure of <i>CRY2</i> Pre-mRNA. <i>Molecular and Cellular Biology</i> , 1995, 15, 6454-6464. | 1.1 | 45 |
| 2537 | Yeast MEK-Dependent Signal Transduction: Response Thresholds and Parameters Affecting Fidelity. <i>Molecular and Cellular Biology</i> , 1995, 15, 6545-6553. | 1.1 | 65 |
| 2538 | <i>AFR1</i> Promotes Polarized Apical Morphogenesis in <i>Saccharomyces cerevisiae</i> . <i>Molecular and Cellular Biology</i> , 1995, 15, 723-730. | 1.1 | 46 |

| # | ARTICLE | IF | CITATIONS |
|------|--|-----|-----------|
| 2539 | A complex unidirectional signal element mediates GCN4 mRNA 3' end formation in <i>Saccharomyces cerevisiae</i> . <i>Molecular and Cellular Biology</i> , 1995, 15, 2466-2473. | 1.1 | 30 |
| 2540 | Yeast Virus Propagation Depends Critically on Free 60S Ribosomal Subunit Concentration. <i>Molecular and Cellular Biology</i> , 1995, 15, 2772-2781. | 1.1 | 109 |
| 2541 | Synergistic Activation of <i>ADH2</i> Expression Is Sensitive to Upstream Activation Sequence 2 (UAS2) Orientation, Copy Number, and UAS1-UAS2 Helical Phasing. <i>Molecular and Cellular Biology</i> , 1995, 15, 3442-3449. | 1.1 | 20 |
| 2542 | Genetic and Biochemical Interactions between an Essential Kinetochores Protein, Cbf2p/Ndc10p, and the <i>CDC34</i> Ubiquitin-Conjugating Enzyme. <i>Molecular and Cellular Biology</i> , 1995, 15, 4835-4842. | 1.1 | 34 |
| 2543 | Negative Regulation of G ₁ and G ₂ by S-Phase Cyclins of <i>Saccharomyces cerevisiae</i> . <i>Molecular and Cellular Biology</i> , 1995, 15, 5030-5042. | 1.1 | 36 |
| 2544 | The <i>maf</i> Proto-oncogene Stimulates Transcription from Multiple Sites in a Promoter That Directs Purkinje Neuron-Specific Gene Expression. <i>Molecular and Cellular Biology</i> , 1995, 15, 246-254. | 1.1 | 76 |
| 2545 | Cloning and expression of the carboxypeptidase gene from <i>Aspergillus saitoi</i> and determination of the catalytic residues by site-directed mutagenesis. <i>Biochemical Journal</i> , 1995, 308, 405-409. | 1.7 | 22 |
| 2546 | A subunit interface mutant of yeast pyruvate kinase requires the allosteric activator fructose 1,6-bisphosphate for activity. <i>Biochemical Journal</i> , 1995, 310, 117-123. | 1.7 | 26 |
| 2547 | [15] Analysis of interaction between Ras and CDC25 guanine nucleotide exchange factor using yeast GAL4 two-hybrid system. <i>Methods in Enzymology</i> , 1995, 255, 135-148. | 0.4 | 6 |
| 2548 | A practical method for fission yeast transformation by electroporation.. <i>Japanese Journal of Genetics</i> , 1995, 70, 1-6. | 1.0 | 13 |
| 2549 | Papulacandin B resistance in budding and fission yeasts: isolation and characterization of a gene involved in (1,3)beta-D-glucan synthesis in <i>Saccharomyces cerevisiae</i> . <i>Journal of Bacteriology</i> , 1995, 177, 5732-5739. | 1.0 | 81 |
| 2550 | Regulation of <i>Saccharomyces cerevisiae</i> Flavohemoglobin Gene Expression. <i>Journal of Biological Chemistry</i> , 1995, 270, 6991-6996. | 1.6 | 44 |
| 2551 | The <i>Candida boidinii</i> peroxisomal membrane protein Pmp30 has a role in peroxisomal proliferation and is functionally homologous to Pmp27 from <i>Saccharomyces cerevisiae</i> . <i>Journal of Bacteriology</i> , 1995, 177, 6773-6781. | 1.0 | 62 |
| 2552 | [15] High expression cloning, purification, and assay of Ypt ^{GTPase-activating} proteins. <i>Methods in Enzymology</i> , 1995, 257, 118-128. | 0.4 | 27 |
| 2553 | MOP2 (SLA2) Affects the Abundance of the Plasma Membrane H ⁺ -ATPase of <i>Saccharomyces cerevisiae</i> . <i>Journal of Biological Chemistry</i> , 1995, 270, 6815-6823. | 1.6 | 58 |
| 2554 | Mutant rat phosphatidylinositol/phosphatidylcholine transfer proteins specifically defective in phosphatidylinositol transfer: implications for the regulation of phospholipid transfer activity.. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 1995, 92, 8826-8830. | 3.3 | 60 |
| 2555 | Sug1 Modulates Yeast Transcription Activation by Cdc68. <i>Molecular and Cellular Biology</i> , 1995, 15, 6025-6035. | 1.1 | 56 |
| 2556 | <i>SOK2</i> May Regulate Cyclic AMP-Dependent Protein Kinase-Stimulated Growth and Pseudohyphal. <i>Molecular and Cellular Biology</i> , 1995, 15, 6854-6863. | 1.1 | 197 |

| # | ARTICLE | IF | CITATIONS |
|------|---|-----|-----------|
| 2557 | Mutations Altering the Mitochondrial-Cytoplasmic Distribution of Mod5p Implicate the Actin Cytoskeleton and mRNA 3' Ends and/or Protein Synthesis in Mitochondrial Delivery. <i>Molecular and Cellular Biology</i> , 1995, 15, 6884-6894. | 1.1 | 79 |
| 2558 | Identification and Characterization of a Sequence Motif Involved in Nonsense-Mediated mRNA Decay. <i>Molecular and Cellular Biology</i> , 1995, 15, 2231-2244. | 1.1 | 126 |
| 2559 | Analysis of the <i>Saccharomyces cerevisiae</i> Mitochondrial COX3 mRNA 5' Untranslated Leader: Translational Activation and mRNA Processing. <i>Molecular and Cellular Biology</i> , 1995, 15, 3291-3300. | 1.1 | 42 |
| 2560 | Mutational Analysis of Hsp90 Function: Interactions with a Steroid Receptor and a Protein Kinase. <i>Molecular and Cellular Biology</i> , 1995, 15, 3917-3925. | 1.1 | 407 |
| 2561 | Functional Differences and Interactions among the Putative RecA Homologs Rad51, Rad55, and Rad57. <i>Molecular and Cellular Biology</i> , 1995, 15, 4843-4850. | 1.1 | 229 |
| 2562 | Functional expression of a vertebrate inwardly rectifying K ⁺ channel in yeast.. <i>Molecular Biology of the Cell</i> , 1995, 6, 1231-1240. | 0.9 | 60 |
| 2563 | Isolation and characterization of pokeweed antiviral protein mutations in <i>Saccharomyces cerevisiae</i> : identification of residues important for toxicity.. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 1995, 92, 8448-8452. | 3.3 | 60 |
| 2564 | Modulation of tRNA ^{Met} , eIF-2, and eIF-2B Expression Shows that GCN4 Translation Is Inversely Coupled to the Level of eIF-2·GTP·Met-tRNA ^{Met} Ternary Complexes. <i>Molecular and Cellular Biology</i> , 1995, 15, 6351-6363. | 1.1 | 126 |
| 2565 | Characterization of cis-Acting Sequences and Decay Intermediates Involved in Nonsense-Mediated mRNA Turnover. <i>Molecular and Cellular Biology</i> , 1995, 15, 809-823. | 1.1 | 144 |
| 2566 | Association of the vav proto-oncogene product with poly(rC)-specific RNA-binding proteins. <i>Molecular and Cellular Biology</i> , 1995, 15, 1324-1332. | 1.1 | 92 |
| 2567 | Role of the <i>Saccharomyces cerevisiae</i> general regulatory factor CP1 in methionine biosynthetic gene transcription. <i>Molecular and Cellular Biology</i> , 1995, 15, 1879-1888. | 1.1 | 53 |
| 2568 | The <i>Saccharomyces SHP1</i> Gene, Which Encodes a Regulator of Phosphoprotein Phosphatase 1 with Differential Effects on Glycogen Metabolism, Meiotic Differentiation, and Mitotic Cell Cycle Progression. <i>Molecular and Cellular Biology</i> , 1995, 15, 2037-2050. | 1.1 | 76 |
| 2569 | Functional Characterization of the MKC1 Gene of <i>Candida albicans</i> , Which Encodes a Mitogen-Activated Protein Kinase Homolog Related to Cell Integrity. <i>Molecular and Cellular Biology</i> , 1995, 15, 2197-2206. | 1.1 | 190 |
| 2570 | RAD1 and RAD10, but Not Other Excision Repair Genes, Are Required for Double-Strand Break-Induced Recombination in <i>Saccharomyces cerevisiae</i> . <i>Molecular and Cellular Biology</i> , 1995, 15, 2245-2251. | 1.1 | 233 |
| 2571 | Disturbance of Normal Cell Cycle Progression Enhances the Establishment of Transcriptional Silencing in <i>Saccharomyces cerevisiae</i> . <i>Molecular and Cellular Biology</i> , 1995, 15, 3608-3617. | 1.1 | 58 |
| 2572 | NPI-1, the human homolog of SRP-1, Interacts with influenza virus nucleoprotein. <i>Virology</i> , 1995, 206, 116-125. | 1.1 | 158 |
| 2573 | Development of metabolically engineered <i>Saccharomyces cerevisiae</i> cells for the production of lactic acid. <i>Biotechnology Progress</i> , 1995, 11, 294-298. | 1.3 | 114 |
| 2574 | Role of maltose permease and α -glucosidase in maltose fermentation and dough-leavening of <i>Torulasporea pretoriensis</i> YK-1. <i>Food Microbiology</i> , 1995, 12, 405-411. | 2.1 | 3 |

| # | ARTICLE | IF | CITATIONS |
|------|--|-----|-----------|
| 2575 | Human Dioxin Receptor Chimera Transactivation in a Yeast Model System and Studies on Receptor Agonists and Antagonists. <i>Basic and Clinical Pharmacology and Toxicology</i> , 1995, 76, 328-333. | 0.0 | 5 |
| 2576 | Inter-Species DNA Polymerase delta Chimeras are Functional in <i>Saccharomyces Cerevisiae</i> . <i>FEBS Journal</i> , 1995, 231, 45-49. | 0.2 | 3 |
| 2577 | Site-Directed Mutagenesis and Sugar-Binding Properties of the Wheat Germ Agglutinin Mutants Tyr73Phe and Phe116Tyr. <i>FEBS Journal</i> , 1995, 233, 27-34. | 0.2 | 10 |
| 2578 | Limitations to in vivo Import of Hydrophobic Proteins into Yeast Mitochondria. The Case of a Cytoplasmically Synthesized Apocytochrome b. <i>FEBS Journal</i> , 1995, 228, 762-771. | 0.2 | 141 |
| 2579 | Expression and Characterization of <i>Geotrichum candidum</i> Lipase I Gene. Comparison of Specificity Profile with Lipase II. <i>FEBS Journal</i> , 1995, 228, 863-869. | 0.2 | 42 |
| 2580 | The properties of the multicopy suppressor of theogdl mutation in yeast. <i>Journal of Basic Microbiology</i> , 1995, 35, 229-232. | 1.8 | 2 |
| 2581 | Deletion analysis of protein kinase c inactivation by calphostin C. <i>Molecular Carcinogenesis</i> , 1995, 12, 42-49. | 1.3 | 42 |
| 2582 | Differential protein kinase C ligand regulation detected in vivo by a phenotypic yeast assay. <i>Molecular Carcinogenesis</i> , 1995, 12, 166-176. | 1.3 | 32 |
| 2583 | A strategy for construction of industrial strains of distiller's yeast. <i>Biotechnology and Bioengineering</i> , 1995, 46, 285-290. | 1.7 | 17 |
| 2584 | Positive and negative feedback loops affect the transcription ofIME1, a positive regulator of meiosis in <i>Saccharomyces cerevisiae</i> . <i>Genesis</i> , 1995, 16, 219-228. | 3.1 | 28 |
| 2585 | ?-Tubulin mutation suppresses microtubule dynamics in vitro and slows mitosis in vivo. <i>Cytoskeleton</i> , 1995, 30, 285-300. | 4.4 | 23 |
| 2586 | Regulation of cell wall β -glucan assembly:PTC1 Negatively affectsPBS2 Action in a pathway that includes modulation ofEXG1 transcription. <i>Molecular Genetics and Genomics</i> , 1995, 248, 260-269. | 2.4 | 97 |
| 2587 | A temperature-sensitive lambda cl repressor functions on a modified operator in yeast cells by masking the TATA element. <i>Molecular Genetics and Genomics</i> , 1995, 248, 499-505. | 2.4 | 3 |
| 2588 | MMK2, a novel alfalfa MAP kinase, specifically complements the yeast MPK1 function. <i>Molecular Genetics and Genomics</i> , 1995, 248, 686-694. | 2.4 | 61 |
| 2589 | Dosage suppressors of the dominant G1 cyclin mutantCLN3-2: Identification of a yeast gene encoding a putative RNA/ssDNA binding protein. <i>Molecular Genetics and Genomics</i> , 1995, 248, 712-718. | 2.4 | 17 |
| 2590 | Stable expression of vicilin from <i>Vicia faba</i> with eight additional single methionine residues but failure of accumulation of legumin with an attached peptide segment in tobacco seeds. <i>Molecular Breeding</i> , 1995, 1, 245-258. | 1.0 | 18 |
| 2591 | Characteristics comparison between plasmid-harboring and chromosome-integrated recombinant <i>Saccharomyces ceremsiae</i> cultures. <i>Korean Journal of Chemical Engineering</i> , 1995, 12, 567-571. | 1.2 | 4 |
| 2592 | Industrial mycology and the new genetics. <i>Journal of Industrial Microbiology</i> , 1995, 14, 355-364. | 0.9 | 8 |

| # | ARTICLE | IF | CITATIONS |
|------|---|-----|-----------|
| 2593 | A complex interplay of positive and negative elements is responsible for the different transcriptional activity of liver NF1 variants. <i>Molecular Biology Reports</i> , 1995, 21, 147-158. | 1.0 | 7 |
| 2594 | The role of <i>Saccharomyces cerevisiae</i> Cdc40p in DNA replication and mitotic spindle formation and/or maintenance. <i>Molecular Genetics and Genomics</i> , 1995, 247, 123-136. | 2.4 | 46 |
| 2595 | Gain-of-function mutations in a human calmodulin-like protein identify residues critical for calmodulin action in yeast. <i>Molecular Genetics and Genomics</i> , 1995, 247, 137-147. | 2.4 | 8 |
| 2596 | Isolation and characterization of temperature-sensitive <i>plc1</i> mutants of the yeast <i>Saccharomyces cerevisiae</i> . <i>Molecular Genetics and Genomics</i> , 1995, 247, 148-156. | 2.4 | 26 |
| 2597 | Expression of arginine decarboxylase is induced during early fruit development and in young tissues of <i>Pisum sativum</i> (L.). <i>Plant Molecular Biology</i> , 1995, 28, 997-1009. | 2.0 | 81 |
| 2598 | Inactivation of the UAS1 of STA1 by glucose and STA10 and identification of two loci, SNS1 and MSS1, involved in STA10-dependent repression in <i>Saccharomyces cerevisiae</i> . <i>Molecular Genetics and Genomics</i> , 1995, 246, 529-537. | 2.4 | 15 |
| 2599 | Cis and trans-acting regulatory elements required for regulation of the CPS1 gene in <i>Saccharomyces cerevisiae</i> . <i>Molecular Genetics and Genomics</i> , 1995, 246, 580-589. | 2.4 | 8 |
| 2600 | RAD58 (XRS4)? A new gene in the RAD52 epistasis group. <i>Current Genetics</i> , 1995, 28, 274-279. | 0.8 | 20 |
| 2601 | NCA3, a nuclear gene involved in the mitochondrial expression of subunits 6 and 8 of the Fo-F1 ATP synthase of <i>S. cerevisiae</i> . <i>Current Genetics</i> , 1995, 27, 409-416. | 0.8 | 58 |
| 2602 | A mutant allele of the SUP45 (SAL4) gene of <i>Saccharomyces cerevisiae</i> shows temperature-dependent allosuppressor and omnipotent suppressor phenotypes. <i>Current Genetics</i> , 1995, 27, 417-426. | 0.8 | 75 |
| 2603 | Cloning and sequencing of cellulase cDNA from <i>Aspergillus kawachii</i> and its expression in <i>Saccharomyces cerevisiae</i> . <i>Current Genetics</i> , 1995, 27, 435-439. | 0.8 | 33 |
| 2604 | Molecular cloning of the meiosis-induced <i>rec10</i> gene of <i>Schizosaccharomyces pombe</i> . <i>Current Genetics</i> , 1995, 27, 440-446. | 0.8 | 31 |
| 2605 | Mating-type suppression of the DNA-repair defect of the yeast <i>rad6?</i> mutation requires the activity of genes in the RAD52 epistasis group. <i>Current Genetics</i> , 1995, 28, 12-18. | 0.8 | 18 |
| 2606 | The evolutionary relationships between homologs of ribosomal YL8 protein and YL8-like proteins. <i>Current Genetics</i> , 1995, 28, 19-25. | 0.8 | 10 |
| 2607 | Molecular genetic properties of the yeast <i>Torulaspota pretoriensis</i> : Characterization of chromosomal DNA and genetic transformation by <i>Saccharomyces cerevisiae</i> -based plasmids. <i>Current Genetics</i> , 1995, 27, 131-134. | 0.8 | 11 |
| 2608 | Molecular analysis of the yeast SER1 gene encoding 3-phosphoserine aminotransferase: regulation by general control and serine repression. <i>Current Genetics</i> , 1995, 27, 501-508. | 0.8 | 25 |
| 2609 | A <i>Kluyveromyces lactis</i> gene homologue to AAC2 complements the <i>Saccaromyces cerevisiae</i> <i>op1</i> mutation. <i>Current Genetics</i> , 1995, 27, 229-233. | 0.8 | 19 |
| 2610 | Identification of the heterothallic mutation in HO-endonuclease of <i>S. cerevisiae</i> using HO/ho chimeric genes. <i>Current Genetics</i> , 1995, 28, 367-373. | 0.8 | 41 |

| # | ARTICLE | IF | CITATIONS |
|------|---|-----|-----------|
| 2611 | Construction of a marker gene cassette which is repeatedly usable for gene disruption in yeast. <i>Current Genetics</i> , 1995, 27, 293-297. | 0.8 | 31 |
| 2612 | Strategy for deletion of complete open reading frames in <i>Saccharomyces cerevisiae</i> . <i>Current Genetics</i> , 1995, 27, 306-308. | 0.8 | 36 |
| 2613 | DNA insertion system for complex yeast shuttle vectors. <i>Current Genetics</i> , 1995, 27, 309-311. | 0.8 | 9 |
| 2614 | Molecular cloning and characterization of a novel gene of <i>Candida albicans</i> , CDR1, conferring multiple resistance to drugs and antifungals. <i>Current Genetics</i> , 1995, 27, 320-329. | 0.8 | 475 |
| 2615 | Molecular and biochemical characterization of the hexokinase from the starch-utilizing yeast <i>Schwanniomyces occidentalis</i> . <i>Current Genetics</i> , 1995, 27, 330-338. | 0.8 | 21 |
| 2616 | Expression of a yeast gene can be blocked by insertion of short yeast DNA fragments between a UAS and the TATA box. <i>Current Genetics</i> , 1995, 27, 387-389. | 0.8 | 9 |
| 2617 | Heat shock protein HSP60 can alleviate the phenotype of mitochondrial RNA-deficient temperature-sensitive <i>mna2</i> pet mutants. <i>Molecular Genetics and Genomics</i> , 1995, 246, 56-64. | 2.4 | 15 |
| 2618 | Chromosome polymorphisms close to the <i>cm-ADE1</i> locus of <i>Candida maltosa</i> . <i>Molecular Genetics and Genomics</i> , 1995, 247, 591-602. | 2.4 | 7 |
| 2619 | Molecular characterization and heterologous expression of an endo- β -1,6-glucanase gene from the mycoparasitic fungus <i>Trichoderma harzianum</i> . <i>Molecular Genetics and Genomics</i> , 1995, 247, 639-645. | 2.4 | 66 |
| 2620 | Cloning and characterization of seven cDNAs for hyperosmolarity-responsive (HOR) genes of <i>Saccharomyces cerevisiae</i> . <i>Molecular Genetics and Genomics</i> , 1995, 249, 127-138. | 2.4 | 103 |
| 2621 | A large circular minichromosome of <i>Schizosaccharomyces pombe</i> requires a high dose of type II DNA topoisomerase for its stabilization. <i>Molecular Genetics and Genomics</i> , 1995, 246, 671-679. | 2.4 | 21 |
| 2622 | Genetic evidence for the interaction of the yeast transcriptional co-activator proteins GCN5 and ADA2. <i>Molecular Genetics and Genomics</i> , 1995, 246, 723-728. | 2.4 | 65 |
| 2623 | Isolation and characterization of extragenic mutations affecting the expression of the uroporphyrinogen decarboxylase gene (<i>HEM12</i>) in <i>Saccharomyces cerevisiae</i> . <i>Molecular Genetics and Genomics</i> , 1995, 247, 471-481. | 2.4 | 8 |
| 2624 | Cloning and expression of a cDNA copy of the viral K28 killer toxin gene in yeast. <i>Molecular Genetics and Genomics</i> , 1995, 246, 236-246. | 2.4 | 32 |
| 2625 | Evidence that two zinc fingers in the methionine aminopeptidase from <i>Saccharomyces cerevisiae</i> are important for normal growth. <i>Molecular Genetics and Genomics</i> , 1995, 246, 247-253. | 2.4 | 57 |
| 2626 | Identification and characterization of <i>CEN12</i> in the budding yeast <i>Saccharomyces cerevisiae</i> . <i>Current Genetics</i> , 1995, 28, 512-516. | 0.8 | 3 |
| 2627 | Inactivation of the yeast <i>Sen1</i> protein affects the localization of nucleolar proteins. <i>Molecular Genetics and Genomics</i> , 1995, 249, 571-584. | 2.4 | 40 |
| 2628 | Constitutive activation of the <i>Saccharomyces cerevisiae</i> mating response pathway by a MAP kinase kinase from <i>Candida albicans</i> . <i>Molecular Genetics and Genomics</i> , 1995, 249, 609-621. | 2.4 | 40 |

| # | ARTICLE | IF | CITATIONS |
|------|--|-----|-----------|
| 2629 | Mode of action of the <i>gcr9</i> and <i>cat3</i> mutations in restoring the ability of <i>Saccharomyces cerevisiae</i> <i>tps1</i> mutants to grow on glucose. <i>Molecular Genetics and Genomics</i> , 1995, 249, 655-664. | 2.4 | 23 |
| 2630 | The glucose repression and RAS-cAMP signal transduction pathways of <i>Saccharomyces cerevisiae</i> each affect RNA processing and the synthesis of a reporter protein. <i>Molecular Genetics and Genomics</i> , 1995, 247, 48-54. | 2.4 | 18 |
| 2631 | Influence of non-homology between recombining DNA sequences on double-strand break repair in <i>Saccharomyces cerevisiae</i> . <i>Molecular Genetics and Genomics</i> , 1995, 247, 55-60. | 2.4 | 4 |
| 2632 | Molecular cloning of the <i>plc1+</i> gene of <i>Schizosaccharomyces pombe</i> , which encodes a putative phosphoinositide-specific phospholipase C. <i>Yeast</i> , 1995, 11, 179-185. | 0.8 | 27 |
| 2633 | Localization of a protein a-tagged <i>kex2</i> protein to the vacuole of <i>Saccharomyces cerevisiae</i> allows rapid purification of vacuolar membranes. <i>Yeast</i> , 1995, 11, 201-210. | 0.8 | 9 |
| 2634 | Molecular analysis of the <i>SNF8</i> gene of <i>Saccharomyces cerevisiae</i> . <i>Yeast</i> , 1995, 11, 219-224. | 0.8 | 22 |
| 2635 | Genetic and carbon source regulation of phosphorylation of <i>Sip1p</i> , a <i>Snf1p</i> -associated protein involved in carbon response in <i>Saccharomyces cerevisiae</i> . <i>Yeast</i> , 1995, 11, 233-246. | 0.8 | 4 |
| 2636 | <i>UASNTR</i> functioning in combination with other <i>UAS</i> elements underlies exceptional patterns of nitrogen regulation in <i>Saccharomyces cerevisiae</i> . <i>Yeast</i> , 1995, 11, 247-260. | 0.8 | 30 |
| 2637 | Plasmid reorganization during integrative transformation in <i>Hansenula polymorpha</i> . <i>Yeast</i> , 1995, 11, 343-353. | 0.8 | 59 |
| 2638 | Studies on the transformation of intact yeast cells by the LiAc/SS-DNA/PEG procedure. <i>Yeast</i> , 1995, 11, 355-360. | 0.8 | 1,939 |
| 2639 | Nucleotide sequence and characterization of the <i>Saccharomyces cerevisiae</i> <i>RPL19A</i> gene encoding a homolog of the mammalian ribosomal protein l19. <i>Yeast</i> , 1995, 11, 383-389. | 0.8 | 5 |
| 2640 | Induced expression of bacterial β -glucosidase activity in <i>Saccharomyces</i> . <i>Yeast</i> , 1995, 11, 395-406. | 0.8 | 35 |
| 2641 | Transcriptional regulation of flocculation genes in <i>Saccharomyces cerevisiae</i> . <i>Yeast</i> , 1995, 11, 435-446. | 0.8 | 86 |
| 2642 | <i>Saccharomyces cerevisiae</i> mRNA 3' end forming signals are also involved in transcription termination. <i>Yeast</i> , 1995, 11, 447-453. | 0.8 | 26 |
| 2643 | Host cell properties and external pH affect proinsulin production by <i>Saccharomyces</i> yeast. <i>Yeast</i> , 1995, 11, 713-724. | 0.8 | 11 |
| 2644 | A regulated <i>MET3-GLC7</i> gene fusion provides evidence of a mitotic role for <i>Saccharomyces cerevisiae</i> protein phosphatase 1. <i>Yeast</i> , 1995, 11, 747-759. | 0.8 | 43 |
| 2645 | <i>Scp160p</i> , a new yeast protein associated with the nuclear membrane and the endoplasmic reticulum, is necessary for maintenance of exact ploidy. <i>Yeast</i> , 1995, 11, 929-944. | 0.8 | 70 |
| 2646 | Molecular cloning of the GTP-cyclohydrolase structural gene <i>RIB1</i> of <i>Pichia guilliermondii</i> involved in riboflavin biosynthesis. <i>Yeast</i> , 1995, 11, 945-952. | 0.8 | 20 |

| # | ARTICLE | IF | CITATIONS |
|------|--|-----|-----------|
| 2647 | Isolation and characterization of a novel yeast gene,ATH1, that is required for vacuolar acid trehalase activity. <i>Yeast</i> , 1995, 11, 1015-1025. | 0.8 | 72 |
| 2648 | Transcriptional regulation by an upstream repression sequence from the yeast enolase geneENO1. <i>Yeast</i> , 1995, 11, 1031-1043. | 0.8 | 9 |
| 2649 | UVS112â€”A gene involved in excision repair of yeast. <i>Yeast</i> , 1995, 11, 1129-1138. | 0.8 | 5 |
| 2650 | Molecular characterization of thePEL1 gene encoding a putative phosphatidylserine synthase. <i>Yeast</i> , 1995, 11, 1223-1231. | 0.8 | 27 |
| 2651 | DOGR1 andDOGR2: Two genes from <i>Saccharomyces cerevisiae</i> that confer 2-deoxyglucose resistance when overexpressed. <i>Yeast</i> , 1995, 11, 1233-1240. | 0.8 | 46 |
| 2652 | A disruption-replacement approach for the targeted integration of foreign genes in <i>Hansenula polymorpha</i> . <i>Yeast</i> , 1995, 11, 1241-1247. | 0.8 | 39 |
| 2653 | XV. Yeast sequencing reports. Nucleotide sequence of theGDS1 gene of <i>Saccharomyces cerevisiae</i> . <i>Yeast</i> , 1995, 11, 1513-1518. | 0.8 | 4 |
| 2654 | Cold-shock induction of a family of TIP1-related proteins associated with the membrane in <i>Saccharomyces cerevisiae</i> . <i>Molecular Microbiology</i> , 1995, 15, 341-353. | 1.2 | 87 |
| 2655 | The role of the YAP1 and YAP2 genes in the regulation of the adaptive oxidative stress responses of <i>Saccharomyces cerevisiae</i> . <i>Molecular Microbiology</i> , 1995, 16, 415-423. | 1.2 | 198 |
| 2656 | Functional analysis of mutations in thePISgene, which encodes <i>Saccharomyces cerevisiae</i> phosphatidylinositol synthase. <i>FEMS Microbiology Letters</i> , 1995, 126, 81-84. | 0.7 | 9 |
| 2657 | Functional role of <i>Aspergillus oryzae</i> glucoamylase C-terminal domain investigated using its cDNA. <i>Journal of Bioscience and Bioengineering</i> , 1995, 80, 280-282. | 0.9 | 6 |
| 2658 | Development of an integrative DNA transformation system for the yeast <i>Hansenula anomala</i> . <i>Journal of Bioscience and Bioengineering</i> , 1995, 79, 1-5. | 0.9 | 11 |
| 2659 | Effects of elevated expression of the CYP51 (P45014DM) gene on the sterol contents of <i>Saccharomyces cerevisiae</i> . <i>Journal of Bioscience and Bioengineering</i> , 1995, 79, 64-66. | 0.9 | 2 |
| 2660 | Secretive expression of the <i>Aspergillus aculeatus</i> cellulase (FI-CMCase) by <i>Saccharomyces cerevisiae</i> . <i>Journal of Bioscience and Bioengineering</i> , 1995, 79, 363-366. | 0.9 | 17 |
| 2661 | Effect of yeast fumarase gene (FUM1) disruption on production of malic, fumaric and succinic acids in sake mash. <i>Journal of Bioscience and Bioengineering</i> , 1995, 80, 355-361. | 0.9 | 32 |
| 2662 | Production of HM-1 killer toxin in <i>Saccharomyces cerevisiae</i> transformed with the PDR4 gene and Î—-sequence-mediated multi-integration system. <i>Journal of Bioscience and Bioengineering</i> , 1995, 80, 423-428. | 0.9 | 11 |
| 2663 | A high dose of theSTM1 gene suppresses the temperature sensitivity of thetom1 andhtr1 mutants in <i>Saccharomyces cerevisiae</i> . <i>Biochimica Et Biophysica Acta Gene Regulatory Mechanisms</i> , 1995, 1263, 285-288. | 2.4 | 24 |
| 2664 | Diminished activity of the first N-glycosylation enzyme, dolichol-P-dependent N-acetylglucosamine-1-P transferase (GPT), gives rise to mutant phenotypes in yeast. <i>BBA - Proteins and Proteomics</i> , 1995, 1247, 51-59. | 2.1 | 34 |

| # | ARTICLE | IF | CITATIONS |
|------|--|-----|-----------|
| 2665 | Preparation by site-directed mutagenesis and characterization of the E211Q mutant of yeast enolase 1. BBA - Proteins and Proteomics, 1995, 1251, 23-31. | 2.1 | 15 |
| 2666 | Molecular cloning and nucleotide sequence of the 1,2- α -D-mannosidase gene, <i>msdS</i> , from <i>Aspergillus saitoi</i> and expression of the gene in yeast cells. BBA - Proteins and Proteomics, 1995, 1253, 141-145. | 2.1 | 40 |
| 2667 | A Potyvirus Polymerase Interacts with the Viral Coat Protein and VPg in Yeast Cells. Virology, 1995, 214, 159-166. | 1.1 | 92 |
| 2668 | Rapid transfer of low copy-number episomal plasmids from <i>Saccharomyces cerevisiae</i> to <i>Escherichia coli</i> by electroporation. Molecular Biotechnology, 1995, 3, 79-84. | 1.3 | 15 |
| 2669 | A method for rapid mapping of mutations by plasmid rescue strategy in <i>Saccharomyces cerevisiae</i> . Journal of Genetics, 1995, 74, 127-138. | 0.4 | 1 |
| 2670 | A downstream target of RHO1 small GTP-binding protein is PKC1, a homolog of protein kinase C, which leads to activation of the MAP kinase cascade in <i>Saccharomyces cerevisiae</i> .. EMBO Journal, 1995, 14, 5931-5938. | 3.5 | 303 |
| 2671 | Molecular cloning of the plasma membrane H(+)-ATPase from <i>Kluyveromyces lactis</i> : a single nucleotide substitution in the gene confers ethidium bromide resistance and deficiency in K ⁺ uptake. Journal of Bacteriology, 1995, 177, 2360-2367. | 1.0 | 21 |
| 2672 | REG1 binds to protein phosphatase type 1 and regulates glucose repression in <i>Saccharomyces cerevisiae</i> .. EMBO Journal, 1995, 14, 5939-5946. | 3.5 | 207 |
| 2673 | Functionality and specific membrane localization of transport GTPases carrying C-terminal membrane anchors of synaptobrevin-like proteins.. EMBO Journal, 1995, 14, 3645-3653. | 3.5 | 41 |
| 2674 | Recognition of nitrogen-responsive upstream activation sequences of <i>Saccharomyces cerevisiae</i> by the product of the <i>GLN3</i> gene. Journal of Bacteriology, 1995, 177, 4190-4193. | 1.0 | 69 |
| 2675 | <i>bfr1+</i> , a novel gene of <i>Schizosaccharomyces pombe</i> which confers brefeldin A resistance, is structurally related to the ATP-binding cassette superfamily. Journal of Bacteriology, 1995, 177, 1536-1543. | 1.0 | 67 |
| 2676 | Mutations in <i>GCD11</i> , the structural gene for eIF-2 gamma in yeast, alter translational regulation of <i>GCN4</i> and the selection of the start site for protein synthesis.. EMBO Journal, 1995, 14, 2239-2249. | 3.5 | 60 |
| 2677 | <i>Nin1p</i> , a regulatory subunit of the 26S proteasome, is necessary for activation of <i>Cdc28p</i> kinase of <i>Saccharomyces cerevisiae</i> .. EMBO Journal, 1995, 14, 3105-3115. | 3.5 | 88 |
| 2678 | Translation and M1 double-stranded RNA propagation: MAK18 = RPL41B and cycloheximide curing. Journal of Bacteriology, 1995, 177, 2887-2891. | 1.0 | 38 |
| 2679 | Structurally Related but Functionally Distinct Yeast Sm D Core Small Nuclear Ribonucleoprotein Particle Proteins. Molecular and Cellular Biology, 1995, 15, 445-455. | 1.1 | 66 |
| 2680 | Pulling the Ribosome out of Frame by 11 at a Programmed Frameshift Site by Cognate Binding of Aminoacyl-tRNA. Molecular and Cellular Biology, 1995, 15, 298-304. | 1.1 | 56 |
| 2681 | Sm and Sm-like proteins belong to a large family: identification of proteins of the U6 as well as the U1, U2, U4 and U5 snRNPs.. EMBO Journal, 1995, 14, 2089-2098. | 3.5 | 250 |
| 2682 | Transcriptional and posttranslational regulation of the general amino acid permease of <i>Saccharomyces cerevisiae</i> . Journal of Bacteriology, 1995, 177, 94-102. | 1.0 | 166 |

| # | ARTICLE | IF | CITATIONS |
|------|---|-----|-----------|
| 2683 | Improvement of Maltose Fermentation Efficiency: Constitutive Expression of MAL Genes in Brewing Yeasts. <i>Journal of the American Society of Brewing Chemists</i> , 1995, 53, 24-29. | 0.8 | 65 |
| 2684 | Identification of the cis-elements mediating the autogenous control of ribosomal protein L2 mRNA stability in yeast.. <i>EMBO Journal</i> , 1995, 14, 4022-4030. | 3.5 | 26 |
| 2685 | Genetic evidence for Gln3p-independent, nitrogen catabolite repression-sensitive gene expression in <i>Saccharomyces cerevisiae</i> . <i>Journal of Bacteriology</i> , 1995, 177, 6910-6918. | 1.0 | 67 |
| 2686 | <i>Schizosaccharomyces pombe atf1+</i> encodes a transcription factor required for sexual development and entry into stationary phase.. <i>EMBO Journal</i> , 1995, 14, 6193-6208. | 3.5 | 229 |
| 2687 | The bovine papillomavirus 1 E2 protein contains two activation domains: one that interacts with TBP and another that functions after TBP binding.. <i>EMBO Journal</i> , 1995, 14, 329-340. | 3.5 | 67 |
| 2688 | GCN20, a novel ATP binding cassette protein, and GCN1 reside in a complex that mediates activation of the eIF-2 alpha kinase GCN2 in amino acid-starved cells.. <i>EMBO Journal</i> , 1995, 14, 3184-3199. | 3.5 | 155 |
| 2689 | A novel protein kinase gene <i>ssp1+</i> is required for alteration of growth polarity and actin localization in fission yeast.. <i>EMBO Journal</i> , 1995, 14, 3325-3338. | 3.5 | 57 |
| 2690 | The <i>Saccharomyces cerevisiae</i> translation initiation factor Tif3 and its mammalian homologue, eIF-4B, have RNA annealing activity.. <i>EMBO Journal</i> , 1995, 14, 3820-3827. | 3.5 | 89 |
| 2691 | Characterization of Physical Interactions of the Putative Transcriptional Adaptor, ADA2, with Acidic Activation Domains and TATA-binding Protein. <i>Journal of Biological Chemistry</i> , 1995, 270, 19337-19344. | 1.6 | 174 |
| 2692 | Termination of translation in eukaryotes is governed by two interacting polypeptide chain release factors, eRF1 and eRF3.. <i>EMBO Journal</i> , 1995, 14, 4065-4072. | 3.5 | 523 |
| 2693 | Control of Vicinal Diketone Production by Brewer's Yeast. I. Effects of <i>ILV5</i> and <i>ILV3</i> Gene Amplification on Vicinal Diketone Production and <i>ILV</i> Enzyme Activity. <i>Journal of the American Society of Brewing Chemists</i> , 1995, 53, 49-53. | 0.8 | 21 |
| 2694 | A Direct Interaction between G-Protein $\beta\gamma$ Subunits and the Raf-1 Protein Kinase. <i>Journal of Biological Chemistry</i> , 1995, 270, 14251-14254. | 1.6 | 74 |
| 2695 | Different Repair Kinetics for Short and Long DNA Double-strand Gaps in <i>Saccharomyces Cerevisiae</i> . <i>International Journal of Radiation Biology</i> , 1995, 68, 421-428. | 1.0 | 2 |
| 2696 | Characterization of Rat Neuronal Nitric Oxide Synthase Expressed in <i>Saccharomyces cerevisiae</i> . <i>DNA and Cell Biology</i> , 1995, 14, 789-794. | 0.9 | 18 |
| 2697 | Protein O-Glycosylation in Yeast. <i>Journal of Biological Chemistry</i> , 1995, 270, 2770-2775. | 1.6 | 88 |
| 2698 | A bovine cDNA and a yeast gene (<i>VMA8</i>) encoding the subunit D of the vacuolar H(+)-ATPase.. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 1995, 92, 497-501. | 3.3 | 82 |
| 2699 | Zip1-induced changes in synaptonemal complex structure and polycomplex assembly.. <i>Journal of Cell Biology</i> , 1995, 128, 455-466. | 2.3 | 174 |
| 2700 | C-terminal deletion analysis of plant plasma membrane H(+)-ATPase: yeast as a model system for solute transport across the plant plasma membrane.. <i>Plant Cell</i> , 1995, 7, 1655-1666. | 3.1 | 121 |

| # | ARTICLE | IF | CITATIONS |
|------|---|-----|-----------|
| 2701 | The origin recognition complex in silencing, cell cycle progression, and DNA replication.. Molecular Biology of the Cell, 1995, 6, 741-756. | 0.9 | 204 |
| 2702 | The alpha subunit of the Saccharomyces cerevisiae oligosaccharyltransferase complex is essential for vegetative growth of yeast and is homologous to mammalian ribophorin I.. Journal of Cell Biology, 1995, 128, 525-536. | 2.3 | 67 |
| 2703 | Hormone-dependent Transactivation by the Human Androgen Receptor Is Regulated by a dnaJ Protein. Journal of Biological Chemistry, 1995, 270, 5251-5257. | 1.6 | 123 |
| 2704 | Role of Bud3p in producing the axial budding pattern of yeast.. Journal of Cell Biology, 1995, 129, 767-778. | 2.3 | 188 |
| 2705 | The Cellular Content of Cdc25p, the Ras Exchange Factor in Saccharomyces cerevisiae, Is Regulated by Destabilization Through a Cyclin Destruction Box. Journal of Biological Chemistry, 1995, 270, 20742-20747. | 1.6 | 37 |
| 2706 | Amino acid substitutions in the structured domains of histones H3 and H4 partially relieve the requirement of the yeast SWI/SNF complex for transcription.. Genes and Development, 1995, 9, 2770-2779. | 2.7 | 241 |
| 2707 | Proteolysis Prevents in Vivo Chimeric Fusion Protein Import into Yeast Mitochondria.. Journal of Biological Chemistry, 1995, 270, 16689-16693. | 1.6 | 12 |
| 2708 | Histone H4 and the maintenance of genome integrity.. Genes and Development, 1995, 9, 1716-1727. | 2.7 | 153 |
| 2709 | The Saccharomyces cerevisiae RIB4 Gene Codes for 6,7-Dimethyl- 8-ribityllumazine Synthase Involved in Riboflavin Biosynthesis. Journal of Biological Chemistry, 1995, 270, 23801-23807. | 1.6 | 44 |
| 2710 | Expression and Function of a Mislocalized Form of Peroxisomal Malate Dehydrogenase (MDH3) in Yeast. Journal of Biological Chemistry, 1995, 270, 21220-21225. | 1.6 | 28 |
| 2711 | Sec6, Sec8, and Sec15 are components of a multisubunit complex which localizes to small bud tips in Saccharomyces cerevisiae.. Journal of Cell Biology, 1995, 130, 299-312. | 2.3 | 287 |
| 2712 | Partial Truncation of the Yeast RNA Polymerase II Carboxyl-terminal Domain Preferentially Reduces Expression of Glycolytic Genes. Journal of Biological Chemistry, 1995, 270, 31255-31261. | 1.6 | 14 |
| 2713 | Functional Complementation of a Null Mutation of the Yeast Saccharomyces cerevisiae Plasma Membrane H ⁺ -ATPase by a Plant H ⁺ -ATPase Gene. Journal of Biological Chemistry, 1995, 270, 23828-23837. | 1.6 | 84 |
| 2714 | Random Substitution of Large Parts of the Propeptide of Yeast Proteinase A. Journal of Biological Chemistry, 1995, 270, 8602-8609. | 1.6 | 28 |
| 2715 | The majority of yeast UPF1 co-localizes with polyribosomes in the cytoplasm.. Molecular Biology of the Cell, 1995, 6, 611-625. | 0.9 | 126 |
| 2716 | The Saccharomyces cerevisiae VMA10 Is an Intron-containing Gene Encoding a Novel 13-kDa Subunit of Vacuolar H ⁺ -ATPase. Journal of Biological Chemistry, 1995, 270, 13726-13732. | 1.6 | 90 |
| 2717 | A Camptothecin-resistant DNA Topoisomerase I Mutant Exhibits Altered Sensitivities to Other DNA Topoisomerase Poisons. Journal of Biological Chemistry, 1995, 270, 6141-6148. | 1.6 | 47 |
| 2718 | Cloning and Expression of a Yeast Gene Encoding a Protein with ATPase Activity and High Identity to the Subunit 4 of the Human 26 S Protease. Journal of Biological Chemistry, 1995, 270, 9178-9184. | 1.6 | 19 |

| # | ARTICLE | IF | CITATIONS |
|------|--|-----|-----------|
| 2719 | Multidrug resistance in <i>Candida albicans</i> : disruption of the BENr gene. <i>Antimicrobial Agents and Chemotherapy</i> , 1995, 39, 422-426. | 1.4 | 91 |
| 2720 | The protein kinase C-activated MAP kinase pathway of <i>Saccharomyces cerevisiae</i> mediates a novel aspect of the heat shock response.. <i>Genes and Development</i> , 1995, 9, 1559-1571. | 2.7 | 459 |
| 2721 | Integrative modification of YAC clones in a 96-well microplate format. <i>Nucleic Acids Research</i> , 1995, 23, 1830-1831. | 6.5 | 1 |
| 2722 | Cis- and trans-acting elements determining induction of the genes of the \hat{I}^3 -aminobutyrate (GABA) utilization pathway in <i>Saccharomyces cerevisiae</i> . <i>Nucleic Acids Research</i> , 1995, 23, 550-557. | 6.5 | 59 |
| 2723 | The essential OST2 gene encodes the 16-kD subunit of the yeast oligosaccharyltransferase, a highly conserved protein expressed in diverse eukaryotic organisms.. <i>Journal of Cell Biology</i> , 1995, 131, 371-383. | 2.3 | 113 |
| 2724 | Mutational Analysis of <i>Saccharomyces cerevisiae</i> ARF1. <i>Journal of Biological Chemistry</i> , 1995, 270, 143-150. | 1.6 | 81 |
| 2725 | Functional Dissection of P-glycoprotein Nucleotide-binding Domains in Chimeric and Mutant Proteins. <i>Journal of Biological Chemistry</i> , 1995, 270, 17159-17170. | 1.6 | 76 |
| 2726 | Dual Retinoblastoma-binding Proteins with Properties Related to a Negative Regulator of Ras in Yeast. <i>Journal of Biological Chemistry</i> , 1995, 270, 25507-25513. | 1.6 | 125 |
| 2727 | Microtubule stability in budding yeast: characterization and dosage suppression of a benomyl-dependent tubulin mutant.. <i>Molecular Biology of the Cell</i> , 1995, 6, 1241-1259. | 0.9 | 49 |
| 2728 | The Protooncogene c-jun Contains an Unusual Estrogen-inducible Enhancer within the Coding Sequence. <i>Journal of Biological Chemistry</i> , 1995, 270, 8506-8513. | 1.6 | 83 |
| 2729 | GCD10, a translational repressor of GCN4, is the RNA-binding subunit of eukaryotic translation initiation factor-3.. <i>Genes and Development</i> , 1995, 9, 1781-1796. | 2.7 | 70 |
| 2730 | A Mutation in an ATP-binding Loop of <i>Saccharomyces cerevisiae</i> Actin (S14A) Causes a Temperature-sensitive Phenotype in Vivo and in Vitro. <i>Journal of Biological Chemistry</i> , 1995, 270, 11406-11414. | 1.6 | 42 |
| 2731 | Human RNA polymerase II subunit hsRPB7 functions in yeast and influences stress survival and cell morphology.. <i>Molecular Biology of the Cell</i> , 1995, 6, 759-775. | 0.9 | 71 |
| 2732 | The <i>GCR1</i> Requirement for Yeast Glycolytic Gene Expression Is Suppressed by Dominant Mutations in the <i>SGC1</i> Gene, Which Encodes a Novel Basic-Helix-Loop-Helix Protein. <i>Molecular and Cellular Biology</i> , 1995, 15, 2646-2653. | 1.1 | 48 |
| 2733 | Expression of an ATP-Binding Cassette Transporter-Encoding Gene (<i>YOR1</i>) Is Required for Oligomycin Resistance in <i>Saccharomyces cerevisiae</i> . <i>Molecular and Cellular Biology</i> , 1995, 15, 6875-6883. | 1.1 | 210 |
| 2734 | AH/PH Domain-Mediated Interaction between Akt Molecules and Its Potential Role in Akt Regulation. <i>Molecular and Cellular Biology</i> , 1995, 15, 2304-2310. | 1.1 | 160 |
| 2735 | Regulation of Cation Transport in <i>Saccharomyces cerevisiae</i> by the Salt Tolerance Gene <i>HAL3</i> . <i>Molecular and Cellular Biology</i> , 1995, 15, 5470-5481. | 1.1 | 159 |
| 2736 | Yeast N-terminal Amidase. <i>Journal of Biological Chemistry</i> , 1995, 270, 12065-12074. | 1.6 | 90 |

| # | ARTICLE | IF | CITATIONS |
|------|---|-----|-----------|
| 2737 | Successful transformation of yeast mitochondria with RPM1: an approach for in vivo studies of mitochondrial RNase P RNA structure, function and biosynthesis. <i>Nucleic Acids Research</i> , 1995, 23, 856-860. | 6.5 | 15 |
| 2738 | Novel PI(4)P 5-kinase homologue, Fab1p, essential for normal vacuole function and morphology in yeast.. <i>Molecular Biology of the Cell</i> , 1995, 6, 525-539. | 0.9 | 283 |
| 2739 | gar2 is a nucleolar protein from <i>Schizosaccharomyces pombe</i> required for 18S rRNA and 40S ribosomal subunit accumulation. <i>Nucleic Acids Research</i> , 1995, 23, 1912-1918. | 6.5 | 60 |
| 2740 | Analysis of Positive Elements Sensitive to Glucose in the Promoter of the FBP1 Gene from Yeast. <i>Journal of Biological Chemistry</i> , 1995, 270, 12832-12838. | 1.6 | 25 |
| 2741 | Site-directed Mutagenesis of the Yeast PMA1 H ⁺ -ATPase STRUCTURAL AND FUNCTIONAL ROLE OF CYSTEINE RESIDUES. <i>Journal of Biological Chemistry</i> , 1995, 270, 28535-28540. | 1.6 | 39 |
| 2742 | Functional characterization of the 180-kD ribosome receptor in vivo.. <i>Journal of Cell Biology</i> , 1995, 130, 29-39. | 2.3 | 84 |
| 2743 | The cytoplasmic tail domain of the vacuolar protein sorting receptor Vps10p and a subset of VPS gene products regulate receptor stability, function, and localization.. <i>Molecular Biology of the Cell</i> , 1995, 6, 1089-1102. | 0.9 | 151 |
| 2744 | A recombinatorial method useful for cloning dominant alleles in <i>Saccharomyces cerevisiae</i> . <i>Nucleic Acids Research</i> , 1995, 23, 537-538. | 6.5 | 1 |
| 2745 | Processing of the Papain Precursor. <i>Journal of Biological Chemistry</i> , 1995, 270, 10838-10846. | 1.6 | 130 |
| 2746 | HYS2, an essential gene required for DNA replication in <i>Saccharomyces cerevisiae</i> . <i>Nucleic Acids Research</i> , 1995, 23, 3493-3500. | 6.5 | 46 |
| 2747 | Riboflavin Biosynthesis in <i>Saccharomyces cerevisiae</i> . <i>Journal of Biological Chemistry</i> , 1995, 270, 437-444. | 1.6 | 40 |
| 2748 | In Vivo Mutational Analysis of the DNA Binding Domain of the Tissue-specific Transcription Factor, Pit-1. <i>Journal of Biological Chemistry</i> , 1995, 270, 25520-25525. | 1.6 | 24 |
| 2749 | Deletion analysis of the starch-binding domain of <i>Aspergillus glucoamylase</i> . <i>Protein Engineering, Design and Selection</i> , 1995, 8, 1049-1055. | 1.0 | 27 |
| 2750 | Cloning and Characterization of a <i>Saccharomyces cerevisiae</i> Gene Encoding the Low Molecular Weight Protein-tyrosine Phosphatase. <i>Journal of Biological Chemistry</i> , 1995, 270, 18491-18499. | 1.6 | 39 |
| 2751 | Evidence that the MIF2 gene of <i>Saccharomyces cerevisiae</i> encodes a centromere protein with homology to the mammalian centromere protein CENP-C.. <i>Molecular Biology of the Cell</i> , 1995, 6, 793-807. | 0.9 | 386 |
| 2752 | Functional characterization of Ost3p. Loss of the 34-kD subunit of the <i>Saccharomyces cerevisiae</i> oligosaccharyltransferase results in biased underglycosylation of acceptor substrates.. <i>Journal of Cell Biology</i> , 1995, 130, 567-577. | 2.3 | 74 |
| 2753 | Growth impairment resulting from expression of influenza virus M2 protein in <i>Saccharomyces cerevisiae</i> : identification of a novel inhibitor of influenza virus. <i>Antimicrobial Agents and Chemotherapy</i> , 1995, 39, 2204-2209. | 1.4 | 77 |
| 2754 | Yeast Gaa1p is required for attachment of a completed GPI anchor onto proteins.. <i>Journal of Cell Biology</i> , 1995, 129, 629-639. | 2.3 | 172 |

| # | ARTICLE | IF | CITATIONS |
|------|--|-----|-----------|
| 2755 | A method for performing precise alterations in the yeast genome using a recyclable selectable marker. <i>Nucleic Acids Research</i> , 1995, 23, 3079-3081. | 6.5 | 95 |
| 2756 | <i>Saccharomyces cerevisiae</i> CNE1 Encodes an Endoplasmic Reticulum (ER) Membrane Protein with Sequence Similarity to Calnexin and Calreticulin and Functions as a Constituent of the ER Quality Control Apparatus. <i>Journal of Biological Chemistry</i> , 1995, 270, 244-253. | 1.6 | 160 |
| 2757 | Mas37p, a novel receptor subunit for protein import into mitochondria.. <i>Journal of Cell Biology</i> , 1995, 129, 25-34. | 2.3 | 172 |
| 2758 | The role of Myo2, a yeast class V myosin, in vesicular transport.. <i>Journal of Cell Biology</i> , 1995, 128, 1055-1068. | 2.3 | 316 |
| 2759 | [16] Analyzing protein-protein interactions using two-hybrid system. <i>Methods in Enzymology</i> , 1995, 254, 241-263. | 0.4 | 330 |
| 2760 | A late Golgi sorting function for <i>Saccharomyces cerevisiae</i> Apm1p, but not for Apm2p, a second yeast clathrin AP medium chain-related protein.. <i>Molecular Biology of the Cell</i> , 1995, 6, 41-58. | 0.9 | 87 |
| 2761 | The rho-GAP encoded by BEM2 regulates cytoskeletal structure in budding yeast.. <i>Molecular Biology of the Cell</i> , 1995, 6, 1011-1024. | 0.9 | 69 |
| 2762 | The Essential Function of Protein-disulfide Isomerase Is to Unscramble Non-native Disulfide Bonds. <i>Journal of Biological Chemistry</i> , 1995, 270, 28006-28009. | 1.6 | 192 |
| 2763 | The GLFG repetitive region of the nucleoporin Nup116p interacts with Kap95p, an essential yeast nuclear import factor.. <i>Journal of Cell Biology</i> , 1995, 131, 1699-1713. | 2.3 | 197 |
| 2764 | A novel cis-acting centromeric DNA element affects <i>S. pombe</i> centromeric chromatin structure at a distance.. <i>Journal of Cell Biology</i> , 1995, 128, 445-454. | 2.3 | 48 |
| 2765 | Control of RAS mRNA level by the mevalonate pathway.. <i>Molecular Biology of the Cell</i> , 1995, 6, 59-70. | 0.9 | 23 |
| 2766 | Human Mitogen-activated Protein Kinase CSBP1, but Not CSBP2, Complements a hog1 Deletion in Yeast. <i>Journal of Biological Chemistry</i> , 1995, 270, 29043-29046. | 1.6 | 47 |
| 2767 | VMA8 Encodes a 32-kDa V1 Subunit of the <i>Saccharomyces cerevisiae</i> Vacuolar H ⁺ -ATPase Required for Function and Assembly of the Enzyme Complex. <i>Journal of Biological Chemistry</i> , 1995, 270, 15037-15044. | 1.6 | 55 |
| 2768 | DNM1, a dynamin-related gene, participates in endosomal trafficking in yeast.. <i>Journal of Cell Biology</i> , 1995, 130, 553-566. | 2.3 | 117 |
| 2769 | Genetic Analysis of the Fluorescein Isothiocyanate Binding Site of the Yeast Plasma Membrane H ⁺ -ATPase. <i>Journal of Biological Chemistry</i> , 1995, 270, 8655-8659. | 1.6 | 16 |
| 2770 | A homeo domain protein lacking specific side chains of helix 3 can still bind DNA and direct transcriptional repression.. <i>Genes and Development</i> , 1995, 9, 182-192. | 2.7 | 62 |
| 2771 | A Transient GCN4 mRNA Destabilization Follows GCN4 Translational Derepression. <i>Journal of Biological Chemistry</i> , 1995, 270, 17317-17320. | 1.6 | 4 |
| 2772 | Fus2 localizes near the site of cell fusion and is required for both cell fusion and nuclear alignment during zygote formation.. <i>Journal of Cell Biology</i> , 1995, 130, 1283-1296. | 2.3 | 68 |

| # | ARTICLE | IF | CITATIONS |
|------|---|-----|-----------|
| 2773 | The Highly Conserved Protein PO Carboxyl End Is Essential for Ribosome Activity Only in the Absence of Proteins P1 and P2. <i>Journal of Biological Chemistry</i> , 1995, 270, 20608-20614. | 1.6 | 82 |
| 2774 | Changing Stereochemistry for a Metabolic Pathway in Vivo. <i>Journal of Biological Chemistry</i> , 1995, 270, 27453-27457. | 1.6 | 54 |
| 2775 | Tropomyosin is essential in yeast, yet the TPM1 and TPM2 products perform distinct functions.. <i>Journal of Cell Biology</i> , 1995, 128, 383-392. | 2.3 | 140 |
| 2776 | p93dis1, which is required for sister chromatid separation, is a novel microtubule and spindle pole body-associating protein phosphorylated at the Cdc2 target sites.. <i>Genes and Development</i> , 1995, 9, 1572-1585. | 2.7 | 149 |
| 2777 | Two microtubule-associated proteins required for anaphase spindle movement in <i>Saccharomyces cerevisiae</i> [published erratum appears in <i>J Cell Biol</i> 1995 Oct;131(2):561]. <i>Journal of Cell Biology</i> , 1995, 130, 1373-1385. | 2.3 | 162 |
| 2778 | Lethal Mutations in a Yeast U6 RNA Gene B Block Promoter Element Identify Essential Contacts with Transcription Factor-IIIc. <i>Journal of Biological Chemistry</i> , 1995, 270, 11398-11405. | 1.6 | 22 |
| 2779 | The Influence of Subunit Structure on the Interaction of Na/K-ATPase Complexes with Na.. <i>Journal of Biological Chemistry</i> , 1995, 270, 13937-13947. | 1.6 | 46 |
| 2780 | Vesicle-mediated protein transport: regulatory interactions between the Vps15 protein kinase and the Vps34 PtdIns 3-kinase essential for protein sorting to the vacuole in yeast.. <i>Journal of Cell Biology</i> , 1995, 129, 321-334. | 2.3 | 226 |
| 2781 | Membrane protein retrieval from the Golgi apparatus to the endoplasmic reticulum (ER): characterization of the RER1 gene product as a component involved in ER localization of Sec12p.. <i>Molecular Biology of the Cell</i> , 1995, 6, 1459-1477. | 0.9 | 98 |
| 2782 | Phosphorylation of CTP Synthetase from <i>Saccharomyces cerevisiae</i> by Protein Kinase C. <i>Journal of Biological Chemistry</i> , 1995, 270, 14983-14988. | 1.6 | 60 |
| 2783 | Pmp27 promotes peroxisomal proliferation.. <i>Journal of Cell Biology</i> , 1995, 129, 345-355. | 2.3 | 203 |
| 2784 | <i>Saccharomyces cerevisiae</i> Phosphoenolpyruvate Carboxykinase: Revised Amino Acid Sequence, Site-Directed Mutagenesis, and Microenvironment Characteristics of Cysteines 365 and 458. <i>Biochemistry</i> , 1995, 34, 6382-6388. | 1.2 | 32 |
| 2785 | Site-Directed Mutagenesis of Putative GTP-Binding Sites of Yeast .beta.-Tubulin: Evidence That .alpha.-, .beta.-, and .gamma.-Tubulins Are Atypical GTPases. <i>Biochemistry</i> , 1995, 34, 7409-7419. | 1.2 | 30 |
| 2786 | Glucose-dependent turnover of the mRNAs encoding succinate dehydrogenase peptides in <i>Saccharomyces cerevisiae</i> : sequence elements in the 5' untranslated region of the Ip mRNA play a dominant role.. <i>Molecular Biology of the Cell</i> , 1995, 6, 1125-1143. | 0.9 | 71 |
| 2787 | Expression of the human MGMT O6-methylguanine DNA methyltransferase gene in a yeast alkylation-sensitive mutant: its effects on both exogenous and endogenous DNA alkylation damage. <i>Mutation Research DNA Repair</i> , 1995, 336, 133-142. | 3.8 | 10 |
| 2788 | Cloning and sequence of a gene encoding the pyruvate dehydrogenase E1 β subunit of <i>Schizosaccharomyces pombe</i> . <i>Gene</i> , 1995, 152, 117-120. | 1.0 | 4 |
| 2789 | Expression of the <i>Arabidopsis thaliana</i> 2S albumin gene 3 in <i>Saccharomyces cerevisiae</i> . <i>Gene</i> , 1995, 153, 175-178. | 1.0 | 2 |
| 2790 | Allele shuffling: conjugational transfer, plasmid shuffling and suppressor analysis in <i>saccharomyces cerevisiae</i> . <i>Gene</i> , 1995, 155, 51-59. | 1.0 | 12 |

| # | ARTICLE | IF | CITATIONS |
|------|--|------|-----------|
| 2791 | Isolation and characterization of ten mutator alleles of the mitochondrial DNA polymerase-encoding MIP1 gene from <i>Saccharomyces cerevisiae</i> . <i>Gene</i> , 1995, 160, 105-110. | 1.0 | 42 |
| 2792 | Role of Ser-53 phosphorylation in the activity of human translation initiation factor eIF-4E in mammalian and yeast cells. <i>Gene</i> , 1995, 163, 283-288. | 1.0 | 11 |
| 2793 | Cloning and characterization of the <i>Saccharomyces cerevisiae</i> SVS1 gene which encodes a serine- and threonine-rich protein required for vanadate resistance. <i>Gene</i> , 1995, 165, 25-29. | 1.0 | 19 |
| 2794 | Cloning of the <i>Candida glabrata</i> TRP1 and HIS3 genes, and construction of their disruptant strains by sequential integrative transformation. <i>Gene</i> , 1995, 165, 203-206. | 1.0 | 144 |
| 2795 | Construction of baker's yeast strains that secrete <i>Aspergillus oryzae</i> alpha-amylase and their use in bread making. <i>Journal of Cereal Science</i> , 1995, 21, 185-193. | 1.8 | 39 |
| 2796 | Identification of a novel nuclear pore-associated protein as a functional target of the HIV-1 Rev protein in yeast. <i>Cell</i> , 1995, 82, 495-506. | 13.5 | 247 |
| 2797 | TEL1, an <i>S. cerevisiae</i> homolog of the human gene mutated in ataxia telangiectasia, is functionally related to the yeast checkpoint gene MEC1. <i>Cell</i> , 1995, 82, 831-840. | 13.5 | 372 |
| 2798 | Improvement of S-adenosylmethionine production by integration of the ethionine-resistance gene into chromosomes of the yeast <i>Saccharomyces cerevisiae</i> . <i>Applied Microbiology and Biotechnology</i> , 1995, 42, 730-733. | 1.7 | 33 |
| 2799 | <i>Schizosaccharomyces pombe</i> : A Model for Molecular Studies of Eukaryotic Genes. <i>DNA and Cell Biology</i> , 1995, 14, 359-371. | 0.9 | 89 |
| 2800 | Production of Rat Protein Disulfide Isomerase in <i>Saccharomyces cerevisiae</i> . <i>Protein Expression and Purification</i> , 1995, 6, 700-706. | 0.6 | 16 |
| 2801 | Induction in the gene RNR3 in <i>saccharomyces cerevisiae</i> upon exposure to different agents related to carcinogenesis. <i>Biochemical Pharmacology</i> , 1995, 50, 1695-1699. | 2.0 | 13 |
| 2802 | Identification of two cell cycle regulated genes affecting the β -1,3-glucan content of cell walls in <i>Saccharomyces cerevisiae</i> . <i>FEBS Letters</i> , 1995, 358, 165-170. | 1.3 | 134 |
| 2803 | Transcriptional regulation of the yeast vacuolar aminopeptidase <i>ysc1</i> encoding gene (APE1) by carbon sources. <i>FEBS Letters</i> , 1995, 364, 13-16. | 1.3 | 8 |
| 2804 | Protein disulfide isomerase mutant lacking its isomerase activity accelerates protein folding in the cell. <i>FEBS Letters</i> , 1995, 377, 505-511. | 1.3 | 85 |
| 2805 | One-step enzymatic hydrolysis of starch using a recombinant strain of <i>Saccharomyces cerevisiae</i> producing α -amylase, glucoamylase and pullulanase. <i>Applied Microbiology and Biotechnology</i> , 1995, 42, 878-883. | 1.7 | 42 |
| 2806 | Yeast Transformation and the Preparation of Frozen Spheroplasts for Electroporation. , 1995, 47, 261-268. | | 11 |
| 2807 | Sequences 5' of the first upstream open reading frame in GCN4 mRNA are required for efficient translational reinitiation. <i>Nucleic Acids Research</i> , 1995, 23, 3980-3988. | 6.5 | 44 |
| 2808 | The upstream region of the isocitrate lyase gene (UPR-ICL) of <i>Candida tropicalis</i> induces gene expression in both <i>Saccharomyces cerevisiae</i> and <i>Escherichia coli</i> by acetate via two distinct promoters. <i>Archives of Microbiology</i> , 1995, 163, 322-328. | 1.0 | 13 |

| # | ARTICLE | IF | CITATIONS |
|------|--|-----|-----------|
| 2809 | Cloning of <i>Corticium rolfii</i> glucoamylase cDNA and its expression in <i>Saccharomyces cerevisiae</i> . <i>Applied Microbiology and Biotechnology</i> , 1995, 44, 451-458. | 1.7 | 8 |
| 2810 | Development of yeast strains for the efficient utilisation of starch: evaluation of constructs that express α -amylase and glucoamylase separately or as bifunctional fusion proteins. <i>Applied Microbiology and Biotechnology</i> , 1995, 43, 1067-1076. | 1.7 | 90 |
| 2811 | CEN14 sequences cause slower proliferation, reduced cell size and asporogeny in <i>Saccharomyces cerevisiae</i> . <i>Applied Microbiology and Biotechnology</i> , 1995, 43, 877-879. | 1.7 | 2 |
| 2812 | A novel promoter, derived from the isocitrate lyase gene of <i>Candida tropicalis</i> , inducible with acetate in <i>Saccharomyces cerevisiae</i> . <i>Applied Microbiology and Biotechnology</i> , 1995, 43, 489-492. | 1.7 | 41 |
| 2813 | Secretion by <i>Saccharomyces cerevisiae</i> of rat apolipoprotein E as a fusion to <i>Mucor</i> rennin. <i>Applied Microbiology and Biotechnology</i> , 1995, 42, 865-870. | 1.7 | 1 |
| 2814 | Generation of a stable non-reverting <i>Leu⁻</i> mutant of <i>Kluyveromyces lactis</i> by gene disruption. <i>Applied Microbiology and Biotechnology</i> , 1995, 42, 734-737. | 1.7 | 4 |
| 2815 | An Essential Gene Pair in <i>Saccharomyces cerevisiae</i> with a Potential Role in Mating. <i>DNA and Cell Biology</i> , 1995, 14, 411-418. | 0.9 | 32 |
| 2816 | NCA2, a second nuclear gene required for the control of mitochondrial synthesis of subunits 6 and 8 of ATP synthase in <i>Saccharomyces cerevisiae</i> . <i>Journal of Molecular Biology</i> , 1995, 247, 588-596. | 2.0 | 27 |
| 2817 | Functional interaction between the CSE2 gene product and centromeres in <i>Saccharomyces cerevisiae</i> . <i>Journal of Molecular Biology</i> , 1995, 248, 255-263. | 2.0 | 5 |
| 2818 | Two mutually exclusive regulatory systems inhibit UASGATA, a cluster of 5'-GAT(A/T)A-3' upstream from the UGA4 gene of <i>Saccharomyces cerevisiae</i> . <i>Nucleic Acids Research</i> , 1995, 23, 558-564. | 6.5 | 55 |
| 2819 | Structural and functional analysis of N-terminal point mutants of the human estrogen receptor. <i>Journal of Steroid Biochemistry and Molecular Biology</i> , 1996, 57, 293-300. | 1.2 | 23 |
| 2820 | Synaptotagmin II expression partially rescues the growth defect of the yeast <i>sec15</i> secretory mutant. <i>Biology of the Cell</i> , 1996, 88, 55-63. | 0.7 | 1 |
| 2821 | Overexpression of binding protein and disruption of the <i>PMR1</i> gene synergistically stimulate secretion of bovine prochymosin but not plant Thaumatin in yeast. <i>Applied Microbiology and Biotechnology</i> , 1996, 46, 365-370. | 1.7 | 95 |
| 2822 | High-level secretion of human α 1-antitrypsin from <i>Saccharomyces cerevisiae</i> using inulinase signal sequence. <i>Journal of Biotechnology</i> , 1996, 48, 15-24. | 1.9 | 38 |
| 2823 | The yeast splicing factor Mud13p is a commitment complex component and corresponds to CBP20, the small subunit of the nuclear cap-binding complex. <i>Genes and Development</i> , 1996, 10, 1699-1708. | 2.7 | 137 |
| 2824 | A lea-class gene of tomato confers salt and freezing tolerance when expressed in <i>Saccharomyces cerevisiae</i> . <i>Gene</i> , 1996, 170, 243-248. | 1.0 | 170 |
| 2825 | Isolation of a <i>Candida glabrata</i> centromere and its use in construction of plasmid vectors. <i>Gene</i> , 1996, 175, 105-108. | 1.0 | 74 |
| 2826 | Cloning of <i>Schizosaccharomyces pombe</i> <i>rph16+</i> , a gene homologous to the <i>Saccharomyces cerevisiae</i> <i>RAD16</i> gene. <i>Mutation Research DNA Repair</i> , 1996, 364, 57-71. | 3.8 | 11 |

| # | ARTICLE | IF | CITATIONS |
|------|---|------|-----------|
| 2827 | Maintenance of the peroxisomal compartment in glucose-repressed and anaerobically grown <i>Saccharomyces cerevisiae</i> cells. <i>Biochimie</i> , 1996, 78, 95-102. | 1.3 | 11 |
| 2828 | <i>Schizosaccharomyces pombe</i> possesses an unusual and a conventional hexokinase: biochemical and molecular characterization of both hexokinases. <i>FEBS Letters</i> , 1996, 378, 185-189. | 1.3 | 32 |
| 2829 | Site-directed mutagenesis of <i>Saccharomyces cerevisiae</i> β -tubulin: interaction between residue 167 and benzimidazole compounds. <i>FEBS Letters</i> , 1996, 385, 7-10. | 1.3 | 42 |
| 2830 | Positive and negative regulation of a sterol biosynthetic gene (ERG3) in the post-squalene portion of the yeast ergosterol pathway. <i>FEBS Letters</i> , 1996, 392, 161-165. | 1.3 | 75 |
| 2831 | Molecular cloning of rat phosphatidylinositol synthase cDNA by functional complementation of the yeast <i>Saccharomyces cerevisiae</i> pismutation. <i>FEBS Letters</i> , 1996, 393, 89-92. | 1.3 | 38 |
| 2832 | Determination of the disulfide array of the first inducible antifungal peptide from insects: drosomycin from <i>Drosophila melanogaster</i> . <i>FEBS Letters</i> , 1996, 395, 6-10. | 1.3 | 50 |
| 2833 | Polyprenyl diphosphate synthase essentially defines the length of the side chain of ubiquinone. <i>Lipids and Lipid Metabolism</i> , 1996, 1302, 217-223. | 2.6 | 103 |
| 2834 | Interactions of the F1-ATPase subunits from <i>Escherichia coli</i> detected by the yeast two-hybrid system. <i>Biochimica Et Biophysica Acta - Bioenergetics</i> , 1996, 1274, 67-72. | 0.5 | 6 |
| 2835 | FOG1 and FOG2 genes, required for the transcriptional activation of glucose-repressible genes of <i>Kluyveromyces lactis</i> , are homologous to GAL83 and SNF1 of <i>Saccharomyces cerevisiae</i> . <i>Current Genetics</i> , 1996, 29, 316-326. | 0.8 | 13 |
| 2836 | Construction of the complete rat fatty acid synthase cDNA and its expression in <i>Saccharomyces cerevisiae</i> . <i>Current Genetics</i> , 1996, 29, 219-226. | 0.8 | 2 |
| 2837 | yAP-1- and yAP-2-mediated, heat shock-induced transcriptional activation of the multidrug resistance ABC transporter genes in <i>Saccharomyces cerevisiae</i> . <i>Current Genetics</i> , 1996, 29, 103-105. | 0.8 | 60 |
| 2838 | Detection of antagonistic cellular regulatory functions by the gene-gene interference method in yeast. <i>Current Genetics</i> , 1996, 29, 114-121. | 0.8 | 7 |
| 2839 | Molecular and functional characterization of a mutant allele of the mitogen-activated protein-kinase gene SLT2 (MPK1) rescued from yeast autolytic mutants. <i>Current Genetics</i> , 1996, 29, 516-522. | 0.8 | 50 |
| 2840 | Budding Yeast SKP1 Encodes an Evolutionarily Conserved Kinetochores Protein Required for Cell Cycle Progression. <i>Cell</i> , 1996, 86, 275-285. | 13.5 | 291 |
| 2841 | A Novel Mechanism for Regulating Activity of a Transcription Factor That Controls the Unfolded Protein Response. <i>Cell</i> , 1996, 87, 391-404. | 13.5 | 923 |
| 2842 | Glutamic acid decarboxylase autoantibody assay using ¹²⁵ I-labelled recombinant GAD65 produced in yeast. <i>Clinica Chimica Acta</i> , 1996, 256, 175-188. | 0.5 | 75 |
| 2843 | Two isozymes of P450nor of <i>Cylindrocarpus tonkinensis</i> : Molecular cloning of the cDNAs and genes, expressions in the yeast, and the putative NAD(P)H-binding site. <i>Biochimie</i> , 1996, 78, 792-799. | 1.3 | 56 |
| 2844 | An Active Sucrose Carrier (Scr1) that is Predominantly Expressed in the Seedling of <i>Ricinus communis</i> L.. <i>Journal of Plant Physiology</i> , 1996, 147, 685-690. | 1.6 | 47 |

| # | ARTICLE | IF | CITATIONS |
|------|--|-----|-----------|
| 2845 | Base analog 6-N-hydroxylaminopurine mutagenesis in the yeast <i>Saccharomyces cerevisiae</i> is controlled by replicative DNA polymerases. <i>Mutation Research - Genetic Toxicology Testing and Biomonitoring of Environmental Or Occupational Exposure</i> , 1996, 369, 33-44. | 1.2 | 49 |
| 2846 | Sterol Esterification in Yeast: A Two-Gene Process. <i>Science</i> , 1996, 272, 1353-1356. | 6.0 | 256 |
| 2847 | Application of Ty1 for cloned gene insertion: amplification of a large regulated expression cassette in <i>Saccharomyces cerevisiae</i> . <i>Applied Microbiology and Biotechnology</i> , 1996, 44, 620-623. | 1.7 | 10 |
| 2848 | Transformation of <i>Saccharomyces cerevisiae</i> . , 1996, , 97-103. | | 1 |
| 2849 | Reading two bases twice: mammalian antizyme frameshifting in yeast.. <i>EMBO Journal</i> , 1996, 15, 1360-1370. | 3.5 | 72 |
| 2850 | Copper-dependent degradation of the <i>Saccharomyces cerevisiae</i> plasma membrane copper transporter Ctr1p in the apparent absence of endocytosis.. <i>EMBO Journal</i> , 1996, 15, 3515-3523. | 3.5 | 179 |
| 2851 | Human platelet cGI-PDE: expression in yeast and localization of the catalytic domain by deletion mutagenesis. <i>Blood</i> , 1996, 88, 1321-1329. | 0.6 | 45 |
| 2852 | Ras2 signals via the Cdc42/Ste20/mitogen-activated protein kinase module to induce filamentous growth in <i>Saccharomyces cerevisiae</i> .. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 1996, 93, 5352-5356. | 3.3 | 345 |
| 2853 | A yeast manganese transporter related to the macrophage protein involved in conferring resistance to mycobacteria.. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 1996, 93, 5105-5110. | 3.3 | 333 |
| 2854 | Mutations within the first LSGGQ motif of Ste6p cause defects in a-factor transport and mating in <i>Saccharomyces cerevisiae</i> . <i>Journal of Bacteriology</i> , 1996, 178, 1712-1719. | 1.0 | 30 |
| 2855 | Identification and Characterization of Mutations in the <i>UPF1</i> Gene That Affect Nonsense Suppression and the Formation of the Upf Protein Complex but Not mRNA Turnover. <i>Molecular and Cellular Biology</i> , 1996, 16, 5491-5506. | 1.1 | 146 |
| 2856 | Genetic analysis of glucose regulation in <i>saccharomyces cerevisiae</i> : control of transcription versus mRNA turnover.. <i>EMBO Journal</i> , 1996, 15, 363-374. | 3.5 | 70 |
| 2857 | Single point mutations in various domains of a plant plasma membrane H(+)-ATPase expressed in <i>Saccharomyces cerevisiae</i> increase H(+)-pumping and permit yeast growth at low pH.. <i>EMBO Journal</i> , 1996, 15, 5513-5526. | 3.5 | 122 |
| 2858 | Functional analysis of the interaction between the small GTP binding protein Cdc42 and the Ste20 protein kinase in yeast.. <i>EMBO Journal</i> , 1996, 15, 7046-7059. | 3.5 | 222 |
| 2859 | Amino acid substitutions in membrane-spanning domains of Hol1, a member of the major facilitator superfamily of transporters, confer nonselective cation uptake in <i>Saccharomyces cerevisiae</i> . <i>Journal of Bacteriology</i> , 1996, 178, 7197-7205. | 1.0 | 29 |
| 2860 | CWH41 encodes a novel endoplasmic reticulum membrane N-glycoprotein involved in beta 1,6-glucan assembly. <i>Journal of Bacteriology</i> , 1996, 178, 1162-1171. | 1.0 | 66 |
| 2861 | Two transcription factors, Gln3p and Nil1p, use the same GATAAG sites to activate the expression of GAP1 of <i>Saccharomyces cerevisiae</i> . <i>Journal of Bacteriology</i> , 1996, 178, 2465-2468. | 1.0 | 62 |
| 2862 | Pip2p: a transcriptional regulator of peroxisome proliferation in the yeast <i>Saccharomyces cerevisiae</i> .. <i>EMBO Journal</i> , 1996, 15, 2924-2934. | 3.5 | 100 |

| # | ARTICLE | IF | CITATIONS |
|------|---|-----|-----------|
| 2863 | Functional Expression of the Multidrug Resistance-associated Protein in the Yeast <i>Saccharomyces cerevisiae</i> . <i>Journal of Biological Chemistry</i> , 1996, 271, 4154-4160. | 1.6 | 34 |
| 2864 | Fission yeast <i>cdc21</i> , a member of the MCM protein family, is required for onset of S phase and is located in the nucleus throughout the cell cycle.. <i>EMBO Journal</i> , 1996, 15, 861-872. | 3.5 | 86 |
| 2865 | A suppressor of mutations in the class III transcription system encodes a component of yeast TFIIIB.. <i>EMBO Journal</i> , 1996, 15, 1941-1949. | 3.5 | 81 |
| 2866 | A novel Hsp70 of the yeast ER lumen is required for the efficient translocation of a number of protein precursors.. <i>EMBO Journal</i> , 1996, 15, 2640-2650. | 3.5 | 150 |
| 2867 | The Exocyst is a multiprotein complex required for exocytosis in <i>Saccharomyces cerevisiae</i> .. <i>EMBO Journal</i> , 1996, 15, 6483-6494. | 3.5 | 757 |
| 2868 | Requirement for PP1 phosphatase and 20S cyclosome/APC for the onset of anaphase is lessened by the dosage increase of a novel gene <i>sds23+</i> .. <i>EMBO Journal</i> , 1996, 15, 6629-6640. | 3.5 | 73 |
| 2869 | Multiple Pdr1p/Pdr3p Binding Sites Are Essential for Normal Expression of the ATP Binding Cassette Transporter Protein-encoding Gene. <i>Journal of Biological Chemistry</i> , 1996, 271, 23049-23054. | 1.6 | 121 |
| 2870 | Role of three chitin synthase genes in the growth of <i>Candida albicans</i> . <i>Journal of Bacteriology</i> , 1996, 178, 2416-2419. | 1.0 | 114 |
| 2871 | The HeLa 200 kDa U5 snRNP-specific protein and its homologue in <i>Saccharomyces cerevisiae</i> are members of the DEXH-box protein family of putative RNA helicases.. <i>EMBO Journal</i> , 1996, 15, 4001-4015. | 3.5 | 107 |
| 2872 | Kes1p shares homology with human oxysterol binding protein and participates in a novel regulatory pathway for yeast Golgi-derived transport vesicle biogenesis.. <i>EMBO Journal</i> , 1996, 15, 6447-6459. | 3.5 | 183 |
| 2873 | Competition between folding and glycosylation in the endoplasmic reticulum.. <i>EMBO Journal</i> , 1996, 15, 3538-3546. | 3.5 | 85 |
| 2874 | A pathway in the yeast cell division cycle linking protein kinase C (Pkc1) to activation of Cdc28 at START.. <i>EMBO Journal</i> , 1996, 15, 3040-3052. | 3.5 | 66 |
| 2875 | The Yeast Homolog of Mammalian Ribosomal Protein S30 Is Expressed from a Duplicated Gene without a Ubiquitin-like Protein Fusion Sequence. <i>Journal of Biological Chemistry</i> , 1996, 271, 13549-13555. | 1.6 | 41 |
| 2876 | Utilizing the GCN4 leader region to investigate the role of the sequence determinants in nonsense-mediated mRNA decay.. <i>EMBO Journal</i> , 1996, 15, 2810-2819. | 3.5 | 47 |
| 2877 | Mutational Analysis of the Vacuolar Sorting Signal of Procarboxypeptidase Y in Yeast Shows a Low Requirement for Sequence Conservation. <i>Journal of Biological Chemistry</i> , 1996, 271, 841-846. | 1.6 | 18 |
| 2878 | Two GTPase isoforms, Ypt31p and Ypt32p, are essential for Golgi function in yeast.. <i>EMBO Journal</i> , 1996, 15, 6460-6475. | 3.5 | 153 |
| 2879 | Isolation and Analysis of the Yeast <i>TEA1</i> Gene, Which Encodes a Zinc Cluster Tyrosine Enhancer-Binding Protein. <i>Molecular and Cellular Biology</i> , 1996, 16, 347-358. | 1.1 | 28 |
| 2880 | The mating-specific G(alpha) protein of <i>Saccharomyces cerevisiae</i> downregulates the mating signal by a mechanism that is dependent on pheromone and independent of G(beta)(gamma) sequestration. <i>Molecular and Cellular Biology</i> , 1996, 16, 6325-6337. | 1.1 | 38 |

| # | ARTICLE | IF | CITATIONS |
|------|--|-----|-----------|
| 2881 | A Role for Nucleoporin FG Repeat Domains in Export of Human Immunodeficiency Virus Type 1 Rev Protein and RNA from the Nucleus. <i>Molecular and Cellular Biology</i> , 1996, 16, 7144-7150. | 1.1 | 97 |
| 2882 | A Yeast Acetyl Coenzyme A Carboxylase Mutant Links Very-Long-Chain Fatty Acid Synthesis to the Structure and Function of the Nuclear Membrane-Pore Complex. <i>Molecular and Cellular Biology</i> , 1996, 16, 7161-7172. | 1.1 | 184 |
| 2883 | Genetic Analysis of the Bipolar Pattern of Bud Site Selection in the Yeast <i>Saccharomyces cerevisiae</i> . <i>Molecular and Cellular Biology</i> , 1996, 16, 1857-1870. | 1.1 | 220 |
| 2884 | Bul1, a New Protein That Binds to the Rsp5 Ubiquitin Ligase in <i>Saccharomyces cerevisiae</i> . <i>Molecular and Cellular Biology</i> , 1996, 16, 3255-3263. | 1.1 | 105 |
| 2885 | Functional Interaction of Cytosolic hsp70 and a DnaJ-Related Protein, Ydj1p, in Protein Translocation In Vivo. <i>Molecular and Cellular Biology</i> , 1996, 16, 4378-4386. | 1.1 | 220 |
| 2886 | Yeast Nle3p/Nup170p Is Required for Normal Stoichiometry of FG Nucleoporins within the Nuclear Pore Complex. <i>Molecular and Cellular Biology</i> , 1996, 16, 2025-2036. | 1.1 | 53 |
| 2887 | <i>SSN</i> Genes That Affect Transcriptional Repression in <i>Saccharomyces cerevisiae</i> Encode SIN4, ROX3, and SRB Proteins Associated with RNA Polymerase II. <i>Molecular and Cellular Biology</i> , 1996, 16, 115-120. | 1.1 | 123 |
| 2888 | Identification of a Regulatory Subcomplex in the Guanine Nucleotide Exchange Factor eIF2B That Mediates Inhibition by Phosphorylated eIF2. <i>Molecular and Cellular Biology</i> , 1996, 16, 6603-6616. | 1.1 | 89 |
| 2889 | Invariant U2 RNA Sequences Bordering the Branchpoint Recognition Region Are Essential for Interaction with Yeast SF3a and SF3b Subunits. <i>Molecular and Cellular Biology</i> , 1996, 16, 818-828. | 1.1 | 62 |
| 2890 | Sex-Specific and Non-Sex-Specific Oligomerization Domains in Both of the <i>doublesex</i> Transcription Factors from <i>Drosophila melanogaster</i> . <i>Molecular and Cellular Biology</i> , 1996, 16, 3106-3111. | 1.1 | 73 |
| 2891 | Pho85p, a Cyclin-Dependent Protein Kinase, and the Snf1p Protein Kinase Act Antagonistically To Control Glycogen Accumulation in <i>Saccharomyces cerevisiae</i> . <i>Molecular and Cellular Biology</i> , 1996, 16, 4357-4365. | 1.1 | 92 |
| 2892 | Signals Sufficient for 3' End Formation of Yeast mRNA. <i>Molecular and Cellular Biology</i> , 1996, 16, 2772-2776. | 1.1 | 68 |
| 2893 | Identification of Ste4 as a Potential Regulator of Byr2 in the Sexual Response Pathway of <i>Schizosaccharomyces pombe</i> . <i>Molecular and Cellular Biology</i> , 1996, 16, 5597-5603. | 1.1 | 75 |
| 2894 | The Cyc8 (Ssn6)-Tup1 Corepressor Complex Is Composed of One Cyc8 and Four Tup1 Subunits. <i>Molecular and Cellular Biology</i> , 1996, 16, 6707-6714. | 1.1 | 126 |
| 2895 | A Novel Role for Cdc5p in DNA Replication. <i>Molecular and Cellular Biology</i> , 1996, 16, 6775-6782. | 1.1 | 86 |
| 2896 | Induction of Meiosis in <i>Saccharomyces cerevisiae</i> Depends on Conversion of the Transcriptional Repressor Ume6 to a Positive Regulator by Its Regulated Association with the Transcriptional Activator Ime1. <i>Molecular and Cellular Biology</i> , 1996, 16, 2518-2526. | 1.1 | 136 |
| 2897 | Inactivation of YME2/RNA12, which encodes an integral inner mitochondrial membrane protein, causes increased escape of DNA from mitochondria to the nucleus in <i>Saccharomyces cerevisiae</i> . <i>Molecular and Cellular Biology</i> , 1996, 16, 2764-2771. | 1.1 | 45 |
| 2898 | Activation Mechanism of the Multifunctional Transcription Factor Repressor-Activator Protein 1 (Rap1p). <i>Molecular and Cellular Biology</i> , 1996, 16, 3187-3196. | 1.1 | 50 |

| # | ARTICLE | IF | CITATIONS |
|------|---|-----|-----------|
| 2899 | DNA sequence preferences of GAL4 and PPR1: how a subset of Zn ²⁺ Cys ⁶ binuclear cluster proteins recognizes DNA. <i>Molecular and Cellular Biology</i> , 1996, 16, 3773-3780. | 1.1 | 97 |
| 2900 | <i>SUM1-1</i> , a Dominant Suppressor of <i>SIR</i> Mutations in <i>Saccharomyces cerevisiae</i> , Increases Transcriptional Silencing at Telomeres and <i>HM</i> Mating-Type Loci and Decreases Chromosome Stability. <i>Molecular and Cellular Biology</i> , 1996, 16, 4281-4294. | 1.1 | 47 |
| 2901 | Identification of the Bud Emergence Gene <i>BEM4</i> and Its Interactions with Rho-Type GTPases in <i>Saccharomyces cerevisiae</i> . <i>Molecular and Cellular Biology</i> , 1996, 16, 4387-4395. | 1.1 | 49 |
| 2902 | Loss of Sustained Fus3p Kinase Activity and the G ₁ Arrest Response in Cells Expressing an Inappropriate Pheromone Receptor. <i>Molecular and Cellular Biology</i> , 1996, 16, 4478-4485. | 1.1 | 18 |
| 2903 | Semidominant mutations in the yeast Rad51 protein and their relationships with the Srs2 helicase. <i>Molecular and Cellular Biology</i> , 1996, 16, 4782-4789. | 1.1 | 91 |
| 2904 | Allele-Specific Suppression of a Defective <i>trans</i> -Golgi Network (TGN) Localization Signal in <i>Kex2p</i> Identifies Three Genes Involved in Localization of TGN Transmembrane Proteins. <i>Molecular and Cellular Biology</i> , 1996, 16, 6208-6217. | 1.1 | 63 |
| 2905 | The Immunosuppressant SR 31747 Blocks Cell Proliferation by Inhibiting a Steroid Isomerase in <i>Saccharomyces cerevisiae</i> . <i>Molecular and Cellular Biology</i> , 1996, 16, 2719-2727. | 1.1 | 68 |
| 2906 | A Yeast Protein Related to a Mammalian Ras-Binding Protein, Vps9p, Is Required for Localization of Vacuolar Proteins. <i>Molecular and Cellular Biology</i> , 1996, 16, 2369-2377. | 1.1 | 78 |
| 2907 | <i>EGT2</i> Gene Transcription Is Induced Predominantly by Swi5 in Early G ₁ . <i>Molecular and Cellular Biology</i> , 1996, 16, 3264-3274. | 1.1 | 79 |
| 2908 | The AD1 Transactivation Domain of E2A Contains a Highly Conserved Helix Which Is Required for Its Activity in both <i>Saccharomyces cerevisiae</i> and Mammalian Cells. <i>Molecular and Cellular Biology</i> , 1996, 16, 121-129. | 1.1 | 96 |
| 2909 | Regulation of the G-Protein-Coupled $\hat{\pm}$ -Factor Pheromone Receptor by Phosphorylation. <i>Molecular and Cellular Biology</i> , 1996, 16, 247-257. | 1.1 | 77 |
| 2910 | Genetic and Biochemical Characterization of Mutations in the ATPase and Helicase Regions of the Upf1 Protein. <i>Molecular and Cellular Biology</i> , 1996, 16, 5477-5490. | 1.1 | 202 |
| 2911 | Mechanism of <i>MAT</i> Donor Preference during Mating-Type Switching of <i>Saccharomyces cerevisiae</i> . <i>Molecular and Cellular Biology</i> , 1996, 16, 657-668. | 1.1 | 45 |
| 2912 | Autoactivation by a <i>Candida glabrata</i> Copper Metalloregulatory Transcription Factor Requires Critical Minor Groove Interactions. <i>Molecular and Cellular Biology</i> , 1996, 16, 724-734. | 1.1 | 61 |
| 2913 | <i>BRO1</i> , a Novel Gene That Interacts with Components of the Plc1p Mitogen-Activated Protein Kinase Pathway in <i>Saccharomyces cerevisiae</i> . <i>Molecular and Cellular Biology</i> , 1996, 16, 2585-2593. | 1.1 | 75 |
| 2914 | Faithful Chromosome Transmission Requires Spt4p, a Putative Regulator of Chromatin Structure in <i>Saccharomyces cerevisiae</i> . <i>Molecular and Cellular Biology</i> , 1996, 16, 2838-2847. | 1.1 | 67 |
| 2915 | Analysis of Functional Domain Organization in DNA Topoisomerase II from Humans and <i>Saccharomyces cerevisiae</i> . <i>Molecular and Cellular Biology</i> , 1996, 16, 3866-3877. | 1.1 | 48 |
| 2916 | The EH-Domain-Containing Protein Pan1 Is Required for Normal Organization of the Actin Cytoskeleton in <i>Saccharomyces cerevisiae</i> . <i>Molecular and Cellular Biology</i> , 1996, 16, 4897-4914. | 1.1 | 101 |

| # | ARTICLE | IF | CITATIONS |
|------|---|-----|-----------|
| 2917 | CDC37 is required for p60v-src activity in yeast.. Molecular Biology of the Cell, 1996, 7, 1405-1417. | 0.9 | 76 |
| 2918 | Multilamellar endosome-like compartment accumulates in the yeast vps28 vacuolar protein sorting mutant.. Molecular Biology of the Cell, 1996, 7, 985-999. | 0.9 | 263 |
| 2919 | Dual Requirement for the Yeast <i>MMS19</i> Gene in DNA Repair and RNA Polymerase II Transcription. Molecular and Cellular Biology, 1996, 16, 6783-6793. | 1.1 | 58 |
| 2920 | The Fission Yeast <i>pmk1⁺</i> Gene Encodes a Novel Mitogen-Activated Protein Kinase Homolog Which Regulates Cell Integrity and Functions Coordinately with the Protein Kinase C Pathway. Molecular and Cellular Biology, 1996, 16, 6752-6764. | 1.1 | 201 |
| 2921 | Mitotic Crossovers between Diverged Sequences Are Regulated by Mismatch Repair Proteins in <i>Saccharomyces cerevisiae</i> . Molecular and Cellular Biology, 1996, 16, 1085-1093. | 1.1 | 221 |
| 2922 | Effects of Terminal Nonhomology and Homeology on Double-Strand-Break-Induced Gene Conversion Tract Directionality. Molecular and Cellular Biology, 1996, 16, 2951-2957. | 1.1 | 32 |
| 2923 | The CCAAT box-binding factor stimulates ammonium assimilation in <i>Saccharomyces cerevisiae</i> , defining a new cross-pathway regulation between nitrogen and carbon metabolisms. Journal of Bacteriology, 1996, 178, 1842-1849. | 1.0 | 59 |
| 2924 | Isolation and characterization of the GFA1 gene encoding the glutamine:fructose-6-phosphate amidotransferase of <i>Candida albicans</i> . Journal of Bacteriology, 1996, 178, 2320-2327. | 1.0 | 42 |
| 2925 | Glucose uptake in <i>Kluyveromyces lactis</i> : role of the HGT1 gene in glucose transport. Journal of Bacteriology, 1996, 178, 5860-5866. | 1.0 | 101 |
| 2926 | The <i>Candida albicans</i> HYR1 gene, which is activated in response to hyphal development, belongs to a gene family encoding yeast cell wall proteins. Journal of Bacteriology, 1996, 178, 5353-5360. | 1.0 | 231 |
| 2927 | Double-strand break repair in the absence of RAD51 in yeast: a possible role for break-induced DNA replication.. Proceedings of the National Academy of Sciences of the United States of America, 1996, 93, 7131-7136. | 3.3 | 409 |
| 2928 | Structure and function in rhodopsin: expression of functional mammalian opsin in <i>Saccharomyces cerevisiae</i> .. Proceedings of the National Academy of Sciences of the United States of America, 1996, 93, 11482-11486. | 3.3 | 40 |
| 2929 | TOR2 is required for organization of the actin cytoskeleton in yeast. Proceedings of the National Academy of Sciences of the United States of America, 1996, 93, 13780-13785. | 3.3 | 238 |
| 2930 | Interaction between human tRNA synthetases involves repeated sequence elements.. Proceedings of the National Academy of Sciences of the United States of America, 1996, 93, 10128-10133. | 3.3 | 81 |
| 2931 | E2 proteins: Modulators of papillomavirus transcription and replication. Methods in Enzymology, 1996, 274, 173-185. | 0.4 | 14 |
| 2932 | Characterization of <i>Saccharomyces cerevisiae</i> deficient in expression of phospholipase D. Biochemical Journal, 1996, 314, 15-19. | 1.7 | 78 |
| 2933 | Cloning and functional expression of glycosyltransferases from parasitic protozoans by heterologous complementation in yeast: the dolichol phosphate mannose synthase from <i>Trypanosoma brucei brucei</i> . Biochemical Journal, 1996, 316, 853-858. | 1.7 | 53 |
| 2934 | Localization and targeting of isocitrate lyases in <i>Saccharomyces cerevisiae</i> . Biochemical Journal, 1996, 319, 255-262. | 1.7 | 57 |

| # | ARTICLE | IF | CITATIONS |
|------|---|-----|-----------|
| 2935 | Characterization of rat 5 α -reductases type 1 and type 2 expressed in <i>Saccharomyces cerevisiae</i> . <i>Biochemical Journal</i> , 1996, 314, 1047-1052. | 1.7 | 22 |
| 2936 | Active heterodimers are formed from human DNA topoisomerase II alpha and II beta isoforms.. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 1996, 93, 8288-8293. | 3.3 | 45 |
| 2937 | Complete replication of an animal virus and maintenance of expression vectors derived from it in <i>Saccharomyces cerevisiae</i> .. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 1996, 93, 9465-9470. | 3.3 | 109 |
| 2938 | The <i>pfmdr1</i> gene of <i>Plasmodium falciparum</i> confers cellular resistance to antimalarial drugs in yeast cells.. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 1996, 93, 9942-9947. | 3.3 | 44 |
| 2939 | ECA39, a conserved gene regulated by c-Myc in mice, is involved in G1/S cell cycle regulation in yeast.. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 1996, 93, 7143-7148. | 3.3 | 61 |
| 2940 | Msn2p, a zinc finger DNA-binding protein, is the transcriptional activator of the multistress response in <i>Saccharomyces cerevisiae</i> .. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 1996, 93, 5777-5782. | 3.3 | 440 |
| 2941 | A simple assay for optimizing yeast-mammalian cell fusion conditions. <i>Molecular Biotechnology</i> , 1996, 6, 99-104. | 1.3 | 6 |
| 2942 | Acetyl-CoA hydrolase involved in acetate utilization in <i>Saccharomyces cerevisiae</i> . <i>BBA - Proteins and Proteomics</i> , 1996, 1297, 105-109. | 2.1 | 21 |
| 2943 | Expression of lipase cDNA from <i>Fusarium heterosporum</i> by <i>Saccharomyces cerevisiae</i> : High-level production and purification. <i>Journal of Bioscience and Bioengineering</i> , 1996, 81, 488-492. | 0.9 | 18 |
| 2944 | Strong expression of peroxisomal malate synthase genes from n-alkane-utilizable yeast <i>Candida tropicalis</i> in <i>Saccharomyces cerevisiae</i> . <i>Journal of Bioscience and Bioengineering</i> , 1996, 81, 61-63. | 0.9 | 3 |
| 2945 | Application of artificial neural network and fuzzy control for fed-batch cultivation of recombinant <i>Saccharomyces cerevisiae</i> . <i>Journal of Bioscience and Bioengineering</i> , 1996, 81, 412-421. | 0.9 | 24 |
| 2946 | Breeding of a new type of baker's yeast by $\hat{\Gamma}$ -integration for overproduction of glucoamylase using a homothallic yeast. <i>Journal of Bioscience and Bioengineering</i> , 1996, 81, 98-103. | 0.9 | 30 |
| 2947 | One-step splitting of a chromosome in haploid cells of <i>Saccharomyces cerevisiae</i> and its effect on the cell proliferation. <i>Journal of Bioscience and Bioengineering</i> , 1996, 82, 199-204. | 0.9 | 11 |
| 2948 | Molecular cloning and nucleotide sequence of the isoamyl acetate-hydrolyzing esterase gene (EST2) from <i>Saccharomyces cerevisiae</i> . <i>Journal of Bioscience and Bioengineering</i> , 1996, 82, 8-15. | 0.9 | 37 |
| 2949 | The construction of novel mobilizable YAC plasmids and their behavior during tranks-kingdom conjugation between bacteria and yeasts. <i>Genetic Analysis, Techniques and Applications</i> , 1996, 13, 25-31. | 1.5 | 7 |
| 2950 | Mutational analysis of <i>Vam4/Ypt7p</i> function in the vacuolar biogenesis and morphogenesis in the yeast, <i>Saccharomyces cerevisiae</i> . <i>Protoplasma</i> , 1996, 191, 126-135. | 1.0 | 15 |
| 2951 | Characterization of a promoter mutation in the <i>CYP3</i> gene of <i>Saccharomyces cerevisiae</i> which cancels regulation by <i>Cyp1p</i> (<i>Hap1p</i>) without affecting its binding site. <i>Molecular Genetics and Genomics</i> , 1996, 253, 103-110. | 2.4 | 2 |
| 2952 | Measuring the toxic effects of high gene dosage on yeast cells. <i>Molecular Genetics and Genomics</i> , 1996, 253, 393-396. | 2.4 | 9 |

| # | ARTICLE | IF | CITATIONS |
|------|---|-----|-----------|
| 2953 | Genetic interactions indicate a role for Mdg1p and the SH3 domain protein Bem1p in linking the G-protein mediated yeast pheromone signalling pathway to regulators of cell polarity. <i>Molecular Genetics and Genomics</i> , 1996, 252, 608-621. | 2.4 | 23 |
| 2954 | Dominant mutant alleles of yeast protein kinase gene CDC15 suppress the tel1 defect in termination of M phase and genetically interact with CDC14. <i>Molecular Genetics and Genomics</i> , 1996, 251, 176-185. | 2.4 | 25 |
| 2955 | Yeast chitin synthases 1 and 2 consist of a non-homologous and dispensable N-terminal region and of a homologous moiety essential for function. <i>Molecular Genetics and Genomics</i> , 1996, 252, 420-428. | 2.4 | 22 |
| 2956 | Mutational analysis of the β -subunit of yeast geranylgeranyl transferase I. <i>Molecular Genetics and Genomics</i> , 1996, 252, 1-10. | 2.4 | 15 |
| 2957 | The mus-8 gene of <i>Neurospora crassa</i> encodes a structural and functional homolog of the Rad6 protein of <i>Saccharomyces cerevisiae</i> . <i>Current Genetics</i> , 1996, 30, 224-231. | 0.8 | 19 |
| 2958 | The repair of DNA methylation damage in <i>Saccharomyces cerevisiae</i> . <i>Current Genetics</i> , 1996, 30, 461-468. | 0.8 | 98 |
| 2959 | Immunogold labeling of yeast cells: an efficient tool for the study of protein targeting and morphological alterations due to overexpression and inactivation of genes. <i>Histochemistry and Cell Biology</i> , 1996, 106, 115-130. | 0.8 | 9 |
| 2960 | cAMP inhibits bud growth in a yeast strain compromised for Ca ²⁺ influx into the Golgi. <i>Molecular Genetics and Genomics</i> , 1996, 251, 556-564. | 2.4 | 6 |
| 2961 | Genetic evidence for functional specificity of the yeast GCN2 kinase. <i>Molecular Genetics and Genomics</i> , 1996, 251, 613-618. | 2.4 | 5 |
| 2962 | In vitro mutagenesis of the mitochondrial leucyl tRNA synthetase of <i>Saccharomyces cerevisiae</i> shows that the suppressor activity of the mutant proteins is related to the splicing function of the wild-type protein. <i>Molecular Genetics and Genomics</i> , 1996, 252, 667-675. | 2.4 | 20 |
| 2963 | An extragenic suppressor of prp24-1 defines genetic interaction between PRP24 and PRP21 gene products of <i>Saccharomyces cerevisiae</i> . <i>Molecular Genetics and Genomics</i> , 1996, 250, 267-276. | 2.4 | 7 |
| 2964 | The CBP2 gene from <i>Saccharomyces douglasii</i> is a functional homologue of the <i>Saccharomyces cerevisiae</i> gene and is essential for respiratory growth in the presence of a wild-type (intron-containing) mitochondrial genome. <i>Molecular Genetics and Genomics</i> , 1996, 250, 316-322. | 2.4 | 3 |
| 2965 | Caffeine-resistance in fission yeast is caused by mutations in a single essential gene, crm1+. <i>Molecular Genetics and Genomics</i> , 1996, 250, 59-68. | 2.4 | 25 |
| 2966 | AgTHR4, a new selection marker for transformation of the filamentous fungus <i>Ashbya gossypii</i> , maps in a four-gene cluster that is conserved between <i>A. gossypii</i> and <i>Saccharomyces cerevisiae</i> . <i>Molecular Genetics and Genomics</i> , 1996, 250, 69-80. | 2.4 | 45 |
| 2967 | Exchange of Regions of the Carboxypeptidase Y Propeptide. Sequence Specificity and Function in Folding in Vivo. <i>FEBS Journal</i> , 1996, 242, 29-35. | 0.2 | 9 |
| 2968 | Analysis of the Mechanism of Activation of cAMP-Dependent Protein Kinase Through the Study of Mutants of the Yeast Regulatory Subunit. <i>FEBS Journal</i> , 1996, 237, 136-142. | 0.2 | 12 |
| 2969 | Recombinant Soluble beta-1,4-Galactosyltransferases Expressed in <i>Saccharomyces cerevisiae</i> . Purification, Characterization and Comparison with Human Enzyme. <i>FEBS Journal</i> , 1996, 239, 340-348. | 0.2 | 47 |
| 2970 | Transcription of the AAC1 Gene Encoding an Isoform of Mitochondrial ADP/ATP Carrier in <i>Saccharomyces cerevisiae</i> is Regulated by Oxygen in a Heme-Independent Manner. <i>FEBS Journal</i> , 1996, 239, 759-763. | 0.2 | 23 |

| # | ARTICLE | IF | CITATIONS |
|------|--|-----|-----------|
| 2971 | cDNA Cloning and Functional Analysis of the p97 Subunit of the 26S Proteasome, a Polypeptide Identical to the Type-1 Tumor-Necrosis-Factor-Receptor-Associated Protein-2/55.11. FEBS Journal, 1996, 239, 912-921. | 0.2 | 54 |
| 2972 | Mutations in yeast ribosomal proteins S28 and S4 affect the accuracy of translation and alter the sensitivity of the ribosomes to paromomycin. Biochimica Et Biophysica Acta Gene Regulatory Mechanisms, 1996, 1309, 156-166. | 2.4 | 47 |
| 2973 | Yeast expression of the cytokine receptor domain of the soluble interleukin-6 receptor. Journal of Immunological Methods, 1996, 199, 47-54. | 0.6 | 31 |
| 2974 | Expression and characterization of rat pancreatic secretory trypsin inhibitor-I and -II in <i>Saccharomyces cerevisiae</i> , and simple purification by affinity chromatography. Journal of Bioscience and Bioengineering, 1996, 82, 444-447. | 0.9 | 1 |
| 2975 | Bud10p directs axial cell polarization in budding yeast and resembles a transmembrane receptor. Current Biology, 1996, 6, 570-579. | 1.8 | 82 |
| 2976 | GFP tagging of budding yeast chromosomes reveals that protein-protein interactions can mediate sister chromatid cohesion. Current Biology, 1996, 6, 1599-1608. | 1.8 | 580 |
| 2977 | HAM1, the gene controlling 6-N-hydroxylaminopurine sensitivity and mutagenesis in the yeast <i>Saccharomyces cerevisiae</i> . Yeast, 1996, 12, 17-29. | 0.8 | 45 |
| 2978 | Cloning and characterization of the <i>Pichia pastoris</i> PRC1 gene encoding carboxypeptidase Y. Yeast, 1996, 12, 31-40. | 0.8 | 14 |
| 2979 | Separate roles for N- and C-termini of the STE4 (\hat{I}^2) subunit of the <i>Saccharomyces cerevisiae</i> G protein in the mediation of the growth arrest. Lack of growth-arresting activity of mammalian $\hat{I}^2\hat{I}^3$ complexes. Yeast, 1996, 12, 41-51. | 0.8 | 6 |
| 2980 | Identification of a class of <i>Saccharomyces cerevisiae</i> mutants defective in fatty acid repression of gene transcription and analysis of the <i>frm2</i> Gene. Yeast, 1996, 12, 319-331. | 0.8 | 22 |
| 2981 | Functional characterization of the repeated UASINO element in the promoters of the INO1 and CHO2 genes of yeast. , 1996, 12, 653-665. | | 29 |
| 2982 | The <i>KNH1</i> gene of <i>Saccharomyces cerevisiae</i> is a functional homolog of <i>KRE9</i> . , 1996, 12, 683-692. | | 48 |
| 2983 | The <i>Candida albicans</i> PKC1 gene encodes a protein kinase C homolog necessary for cellular integrity but not dimorphism. , 1996, 12, 741-756. | | 75 |
| 2984 | The isolation of a dol-P-man synthase from <i>Ustilago maydis</i> that functions in <i>Saccharomyces cerevisiae</i> . , 1996, 12, 765-771. | | 32 |
| 2985 | Vacuolar and extracellular maturation of <i>Saccharomyces cerevisiae</i> proteinase A. , 1996, 12, 823-832. | | 27 |
| 2986 | Characterization of <i>cwl1+</i> , a gene from <i>Schizosaccharomyces pombe</i> whose overexpression causes cell lysis. , 1996, 12, 983-990. | | 7 |
| 2987 | Cloning of <i>Candida albicans</i> SEC14 gene homologue coding for a putative essential function. Yeast, 1996, 12, 1097-1105. | 0.8 | 45 |
| 2988 | Isolation of a gene encoding a G protein \hat{I}^{\pm} subunit involved in the regulation of cAMP levels in the yeast <i>Kluyveromyces lactis</i> . Yeast, 1996, 12, 1125-1133. | 0.8 | 22 |

| # | ARTICLE | IF | CITATIONS |
|------|--|-----|-----------|
| 2989 | The <i>S. cerevisiae</i> nitrogen starvation-induced Yvh1p and Ptp2p phosphatases play a role in control of sporulation. , 1996, 12, 1135-1151. | | 24 |
| 2990 | Genetic aspects of carbon catabolite repression of the STA2 glucoamylase gene in <i>Saccharomyces cerevisiae</i> . <i>Yeast</i> , 1996, 12, 1297-1300. | 0.8 | 19 |
| 2991 | CtCdc55p and CtHal3p: Two putative regulatory proteins from <i>Candida tropicalis</i> with long acidic domains. , 1996, 12, 1321-1329. | | 14 |
| 2992 | Understanding <i>Candida albicans</i> at the Molecular Level. <i>Yeast</i> , 1996, 12, 1677-1702. | 0.8 | 66 |
| 2993 | Target gene identification: Target specific transcriptional activation by three murine Homeodomain/VP16 hybrid proteins in <i>Saccharomyces cerevisiae</i> . , 1996, 274, 145-156. | | 6 |
| 2994 | Requirements for perpendicular helix pairing. , 1996, 26, 95-107. | | 4 |
| 2995 | Regulatory domain of human protein kinase C δ dominantly inhibits protein kinase C δ -I regulated growth and morphology in <i>Saccharomyces cerevisiae</i> . , 1996, 166, 609-617. | | 13 |
| 2996 | Site-directed replacement of the invariant lysine 73 of <i>Saccharomyces cerevisiae</i> iso-1-cytochrome c with all ribosomally encoded amino acids. <i>Inorganica Chimica Acta</i> , 1996, 242, 97-103. | 1.2 | 13 |
| 2997 | Dissection of fission yeast microtubule associating protein p93 Dis1 : regions implicated in regulated localization and microtubule interaction. <i>Genes To Cells</i> , 1996, 1, 633-644. | 0.5 | 47 |
| 2998 | PAT1 , an evolutionarily conserved acetyltransferase homologue, is required for multiple steps in the cell cycle. <i>Genes To Cells</i> , 1996, 1, 923-942. | 0.5 | 21 |
| 2999 | Signalling from endoplasmic reticulum to nucleus: transcription factor with a basic-leucine zipper motif is required for the unfolded protein-response pathway. <i>Genes To Cells</i> , 1996, 1, 803-817. | 0.5 | 352 |
| 3000 | The GalF protein of <i>Escherichia coli</i> is not a UDP-glucose pyrophosphorylase but interacts with the GalU protein possibly to regulate cellular levels of UDP-glucose. <i>Molecular Microbiology</i> , 1996, 22, 827-840. | 1.2 | 47 |
| 3001 | Isolation of an additional soybean cDNA encoding Ypt/Rab-related small GTP-binding protein and its functional comparison to Sypt using a yeast ypt1-1 mutant. <i>Plant Molecular Biology</i> , 1996, 31, 783-792. | 2.0 | 20 |
| 3002 | The use of inulinase pre-pro leader peptide for secretion of heterologous proteins in <i>Saccharomyces cerevisiae</i> . <i>Biotechnology Letters</i> , 1996, 18, 627-632. | 1.1 | 3 |
| 3003 | Retention of human apolipoprotein A-II by a mutant of <i>Saccharomyces cerevisiae</i> . <i>Biotechnology Letters</i> , 1996, 18, 355-360. | 1.1 | 0 |
| 3004 | The null mutant of <i>Saccharomyces cerevisiae</i> is deficient in cardiolipin and does not survive the disruption of the CHO1 gene encoding phosphatidylserine synthase. <i>FEMS Microbiology Letters</i> , 1996, 140, 43-47. | 0.7 | 29 |
| 3005 | Glutathione is an important antioxidant molecule in the yeast <i>Saccharomyces cerevisiae</i> . <i>FEMS Microbiology Letters</i> , 1996, 141, 207-212. | 0.7 | 126 |
| 3006 | Molecular cloning and characterization of tumour necrosis factor alpha (TNF- α) from the Australian common brushtail possum, <i>Trichosurus vulpecula</i> . <i>Immunology and Cell Biology</i> , 1996, 74, 151-158. | 1.0 | 27 |

| # | ARTICLE | IF | CITATIONS |
|------|--|------|-----------|
| 3007 | A new group of conserved coactivators that increase the specificity of AP-1 transcription factors. <i>Nature</i> , 1996, 383, 453-457. | 13.7 | 441 |
| 3008 | Mouse RNA Polymerase I 16-kDa Subunit Able to Associate with 40-kDa Subunit Is a Homolog of Yeast AC19 Subunit of RNA Polymerases I and III. <i>Journal of Biological Chemistry</i> , 1996, 271, 32881-32885. | 1.6 | 20 |
| 3009 | SCD5, a suppressor of clathrin deficiency, encodes a novel protein with a late secretory function in yeast.. <i>Molecular Biology of the Cell</i> , 1996, 7, 245-260. | 0.9 | 33 |
| 3010 | Actin organization, bristle morphology, and viability are affected by actin capping protein mutations in <i>Drosophila</i> .. <i>Journal of Cell Biology</i> , 1996, 133, 1293-1305. | 2.3 | 98 |
| 3011 | Six novel genes necessary for pre-mRNA splicing in <i>Saccharomyces cerevisiae</i> . <i>Nucleic Acids Research</i> , 1996, 24, 1037-1044. | 6.5 | 21 |
| 3012 | <i>Saccharomyces cerevisiae</i> IRE2/HAC1 is involved in IRE1-mediated KAR2 expression. <i>Nucleic Acids Research</i> , 1996, 24, 4222-4226. | 6.5 | 99 |
| 3013 | SYR2, A Gene Necessary for Syringomycin Growth Inhibition of <i>Saccharomyces Cerevisiae</i> . <i>Microbiology (United Kingdom)</i> , 1996, 142, 477-484. | 0.7 | 48 |
| 3014 | <i>Candida maltosa</i> . , 1996, , 411-580. | | 34 |
| 3015 | Transmembrane Topology of $\hat{1}\pm$ - and $\hat{1}^2$ -Subunits of Na ⁺ ,K ⁺ -ATPase Derived from $\hat{1}^2$ -Galactosidase Fusion Proteins Expressed in Yeast. <i>Journal of Biological Chemistry</i> , 1996, 271, 29312-29320. | 1.6 | 20 |
| 3016 | Isolation and Characterization of SUG2. <i>Journal of Biological Chemistry</i> , 1996, 271, 32810-32817. | 1.6 | 60 |
| 3017 | Aberrant mitosis in fission yeast mutants defective in fatty acid synthetase and acetyl CoA carboxylase.. <i>Journal of Cell Biology</i> , 1996, 134, 949-961. | 2.3 | 105 |
| 3018 | Probing Conserved Regions of the Cytoplasmic LOOP1 Segment Linking Transmembrane Segments 2 and 3 of the <i>Saccharomyces cerevisiae</i> Plasma Membrane H ⁺ -ATPase. <i>Journal of Biological Chemistry</i> , 1996, 271, 25438-25445. | 1.6 | 28 |
| 3019 | The 110-kD spindle pole body component of <i>Saccharomyces cerevisiae</i> is a phosphoprotein that is modified in a cell cycle-dependent manner.. <i>Journal of Cell Biology</i> , 1996, 132, 903-914. | 2.3 | 45 |
| 3020 | Functionally redundant interactions between U2 and U6 spliceosomal snRNAs.. <i>Genes and Development</i> , 1996, 10, 489-501. | 2.7 | 37 |
| 3021 | A Novel RING Finger Protein, Vps8p, Functionally Interacts with the Small GTPase, Vps21p, to Facilitate Soluble Vacuolar Protein Localization. <i>Journal of Biological Chemistry</i> , 1996, 271, 33607-33615. | 1.6 | 73 |
| 3022 | Regulatory Elements That Control Transcription Activation and Unsaturated Fatty Acid-mediated Repression of the <i>Saccharomyces cerevisiae</i> OLE1 Gene. <i>Journal of Biological Chemistry</i> , 1996, 271, 3581-3589. | 1.6 | 144 |
| 3023 | The Yeast VRG4 Gene Is Required for Normal Golgi Functions and Defines a New Family of Related Genes. <i>Journal of Biological Chemistry</i> , 1996, 271, 3837-3845. | 1.6 | 63 |
| 3024 | TOR controls translation initiation and early G1 progression in yeast.. <i>Molecular Biology of the Cell</i> , 1996, 7, 25-42. | 0.9 | 680 |

| # | ARTICLE | IF | CITATIONS |
|------|---|-----|-----------|
| 3025 | Site-directed Mutagenesis of the Cysteine Residues in the Neurospora crassa Plasma Membrane H ⁺ -ATPase. Journal of Biological Chemistry, 1996, 271, 367-371. | 1.6 | 17 |
| 3026 | Sequence Requirements for Mitochondrial Import of Yeast Cytochrome c. Journal of Biological Chemistry, 1996, 271, 6594-6604. | 1.6 | 36 |
| 3027 | ADR1 Activation Domains Contact the Histone Acetyltransferase GCN5 and the Core Transcriptional Factor TFIIB. Journal of Biological Chemistry, 1996, 271, 32359-32365. | 1.6 | 85 |
| 3028 | Biochemical Studies of Saccharomyces cerevisiae Myristoyl-coenzyme A:Protein N-Myristoyltransferase Mutants. Journal of Biological Chemistry, 1996, 271, 33131-33140. | 1.6 | 23 |
| 3029 | ROD1, a Novel Gene Conferring Multiple Resistance Phenotypes in Saccharomyces cerevisiae. Journal of Biological Chemistry, 1996, 271, 2914-2920. | 1.6 | 19 |
| 3030 | Identification, localization, and functional implications of an abundant nematode annexin.. Journal of Cell Biology, 1996, 132, 1079-1092. | 2.3 | 35 |
| 3031 | Cell fusion during yeast mating requires high levels of a-factor mating pheromone.. Journal of Cell Biology, 1996, 135, 1727-1739. | 2.3 | 70 |
| 3032 | The Ydj1 molecular chaperone facilitates formation of active p60v-src in yeast.. Molecular Biology of the Cell, 1996, 7, 91-100. | 0.9 | 66 |
| 3033 | Nuclear transport defects and nuclear envelope alterations are associated with mutation of the Saccharomyces cerevisiae NPL4 gene.. Molecular Biology of the Cell, 1996, 7, 1835-1855. | 0.9 | 61 |
| 3034 | Identification of an overexpressed yeast gene which prevents aminoglycoside toxicity. Microbiology (United Kingdom), 1996, 142, 3407-3414. | 0.7 | 47 |
| 3035 | Functional analysis in Saccharomyces cerevisiae of naturally occurring amino acid substitutions in human dihydrolipoamide dehydrogenase. Human Molecular Genetics, 1996, 5, 1643-1648. | 1.4 | 17 |
| 3036 | Alanine-scanning Mutagenesis along Membrane Segment 4 of the Yeast Plasma Membrane H ⁺ -ATPase EFFECTS ON STRUCTURE AND FUNCTION. Journal of Biological Chemistry, 1996, 271, 22999-23005. | 1.6 | 40 |
| 3037 | Assembly and Function of a Cytosolic Form of NADH-specific Isocitrate Dehydrogenase in Yeast. Journal of Biological Chemistry, 1996, 271, 10347-10352. | 1.6 | 19 |
| 3038 | Transcriptional Activation by Yeast PDR1p Is Inhibited by Its Association with NGG1p/ADA3p. Journal of Biological Chemistry, 1996, 271, 15884-15890. | 1.6 | 59 |
| 3039 | Yeast TAF(II)90 is required for cell-cycle progression through G2/M but not for general transcription activation.. Genes and Development, 1996, 10, 2368-2380. | 2.7 | 143 |
| 3040 | Selection of axial growth sites in yeast requires Axl2p, a novel plasma membrane glycoprotein.. Genes and Development, 1996, 10, 777-793. | 2.7 | 132 |
| 3041 | The PRP31 gene encodes a novel protein required for pre-mRNA splicing in Saccharomyces cerevisiae. Nucleic Acids Research, 1996, 24, 1164-1170. | 6.5 | 39 |
| 3042 | MIG1 overexpression causes flocculation in Saccharomyces cerevisiae. Microbiology (United Kingdom) Tj ETQq1 1 0.784314 rgBT /Overlock 10 | 0.9 | 9 |

| # | ARTICLE | IF | CITATIONS |
|------|--|-----|-----------|
| 3043 | Isolation of the mRNA-capping enzyme and ferric-reductase-related genes from <i>Candida albicans</i> . <i>Microbiology (United Kingdom)</i> , 1996, 142, 2515-2523. | 0.7 | 51 |
| 3044 | HDF2, the Second Subunit of the Ku Homologue from <i>Saccharomyces cerevisiae</i> . <i>Journal of Biological Chemistry</i> , 1996, 271, 27765-27769. | 1.6 | 76 |
| 3045 | Novel syntaxin homologue, Pep12p, required for the sorting of luminal hydrolases to the lysosome-like vacuole in yeast.. <i>Molecular Biology of the Cell</i> , 1996, 7, 579-594. | 0.9 | 282 |
| 3046 | The sequence NPFXD defines a new class of endocytosis signal in <i>Saccharomyces cerevisiae</i> .. <i>Journal of Cell Biology</i> , 1996, 135, 1789-1800. | 2.3 | 105 |
| 3047 | A new pathway for protein export in <i>Saccharomyces cerevisiae</i> .. <i>Journal of Cell Biology</i> , 1996, 133, 1017-1026. | 2.3 | 202 |
| 3048 | Yeast src homology region 3 domain-binding proteins involved in bud formation.. <i>Journal of Cell Biology</i> , 1996, 133, 865-878. | 2.3 | 76 |
| 3049 | CDNA cloning of p112, the largest regulatory subunit of the human 26s proteasome, and functional analysis of its yeast homologue, sen3p.. <i>Molecular Biology of the Cell</i> , 1996, 7, 853-870. | 0.9 | 50 |
| 3050 | Differential Structural Requirements for Interaction of Ras Protein with Its Distinct Downstream Effectors. <i>Journal of Biological Chemistry</i> , 1996, 271, 5353-5360. | 1.6 | 64 |
| 3051 | Directed mutagenesis of YAC-cloned DNA using a rapid, PCR-based screening protocol. <i>Nucleic Acids Research</i> , 1996, 24, 3467-3468. | 6.5 | 4 |
| 3052 | Acid Proteinase Secreted by <i>Candida Tropicalis</i> : Functional Analysis of Preproregion Cleavages in <i>C. Tropicalis</i> and <i>Saccharomyces Cerevisiae</i> . <i>Microbiology (United Kingdom)</i> , 1996, 142, 493-503. | 0.7 | 48 |
| 3053 | The Light Chain Subunit Is Required for Clathrin Function in <i>Saccharomyces cerevisiae</i> . <i>Journal of Biological Chemistry</i> , 1996, 271, 33123-33130. | 1.6 | 67 |
| 3054 | Two Yeast Homologs of ECA39, a Target for c-Myc Regulation, Code for Cytosolic and Mitochondrial Branched-chain Amino Acid Aminotransferases. <i>Journal of Biological Chemistry</i> , 1996, 271, 20242-20245. | 1.6 | 103 |
| 3055 | Involvement of the <i>Saccharomyces cerevisiae</i> HDF1 Gene in DNA Double-strand Break Repair and Recombination. <i>Journal of Biological Chemistry</i> , 1996, 271, 7910-7915. | 1.6 | 60 |
| 3056 | ATP Synthase of Yeast Mitochondria. <i>Journal of Biological Chemistry</i> , 1996, 271, 20284-20290. | 1.6 | 86 |
| 3057 | The OST4 Gene of <i>Saccharomyces cerevisiae</i> Encodes an Unusually Small Protein Required for Normal Levels of Oligosaccharyltransferase Activity. <i>Journal of Biological Chemistry</i> , 1996, 271, 3132-3140. | 1.6 | 61 |
| 3058 | The NH2-terminal Extension of Protein Phosphatase PPZ1 Has an Essential Functional Role. <i>Journal of Biological Chemistry</i> , 1996, 271, 26349-26355. | 1.6 | 59 |
| 3059 | The PAL1 gene product is a peroxisomal ATP-binding cassette transporter in the yeast <i>Saccharomyces cerevisiae</i> .. <i>Journal of Cell Biology</i> , 1996, 132, 549-563. | 2.3 | 74 |
| 3060 | Selective Activation of MEK1 but Not MEK2 by A-Raf from Epidermal Growth Factor-stimulated Hela Cells. <i>Journal of Biological Chemistry</i> , 1996, 271, 3265-3271. | 1.6 | 104 |

| # | ARTICLE | IF | CITATIONS |
|------|--|-----|-----------|
| 3061 | Expression of Rat Aspartyl-tRNA Synthetase in <i>Saccharomyces cerevisiae</i> . <i>Journal of Biological Chemistry</i> , 1996, 271, 29295-29303. | 1.6 | 26 |
| 3062 | The ARG11 Gene of <i>Saccharomyces cerevisiae</i> Encodes a Mitochondrial Integral Membrane Protein Required for Arginine Biosynthesis. <i>Journal of Biological Chemistry</i> , 1996, 271, 25011-25018. | 1.6 | 53 |
| 3063 | Degradation of Aspartic Proteinase-I with Mutated Prosequences Occurs in the Endoplasmic Reticulum of. <i>Journal of Biological Chemistry</i> , 1996, 271, 14252-14255. | 1.6 | 12 |
| 3064 | A widespread transposable element masks expression of a yeast copper transport gene.. <i>Genes and Development</i> , 1996, 10, 1917-1929. | 2.7 | 239 |
| 3065 | <i>Kluyveromyces lactis</i> killer system: analysis of cytoplasmic promoters of the linear plasmids. <i>Nucleic Acids Research</i> , 1996, 24, 1879-1886. | 6.5 | 39 |
| 3066 | Structural and Functional Analysis of Yeast Putative Adaptors. <i>Journal of Biological Chemistry</i> , 1996, 271, 5237-5245. | 1.6 | 113 |
| 3067 | Amino Acid and Adenine Cross-pathway Regulation Act through the Same 5â€²-TGACTC-3â€² Motif in the Yeast HIS7 Promoter. <i>Journal of Biological Chemistry</i> , 1996, 271, 29637-29643. | 1.6 | 35 |
| 3068 | Function and Expression of Flavohemoglobin in <i>Saccharomyces cerevisiae</i> . <i>Journal of Biological Chemistry</i> , 1996, 271, 25131-25138. | 1.6 | 73 |
| 3069 | Structure/Function Properties of the Yeast Dual Regulator Protein NGG1 That Are Required for Glucose Repression. <i>Journal of Biological Chemistry</i> , 1996, 271, 9298-9306. | 1.6 | 34 |
| 3070 | Cloning of a cDNA Encoded by a Member of the <i>Arabidopsis thaliana</i> ATP Sulfurylase Multigene Family. <i>Journal of Biological Chemistry</i> , 1996, 271, 12227-12233. | 1.6 | 90 |
| 3071 | A Recombinant Sickle Hemoglobin Triple Mutant with Independent Inhibitory Effects on Polymerization. <i>Journal of Biological Chemistry</i> , 1996, 271, 25152-25156. | 1.6 | 16 |
| 3072 | Different Glycosylation Requirements for the Synthesis of Enzymatically Active Angiotensin-converting Enzyme in Mammalian Cells and Yeast. <i>Journal of Biological Chemistry</i> , 1996, 271, 6429-6434. | 1.6 | 30 |
| 3073 | A connection between pre-mRNA splicing and the cell cycle in fission yeast: <i>cdc28+</i> is allelic with <i>prp8+</i> and encodes an RNA-dependent ATPase/helicase.. <i>Molecular Biology of the Cell</i> , 1996, 7, 1083-1094. | 0.9 | 71 |
| 3074 | GLE2, a <i>Saccharomyces cerevisiae</i> homologue of the <i>Schizosaccharomyces pombe</i> export factor RAE1, is required for nuclear pore complex structure and function.. <i>Molecular Biology of the Cell</i> , 1996, 7, 1921-1937. | 0.9 | 173 |
| 3075 | A Novel Family of TRF (DNA Topoisomerase I-Related Function) Genes Required for Proper Nuclear Segregation. <i>Nucleic Acids Research</i> , 1996, 24, 2404-2410. | 6.5 | 77 |
| 3076 | The effects of clathrin inactivation on localization of Kex2 protease are independent of the TGN localization signal in the cytosolic tail of Kex2p.. <i>Molecular Biology of the Cell</i> , 1996, 7, 1667-1677. | 0.9 | 45 |
| 3077 | Immunodominant epitopes defined by a yeast-expressed library of random fragments of the rabies virus glycoprotein map outside major antigenic sites. <i>Journal of General Virology</i> , 1996, 77, 339-346. | 1.3 | 39 |
| 3078 | Gat1p, a GATA Family Protein Whose Production Is Sensitive to Nitrogen Catabolite Repression, Participates in Transcriptional Activation of Nitrogen-Catabolic Genes in <i>Saccharomyces cerevisiae</i> . <i>Molecular and Cellular Biology</i> , 1996, 16, 847-858. | 1.1 | 137 |

| # | ARTICLE | IF | CITATIONS |
|------|---|-----|-----------|
| 3079 | CHARACTERISATION OF AMYLOLYTIC BREWING YEAST. <i>Journal of the Institute of Brewing</i> , 1996, 102, 27-32. | 0.8 | 6 |
| 3080 | Transcriptional Regulation of a Sterol-Biosynthetic Enzyme by Sterol Levels in <i>Saccharomyces cerevisiae</i> . <i>Molecular and Cellular Biology</i> , 1996, 16, 3981-3989. | 1.1 | 55 |
| 3081 | Sst2, a Negative Regulator of Pheromone Signaling in the Yeast <i>Saccharomyces cerevisiae</i> : Expression, Localization, and Genetic Interaction and Physical Association with Gpa1 (the G-Protein β) Tj ETQq0 0.1gBT /Overb 10 T | 0.1 | 10 |
| 3082 | Transformation of Lithium-Treated Yeast Cells and the Selection of Auxotrophic and Dominant Markers. , 1996, 53, 139-146. | | 15 |
| 3083 | Steroid 21-Hydroxylase Autoantibodies: Measurements with a New Immunoprecipitation Assay ¹ . <i>Journal of Clinical Endocrinology and Metabolism</i> , 1997, 82, 1440-1446. | 1.8 | 80 |
| 3084 | Estrogen Receptor Residues Required for Stereospecific Ligand Recognition and Activation. <i>Molecular Endocrinology</i> , 1997, 11, 587-594. | 3.7 | 16 |
| 3085 | Cellular distribution of mammalian DNA topoisomerase II is determined by its catalytically dispensable C-terminal domain. <i>Nucleic Acids Research</i> , 1997, 25, 3135-3142. | 6.5 | 49 |
| 3086 | A Multispecificity Syntaxin Homologue, Vam3p, Essential for Autophagic and Biosynthetic Protein Transport to the Vacuole. <i>Journal of Cell Biology</i> , 1997, 138, 517-529. | 2.3 | 332 |
| 3087 | The Yeast Gene, MDM20, Is Necessary for Mitochondrial Inheritance and Organization of the Actin Cytoskeleton. <i>Journal of Cell Biology</i> , 1997, 137, 141-153. | 2.3 | 130 |
| 3088 | Homeodomain-DNA interactions of the Pho2 protein are promoter-dependent. <i>Nucleic Acids Research</i> , 1997, 25, 4730-4739. | 6.5 | 10 |
| 3089 | Active Site Mutations in Yeast Protein Disulfide Isomerase Cause Dithiothreitol Sensitivity and a Reduced Rate of Protein Folding in the Endoplasmic Reticulum. <i>Journal of Cell Biology</i> , 1997, 138, 1229-1238. | 2.3 | 76 |
| 3090 | Evidence for physical and functional interactions among two <i>Saccharomyces cerevisiae</i> SH3 domain proteins, an adenyl cyclase-associated protein and the actin cytoskeleton.. <i>Molecular Biology of the Cell</i> , 1997, 8, 367-385. | 0.9 | 162 |
| 3091 | The unique hetero-oligomeric nature of the subunits in the catalytic cooperativity of the yeast Cct chaperonin complex. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 1997, 94, 10780-10785. | 3.3 | 70 |
| 3092 | The VRG4 Gene Is Required for GDP-mannose Transport into the Lumen of the Golgi in the Yeast, <i>Saccharomyces cerevisiae</i> . <i>Journal of Biological Chemistry</i> , 1997, 272, 31908-31914. | 1.6 | 123 |
| 3093 | Identification of three regions essential for interaction between a β -like factor and core RNA polymerase. <i>Genes and Development</i> , 1997, 11, 2897-2909. | 2.7 | 45 |
| 3094 | Mdm12p, a Component Required for Mitochondrial Inheritance That Is Conserved between Budding and Fission Yeast. <i>Journal of Cell Biology</i> , 1997, 136, 545-553. | 2.3 | 185 |
| 3095 | Marmoset CYP1A2: primary structure and constitutive expression in livers. <i>Carcinogenesis</i> , 1997, 18, 1985-1991. | 1.3 | 22 |
| 3096 | Affinity Purification and Kinetic Analysis of Mutant Forms of Yeast NAD ⁺ -specific Isocitrate Dehydrogenase. <i>Journal of Biological Chemistry</i> , 1997, 272, 21811-21817. | 1.6 | 32 |

| # | ARTICLE | IF | CITATIONS |
|------|--|-----|-----------|
| 3097 | Aip3p/Bud6p, a yeast actin-interacting protein that is involved in morphogenesis and the selection of bipolar budding sites.. <i>Molecular Biology of the Cell</i> , 1997, 8, 729-753. | 0.9 | 169 |
| 3098 | Ras2 and Ras1 Protein Phosphorylation in <i>Saccharomyces cerevisiae</i> . <i>Journal of Biological Chemistry</i> , 1997, 272, 18790-18800. | 1.6 | 17 |
| 3099 | A Nuclear Export Signal in Kap95p Is Required for Both Recycling the Import Factor and Interaction with the Nucleoporin GLFG Repeat Regions of Nup116p and Nup100p. <i>Journal of Cell Biology</i> , 1997, 137, 797-811. | 2.3 | 84 |
| 3100 | A <i>Candida albicans</i> RAS-related gene (CaRSRI) is involved in budding, cell morphogenesis and hypha development. <i>Microbiology (United Kingdom)</i> , 1997, 143, 3033-3044. | 0.7 | 74 |
| 3101 | Human Bcl-2 Reverses Survival Defects in Yeast Lacking Superoxide Dismutase and Delays Death of Wild-Type Yeast. <i>Journal of Cell Biology</i> , 1997, 137, 1581-1588. | 2.3 | 203 |
| 3102 | Osmotic Balance Regulates Cell Fusion during Mating in <i>Saccharomyces cerevisiae</i> . <i>Journal of Cell Biology</i> , 1997, 138, 961-974. | 2.3 | 125 |
| 3103 | A novel Sec18p/NSF-dependent complex required for Golgi-to-endosome transport in yeast.. <i>Molecular Biology of the Cell</i> , 1997, 8, 1089-1104. | 0.9 | 153 |
| 3104 | Intracellular Sequestration of Sodium by a Novel Na ⁺ /H ⁺ Exchanger in Yeast Is Enhanced by Mutations in the Plasma Membrane H ⁺ -ATPase. <i>Journal of Biological Chemistry</i> , 1997, 272, 26145-26152. | 1.6 | 233 |
| 3105 | Tubulin Subunits Exist in an Activated Conformational State Generated and Maintained by Protein Cofactors. <i>Journal of Cell Biology</i> , 1997, 138, 821-832. | 2.3 | 190 |
| 3106 | <i>Saccharomyces cerevisiae</i> Homologs of Mammalian B and B ² Subunits of Protein Phosphatase 2A Direct the Enzyme to Distinct Cellular Functions. <i>Journal of Biological Chemistry</i> , 1997, 272, 8256-8262. | 1.6 | 97 |
| 3107 | Endoplasmic Reticulum Stress-induced mRNA Splicing Permits Synthesis of Transcription Factor Hac1p/Ern4p That Activates the Unfolded Protein Response. <i>Molecular Biology of the Cell</i> , 1997, 8, 1845-1862. | 0.9 | 263 |
| 3108 | A Novel RING Finger Protein Complex Essential for a Late Step in Protein Transport to the Yeast Vacuole. <i>Molecular Biology of the Cell</i> , 1997, 8, 2307-2327. | 0.9 | 290 |
| 3109 | A sorting nexin-1 homologue, Vps5p, forms a complex with Vps17p and is required for recycling the vacuolar protein-sorting receptor.. <i>Molecular Biology of the Cell</i> , 1997, 8, 1529-1541. | 0.9 | 209 |
| 3110 | Targeting of Chitin Synthase 3 to Polarized Growth Sites in Yeast Requires Chs5p and Myo2p. <i>Journal of Cell Biology</i> , 1997, 136, 95-110. | 2.3 | 196 |
| 3111 | Endosome to Golgi Retrieval of the Vacuolar Protein Sorting Receptor, Vps10p, Requires the Function of the VPS29, VPS30, and VPS35 Gene Products. <i>Journal of Cell Biology</i> , 1997, 137, 79-92. | 2.3 | 368 |
| 3112 | Phosphorylation- and ubiquitin-dependent degradation of the cyclin-dependent kinase inhibitor Far1p in budding yeast. <i>Genes and Development</i> , 1997, 11, 3046-3060. | 2.7 | 191 |
| 3113 | Molecular Cloning and Cell Cycle-dependent Expression of Mammalian CRM1, a Protein Involved in Nuclear Export of Proteins. <i>Journal of Biological Chemistry</i> , 1997, 272, 29742-29751. | 1.6 | 188 |
| 3114 | Characterization of an ADP-ribosylation Factor-like 1 Protein in <i>Saccharomyces cerevisiae</i> . <i>Journal of Biological Chemistry</i> , 1997, 272, 30998-31005. | 1.6 | 82 |

| # | ARTICLE | IF | CITATIONS |
|------|--|-----|-----------|
| 3115 | Sec3p is involved in secretion and morphogenesis in <i>Saccharomyces cerevisiae</i> . <i>Molecular Biology of the Cell</i> , 1997, 8, 647-662. | 0.9 | 98 |
| 3116 | The AtVAM3 Encodes a Syntaxin-related Molecule Implicated in the Vacuolar Assembly in <i>Arabidopsis thaliana</i> . <i>Journal of Biological Chemistry</i> , 1997, 272, 24530-24535. | 1.6 | 109 |
| 3117 | Sequence requirements of the bidirectional yeast TRP4 mRNA 3'-end formation signal. <i>Nucleic Acids Research</i> , 1997, 25, 417-422. | 6.5 | 10 |
| 3118 | The Carboxyl Terminus of the <i>Saccharomyces cerevisiae</i> Succinate Dehydrogenase Membrane Subunit, SDH4p, Is Necessary for Ubiquinone Reduction and Enzyme Stability. <i>Journal of Biological Chemistry</i> , 1997, 272, 31382-31388. | 1.6 | 28 |
| 3119 | A Novel Membrane-associated Metalloprotease, Ste24p, Is Required for the First Step of NH ₂ -terminal Processing of the Yeast α -Factor Precursor. <i>Journal of Cell Biology</i> , 1997, 136, 271-285. | 2.3 | 149 |
| 3120 | The Rho-GEF Rom2p Localizes to Sites of Polarized Cell Growth and Participates in Cytoskeletal Functions in <i>Saccharomyces cerevisiae</i> . <i>Molecular Biology of the Cell</i> , 1997, 8, 1829-1844. | 0.9 | 94 |
| 3121 | Characterization of <i>Saccharomyces cerevisiae</i> dna2 Mutants Suggests a Role for the Helicase Late in S Phase. <i>Molecular Biology of the Cell</i> , 1997, 8, 2519-2537. | 0.9 | 58 |
| 3122 | Tam1, a telomere-associated meiotic protein, functions in chromosome synapsis and crossover interference.. <i>Genes and Development</i> , 1997, 11, 1786-1800. | 2.7 | 166 |
| 3123 | MNN6, a Member of the KRE2/MNT1 Family, Is the Gene for Mannosylphosphate Transfer in <i>Saccharomyces cerevisiae</i> . <i>Journal of Biological Chemistry</i> , 1997, 272, 18117-18124. | 1.6 | 76 |
| 3124 | AP1-mediated Multidrug Resistance in <i>Saccharomyces cerevisiae</i> Requires FLR1 Encoding a Transporter of the Major Facilitator Superfamily. <i>Journal of Biological Chemistry</i> , 1997, 272, 19304-19313. | 1.6 | 188 |
| 3125 | Evidence That Intramolecular Interactions Are Involved in Masking the Activation Domain of Transcriptional Activator Leu3p. <i>Journal of Biological Chemistry</i> , 1997, 272, 19383-19392. | 1.6 | 34 |
| 3126 | Glucose-6-P Control of Glycogen Synthase Phosphorylation in Yeast. <i>Journal of Biological Chemistry</i> , 1997, 272, 22495-22501. | 1.6 | 36 |
| 3127 | The Yeast Nucleolar Protein Nop4p Contains Four RNA Recognition Motifs Necessary for Ribosome Biogenesis. <i>Journal of Biological Chemistry</i> , 1997, 272, 25345-25352. | 1.6 | 28 |
| 3128 | The Adjacent Yeast Genes ARO4 and HIS7 Carry No Intergenic Region. <i>Journal of Biological Chemistry</i> , 1997, 272, 26318-26324. | 1.6 | 15 |
| 3129 | V1-situated Stalk Subunits of the Yeast Vacuolar Proton-translocating ATPase. <i>Journal of Biological Chemistry</i> , 1997, 272, 26787-26793. | 1.6 | 94 |
| 3130 | Diazaborine Resistance in the Yeast <i>Saccharomyces cerevisiae</i> Reveals a Link between YAP1 and the Pleiotropic Drug Resistance Genes PDR1 and PDR3. <i>Journal of Biological Chemistry</i> , 1997, 272, 27091-27098. | 1.6 | 75 |
| 3131 | Functional Expression of a cDNA to Human Acyl-coenzyme A:Cholesterol Acyltransferase in Yeast. <i>Journal of Biological Chemistry</i> , 1997, 272, 3980-3985. | 1.6 | 56 |
| 3132 | <i>BIM1</i> Encodes a Microtubule-binding Protein in Yeast. <i>Molecular Biology of the Cell</i> , 1997, 8, 2677-2691. | 0.9 | 179 |

| # | ARTICLE | IF | CITATIONS |
|------|---|-----|-----------|
| 3133 | Overexpression of the G1-cyclin Gene CLN2 Represses the Mating Pathway in <i>Saccharomyces cerevisiae</i> at the Level of the MEK1. <i>Journal of Biological Chemistry</i> , 1997, 272, 13180-13188. | 1.6 | 33 |
| 3134 | A Key Amino Acid Responsible for Substrate Selectivity of Monoamine Oxidase A and B. <i>Journal of Biological Chemistry</i> , 1997, 272, 14033-14036. | 1.6 | 75 |
| 3135 | Reconstitution of a MEK1-Independent Checkpoint in Yeast by Expression of a Novel Human Forkhead cDNA. <i>Molecular and Cellular Biology</i> , 1997, 17, 3037-3046. | 1.1 | 98 |
| 3136 | EXPRESSION OF THE STA2 GLUCOAMYLASE GENE OF <i>SACCHAROMYCES CEREVISIAE</i> IN BREWERS' YEAST. <i>Journal of the Institute of Brewing</i> , 1997, 103, 35-39. | 0.8 | 1 |
| 3137 | The <i>Saccharomyces cerevisiae</i> ACR3 Gene Encodes a Putative Membrane Protein Involved in Arsenite Transport. <i>Journal of Biological Chemistry</i> , 1997, 272, 30061-30066. | 1.6 | 223 |
| 3138 | Fluorescence Probing of Yeast Actin Subdomain 3/4 Hydrophobic Loop 262-274. <i>Journal of Biological Chemistry</i> , 1997, 272, 16829-16837. | 1.6 | 75 |
| 3139 | <i>Saccharomyces cerevisiae</i> Basic Region-Leucine Zipper Protein Regulatory Networks Converge at the ATR1 Structural Gene. <i>Journal of Biological Chemistry</i> , 1997, 272, 23224-23230. | 1.6 | 64 |
| 3140 | Pyridinyl Imidazole Inhibitors of p38 Mitogen-activated Protein Kinase Bind in the ATP Site. <i>Journal of Biological Chemistry</i> , 1997, 272, 12116-12121. | 1.6 | 519 |
| 3141 | The <i>Saccharomyces cerevisiae</i> AP-1 Protein Discriminates between Oxidative Stress Elicited by the Oxidants H ₂ O ₂ and Diamide. <i>Journal of Biological Chemistry</i> , 1997, 272, 7908-7914. | 1.6 | 97 |
| 3142 | Protein-protein interactions in the synaptonemal complex. <i>Molecular Biology of the Cell</i> , 1997, 8, 1405-1414. | 0.9 | 54 |
| 3143 | Yeast counterparts of subunits S5a and p58 (S3) of the human 26S proteasome are encoded by two multicopy suppressors of nin1-1. <i>Molecular Biology of the Cell</i> , 1997, 8, 171-187. | 0.9 | 98 |
| 3144 | A Far Upstream Cis-element Is Required for Wilms' Tumor-1 (WT1) Gene Expression in Renal Cell Culture. <i>Journal of Biological Chemistry</i> , 1997, 272, 32836-32846. | 1.6 | 20 |
| 3145 | Biochemical Characterization of the Wilson Disease Protein and Functional Expression in the Yeast <i>Saccharomyces cerevisiae</i> . <i>Journal of Biological Chemistry</i> , 1997, 272, 21461-21466. | 1.6 | 296 |
| 3146 | p32 Protein, a Splicing Factor 2-associated Protein, Is Localized in Mitochondrial Matrix and Is Functionally Important in Maintaining Oxidative Phosphorylation. <i>Journal of Biological Chemistry</i> , 1997, 272, 24363-24370. | 1.6 | 196 |
| 3147 | Heterologous Expression of Human Cholecystokinin in <i>Saccharomyces cerevisiae</i> . <i>Journal of Biological Chemistry</i> , 1997, 272, 9720-9727. | 1.6 | 23 |
| 3148 | Novel Cdc42-binding proteins Gic1 and Gic2 control cell polarity in yeast. <i>Genes and Development</i> , 1997, 11, 2972-2982. | 2.7 | 161 |
| 3149 | The Transcriptional Activators BAS1, BAS2, and ABF1 Bind Positive Regulatory Sites as the Critical Elements for Adenine Regulation of ADE5,7. <i>Journal of Biological Chemistry</i> , 1997, 272, 13343-13354. | 1.6 | 32 |
| 3150 | A novel cytosolic regulator, Pianissimo, is required for chemoattractant receptor and G protein-mediated activation of the 12 transmembrane domain adenylyl cyclase in <i>Dictyostelium</i> . <i>Genes and Development</i> , 1997, 11, 3218-3231. | 2.7 | 118 |

| # | ARTICLE | IF | CITATIONS |
|------|--|-----|-----------|
| 3151 | Saccharomyces cerevisiae VIC9 Encodes GDP-mannose Pyrophosphorylase, Which Is Essential for Protein Glycosylation. Journal of Biological Chemistry, 1997, 272, 16308-16314. | 1.6 | 82 |
| 3152 | In Vivo Dynamics of Nuclear Pore Complexes in Yeast. Journal of Cell Biology, 1997, 136, 1185-1199. | 2.3 | 113 |
| 3153 | The unfolded protein response coordinates the production of endoplasmic reticulum protein and endoplasmic reticulum membrane.. Molecular Biology of the Cell, 1997, 8, 1805-1814. | 0.9 | 357 |
| 3154 | The Srp40 Protein Plays a Dose-Sensitive Role in Preribosome Assembly or Transport and Depends on Its Carboxy-Terminal Domain for Proper Localization to the Yeast Nucleoskeleton. DNA and Cell Biology, 1997, 16, 1161-1173. | 0.9 | 11 |
| 3155 | Complementation of the DNA repair-deficient swi10 mutant of fission yeast by the human ERCC1 gene. Nucleic Acids Research, 1997, 25, 2823-2827. | 6.5 | 17 |
| 3156 | The RAD5 gene product is involved in the avoidance of non-homologous end-joining of DNA double strand breaks in the yeast Saccharomyces cerevisiae. Nucleic Acids Research, 1997, 25, 743-749. | 6.5 | 53 |
| 3157 | Genetic and Biochemical Studies on Molecular Breeding of Yeast Cells. Nippon Nogeikagaku Kaishi, 1997, 71, 691-697. | 0.0 | 1 |
| 3158 | A glutamate residue in the catalytic center of the yeast chorismate mutase restricts enzyme activity to acidic conditions. Proceedings of the National Academy of Sciences of the United States of America, 1997, 94, 8491-8496. | 3.3 | 38 |
| 3159 | Hrp1, a sequence-specific RNA-binding protein that shuttles between the nucleus and the cytoplasm, is required for mRNA 3' end formation in yeast. Genes and Development, 1997, 11, 2545-2556. | 2.7 | 213 |
| 3160 | The Ubiquitin-Conjugating Enzyme Rad6 (Ubc2) Is Required for Silencing in <i>Saccharomyces cerevisiae</i> . Molecular and Cellular Biology, 1997, 17, 6693-6699. | 1.1 | 126 |
| 3161 | Effect of association with adenylyl cyclase-associated protein on the interaction of yeast adenylyl cyclase with Ras protein. Molecular and Cellular Biology, 1997, 17, 1057-1064. | 1.1 | 57 |
| 3162 | Template Topology and Transcription: Chromatin Templates Relaxed by Localized Linearization Are Transcriptionally Active in Yeast. Molecular and Cellular Biology, 1997, 17, 2825-2834. | 1.1 | 17 |
| 3163 | Nop2p Is Required for Pre-rRNA Processing and 60S Ribosome Subunit Synthesis in Yeast. Molecular and Cellular Biology, 1997, 17, 378-388. | 1.1 | 142 |
| 3164 | The DNA Binding and Activation Domains of Gal4p Are Sufficient for Conveying Its Regulatory Signals. Molecular and Cellular Biology, 1997, 17, 2538-2549. | 1.1 | 31 |
| 3165 | Structure-Function Analysis of <i>TAF130</i> : Identification and Characterization of a High-Affinity TATA-Binding Protein Interaction Domain in the N Terminus of Yeast TAF ₁₃₀ . Molecular and Cellular Biology, 1997, 17, 3081-3093. | 1.1 | 66 |
| 3166 | Analysis of a Meiosis-Specific URS1 Site: Sequence Requirements and Involvement of Replication Protein A. Molecular and Cellular Biology, 1997, 17, 3536-3546. | 1.1 | 18 |
| 3167 | A Family of Ammonium Transporters in <i>Saccharomyces cerevisiae</i> . Molecular and Cellular Biology, 1997, 17, 4282-4293. | 1.1 | 516 |
| 3168 | Small Heat Shock Protein Suppression of Vpr-Induced Cytoskeletal Defects in Budding Yeast. Molecular and Cellular Biology, 1997, 17, 4033-4042. | 1.1 | 37 |

| # | ARTICLE | IF | CITATIONS |
|------|--|-----|-----------|
| 3169 | Regulation of the Protein Kinase PKR by the Vaccinia Virus Pseudosubstrate Inhibitor K3L Is Dependent on Residues Conserved between the K3L Protein and the PKR Substrate eIF2 γ . <i>Molecular and Cellular Biology</i> , 1997, 17, 4146-4158. | 1.1 | 125 |
| 3170 | Extensive Mutagenesis of a Transcriptional Activation Domain Identifies Single Hydrophobic and Acidic Amino Acids Important for Activation In Vivo. <i>Molecular and Cellular Biology</i> , 1997, 17, 115-122. | 1.1 | 65 |
| 3171 | Multiple Regulatory Domains on the Byr2 Protein Kinase. <i>Molecular and Cellular Biology</i> , 1997, 17, 5876-5887. | 1.1 | 84 |
| 3172 | Physical and Functional Interaction between the Human T-Cell Lymphotropic Virus Type 1 Tax₁ Protein and the CCAAT Binding Protein NF-Y. <i>Molecular and Cellular Biology</i> , 1997, 17, 1236-1243. | 1.1 | 32 |
| 3173 | Homologous Segments in Three Subunits of the Guanine Nucleotide Exchange Factor eIF2B Mediate Translational Regulation by Phosphorylation of eIF2. <i>Molecular and Cellular Biology</i> , 1997, 17, 1298-1313. | 1.1 | 133 |
| 3174 | The <i>Saccharomyces cerevisiae</i> MADS-Box Transcription Factor Rlm1 Is a Target for the Mpk1 Mitogen-Activated Protein Kinase Pathway. <i>Molecular and Cellular Biology</i> , 1997, 17, 1848-1859. | 1.1 | 180 |
| 3175 | Plant Farnesyltransferase Can Restore Yeast Ras Signaling and Mating. <i>Molecular and Cellular Biology</i> , 1997, 17, 1986-1994. | 1.1 | 41 |
| 3176 | The Branchpoint Residue Is Recognized during Commitment Complex Formation before Being Bulged out of the U2 snRNA α pre-mRNA Duplex. <i>Molecular and Cellular Biology</i> , 1997, 17, 3469-3476. | 1.1 | 26 |
| 3177 | Cooperative Binding Interactions Required for Function of the Ty1 Sterile Responsive Element. <i>Molecular and Cellular Biology</i> , 1997, 17, 4330-4337. | 1.1 | 58 |
| 3178 | Coordination of the Mating and Cell Integrity Mitogen-Activated Protein Kinase Pathways in <i>Saccharomyces cerevisiae</i> . <i>Molecular and Cellular Biology</i> , 1997, 17, 6517-6525. | 1.1 | 134 |
| 3179 | <i>Saccharomyces cerevisiae</i> BUR6 Encodes a DRAP1/NC2 γ Homolog That Has both Positive and Negative Roles in Transcription In Vivo. <i>Molecular and Cellular Biology</i> , 1997, 17, 2057-2065. | 1.1 | 86 |
| 3180 | <i>CHS5</i> , a Gene Involved in Chitin Synthesis and Mating in <i>Saccharomyces cerevisiae</i> . <i>Molecular and Cellular Biology</i> , 1997, 17, 2485-2496. | 1.1 | 84 |
| 3181 | Inactivation of the Cyclin-Dependent Kinase Cdc28 Abrogates Cell Cycle Arrest Induced by DNA Damage and Disassembly of Mitotic Spindles in <i>Saccharomyces cerevisiae</i> . <i>Molecular and Cellular Biology</i> , 1997, 17, 2723-2734. | 1.1 | 50 |
| 3182 | The <i>STK2</i> Gene, Which Encodes a Putative Ser/Thr Protein Kinase, Is Required for High-Affinity Spermidine Transport in <i>Saccharomyces cerevisiae</i> . <i>Molecular and Cellular Biology</i> , 1997, 17, 2994-3004. | 1.1 | 47 |
| 3183 | Functional Equivalency and Diversity of <i>cis</i> -Acting Elements among Yeast Replication Origins. <i>Molecular and Cellular Biology</i> , 1997, 17, 5473-5484. | 1.1 | 57 |
| 3184 | In Vivo Analysis of the Hsp90 Cochaperone Sti1 (p60). <i>Molecular and Cellular Biology</i> , 1997, 17, 318-325. | 1.1 | 222 |
| 3185 | A Domain Shared by the <i>Polycomb</i> Group Proteins Scm and ph Mediates Heterotypic and Homotypic Interactions. <i>Molecular and Cellular Biology</i> , 1997, 17, 6683-6692. | 1.1 | 139 |
| 3186 | Hir1p and Hir2p Function as Transcriptional Corepressors To Regulate Histone Gene Transcription in the <i>Saccharomyces cerevisiae</i> Cell Cycle. <i>Molecular and Cellular Biology</i> , 1997, 17, 545-552. | 1.1 | 130 |

| # | ARTICLE | IF | CITATIONS |
|------|--|-----|-----------|
| 3187 | Interaction between the Human Nuclear Cap-Binding Protein Complex and hnRNP F. <i>Molecular and Cellular Biology</i> , 1997, 17, 2587-2597. | 1.1 | 47 |
| 3188 | Plk Is a Functional Homolog of <i>Saccharomyces cerevisiae</i> Cdc5, and Elevated Plk Activity Induces Multiple Septation Structures. <i>Molecular and Cellular Biology</i> , 1997, 17, 3408-3417. | 1.1 | 188 |
| 3189 | A critical amino acid residue, Asp446, in UDP-glucuronosyltransferase. <i>Biochemical Journal</i> , 1997, 325, 587-591. | 1.7 | 21 |
| 3190 | Cloning of the <i>Candida albicans</i> homolog of <i>Saccharomyces cerevisiae</i> GSC1/FKS1 and its involvement in beta-1,3-glucan synthesis. <i>Journal of Bacteriology</i> , 1997, 179, 4096-4105. | 1.0 | 144 |
| 3191 | Isolation of the <i>Candida albicans</i> homologs of <i>Saccharomyces cerevisiae</i> KRE6 and SKN1: expression and physiological function. <i>Journal of Bacteriology</i> , 1997, 179, 2363-2372. | 1.0 | 50 |
| 3192 | Vam2/Vps41p and Vam6/Vps39p Are Components of a Protein Complex on the Vacuolar Membranes and Involved in the Vacuolar Assembly in the Yeast <i>Saccharomyces cerevisiae</i> . <i>Journal of Biological Chemistry</i> , 1997, 272, 11344-11349. | 1.6 | 156 |
| 3193 | Molecular cloning of mouse liver flavin containing monooxygenase (FMO1) cDNA and characterization of the expression product: metabolism of the neurotoxin, 1, 2, 3, 4 - tetrahydroisoquinoline (TIQ).. <i>Journal of Toxicological Sciences</i> , 1997, 22, 45-56. | 0.7 | 14 |
| 3194 | Identification of a mannoprotein present in the inner layer of the cell wall of <i>Saccharomyces cerevisiae</i> . <i>Journal of Bacteriology</i> , 1997, 179, 2154-2162. | 1.0 | 74 |
| 3195 | Conserved Properties between Functionally Distinct MutS Homologs in Yeast. <i>Journal of Biological Chemistry</i> , 1997, 272, 30345-30349. | 1.6 | 117 |
| 3196 | Yeast-enhanced green fluorescent protein (yEGFP): a reporter of gene expression in <i>Candida albicans</i> . <i>Microbiology (United Kingdom)</i> , 1997, 143, 303-311. | 0.7 | 559 |
| 3197 | Alternative Yeast Two-Hybrid Systems: The Interaction Trap and Interaction Mating. , 1997, 63, 197-218. | | 42 |
| 3198 | Inducible Expression Cassettes in Yeast: GAL4. , 1997, 62, 131-148. | | 1 |
| 3199 | Recombination between DNA repeats in yeast hpr1Delta cells is linked to transcription elongation. <i>EMBO Journal</i> , 1997, 16, 2826-2835. | 3.5 | 94 |
| 3200 | Mutagenesis of phospholipase D defines a superfamily including a trans-Golgi viral protein required for poxvirus pathogenicity. <i>EMBO Journal</i> , 1997, 16, 4519-4530. | 3.5 | 341 |
| 3201 | Establishment of a Novel Host, High-Red Yeast That Stably Expresses Hamster NADPH-Cytochrome P450 Oxidoreductase: Usefulness for Examination of the Function of Mammalian Cytochrome P450. <i>Archives of Biochemistry and Biophysics</i> , 1997, 343, 215-224. | 1.4 | 11 |
| 3202 | Metabolic Effects of Altering Redundant Targeting Signals for Yeast Mitochondrial Malate Dehydrogenase. <i>Archives of Biochemistry and Biophysics</i> , 1997, 344, 53-60. | 1.4 | 18 |
| 3203 | Determination of FAD-Binding Domain in Flavin-Containing Monooxygenase 1 (FMO1). <i>Archives of Biochemistry and Biophysics</i> , 1997, 345, 271-277. | 1.4 | 24 |
| 3204 | A Dominant Negative Mutation in <i>Saccharomyces cerevisiae</i> Methionine Aminopeptidase-1 Affects Catalysis and Interferes with the Function of Methionine Aminopeptidase-2. <i>Archives of Biochemistry and Biophysics</i> , 1997, 347, 193-200. | 1.4 | 24 |

| # | ARTICLE | IF | CITATIONS |
|------|--|------|-----------|
| 3205 | Characterization of a Sphingomyelinase Activity in <i>Saccharomyces cerevisiae</i> . <i>Archives of Biochemistry and Biophysics</i> , 1997, 340, 101-110. | 1.4 | 31 |
| 3206 | <i>Saccharomyces cerevisiae</i> Expresses Two Genes Encoding Isozymes of 5-Aminoimidazole-4-carboxamide Ribonucleotide Transformylase. <i>Archives of Biochemistry and Biophysics</i> , 1997, 340, 195-200. | 1.4 | 24 |
| 3207 | An AP1 Element Is Involved in Transcriptional Regulation of Δ^9 -Desaturase Gene of <i>Histoplasma capsulatum</i> . <i>Biochemical and Biophysical Research Communications</i> , 1997, 230, 457-461. | 1.0 | 18 |
| 3208 | Association of Elongation Factor ϵ and Ribosomal Protein L3 with the Proline-Rich Region of Yeast Adenylyl Cyclase-Associated Protein CAP. <i>Biochemical and Biophysical Research Communications</i> , 1997, 232, 503-507. | 1.0 | 13 |
| 3209 | The Interaction between Human Cytomegalovirus Immediate-Early Gene 2 (IE2) Protein and Heterogeneous Ribonucleoprotein A1. <i>Biochemical and Biophysical Research Communications</i> , 1997, 232, 590-594. | 1.0 | 11 |
| 3210 | G-Protein Coupled Receptor from Yeast <i>Saccharomyces cerevisiae</i> . <i>Biochemical and Biophysical Research Communications</i> , 1997, 240, 287-292. | 1.0 | 94 |
| 3211 | The <i>amdA</i> Regulatory Gene of <i>Aspergillus nidulans</i> : Characterization of Gain-of-Function Mutations and Identification of Binding Sites for the Gene Product. <i>Fungal Genetics and Biology</i> , 1997, 21, 50-63. | 0.9 | 8 |
| 3212 | Expression of <i>YWP1</i> , a Gene That Encodes a Specific <i>Yarrowia lipolytica</i> Mycelial Cell Wall Protein, in <i>Saccharomyces cerevisiae</i> . <i>Fungal Genetics and Biology</i> , 1997, 22, 77-83. | 0.9 | 6 |
| 3213 | Physical Mapping of Potassium Channel Gene Clusters on Mouse Chromosomes Three and Six. <i>Genomics</i> , 1997, 44, 110-117. | 1.3 | 7 |
| 3214 | Synergistic transcriptional-activation by the papillomavirus E2 protein occurs after DNA binding and correlates with a change in chromatin structure 1 Edited by J. Karn. <i>Journal of Molecular Biology</i> , 1997, 266, 465-478. | 2.0 | 12 |
| 3215 | A dynamic in Vivo view of the HIV-1 Rev-RRE interaction. <i>Journal of Molecular Biology</i> , 1997, 266, 950-962. | 2.0 | 39 |
| 3216 | An in vivo and in vitro structure-function analysis of the <i>Saccharomyces cerevisiae</i> U3A snoRNP: protein-RNA contacts and base-pair interaction with the pre-ribosomal RNA 1 Edited by M. Yaniv. <i>Journal of Molecular Biology</i> , 1997, 273, 552-571. | 2.0 | 105 |
| 3217 | Using <i>GCN4</i> as a Reporter of eIF2 γ Phosphorylation and Translational Regulation in Yeast. <i>Methods</i> , 1997, 11, 403-417. | 1.9 | 33 |
| 3218 | Yeast as a Model System to Study Drugs Effective against Apicomplexan Proteins. <i>Methods</i> , 1997, 13, 190-207. | 1.9 | 37 |
| 3219 | High-Level Intracellular Expression of Hydroxynitrile Lyase from the Tropical Rubber Tree <i>Hevea brasiliensis</i> in Microbial Hosts. <i>Protein Expression and Purification</i> , 1997, 11, 61-71. | 0.6 | 126 |
| 3220 | Two transcription activation functions in the amino terminus of the mouse estrogen receptor that are affected by the carboxy terminus. <i>Steroids</i> , 1997, 62, 508-515. | 0.8 | 13 |
| 3221 | Mis6, a Fission Yeast Inner Centromere Protein, Acts during G1/S and Forms Specialized Chromatin Required for Equal Segregation. <i>Cell</i> , 1997, 90, 131-143. | 13.5 | 227 |
| 3222 | The Yeast Phosphatidylinositol Kinase Homolog TOR2 Activates RHO1 and RHO2 via the Exchange Factor ROM2. <i>Cell</i> , 1997, 88, 531-542. | 13.5 | 293 |

| # | ARTICLE | IF | CITATIONS |
|------|--|-----|-----------|
| 3223 | Modulation of intracellular protein degradation by SSB1-SIS1 chaperon system in yeast <i>S. cerevisiae</i> . <i>FEBS Letters</i> , 1997, 409, 307-311. | 1.3 | 40 |
| 3224 | High expression of the yeast syntaxin-related Vam3 protein suppresses the protein transport defects of <i>apep12</i> null mutant. <i>FEBS Letters</i> , 1997, 411, 48-52. | 1.3 | 39 |
| 3225 | Glucose activation of the yeast plasma membrane H ⁺ -ATPase requires the ubiquitin-proteasome proteolytic pathway. <i>FEBS Letters</i> , 1997, 411, 308-312. | 1.3 | 31 |
| 3226 | Isolation and characterization of mutations affecting expression of the Δ^9 -fatty acid desaturase gene, <i>OLE1</i> , in <i>Saccharomyces cerevisiae</i> . <i>FEBS Letters</i> , 1997, 413, 226-230. | 1.3 | 23 |
| 3227 | Bactericidal action of lysozymes attached with various sizes of hydrophobic peptides to the C-terminal using genetic modification. <i>FEBS Letters</i> , 1997, 415, 114-118. | 1.3 | 49 |
| 3228 | Functional assessment of the yeast <i>Rvs161</i> and <i>Rvs167</i> protein domains. <i>FEBS Letters</i> , 1997, 417, 21-27. | 1.3 | 36 |
| 3229 | Yeast gene <i>YOR137c</i> is involved in the activation of the yeast plasma membrane H ⁺ -ATPase by glucose. <i>FEBS Letters</i> , 1997, 420, 17-19. | 1.3 | 17 |
| 3230 | Overexpression of the sodium ATPase of <i>Saccharomyces cerevisiae</i> : conditions for phosphorylation from ATP and Pi. <i>Biochimica Et Biophysica Acta - Biomembranes</i> , 1997, 1328, 214-225. | 1.4 | 42 |
| 3231 | An amino terminal prosequence is required for efficient synthesis of <i>S. cerevisiae</i> α -factor. <i>Biochimica Et Biophysica Acta - Molecular Cell Research</i> , 1997, 1356, 23-34. | 1.9 | 3 |
| 3232 | Potato (<i>Solanum tuberosum</i> L.) fructokinase expressed in yeast exhibits inhibition by fructose of both in vitro enzyme activity and rate of cell proliferation. <i>Plant Science</i> , 1997, 128, 191-197. | 1.7 | 14 |
| 3233 | Heterologous <i>Kluyveromyces lactis</i> β -galactosidase production and release by <i>Saccharomyces cerevisiae</i> osmotic-remedial thermosensitive autolytic mutants. <i>Biochimica Et Biophysica Acta - General Subjects</i> , 1997, 1335, 235-241. | 1.1 | 27 |
| 3234 | The <i>P-OLE1</i> gene of <i>Pichia angusta</i> encodes a Δ^9 -fatty acid desaturase and complements the <i>ole1</i> mutation of <i>Saccharomyces cerevisiae</i> . <i>Gene</i> , 1997, 184, 299-306. | 1.0 | 26 |
| 3235 | Spermidine biosynthesis in <i>Saccharomyces cerevisiae</i> : Polyamine requirement of a null mutant of the <i>SPE3</i> gene (spermidine synthase). <i>Gene</i> , 1997, 187, 35-43. | 1.0 | 61 |
| 3236 | Isolation and characterization of cDNA encoding mouse RNA polymerase II subunit <i>RPB14</i> . <i>Gene</i> , 1997, 187, 165-170. | 1.0 | 5 |
| 3237 | <i>RIC1</i> , a novel gene required for ribosome synthesis in <i>Saccharomyces cerevisiae</i> . <i>Gene</i> , 1997, 187, 171-178. | 1.0 | 15 |
| 3238 | Identification of a profilin homologue in <i>Trypanosoma brucei</i> by complementation screening. <i>Gene</i> , 1997, 187, 201-209. | 1.0 | 17 |
| 3239 | Identification of <i>Cdc45p</i> , an essential factor required for DNA replication. <i>Gene</i> , 1997, 187, 239-246. | 1.0 | 55 |
| 3240 | Satellite DNA from the brine shrimp <i>Artemia</i> affects the expression of a flanking gene in yeast. <i>Gene</i> , 1997, 189, 13-18. | 1.0 | 5 |

| # | ARTICLE | IF | CITATIONS |
|------|--|-----|-----------|
| 3241 | Cloning and characterization of the <i>S. pombe</i> gene <i>efc25+</i> , a new putative guanine nucleotide exchange factor. <i>Gene</i> , 1997, 193, 203-210. | 1.0 | 22 |
| 3242 | Two vectors for the insertion of mammalian selectable genes into yeast artificial chromosome cloned DNA. <i>Gene</i> , 1997, 199, 25-30. | 1.0 | 1 |
| 3243 | Suppression of the <i>Saccharomyces cerevisiae</i> <i>hac1/ire15</i> mutation by yeast genes and human cDNAs. <i>Gene</i> , 1997, 201, 5-10. | 1.0 | 8 |
| 3244 | Cloning of porcine neuron growth inhibitory factor (metallothionein III) cDNA and expression of the gene in <i>Saccharomyces cerevisiae</i> . <i>Gene</i> , 1997, 203, 189-197. | 1.0 | 7 |
| 3245 | cDNA cloning and functional analysis of p44.5 and p55, two regulatory subunits of the 26S proteasome. <i>Gene</i> , 1997, 203, 241-250. | 1.0 | 26 |
| 3246 | Identification of the Protein That Interacts with the 3' End of Histone mRNA. , 1997, , 163-193. | | 9 |
| 3247 | Position- and orientation-independent activity of the <i>Schizosaccharomyces pombe</i> meiotic recombination hot spot M26. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 1997, 94, 7446-7451. | 3.3 | 37 |
| 3248 | A conserved cysteine residue in yeast uroporphyrinogen decarboxylase is not essential for enzymatic activity. <i>Canadian Journal of Microbiology</i> , 1997, 43, 792-795. | 0.8 | 2 |
| 3249 | DNA double-strand-break sensitivity, DNA replication, and cell cycle arrest phenotypes of Ku-deficient <i>Saccharomyces cerevisiae</i> . <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 1997, 94, 867-872. | 3.3 | 108 |
| 3250 | Cross regulation of four GATA factors that control nitrogen catabolic gene expression in <i>Saccharomyces cerevisiae</i> . <i>Journal of Bacteriology</i> , 1997, 179, 3416-3429. | 1.0 | 114 |
| 3251 | Cloning of the <i>sdsA</i> gene encoding solanesyl diphosphate synthase from <i>Rhodobacter capsulatus</i> and its functional expression in <i>Escherichia coli</i> and <i>Saccharomyces cerevisiae</i> . <i>Journal of Bacteriology</i> , 1997, 179, 5992-5998. | 1.0 | 61 |
| 3252 | Chromatin Remodeling by Transcriptional Activation Domains in a Yeast Episome. <i>Journal of Biological Chemistry</i> , 1997, 272, 11526-11534. | 1.6 | 33 |
| 3253 | Role of Conserved Histidines in Catalytic Activity and Inhibitor Binding of Human Recombinant Phosphodiesterase 4A. <i>Molecular Pharmacology</i> , 1997, 51, 999-1006. | 1.0 | 30 |
| 3254 | Regulation of the proteinase B structural gene <i>PRB1</i> in <i>Saccharomyces cerevisiae</i> . <i>Journal of Bacteriology</i> , 1997, 179, 1469-1474. | 1.0 | 28 |
| 3255 | Detection of heterozygous truncating mutations in the <i>BRCA1</i> and <i>APC</i> genes by using a rapid screening assay in yeast. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 1997, 94, 2449-2453. | 3.3 | 52 |
| 3256 | Activated Alleles of Yeast <i>SLN1</i> Increase <i>Mcm1</i> -dependent Reporter Gene Expression and Diminish Signaling through the <i>Hog1</i> Osmosensing Pathway. <i>Journal of Biological Chemistry</i> , 1997, 272, 13365-13371. | 1.6 | 49 |
| 3257 | Recombinant Sickle Hemoglobin Containing a Lysine Substitution at Asp-85(Î±): Expression in Yeast, Functional Properties, and Participation in Gel Formation. <i>Blood</i> , 1997, 89, 4196-4203. | 0.6 | 9 |
| 3258 | The yeast peptide-methionine sulfoxide reductase functions as an antioxidant in vivo. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 1997, 94, 9585-9589. | 3.3 | 328 |

| # | ARTICLE | IF | CITATIONS |
|------|--|------|-----------|
| 3259 | A novel Mcm1-dependent element in the SWI4, CLN3, CDC6, and CDC47 promoters activates M/G1-specific transcription.. Genes and Development, 1997, 11, 1277-1288. | 2.7 | 191 |
| 3260 | Investigating Ester Metabolism: Characterization of the <i>ATF1</i> Gene in <i>Saccharomyces Cerevisiae</i> . Journal of the American Society of Brewing Chemists, 1997, 55, 141-146. | 0.8 | 13 |
| 3261 | Safety assessment of genetically engineered yeast: elimination of mutagenicity of the yeast <i>Saccharomyces cerevisiae</i> by decreasing the activity of methylglyoxal synthase. International Journal of Food Science and Technology, 1997, 32, 521-526. | 1.3 | 1 |
| 3262 | Essential role for diacylglycerol in protein transport from the yeast Golgi complex. Nature, 1997, 387, 101-105. | 13.7 | 264 |
| 3263 | Yeast DNA ligase IV mediates non-homologous DNA end joining. Nature, 1997, 388, 495-498. | 13.7 | 381 |
| 3264 | Estimation of the Rate of Cloned Gene Integration via Retrotransposition. Biotechnology Progress, 1997, 13, 123-131. | 1.3 | 2 |
| 3265 | Identification and Characterisation of a Human Calmodulin-Stimulated Phosphodiesterase PDE1B1. Cellular Signalling, 1997, 9, 519-529. | 1.7 | 64 |
| 3266 | Molecular Cloning and Expression of a Human Phosphodiesterase 4C. Cellular Signalling, 1997, 9, 575-585. | 1.7 | 28 |
| 3267 | Enhancement of rice α -amylase production in recombinant <i>Yarrowia lipolytica</i> . Journal of Bioscience and Bioengineering, 1997, 84, 421-427. | 0.9 | 11 |
| 3268 | A method for fusing chromosomes in <i>Saccharomyces cerevisiae</i> . Journal of Bioscience and Bioengineering, 1997, 83, 125-131. | 0.9 | 7 |
| 3269 | An IQGAP-related protein controls actin-ring formation and cytokinesis in yeast. Current Biology, 1997, 7, 921-929. | 1.8 | 176 |
| 3270 | Intracellular Location, Complex Formation, and Function of the Transporter Associated with Antigen Processing in Yeast. FEBS Journal, 1997, 245, 266-272. | 0.2 | 35 |
| 3271 | A Heterodimer of the Zn ₂ Cys ₆ Transcription Factors Pip2p and Oaf1p Controls Induction of Genes Encoding Peroxisomal Proteins in <i>Saccharomyces Cerevisiae</i> . FEBS Journal, 1997, 247, 776-783. | 0.2 | 109 |
| 3272 | The Absence of the Mitochondrial ATP Synthase Subunit Promotes a Slow Growth Phenotype of Rho- Yeast Cells by a Lack of Assembly of the Catalytic Sector F ₁ . FEBS Journal, 1997, 245, 813-818. | 0.2 | 73 |
| 3273 | Mitochondria1 Asparaginyl-tRNA Synthetase Encoded by the Yeast Nuclear Gene YCR24c. FEBS Journal, 1997, 243, 268-273. | 0.2 | 8 |
| 3274 | Analysis in yeast of antimalaria drugs that target the dihydrofolate reductase of <i>Plasmodium falciparum</i> . Molecular and Biochemical Parasitology, 1997, 85, 25-40. | 0.5 | 84 |
| 3275 | Novel Golgi to vacuole delivery pathway in yeast: identification of a sorting determinant and required transport component. EMBO Journal, 1997, 16, 2769-2782. | 3.5 | 187 |
| 3276 | An evolutionarily conserved U5 snRNP-specific protein is a GTP-binding factor closely related to the ribosomal translocase EF-2. EMBO Journal, 1997, 16, 4092-4106. | 3.5 | 194 |

| # | ARTICLE | IF | CITATIONS |
|------|---|-----|-----------|
| 3277 | Suppressors of YCK-encoded yeast casein kinase 1 deficiency define the four subunits of a novel clathrin AP-like complex. <i>EMBO Journal</i> , 1997, 16, 4194-4204. | 3.5 | 137 |
| 3278 | DBF2, a cell cycle-regulated protein kinase, is physically and functionally associated with the CCR4 transcriptional regulatory complex. <i>EMBO Journal</i> , 1997, 16, 5289-5298. | 3.5 | 101 |
| 3279 | Essential functions and actin-binding surfaces of yeast cofilin revealed by systematic mutagenesis. <i>EMBO Journal</i> , 1997, 16, 5520-5530. | 3.5 | 235 |
| 3280 | Yeast spore germination: a requirement for Ras protein activity during re-entry into the cell cycle. <i>EMBO Journal</i> , 1997, 16, 6171-6181. | 3.5 | 114 |
| 3281 | The role of branchpoint-3' splice site spacing and interaction between intron terminal nucleotides in 3' splice site selection in <i>Saccharomyces cerevisiae</i> . <i>EMBO Journal</i> , 1997, 16, 779-792. | 3.5 | 74 |
| 3282 | A novel inhibitor of cap-dependent translation initiation in yeast: p20 competes with eIF4G for binding to eIF4E. <i>EMBO Journal</i> , 1997, 16, 1114-1121. | 3.5 | 154 |
| 3283 | Endosomal transport function in yeast requires a novel AAA-type ATPase, Vps4p. <i>EMBO Journal</i> , 1997, 16, 1820-1831. | 3.5 | 377 |
| 3284 | A novel structural model for regulation of clathrin function. <i>EMBO Journal</i> , 1997, 16, 2227-2239. | 3.5 | 37 |
| 3285 | Site-directed mutagenesis in basic amino acid residues of <i>Saccharomyces cerevisiae</i> phosphoenolpyruvate carboxykinase. <i>The Protein Journal</i> , 1997, 16, 233-236. | 1.1 | 4 |
| 3286 | Minimum Length of Sequence Homology Required for <i>In Vivo</i> Cloning by Homologous Recombination in Yeast. <i>Plasmid</i> , 1997, 38, 91-96. | 0.4 | 110 |
| 3287 | Evaluation of Chemicals with Endocrine Modulating Activity in a Yeast-Based Steroid Hormone Receptor Gene Transcription Assay. <i>Toxicology and Applied Pharmacology</i> , 1997, 143, 205-212. | 1.3 | 635 |
| 3288 | Mutational Analysis of Transcriptional Activation by the Bovine Papillomavirus Type 1 E6. <i>Virology</i> , 1997, 236, 30-36. | 1.1 | 14 |
| 3289 | An Imperfect Correlation between DNA Replication Activity of Epstein-Barr Virus Nuclear Antigen 1 (EBNA1) and Binding to the Nuclear Import Receptor, Rch1/importin β . <i>Virology</i> , 1997, 239, 340-351. | 1.1 | 45 |
| 3290 | The presence of a Sar1 gene family in <i>Brassica campestris</i> that suppresses a yeast vesicular transport mutation Sec12-1. <i>Plant Molecular Biology</i> , 1997, 33, 1025-1035. | 2.0 | 17 |
| 3291 | Title is missing!. <i>Molecular and Cellular Biochemistry</i> , 1997, 172, 67-79. | 1.4 | 117 |
| 3292 | Antigen-specific inhibition of CD4+ T-cell responses to beta-lactoglobulin by its single amino acid-substituted mutant form through T-cell receptor antagonism. <i>Cytotechnology</i> , 1997, 25, 115-126. | 0.7 | 9 |
| 3293 | A small variance in the antigenicity but not function of recombinant β -lactoglobulin purified from the culture supernatant of transformed yeast cells. <i>Cytotechnology</i> , 1997, 23, 133-141. | 0.7 | 4 |
| 3294 | Title is missing!. <i>Biotechnology Letters</i> , 1997, 19, 503-506. | 1.1 | 5 |

| # | ARTICLE | IF | CITATIONS |
|------|--|-----|-----------|
| 3295 | Title is missing!. Biotechnology Letters, 1997, 19, 437-441. | 1.1 | 4 |
| 3296 | Rapid and sensitive detection of b-galactosidase-producing yeasts by using microtiter plate assay. Biotechnology Letters, 1997, 11, 399-402. | 0.5 | 7 |
| 3297 | Title is missing!. Biotechnology Letters, 1997, 19, 1169-1173. | 1.1 | 1 |
| 3298 | Assembly of deletion mutants of the Rieske iron-sulfur protein into the cytochrome bc1 complex of yeast mitochondria. Journal of Bioenergetics and Biomembranes, 1997, 29, 45-54. | 1.0 | 6 |
| 3299 | Asparaginase II of <i>Saccharomyces cerevisiae</i> . Applied Biochemistry and Biotechnology, 1997, 63-65, 203-212. | 1.4 | 21 |
| 3300 | Integration of the multiple controls regulating the expression of the arginase gene CAR1 of <i>Saccharomyces cerevisiae</i> in response to different nitrogen signals: role of Gln3p, ArgRp-Mcm1p, and Ume6p. Molecular Genetics and Genomics, 1997, 253, 568-580. | 2.4 | 30 |
| 3301 | A novel yeast gene, RHK1, is involved in the synthesis of the cell wall receptor for the HM-1 killer toxin that inhibits 1,3-glucan synthesis. Molecular Genetics and Genomics, 1997, 254, 139-147. | 2.4 | 22 |
| 3302 | PRS1 is a key member of the gene family encoding phosphoribosylpyrophosphate synthetase in <i>Saccharomyces cerevisiae</i> . Molecular Genetics and Genomics, 1997, 254, 148-156. | 2.4 | 33 |
| 3303 | A novel fission yeast gene, kms1+, is required for the formation of meiotic prophase-specific nuclear architecture. Molecular Genetics and Genomics, 1997, 254, 238-249. | 2.4 | 127 |
| 3304 | Screening and identification of yeast sequences that cause growth inhibition when overexpressed. Molecular Genetics and Genomics, 1997, 254, 267-274. | 2.4 | 75 |
| 3305 | A simple signal element mediates transcription termination and mRNA 3' end formation in the DEG1 gene of <i>Saccharomyces cerevisiae</i> . Molecular Genetics and Genomics, 1997, 254, 681-688. | 2.4 | 11 |
| 3306 | Yeast silencers create domains of nuclease-resistant chromatin in an SIR4 -dependent manner. Chromosoma, 1997, 106, 136-148. | 1.0 | 7 |
| 3307 | Mutations in SPK1/RAD53 that specifically abolish checkpoint but not growth-related functions. Current Genetics, 1997, 31, 97-105. | 0.8 | 53 |
| 3308 | Heme regulates SOD2 transcription by activation and repression in <i>Saccharomyces cerevisiae</i> . Current Genetics, 1997, 31, 281-291. | 0.8 | 35 |
| 3309 | Additional copies of the mitochondrial Ef-Tu and aspartyl-tRNA synthetase genes can compensate for a mutation affecting the maturation of the mitochondrial tRNA Asp. Current Genetics, 1997, 31, 494-496. | 0.8 | 30 |
| 3310 | A novel nuclear gene, CBT1, essential for mitochondrial cytochrome b formation: terminal processing of mRNA and intron dependence. Current Genetics, 1997, 32, 163-174. | 0.8 | 13 |
| 3311 | A <i>Saccharomyces cerevisiae</i> mitochondrial DNA fragment activates Reg1p-dependent glucose-repressible transcription in the nucleus. Current Genetics, 1997, 32, 389-398. | 0.8 | 3 |
| 3312 | Genetic determination of polygalacturonase production in wild-type and laboratory strains of <i>Saccharomyces cerevisiae</i> . Archives of Microbiology, 1997, 167, 284-288. | 1.0 | 24 |

| # | ARTICLE | IF | CITATIONS |
|------|--|-----|-----------|
| 3313 | A counter-selectable marker for genetic transformation of the yeast <i>Schwanniomyces alluvius</i> . <i>Applied Microbiology and Biotechnology</i> , 1997, 48, 204-207. | 1.7 | 8 |
| 3314 | One-step transformation of the dimorphic yeast <i>Yarrowia lipolytica</i> . <i>Applied Microbiology and Biotechnology</i> , 1997, 48, 232-235. | 1.7 | 148 |
| 3315 | Improved efficiency and stability of multiple cloned gene insertions at the $\hat{\Gamma}$ sequences of <i>Saccharomyces cerevisiae</i> . <i>Applied Microbiology and Biotechnology</i> , 1997, 48, 339-345. | 1.7 | 64 |
| 3316 | Transformation system for a wastewater treatment yeast, <i>Hansenula fabianii</i> J640: isolation of the orotidine-5 $\hat{\epsilon}$ -phosphate decarboxylase gene (<i>URA3</i>) and uracil auxotrophic mutants. <i>Applied Microbiology and Biotechnology</i> , 1997, 48, 621-625. | 1.7 | 11 |
| 3317 | Specific interactions between the K domains of AG and AGLs, members of the MADS domain family of DNA binding proteins. <i>Plant Journal</i> , 1997, 12, 999-1010. | 2.8 | 196 |
| 3318 | Involvement of the <i>MRE2</i> gene of yeast in formation of meiosis-specific double-strand breaks and crossover recombination through RNA splicing. <i>Genes To Cells</i> , 1997, 2, 65-79. | 0.5 | 50 |
| 3319 | Region $\hat{\epsilon}$ -specific meiotic recombination in <i>Schizosaccharomyces pombe</i> : the <i>rec11</i> gene. <i>Molecular Microbiology</i> , 1997, 23, 869-878. | 1.2 | 39 |
| 3320 | Bidirectional regulation of two DNA $\hat{\epsilon}$ damage $\hat{\epsilon}$ inducible genes, <i>MAG1</i> and <i>DDI1</i> , from <i>Saccharomyces cerevisiae</i> . <i>Molecular Microbiology</i> , 1997, 23, 777-789. | 1.2 | 45 |
| 3321 | <i>Gzf3p</i> , a fourth GATA factor involved in nitrogen-regulated transcription in <i>Saccharomyces cerevisiae</i> . <i>Molecular Microbiology</i> , 1997, 23, 1157-1168. | 1.2 | 102 |
| 3322 | The C-terminal Domain of <i>Snf3p</i> is Sufficient to Complement the Growth Defect of <i>snf3</i> Null Mutations in <i>Saccharomyces cerevisiae</i> : <i>SNF3</i> Functions in Glucose Recognition. , 1997, 13, 9-20. | | 58 |
| 3323 | Analysis of the Structure of a Natural Alternating d(TA) n Sequence in Yeast Chromatin. , 1997, 13, 313-326. | | 23 |
| 3324 | Identification of Gene encoding a Putative RNA-Helicase, Homologous to <i>SKI2</i> , in Chromosome VII of <i>Saccharomyces cerevisiae</i> . , 1997, 13, 391-397. | | 19 |
| 3325 | Isolation of a <i>Schizosaccharomyces pombe</i> Gene Which in High Copy Confers Resistance to the Nucleoside Analogue 5-Azacytidine. <i>Yeast</i> , 1997, 13, 463-474. | 0.8 | 8 |
| 3326 | Secretion of Mouse $\hat{\Gamma}$ -Amylase from <i>Kluyveromyces lactis</i> . , 1997, 13, 699-706. | | 28 |
| 3327 | Characterization of the Checkpoint Gene <i>RAD53/MEC2</i> in <i>Saccharomyces cerevisiae</i> . , 1997, 13, 735-745. | | 20 |
| 3328 | Isolation of Three Contiguous Genes, <i>ACR1</i> , <i>ACR2</i> and <i>ACR3</i> , Involved in Resistance to Arsenic Compounds in the Yeast <i>Saccharomyces cerevisiae</i> . , 1997, 13, 819-828. | | 211 |
| 3329 | <i>GCR1</i> -Dependent Transcriptional Activation of Yeast Retrotransposon <i>Ty2-917</i> . , 1997, 13, 917-930. | | 29 |
| 3330 | Isolation and Characterization of the <i>KIHEM1</i> Gene in <i>Kluyveromyces lactis</i> . , 1997, 13, 961-971. | | 16 |

| # | ARTICLE | IF | CITATIONS |
|------|---|-----|-----------|
| 3331 | Phylogenetic Relationships of Fungal Cytochromesc. , 1997, 13, 985-990. | | 12 |
| 3332 | Accumulation of Misfolded Protein Aggregates Leads to the Formation of Russell Body-like Dilated Endoplasmic Reticulum in Yeast. , 1997, 13, 1009-1020. | | 45 |
| 3333 | Molecular Cloning of Chromosome I DNA from <i>Saccharomyces cerevisiae</i> : Characterization of the 54â€‰%kb Right Terminal CDC15-FLO1-PHO11 Region. <i>Yeast</i> , 1997, 13, 1251-1263. | 0.8 | 17 |
| 3334 | Lys80p of <i>Saccharomyces cerevisiae</i> , previously proposed as a specific repressor of <i>LYS</i> genes, is a pleiotropic regulatory factor identical to <i>Mks1p</i> . , 1997, 13, 1337-1346. | | 38 |
| 3335 | Functional differences among the six <i>Saccharomyces cerevisiae</i> tRNA ^{Trp} genes. <i>Yeast</i> , 1997, 13, 1357-1362. | 0.8 | 12 |
| 3336 | Hydroxynitrile lyase from <i>Hevea brasiliensis</i> : Molecular characterization and mechanism of enzyme catalysis. , 1997, 27, 438-449. | | 33 |
| 3337 | Genetic and biochemical analysis of the transmembrane domain of <i>Arabidopsis</i> 3-hydroxy-3-methylglutaryl coenzyme A reductase. <i>Journal of Cellular Biochemistry</i> , 1997, 65, 443-459. | 1.2 | 13 |
| 3338 | Title is missing!. <i>Biotechnology Letters</i> , 1998, 20, 9-13. | 1.1 | 4 |
| 3339 | Conserved Ser residues in the basic region of the bZIP-type transcription factor HBP-1a(17): importance in DNA binding and possible targets for phosphorylation. <i>Plant Molecular Biology</i> , 1998, 36, 125-136. | 2.0 | 23 |
| 3340 | Engineering in vivo instability of firefly luciferase and <i>Escherichia coli</i> beta-glucuronidase in higher plants using recognition elements from the ubiquitin pathway. <i>Plant Molecular Biology</i> , 1998, 37, 337-347. | 2.0 | 45 |
| 3341 | pmp1+, a suppressor of calcineurin deficiency, encodes a novel MAP kinase phosphatase in fission yeast. <i>EMBO Journal</i> , 1998, 17, 140-148. | 3.5 | 109 |
| 3342 | D-type cyclins repress transcriptional activation by the v-Myb but not the c-Myb DNA-binding domain. <i>EMBO Journal</i> , 1998, 17, 255-268. | 3.5 | 90 |
| 3343 | The Vps4p AAA ATPase regulates membrane association of a Vps protein complex required for normal endosome function. <i>EMBO Journal</i> , 1998, 17, 2982-2993. | 3.5 | 684 |
| 3344 | Novel developmentally regulated phosphoinositide binding proteins from soybean whose expression bypasses the requirement for an essential phosphatidylinositol transfer protein in yeast. <i>EMBO Journal</i> , 1998, 17, 4004-4017. | 3.5 | 69 |
| 3345 | ATP binding and hydrolysis are essential to the function of the Hsp90 molecular chaperone in vivo. <i>EMBO Journal</i> , 1998, 17, 4829-4836. | 3.5 | 662 |
| 3346 | Phosphoinositide signaling and turnover: PtdIns(3)P, a regulator of membrane traffic, is transported to the vacuole and degraded by a process that requires luminal vacuolar hydrolase activities. <i>EMBO Journal</i> , 1998, 17, 4930-4942. | 3.5 | 139 |
| 3347 | The Cdc42p effector Gic2p is targeted for ubiquitin-dependent degradation by the SCF ^{Grr1} complex. <i>EMBO Journal</i> , 1998, 17, 5360-5373. | 3.5 | 95 |
| 3348 | Apc10 and Ste9/Srw1, two regulators of the APCâ€™“cyclosome, as well as the CDK inhibitor Rum1 are required for G1 cell-cycle arrest in fission yeast. <i>EMBO Journal</i> , 1998, 17, 5388-5399. | 3.5 | 92 |

| # | ARTICLE | IF | CITATIONS |
|------|--|-----|-----------|
| 3349 | Accumulation of mitochondrially synthesized <i>Saccharomyces cerevisiae</i> Cox2p and Cox3p depends on targeting information in untranslated portions of their mRNAs. <i>EMBO Journal</i> , 1998, 17, 5796-5804. | 3.5 | 107 |
| 3350 | Cooperative DNA-binding by Bicoid provides a mechanism for threshold-dependent gene activation in the <i>Drosophila</i> embryo. <i>EMBO Journal</i> , 1998, 17, 5998-6009. | 3.5 | 161 |
| 3351 | The TOR nutrient signalling pathway phosphorylates NPR1 and inhibits turnover of the tryptophan permease. <i>EMBO Journal</i> , 1998, 17, 6924-6931. | 3.5 | 289 |
| 3352 | The yeast histidine protein kinase, Sln1p, mediates phosphotransfer to two response regulators, Ssk1p and Skn7p. <i>EMBO Journal</i> , 1998, 17, 6952-6962. | 3.5 | 162 |
| 3353 | Crm1p mediates regulated nuclear export of a yeast AP-1-like transcription factor. <i>EMBO Journal</i> , 1998, 17, 7416-7429. | 3.5 | 224 |
| 3354 | Distinct functions of calmodulin are required for the uptake step of receptor-mediated endocytosis in yeast: the type I myosin Myo5p is one of the calmodulin targets. <i>EMBO Journal</i> , 1998, 17, 635-647. | 3.5 | 54 |
| 3355 | epsilon-COP is a structural component of coatamer that functions to stabilize alpha-COP. <i>EMBO Journal</i> , 1998, 17, 985-995. | 3.5 | 77 |
| 3356 | GPR1 encodes a putative G protein-coupled receptor that associates with the Gpa2p G β subunit and functions in a Ras-independent pathway. <i>EMBO Journal</i> , 1998, 17, 1996-2007. | 3.5 | 290 |
| 3357 | UTF1, a novel transcriptional coactivator expressed in pluripotent embryonic stem cells and extra-embryonic cells. <i>EMBO Journal</i> , 1998, 17, 2019-2032. | 3.5 | 167 |
| 3358 | A dileucine-like sorting signal directs transport into an AP-3-dependent, clathrin-independent pathway to the yeast vacuole. <i>EMBO Journal</i> , 1998, 17, 2482-2493. | 3.5 | 106 |
| 3359 | Surface-related Changes of Fusing <i>Phytophthora</i> Zoospores. <i>Journal of Phytopathology</i> , 1998, 146, 179-184. | 0.5 | 1 |
| 3360 | Identification of the <i>mgc1</i> mutation which affects mating-pheromone-induced morphogenesis in the yeast <i>Saccharomyces cerevisiae</i> . <i>FEMS Microbiology Letters</i> , 1998, 161, 359-364. | 0.7 | 2 |
| 3361 | Green fluorescent protein-cell wall fusion proteins are covalently incorporated into the cell wall of <i>Saccharomyces cerevisiae</i> . <i>FEMS Microbiology Letters</i> , 1998, 162, 249-255. | 0.7 | 49 |
| 3362 | Cloning, molecular characterization, and expression of an endo-polygalacturonase-encoding gene from <i>Saccharomyces cerevisiae</i> IM1-8b. <i>FEMS Microbiology Letters</i> , 1998, 164, 249-255. | 0.7 | 51 |
| 3363 | Soluble fumarate reductase isoenzymes from <i>Saccharomyces cerevisiae</i> are required for anaerobic growth. <i>FEMS Microbiology Letters</i> , 1998, 165, 111-116. | 0.7 | 40 |
| 3364 | <i>Saccharomyces cerevisiae</i> ACR2 gene encodes an arsenate reductase. <i>FEMS Microbiology Letters</i> , 1998, 168, 127-136. | 0.7 | 118 |
| 3365 | Molecular Cloning and Functional Analysis of <i>Cynomolgus</i> Monkey CYP1A2 11The nucleotide sequence reported in this paper appears in the GSDB, DDBJ, EMBL, and GenBank database with accession number AF086474, definition "Macaca fascicularis adult liver mRNA for cytochrome P-450." <i>Biochemical Pharmacology</i> , 1998, 56, 131-139. | 2.0 | 31 |
| 3366 | Stage-specific expression of the 14-3-3 gene in <i>Echinococcus multilocularis</i> 1Note: sequence information presented in this study is available in the EMBL, GenBank, and DDJB databases under the accession number U63643.1. <i>Molecular and Biochemical Parasitology</i> , 1998, 91, 281-293. | 0.5 | 57 |

| # | ARTICLE | IF | CITATIONS |
|------|---|-----|-----------|
| 3367 | Cloning of a cdc2-related protein kinase from <i>Trypanosoma cruzi</i> that interacts with mammalian cyclins1 Note: The nucleotide sequences reported in this paper has been submitted to the GenBank™/EMBL data bank with accession numbers tzcrk3, U69958; tzcrk3a, U69960; tzcrk3b U69959; tzcrk1, U74762; tzcrk1a, U74763; tzcrk1b, U74764; tzcrk1c, U74765; tzcrk1d, U74766.1. <i>Molecular and Biochemical Parasitology</i> , 1998, 91, 337-351. | 0.5 | 30 |
| 3368 | Structure/function analysis of the domains required for the multimerisation of phenylalanine hydroxylase. <i>BBA - Proteins and Proteomics</i> , 1998, 1382, 295-304. | 2.1 | 19 |
| 3369 | D-Arabinose dehydrogenase and its gene from <i>Saccharomyces cerevisiae</i> . <i>BBA - Proteins and Proteomics</i> , 1998, 1429, 29-39. | 2.1 | 60 |
| 3370 | Transformation of <i>Saccharomyces cerevisiae</i> by the lithium acetate/single-stranded carrier DNA/polyethylene glycol protocol. <i>Technical Tips Online</i> , 1998, 3, 133-137. | 0.2 | 105 |
| 3371 | Brewing properties of sake yeast whose EST2 gene encoding isoamyl acetate-hydrolyzing esterase was disrupted. <i>Journal of Bioscience and Bioengineering</i> , 1998, 85, 101-106. | 0.9 | 43 |
| 3372 | Isolation of oversecreting mutant strains of the yeast <i>Yarrowia lipolytica</i> . <i>Journal of Bioscience and Bioengineering</i> , 1998, 85, 180-184. | 0.9 | 4 |
| 3373 | Expression and subcellular localization of <i>Candida tropicalis</i> catalase in catalase gene disruptants of <i>Saccharomyces cerevisiae</i> . <i>Journal of Bioscience and Bioengineering</i> , 1998, 85, 571-578. | 0.9 | 4 |
| 3374 | Evaluation of the function of arming yeast displaying glucoamylase on its cell surface by direct fermentation of corn to ethanol. <i>Journal of Bioscience and Bioengineering</i> , 1998, 86, 569-572. | 0.9 | 23 |
| 3375 | Fungal protoplasts: relics or modern objects of molecular research?. <i>Microbiological Research</i> , 1998, 153, 97-104. | 2.5 | 8 |
| 3376 | The mammalian small heat-shock protein Hsp20 forms dimers and is a poor chaperone. <i>FEBS Journal</i> , 1998, 258, 1014-1021. | 0.2 | 86 |
| 3377 | Trans-kingdom conjugation offers a powerful gene targeting. <i>Genetic Analysis, Techniques and Applications</i> , 1998, 14, 65-73. | 1.5 | 11 |
| 3378 | Inulase-secreting strain of <i>Saccharomyces cerevisiae</i> produces fructose. , 1998, 60, 492-497. | | 7 |
| 3379 | Cloning of the Multicopy Suppressor Gene SUR7: Evidence for a Functional Relationship between the Yeast Actin-binding Protein Rvs167 and a Putative Membranous Protein. , 1998, 13, 747-761. | | 29 |
| 3380 | Mutations of the CDC28 gene and the radiation sensitivity of <i>Saccharomyces cerevisiae</i> . , 1998, 14, 133-146. | | 21 |
| 3381 | Characterization of a branched-chain amino-acid aminotransferase from <i>Schizosaccharomyces pombe</i> . , 1998, 14, 189-194. | | 18 |
| 3382 | During the initiation of fermentation overexpression of hexokinase PII in yeast transiently causes a similar deregulation of glycolysis as deletion of Tps1. , 1998, 14, 255-269. | | 38 |
| 3383 | Glycosylation of human Î±1-antitrypsin in <i>Saccharomyces cerevisiae</i> and methylotrophic yeasts. , 1998, 14, 371-381. | | 54 |
| 3384 | Identification of YHR019 in <i>Saccharomyces cerevisiae</i> chromosome VIII as the gene for the cytosolic asparaginyl-tRNA synthetase. , 1998, 14, 527-533. | | 5 |

| # | ARTICLE | IF | CITATIONS |
|------|--|-----|-----------|
| 3385 | A novel esterase from <i>Saccharomyces carlsbergensis</i> , a possible function for the yeast TIP1 gene. , 1998, 14, 793-803. | | 18 |
| 3386 | Highly efficient assimilation of lactose by a metabolically engineered strain of <i>Saccharomyces cerevisiae</i> . , 1998, 14, 827-837. | | 44 |
| 3387 | Chromosomal DNA patterns and gene stability of <i>Pichia pastoris</i> . , 1998, 14, 895-903. | | 24 |
| 3388 | Isolation and nucleotide sequence of the gene encoding phosphoenolpyruvate carboxykinase from <i>Kluyveromyces lactis</i> . , 1998, 14, 963-967. | | 7 |
| 3389 | Functional analysis of three adjacent open reading frames from the right arm of yeast chromosome XVI. <i>Yeast</i> , 1998, 14, 1027-1039. | 0.8 | 5 |
| 3390 | New constructs and strategies for efficient PCR-based gene manipulations in yeast. <i>Yeast</i> , 1998, 14, 1139-1146. | 0.8 | 86 |
| 3391 | Molecular cloning and sequence analysis of the <i>Schizosaccharomyces pombe</i> ade10+ gene. <i>Yeast</i> , 1998, 14, 1307-1310. | 0.8 | 1 |
| 3392 | Detection of APC mutations by a yeast-based protein truncation test (YPTT). , 1998, 21, 290-297. | | 7 |
| 3393 | Mms4, a putative transcriptional (co)activator, protects <i>Saccharomyces cerevisiae</i> cells from endogenous and environmental DNA damage. <i>Molecular Genetics and Genomics</i> , 1998, 257, 614-623. | 2.4 | 30 |
| 3394 | Datin, a yeast poly(dA:dT)-binding protein, behaves as an activator of the wild-type ILV1 promoter and interacts synergistically with Reb1p. <i>Molecular Genetics and Genomics</i> , 1998, 258, 95-103. | 2.4 | 12 |
| 3395 | The Cdc14 phosphatase is functionally associated with the Dbf2 protein kinase in <i>Saccharomyces cerevisiae</i> . <i>Molecular Genetics and Genomics</i> , 1998, 258, 104-116. | 2.4 | 23 |
| 3396 | Phosphoinositide-specific phospholipase C forms a complex with 14-3-3 proteins and is involved in expression of UV resistance in fission yeast. <i>Molecular Genetics and Genomics</i> , 1998, 258, 139-147. | 2.4 | 21 |
| 3397 | Isolation of an extragenic suppressor of the rna1-1 mutation in <i>Saccharomyces cerevisiae</i> . <i>Molecular Genetics and Genomics</i> , 1998, 259, 404-413. | 2.4 | 3 |
| 3398 | Involvement of histidine permease (Hip1p) in manganese transport in <i>Saccharomyces cerevisiae</i> . <i>Molecular Genetics and Genomics</i> , 1998, 259, 541-548. | 2.4 | 16 |
| 3399 | Efficient initiation of S-phase in yeast requires Cdc40p, a protein involved in pre-mRNA splicing. <i>Molecular Genetics and Genomics</i> , 1998, 260, 232-241. | 2.4 | 22 |
| 3400 | The MCM16 gene of the yeast <i>Saccharomyces cerevisiae</i> is required for chromosome segregation. <i>Molecular Genetics and Genomics</i> , 1998, 260, 242-250. | 2.4 | 19 |
| 3401 | Improvement of heterologous protein productivity through a selected bioprocess strategy and medium design. <i>Applied Biochemistry and Biotechnology</i> , 1998, 74, 173-189. | 1.4 | 1 |
| 3402 | Simultaneous saccharification of inulin and ethanol fermentation by recombinant <i>Saccharomyces cerevisiae</i> secreting inulinase. <i>Biotechnology and Bioprocess Engineering</i> , 1998, 3, 55-60. | 1.4 | 0 |

| # | ARTICLE | IF | CITATIONS |
|------|---|-----|-----------|
| 3403 | Characterization of expression of a plant oleate desaturase in yeast. <i>JAACS, Journal of the American Oil Chemists' Society</i> , 1998, 75, 77-79. | 0.8 | 10 |
| 3404 | Process development for high-level secretory production of carboxypeptidase Y by <i>Saccharomyces cerevisiae</i> . <i>Applied Microbiology and Biotechnology</i> , 1998, 50, 34-41. | 1.7 | 18 |
| 3405 | A Short Region from the LEU2 Gene of <i>Saccharomyces cerevisiae</i> Functions as an ARS in the Yeast <i>Saccharomyces exiguus</i> Yp74L-3. <i>Current Microbiology</i> , 1998, 37, 426-430. | 1.0 | 3 |
| 3406 | Over-expression of the yeast BFR2 gene partially suppresses the growth defects induced by Brefeldin A and by four ER-to-Golgi mutations. <i>Current Genetics</i> , 1998, 33, 21-28. | 0.8 | 6 |
| 3407 | Synergism between yeast nucleotide and base excision repair pathways in the protection against DNA methylation damage. <i>Current Genetics</i> , 1998, 33, 92-99. | 0.8 | 64 |
| 3408 | The Tup1-Ssn6 general repressor is involved in repression of IME1 encoding a transcriptional activator of meiosis in <i>Saccharomyces cerevisiae</i> . <i>Current Genetics</i> , 1998, 33, 239-247. | 0.8 | 30 |
| 3409 | Exogenous inositol and genes responsible for inositol transport are required for mating and sporulation in <i>Schizosaccharomyces pombe</i> . <i>Current Genetics</i> , 1998, 33, 255-261. | 0.8 | 32 |
| 3410 | Functional analysis of upstream activating elements in the promoter of the FBP1 gene from <i>Saccharomyces cerevisiae</i> . <i>Current Genetics</i> , 1998, 33, 406-411. | 0.8 | 15 |
| 3411 | Role of yeast Rth1 nuclease and its homologs in mutation avoidance, DNA repair, and DNA replication. <i>Current Genetics</i> , 1998, 34, 21-29. | 0.8 | 55 |
| 3412 | The essential <i>Schizosaccharomyces pombe</i> cdc23 DNA replication gene shares structural and functional homology with the <i>Saccharomyces cerevisiae</i> DNA43 (MCM10) gene. <i>Current Genetics</i> , 1998, 34, 164-171. | 0.8 | 38 |
| 3413 | Phosphatidylglycerolphosphate synthase encoded by the PEL1/PCS1 gene in <i>Saccharomyces cerevisiae</i> is localized in mitochondria and its expression is regulated by phospholipid precursors. <i>Current Genetics</i> , 1998, 34, 297-302. | 0.8 | 52 |
| 3414 | The two 3-deoxy- d - arabino -heptulosonate-7-phosphate synthase isoenzymes from <i>Saccharomyces cerevisiae</i> show different kinetic modes of inhibition. <i>Archives of Microbiology</i> , 1998, 169, 517-524. | 1.0 | 31 |
| 3415 | Expression of the yeast BFR2 gene is regulated at the transcriptional level and through degradation of its product. <i>Molecular Genetics and Genomics</i> , 1998, 258, 215-221. | 2.4 | 8 |
| 3416 | Bioconversion using immobilized recombinant flocculent yeast cells carrying a fused enzyme gene in an 'intelligent' bioreactor. <i>Biochemical Engineering Journal</i> , 1998, 2, 229-235. | 1.8 | 29 |
| 3417 | Mutagenesis of histidine 26 demonstrates the importance of loop-loop and loop-protein interactions for the function of iso-cytochrome c. <i>Protein Science</i> , 1998, 7, 994-1005. | 3.1 | 10 |
| 3418 | Isolation and molecular characterization of the <i>Arabidopsis</i> TPS1 gene, encoding trehalose-phosphate synthase. <i>Plant Journal</i> , 1998, 13, 685-689. | 2.8 | 215 |
| 3419 | Identification of sna41 gene, which is the suppressor of nda4 mutation and is involved in DNA replication in <i>Schizosaccharomyces pombe</i> . <i>Genes To Cells</i> , 1998, 3, 157-166. | 0.5 | 32 |
| 3420 | A Ste6p/P-glycoprotein homologue from the asexual yeast <i>Candida albicans</i> transports the a-factor mating pheromone in <i>Saccharomyces cerevisiae</i> . <i>Molecular Microbiology</i> , 1998, 27, 587-598. | 1.2 | 55 |

| # | ARTICLE | IF | CITATIONS |
|------|--|------|-----------|
| 3421 | Probing the active site of mitogillin, a fungal ribotoxin. <i>Molecular Microbiology</i> , 1998, 29, 1019-1027. | 1.2 | 25 |
| 3422 | Characterization of the geranylgeranyl transferase type I from <i>Schizosaccharomyces pombe</i> . <i>Molecular Microbiology</i> , 1998, 29, 1357-1367. | 1.2 | 26 |
| 3423 | Molecular cloning and characterization of a <i>Candida albicans</i> gene coding for cytochrome <i>cha</i> lyase and a cell wall-related protein. <i>Molecular Microbiology</i> , 1998, 30, 67-81. | 1.2 | 15 |
| 3424 | Spermine is not essential for growth of <i>Saccharomyces cerevisiae</i> : identification of the SPE4 gene (spermine synthase) and characterization of a <i>spe4</i> deletion mutant. <i>Gene</i> , 1998, 210, 195-201. | 1.0 | 64 |
| 3425 | cDNA cloning and functional analysis of p28 (Nas6p) and p40.5 (Nas7p), two novel regulatory subunits of the 26S proteasome. <i>Gene</i> , 1998, 216, 113-122. | 1.0 | 83 |
| 3426 | Characterization of <i>alg2</i> encoding a mannosyltransferase in the zygomycete fungus <i>Rhizomucor pusillus</i> . <i>Gene</i> , 1998, 221, 179-184. | 1.0 | 13 |
| 3427 | Isolation of overexpressed yeast genes which prevent aminoglycoside toxicity. <i>Hearing Research</i> , 1998, 120, 62-68. | 0.9 | 4 |
| 3428 | N-Glycosylation does not affect assembly and targeting of proglycinin in yeast. <i>Biochimica Et Biophysica Acta - General Subjects</i> , 1998, 1379, 107-117. | 1.1 | 9 |
| 3429 | 4 Transformation of Yeast by the Lithium Acetate/Single-Stranded Carrier DNA/PEG Method. <i>Methods in Microbiology</i> , 1998, 26, 53-66. | 0.4 | 75 |
| 3430 | Using yeast to study glucocorticoid receptor phosphorylation. <i>Journal of Steroid Biochemistry and Molecular Biology</i> , 1998, 66, 303-318. | 1.2 | 13 |
| 3431 | Regulation of human estrogen receptor by phytoestrogens in yeast and human cells. <i>Journal of Steroid Biochemistry and Molecular Biology</i> , 1998, 67, 421-429. | 1.2 | 82 |
| 3432 | Gametogenesis in Yeast Is Regulated by a Transcriptional Cascade Dependent on Ndt80. <i>Molecular Cell</i> , 1998, 1, 685-696. | 4.5 | 346 |
| 3433 | Broad, but Not Universal, Transcriptional Requirement for γ TAFII17, a Histone H3-like TAFII Present in TFIID and SAGA. <i>Molecular Cell</i> , 1998, 2, 653-661. | 4.5 | 94 |
| 3434 | The Meiosis-Specific Hop2 Protein of <i>S. cerevisiae</i> Ensures Synapsis between Homologous Chromosomes. <i>Cell</i> , 1998, 94, 375-386. | 13.5 | 190 |
| 3435 | Transactivation potential of the C-terminus of human Nm23-H1. <i>FEBS Letters</i> , 1998, 423, 235-238. | 1.3 | 13 |
| 3436 | Yeast putative transcription factors involved in salt tolerance. <i>FEBS Letters</i> , 1998, 425, 323-328. | 1.3 | 129 |
| 3437 | An <i>Arabidopsis</i> cDNA encoding a bifunctional glutamine amidotransferase/cyclase suppresses the histidine auxotrophy of a <i>Saccharomyces cerevisiae</i> <i>his7</i> mutant. <i>FEBS Letters</i> , 1998, 428, 229-234. | 1.3 | 18 |
| 3438 | Heterologous expression of human H1 histones in yeast. <i>FEBS Letters</i> , 1998, 435, 245-250. | 1.3 | 23 |

| # | ARTICLE | IF | CITATIONS |
|------|--|-----|-----------|
| 3439 | phd1+, a histone deacetylase gene of <i>Schizosaccharomyces pombe</i> , is required for the meiotic cell cycle and resistance to trichostatin A1. <i>FEBS Letters</i> , 1998, 436, 193-196. | 1.3 | 16 |
| 3440 | Enhanced secretion of hydrophobic peptide fused lysozyme by the introduction of N-glycosylation signal and the disruption of calnexin gene in <i>Saccharomyces cerevisiae</i> . <i>FEBS Letters</i> , 1998, 440, 89-92. | 1.3 | 32 |
| 3441 | Incorporation of extracellular phospholipids and their effect on the growth and lipid metabolism of the <i>Saccharomyces cerevisiae</i> cho1/pss mutant. <i>Lipids and Lipid Metabolism</i> , 1998, 1394, 23-32. | 2.6 | 11 |
| 3442 | Identification of the Orotidine-5-Phosphate Decarboxylase Gene and Development of a Transformation System in the Yeast <i>Saccharomyces exiguus</i> Yp74L-3. <i>Bioscience, Biotechnology and Biochemistry</i> , 1998, 62, 2280-2282. | 0.6 | 5 |
| 3443 | Morphogenesis beyond Cytokinetic Arrest in <i>Saccharomyces cerevisiae</i> . <i>Journal of Cell Biology</i> , 1998, 143, 1617-1634. | 2.3 | 64 |
| 3444 | Purification and characterization of Hrp1, a Homolog of Mouse CHD1 from the fission yeast <i>Schizosaccharomyces pombe</i> . <i>Korean Journal of Biological Sciences</i> , 1998, 2, 539-543. | 0.1 | 4 |
| 3445 | Acidic Di-leucine Motif Essential for AP-3-dependent Sorting and Restriction of the Functional Specificity of the Vam3p Vacuolar t-SNARE. <i>Journal of Cell Biology</i> , 1998, 142, 913-922. | 2.3 | 130 |
| 3446 | Purification of a Histidine-Tagged Plant Plasma Membrane H ⁺ -ATPase Expressed in Yeast. <i>Protein Expression and Purification</i> , 1998, 12, 29-37. | 0.6 | 32 |
| 3447 | Cloning, Overexpression, and Purification of Cytosine Deaminase from <i>Saccharomyces cerevisiae</i> . <i>Protein Expression and Purification</i> , 1998, 12, 173-184. | 0.6 | 21 |
| 3448 | Cloning of the Fatty Acid Synthetase β^2 Subunit from Fission Yeast, Coexpression with the β^1 Subunit, and Purification of the Intact Multifunctional Enzyme Complex. <i>Protein Expression and Purification</i> , 1998, 13, 403-413. | 0.6 | 8 |
| 3449 | The β^1 -Helix in the Extra-Membranous Domain Contributes to the Stability of the Iron-Sulfur Protein of the Cytochrome bc ₁ Complex. <i>Archives of Biochemistry and Biophysics</i> , 1998, 359, 128-132. | 1.4 | 2 |
| 3450 | Identification of the First Mammalian Sphingosine Phosphate Lyase Gene and Its Functional Expression in Yeast. <i>Biochemical and Biophysical Research Communications</i> , 1998, 242, 502-507. | 1.0 | 170 |
| 3451 | Molecular Cloning of <i>Saccharomyces cerevisiae</i> MLF4/SSH4 Gene Which Confers the Immunosuppressant Leflunomide Resistance. <i>Biochemical and Biophysical Research Communications</i> , 1998, 246, 378-381. | 1.0 | 5 |
| 3452 | A Mutation Ser213/Asn in the Hexokinase 1 from <i>Schizosaccharomyces pombe</i> Increases Its Affinity for Glucose. <i>Biochemical and Biophysical Research Communications</i> , 1998, 251, 714-719. | 1.0 | 7 |
| 3453 | Role of RAS2 in Recovery from Chronic Stress: Effect on Yeast Life Span. <i>Experimental Cell Research</i> , 1998, 245, 368-378. | 1.2 | 46 |
| 3454 | Heat Stress-Induced Life Span Extension in Yeast. <i>Experimental Cell Research</i> , 1998, 245, 379-388. | 1.2 | 130 |
| 3455 | Direct Cloning of Human 10q25 Neocentromere DNA Using Transformation-Associated Recombination (TAR) in Yeast. <i>Genomics</i> , 1998, 47, 399-404. | 1.3 | 37 |
| 3456 | cDNA Cloning and Characterization of a Human Proteasomal Modulator Subunit, p27 (PSMD9). <i>Genomics</i> , 1998, 50, 241-250. | 1.3 | 48 |

| # | ARTICLE | IF | CITATIONS |
|------|---|-----|-----------|
| 3457 | Characterization of a Human Homologue of the <i>Saccharomyces cerevisiae</i> Transcription Factor Spt3 (SUPT3H). <i>Genomics</i> , 1998, 53, 90-96. | 1.3 | 20 |
| 3458 | Genetic analysis of <i>Xenopus</i> transcription factor IIIA 1 Edited by P. E. Wright. <i>Journal of Molecular Biology</i> , 1998, 284, 1307-1322. | 2.0 | 15 |
| 3459 | Yeast Model System for Study of Mammalian Phosphodiesterases. <i>Methods</i> , 1998, 14, 35-42. | 1.9 | 17 |
| 3460 | Essential role of tubulin-folding cofactor D in microtubule assembly and its association with microtubules in fission yeast. <i>EMBO Journal</i> , 1998, 17, 658-666. | 3.5 | 80 |
| 3461 | Transformation. , 1998, 103, 27-40. | | 115 |
| 3462 | The Role of Far1p in Linking the Heterotrimeric G Protein to Polarity Establishment Proteins During Yeast Mating. , 1998, 282, 1511-1516. | | 215 |
| 3463 | Cooperative interaction of branch signals in the actin intron of <i>Saccharomyces cerevisiae</i> . <i>Nucleic Acids Research</i> , 1998, 26, 4137-4145. | 6.5 | 2 |
| 3464 | C-Terminal Peptide of <i>Fusarium heterosporum</i> Lipase Is Necessary for Its Increasing Thermostability. <i>Journal of Biochemistry</i> , 1998, 124, 1124-1129. | 0.9 | 12 |
| 3465 | Isolation and Characterization of the <i>Saccharomyces cerevisiae</i> LPP1 Gene Encoding a Mg ²⁺ -independent Phosphatidate Phosphatase. <i>Journal of Biological Chemistry</i> , 1998, 273, 14331-14338. | 1.6 | 107 |
| 3466 | Pleiotropic effects of the <i>opil</i> regulatory mutation of yeast: its effects on growth and on phospholipid and inositol metabolism. <i>Microbiology (United Kingdom)</i> , 1998, 144, 2739-2748. | 0.7 | 34 |
| 3467 | In Vitro Reconstitution of Glucose-induced Targeting of Fructose-1,6-bisphosphatase into the Vacuole in Semi-intact Yeast Cells. <i>Journal of Biological Chemistry</i> , 1998, 273, 3381-3387. | 1.6 | 34 |
| 3468 | Identification of Kel1p, a Kelch Domain-containing Protein Involved in Cell Fusion and Morphology in <i>Saccharomyces cerevisiae</i> . <i>Journal of Cell Biology</i> , 1998, 143, 375-389. | 2.3 | 88 |
| 3469 | Alteration of the Midpoint Potential and Catalytic Activity of the Rieske Iron-Sulfur Protein by Changes of Amino Acids Forming Hydrogen Bonds to the Iron-Sulfur Cluster. <i>Journal of Biological Chemistry</i> , 1998, 273, 9085-9093. | 1.6 | 165 |
| 3470 | Phospholipase D activity is required for suppression of yeast phosphatidylinositol transfer protein defects. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 1998, 95, 12346-12351. | 3.3 | 159 |
| 3471 | A Novel Fluorescence-based Genetic Strategy Identifies Mutants of <i>Saccharomyces cerevisiae</i> Defective for Nuclear Pore Complex Assembly. <i>Molecular Biology of the Cell</i> , 1998, 9, 2439-2461. | 0.9 | 45 |
| 3472 | Molecular Cloning and Characterization of a <i>Drosophila</i> p38 Mitogen-activated Protein Kinase. <i>Journal of Biological Chemistry</i> , 1998, 273, 369-374. | 1.6 | 94 |
| 3473 | Palindrome with Spacer of One Nucleotide Is Characteristic of the cis-Acting Unfolded Protein Response Element in <i>Saccharomyces cerevisiae</i> . <i>Journal of Biological Chemistry</i> , 1998, 273, 9912-9920. | 1.6 | 125 |
| 3474 | Goosecoid-like, a gene deleted in DiGeorge and velocardiofacial syndromes, recognizes DNA with a bicoid-like specificity and is expressed in the developing mouse brain. <i>Human Molecular Genetics</i> , 1998, 7, 1497-1505. | 1.4 | 19 |

| # | ARTICLE | IF | CITATIONS |
|------|--|-----|-----------|
| 3475 | Nitrogen-regulated Ubiquitination of the Gap1 Permease of <i>Saccharomyces cerevisiae</i> . <i>Molecular Biology of the Cell</i> , 1998, 9, 1253-1263. | 0.9 | 206 |
| 3476 | Cloning and functional expression of a cDNA encoding a pheromone gland-specific acyl-CoA Δ^11 -desaturase of the cabbage looper moth, <i>Trichoplusia ni</i> . <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 1998, 95, 15287-15292. | 3.3 | 109 |
| 3477 | The yeast FBP1 poly(A) signal functions in both orientations and overlaps with a gene promoter. <i>Nucleic Acids Research</i> , 1998, 26, 4588-4596. | 6.5 | 9 |
| 3478 | Cloning and sequencing of the <i>Candida albicans</i> homologue of SRB1/PSA1/MIG9, the essential gene encoding GDP-mannose pyrophosphorylase in <i>Saccharomyces cerevisiae</i> . <i>Microbiology (United Kingdom)</i> 140:107-114 | 0.7 | 10 |
| 3479 | Single Point Mutations Distributed in 10 Soluble and Membrane Regions of the <i>Nicotiana glauca</i> Plasma Membrane PMA2 H ⁺ -ATPase Activate the Enzyme and Modify the Structure of the C-terminal Region. <i>Journal of Biological Chemistry</i> , 1998, 273, 34837-34842. | 1.6 | 120 |
| 3480 | A Human Homologue of the <i>Schizosaccharomyces pombe</i> rad1+ Checkpoint Gene Encodes an Exonuclease. <i>Journal of Biological Chemistry</i> , 1998, 273, 18332-18339. | 1.6 | 92 |
| 3481 | Ste6p Mutants Defective in Exit from the Endoplasmic Reticulum (ER) Reveal Aspects of an ER Quality Control Pathway in <i>Saccharomyces cerevisiae</i> . <i>Molecular Biology of the Cell</i> , 1998, 9, 2767-2784. | 0.9 | 107 |
| 3482 | Tuberous Sclerosis Gene 2 Product Modulates Transcription Mediated by Steroid Hormone Receptor Family Members. <i>Journal of Biological Chemistry</i> , 1998, 273, 20535-20539. | 1.6 | 112 |
| 3483 | Ribosomal P-protein Stalk Function Is Targeted by Sordarin Antifungals. <i>Journal of Biological Chemistry</i> , 1998, 273, 25041-25044. | 1.6 | 47 |
| 3484 | Dominant Negative Alleles of <i>SEC10</i> Reveal Distinct Domains Involved in Secretion and Morphogenesis in Yeast. <i>Molecular Biology of the Cell</i> , 1998, 9, 1725-1739. | 0.9 | 40 |
| 3485 | Effect of CTP Synthetase Regulation by CTP on Phospholipid Synthesis in <i>Saccharomyces cerevisiae</i> . <i>Journal of Biological Chemistry</i> , 1998, 273, 18992-19001. | 1.6 | 112 |
| 3486 | Interaction between Mitochondria and the Actin Cytoskeleton in Budding Yeast Requires Two Integral Mitochondrial Outer Membrane Proteins, Mmm1p and Mdm10p. <i>Journal of Cell Biology</i> , 1998, 141, 1371-1381. | 2.3 | 177 |
| 3487 | Fab1p Is Essential for PtdIns(3)P 5-Kinase Activity and the Maintenance of Vacuolar Size and Membrane Homeostasis. <i>Journal of Cell Biology</i> , 1998, 143, 65-79. | 2.3 | 395 |
| 3488 | The Spermidine Transport System Is Regulated by Ligand Inactivation, Endocytosis, and by the Npr1p Ser/Thr Protein Kinase in <i>Saccharomyces cerevisiae</i> . <i>Journal of Biological Chemistry</i> , 1998, 273, 2109-2117. | 1.6 | 30 |
| 3489 | Relocalization of Phospholipase D Activity Mediates Membrane Formation During Meiosis. <i>Journal of Cell Biology</i> , 1998, 140, 81-90. | 2.3 | 151 |
| 3490 | Identification of Novel Temperature-sensitive Lethal Alleles in Essential β -Tubulin and Nonessential α -Tubulin Genes as Fission Yeast Polarity Mutants. <i>Molecular Biology of the Cell</i> , 1998, 9, 1757-1771. | 0.9 | 87 |
| 3491 | Tropomyosin-containing Actin Cables Direct the Myo2p-dependent Polarized Delivery of Secretory Vesicles in Budding Yeast. <i>Journal of Cell Biology</i> , 1998, 143, 1931-1945. | 2.3 | 310 |
| 3492 | Production of Human Compatible High Mannose-type (Man5GlcNAc2) Sugar Chains in <i>Saccharomyces cerevisiae</i> . <i>Journal of Biological Chemistry</i> , 1998, 273, 26298-26304. | 1.6 | 131 |

| # | ARTICLE | IF | CITATIONS |
|------|---|-----|-----------|
| 3493 | Two-Hybrid Analysis of Ras-Raf Interactions. , 1998, 84, 201-222. | | 14 |
| 3494 | Multidrug Resistance Phenotype Conferred by Overexpressing bfr2+/pad1+/sks1+or pap1+ Genes and Mediated by bfr1+ Gene Product, a Structural and Functional Homologue of P-Glycoprotein in <i>Schizosaccharomyces pombe</i> . <i>Bioscience, Biotechnology and Biochemistry</i> , 1998, 62, 390-392. | 0.6 | 9 |
| 3495 | Identification and Analysis of Frameshift Sites. , 1998, 77, 399-412. | | 4 |
| 3496 | Cloning and Characterization of Two Phosphate Transporters from <i>Medicago truncatula</i> Roots: Regulation in Response to Phosphate and to Colonization by Arbuscular Mycorrhizal (AM) Fungi. <i>Molecular Plant-Microbe Interactions</i> , 1998, 11, 14-22. | 1.4 | 264 |
| 3497 | Structural Similarities Between Hammerhead Ribozymes and the Spliceosomal RNAs Could Be Responsible for Lack of Ribozyme Cleavage in Yeast. <i>Oligonucleotides</i> , 1998, 8, 1-13. | 4.4 | 9 |
| 3498 | Molecular Cloning and Expression in <i>Saccharomyces cerevisiae</i> of Tobacco NADPH-Cytochrome P450 Oxidoreductase cDNA. <i>Bioscience, Biotechnology and Biochemistry</i> , 1998, 62, 1403-1411. | 0.6 | 7 |
| 3499 | Overexpression of peptide-methionine sulfoxide reductase in <i>Saccharomyces cerevisiae</i> and human T cells provides them with high resistance to oxidative stress. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 1998, 95, 14071-14075. | 3.3 | 260 |
| 3500 | HAH1 Is a Copper-binding Protein with Distinct Amino Acid Residues Mediating Copper Homeostasis and Antioxidant Defense. <i>Journal of Biological Chemistry</i> , 1998, 273, 1749-1754. | 1.6 | 130 |
| 3501 | The Role of Charged Amino Acids in the ± 1 - ± 4 Loop of the Iron-Sulfur Protein of the Cytochrome bc ₁ Complex of Yeast Mitochondria. <i>Journal of Biological Chemistry</i> , 1998, 273, 11917-11922. | 1.6 | 7 |
| 3502 | Nip1p Associates with 40 S Ribosomes and the Prt1p Subunit of Eukaryotic Initiation Factor 3 and Is Required for Efficient Translation Initiation. <i>Journal of Biological Chemistry</i> , 1998, 273, 23485-23494. | 1.6 | 22 |
| 3503 | Coiled-coil Interaction of N-terminal 36 Residues of Cyclase-associated Protein with Adenylyl Cyclase Is Sufficient for Its Function in <i>Saccharomyces cerevisiae</i> Ras Pathway. <i>Journal of Biological Chemistry</i> , 1998, 273, 28019-28024. | 1.6 | 45 |
| 3504 | Unconventional Splicing of HAC1/ERN4 mRNA Required for the Unfolded Protein Response. <i>Journal of Biological Chemistry</i> , 1998, 273, 1802-1807. | 1.6 | 84 |
| 3505 | The Mutagenesis Protein MucB Interacts with Single Strand DNA Binding Protein and Induces a Major Conformational Change in Its Complex with Single-stranded DNA. <i>Journal of Biological Chemistry</i> , 1998, 273, 5520-5527. | 1.6 | 31 |
| 3506 | The Gcn5 \hat{A} -Ada Complex Potentiates the Histone Acetyltransferase Activity of Gcn5. <i>Journal of Biological Chemistry</i> , 1998, 273, 24414-24419. | 1.6 | 29 |
| 3507 | A Ribosomal Protein Is Required for Translational Regulation of GCN4 mRNA. <i>Journal of Biological Chemistry</i> , 1998, 273, 32870-32877. | 1.6 | 15 |
| 3508 | Characterization of Two Human Genes Encoding Acyl Coenzyme A:Cholesterol Acyltransferase-related Enzymes. <i>Journal of Biological Chemistry</i> , 1998, 273, 26765-26771. | 1.6 | 341 |
| 3509 | Identification of a CAP (adenylyl-cyclase-associated protein) homologous gene in <i>Lentinus edodes</i> and its functional complementation of yeast CAP mutants. <i>Microbiology (United Kingdom)</i> , 1998, 144, 1085-1093. | 0.7 | 21 |
| 3510 | Phosphorylation Region of the Yeast Plasma-membrane H ⁺ -ATPase. <i>Journal of Biological Chemistry</i> , 1998, 273, 21744-21751. | 1.6 | 49 |

| # | ARTICLE | IF | CITATIONS |
|------|--|-----|-----------|
| 3511 | Radicicol Leads to Selective Depletion of Raf Kinase and Disrupts K-Ras-activated Aberrant Signaling Pathway. <i>Journal of Biological Chemistry</i> , 1998, 273, 822-828. | 1.6 | 87 |
| 3512 | The <i>Saccharomyces cerevisiae</i> SOP1 and SOP2 Genes, Which Act in Cation Homeostasis, Can Be Functionally Substituted by the <i>Drosophila lethal(2)giant</i> larval Tumor Suppressor Gene. <i>Journal of Biological Chemistry</i> , 1998, 273, 33610-33618. | 1.6 | 39 |
| 3513 | Phosphatidylinositol-4-phosphate 5-Kinase Localized on the Plasma Membrane Is Essential for Yeast Cell Morphogenesis. <i>Journal of Biological Chemistry</i> , 1998, 273, 15779-15786. | 1.6 | 164 |
| 3514 | The Integral Membrane Protein Snl1p Is Genetically Linked to Yeast Nuclear Pore Complex Function. <i>Molecular Biology of the Cell</i> , 1998, 9, 355-373. | 0.9 | 51 |
| 3515 | The <i>Saccharomyces cerevisiae</i> Prenylcysteine Carboxyl Methyltransferase Ste14p Is in the Endoplasmic Reticulum Membrane. <i>Molecular Biology of the Cell</i> , 1998, 9, 2231-2247. | 0.9 | 105 |
| 3516 | A Functional GTPase Domain, but not its Transmembrane Domain, is Required for Function of the SRP Receptor β -subunit. <i>Journal of Cell Biology</i> , 1998, 142, 341-354. | 2.3 | 57 |
| 3517 | Functional Interaction of Isr1, a Predicted Protein Kinase, with the Pkc1 Pathway in <i>Saccharomyces cerevisiae</i> . <i>Bioscience, Biotechnology and Biochemistry</i> , 1998, 62, 1376-1380. | 0.6 | 7 |
| 3518 | Expression, Purification, and Characterization of Choline Kinase, Product of the CKI Gene from <i>Saccharomyces cerevisiae</i> . <i>Journal of Biological Chemistry</i> , 1998, 273, 6844-6852. | 1.6 | 51 |
| 3519 | Multiple Functions for Actin during Filamentous Growth of <i>Saccharomyces cerevisiae</i> . <i>Molecular Biology of the Cell</i> , 1998, 9, 1873-1889. | 0.9 | 38 |
| 3520 | Rvs161p Interacts with Fus2p to Promote Cell Fusion in <i>Saccharomyces cerevisiae</i> . <i>Journal of Cell Biology</i> , 1998, 141, 567-584. | 2.3 | 85 |
| 3521 | A Role for the Luminal Domain in Golgi Localization of the <i>Saccharomyces cerevisiae</i> Guanosine Diphosphatase. <i>Molecular Biology of the Cell</i> , 1998, 9, 1351-1365. | 0.9 | 16 |
| 3522 | Divergent Transcriptional Control of Multidrug Resistance Genes in <i>Saccharomyces cerevisiae</i> . <i>Journal of Biological Chemistry</i> , 1998, 273, 2098-2104. | 1.6 | 42 |
| 3523 | The Kinesin-related Proteins, Kip2p and Kip3p, Function Differently in Nuclear Migration in Yeast. <i>Molecular Biology of the Cell</i> , 1998, 9, 2051-2068. | 0.9 | 133 |
| 3524 | Sequence Determinants for Regulated Degradation of Yeast 3-Hydroxy-3-Methylglutaryl-CoA Reductase, an Integral Endoplasmic Reticulum Membrane Protein. <i>Molecular Biology of the Cell</i> , 1998, 9, 2611-2626. | 0.9 | 72 |
| 3525 | Substitutions of Aspartate 378 in the Phosphorylation Domain of the Yeast PMA1 H ⁺ -ATPase Disrupt Protein Folding and Biogenesis. <i>Journal of Biological Chemistry</i> , 1998, 273, 7338-7344. | 1.6 | 43 |
| 3526 | Tyrosine and Tryptophan Act through the Same Binding Site at the Dimer Interface of Yeast Chorismate Mutase. <i>Journal of Biological Chemistry</i> , 1998, 273, 17012-17017. | 1.6 | 27 |
| 3527 | Identification of a Human Homologue of the <i>Schizosaccharomyces pombe</i> rad17 ⁺ Checkpoint Gene. <i>Journal of Biological Chemistry</i> , 1998, 273, 18340-18346. | 1.6 | 64 |
| 3528 | Peroxisomal Δ^3 -cis- Δ^2 -trans-Enoyl-CoA Isomerase Encoded by ECI1 Is Required for Growth of the Yeast <i>Saccharomyces cerevisiae</i> on Unsaturated Fatty Acids. <i>Journal of Biological Chemistry</i> , 1998, 273, 31366-31374. | 1.6 | 56 |

| # | ARTICLE | IF | CITATIONS |
|------|---|-----|-----------|
| 3529 | Evidence for a Salt Bridge between Transmembrane Segments 5 and 6 of the Yeast Plasma-membrane H ⁺ -ATPase. <i>Journal of Biological Chemistry</i> , 1998, 273, 34328-34334. | 1.6 | 36 |
| 3530 | Isolation and Characterization of the <i>Saccharomyces cerevisiae</i> DPP1 Gene Encoding Diacylglycerol Pyrophosphate Phosphatase. <i>Journal of Biological Chemistry</i> , 1998, 273, 3278-3284. | 1.6 | 109 |
| 3531 | Functional Expression of the Menkes Disease Protein Reveals Common Biochemical Mechanisms Among the Copper-transporting P-type ATPases. <i>Journal of Biological Chemistry</i> , 1998, 273, 3765-3770. | 1.6 | 145 |
| 3532 | Time-Lapse Microscopy Reveals Unique Roles for Kinesins during Anaphase in Budding Yeast. <i>Journal of Cell Biology</i> , 1998, 143, 687-694. | 2.3 | 248 |
| 3533 | Distinct Morphological Phenotypes of Cell Fusion Mutants. <i>Molecular Biology of the Cell</i> , 1998, 9, 1395-1410. | 0.9 | 100 |
| 3534 | PCR- and ligation-mediated synthesis of marker cassettes with long flanking homology regions for gene disruption in <i>Saccharomyces cerevisiae</i> . <i>Nucleic Acids Research</i> , 1998, 26, 860-861. | 6.5 | 56 |
| 3535 | The C-terminal silencing domain of Rap1p is essential for the repression of ribosomal protein genes in response to a defect in the secretory pathway. <i>Nucleic Acids Research</i> , 1998, 26, 1063-1069. | 6.5 | 34 |
| 3536 | Functional Evidence for UDP-galactose Transporter in <i>Saccharomyces cerevisiae</i> through the in Vivo Galactosylation and in Vitro Transport Assay. <i>Journal of Biological Chemistry</i> , 1998, 273, 2583-2590. | 1.6 | 26 |
| 3537 | Mammalian Prenylcysteine Carboxyl Methyltransferase Is in the Endoplasmic Reticulum. <i>Journal of Biological Chemistry</i> , 1998, 273, 15030-15034. | 1.6 | 262 |
| 3538 | Structure-Function Relationships in Membrane Segment 5 of the Yeast Pma1 H ⁺ -ATPase. <i>Journal of Biological Chemistry</i> , 1998, 273, 17411-17417. | 1.6 | 26 |
| 3539 | Tomato Fructokinases Exhibit Differential Expression and Substrate Regulation1. <i>Plant Physiology</i> , 1998, 117, 85-90. | 2.3 | 87 |
| 3540 | A Mutation in a Novel Yeast Proteasomal Gene, <i>RPN11/MPR1</i> , Produces a Cell Cycle Arrest, Overreplication of Nuclear and Mitochondrial DNA, and an Altered Mitochondrial Morphology. <i>Molecular Biology of the Cell</i> , 1998, 9, 2917-2931. | 0.9 | 67 |
| 3541 | Myb-Related <i>Schizosaccharomyces pombe</i> <i>cdc5p</i> Is Structurally and Functionally Conserved in Eukaryotes. <i>Molecular and Cellular Biology</i> , 1998, 18, 4097-4108. | 1.1 | 70 |
| 3542 | Identification of <i>APN2</i> , the <i>Saccharomyces cerevisiae</i> homolog of the major human AP endonuclease <i>HAP1</i> , and its role in the repair of abasic sites. <i>Genes and Development</i> , 1998, 12, 3137-3143. | 2.7 | 184 |
| 3543 | Processing of telomeric DNA ends requires the passage of a replication fork. <i>Nucleic Acids Research</i> , 1998, 26, 5365-5371. | 6.5 | 80 |
| 3544 | Cloning and functional expression of the human GlcNAc-1-P transferase, the enzyme for the committed step of the dolichol cycle, by heterologous complementation in <i>Saccharomyces cerevisiae</i> . <i>Glycobiology</i> , 1998, 8, 77-85. | 1.3 | 36 |
| 3545 | Mutation of the polo-box disrupts localization and mitotic functions of the mammalian polo kinase Plk. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 1998, 95, 9301-9306. | 3.3 | 234 |
| 3546 | [7] Identification of ion channel-associated proteins using the yeast two-hybrid system. <i>Methods in Enzymology</i> , 1998, 293, 104-122. | 0.4 | 28 |

| # | ARTICLE | IF | CITATIONS |
|------|--|-----|-----------|
| 3547 | A genetic system based on split-ubiquitin for the analysis of interactions between membrane proteins in vivo. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 1998, 95, 5187-5192. | 3.3 | 567 |
| 3548 | Role for the Ubiquitin-Proteasome System in the Vacuolar Degradation of Ste6p, the <i>Yap1</i> -Factor Transporter in <i>Saccharomyces cerevisiae</i> . <i>Molecular and Cellular Biology</i> , 1998, 18, 779-789. | 1.1 | 66 |
| 3549 | DNA-Binding Activities of Hop1 Protein, a Synaptonemal Complex Component from <i>Saccharomyces cerevisiae</i> . <i>Molecular and Cellular Biology</i> , 1998, 18, 1424-1435. | 1.1 | 78 |
| 3550 | SWI-SNF Complex Participation in Transcriptional Activation at a Step Subsequent to Activator Binding. <i>Molecular and Cellular Biology</i> , 1998, 18, 1774-1782. | 1.1 | 74 |
| 3551 | Spa2p Interacts with Cell Polarity Proteins and Signaling Components Involved in Yeast Cell Morphogenesis. <i>Molecular and Cellular Biology</i> , 1998, 18, 4053-4069. | 1.1 | 218 |
| 3552 | Temperature-Induced Expression of Yeast <i>FKS2</i> Is under the Dual Control of Protein Kinase C and Calcineurin. <i>Molecular and Cellular Biology</i> , 1998, 18, 1013-1022. | 1.1 | 198 |
| 3553 | Carbon Source-Dependent Phosphorylation of Hexokinase PII and Its Role in the Glucose-Signaling Response in Yeast. <i>Molecular and Cellular Biology</i> , 1998, 18, 2940-2948. | 1.1 | 112 |
| 3554 | Identification of a Translation Initiation Factor 3 (eIF3) Core Complex, Conserved in Yeast and Mammals, That Interacts with eIF5. <i>Molecular and Cellular Biology</i> , 1998, 18, 4935-4946. | 1.1 | 173 |
| 3555 | eIF2 independently binds two distinct eIF2B subcomplexes that catalyze and regulate guanine-nucleotide exchange. <i>Genes and Development</i> , 1998, 12, 514-526. | 2.7 | 236 |
| 3556 | Separation of inhibition and activation of the allosteric yeast chorismate mutase. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 1998, 95, 2868-2873. | 3.3 | 18 |
| 3557 | Yeast Ty1 Retrotransposition Is Stimulated by a Synergistic Interaction between Mutations in Chromatin Assembly Factor I and Histone Regulatory Proteins. <i>Molecular and Cellular Biology</i> , 1998, 18, 4783-4792. | 1.1 | 62 |
| 3558 | Inhibition of Double-Stranded RNA-Dependent Protein Kinase PKR by Vaccinia Virus E3: Role of Complex Formation and the E3 N-Terminal Domain. <i>Molecular and Cellular Biology</i> , 1998, 18, 7304-7316. | 1.1 | 176 |
| 3559 | DBF2 Protein Kinase Binds to and Acts through the Cell Cycle-Regulated MOB1 Protein. <i>Molecular and Cellular Biology</i> , 1998, 18, 2100-2107. | 1.1 | 98 |
| 3560 | Prohibitin Family Members Interact Genetically with Mitochondrial Inheritance Components in <i>Saccharomyces cerevisiae</i> . <i>Molecular and Cellular Biology</i> , 1998, 18, 4043-4052. | 1.1 | 179 |
| 3561 | Multiple and Distinct Activation and Repression Sequences Mediate the Regulated Transcription of <i>IME1</i> , a Transcriptional Activator of Meiosis-Specific Genes in <i>Saccharomyces cerevisiae</i> . <i>Molecular and Cellular Biology</i> , 1998, 18, 1985-1995. | 1.1 | 75 |
| 3562 | The Nucleic Acid Binding Activity of Bleomycin Hydrolase Is Involved in Bleomycin Detoxification. <i>Molecular and Cellular Biology</i> , 1998, 18, 3580-3585. | 1.1 | 28 |
| 3563 | Vam7p, a SNAP-25-Like Molecule, and Vam3p, a Syntaxin Homolog, Function Together in Yeast Vacuolar Protein Trafficking. <i>Molecular and Cellular Biology</i> , 1998, 18, 5308-5319. | 1.1 | 187 |
| 3564 | Regulation of Transcription Factor Pdr1p Function by an Hsp70 Protein in <i>Saccharomyces cerevisiae</i> . <i>Molecular and Cellular Biology</i> , 1998, 18, 1147-1155. | 1.1 | 80 |

| # | ARTICLE | IF | CITATIONS |
|------|---|-----|-----------|
| 3565 | myc Boxes, Which Are Conserved in myc Family Proteins, Are Signals for Protein Degradation via the Proteasome. <i>Molecular and Cellular Biology</i> , 1998, 18, 5961-5969. | 1.1 | 117 |
| 3566 | Distinct Subclasses of Small GTPases Interact with Guanine Nucleotide Exchange Factors in a Similar Manner. <i>Molecular and Cellular Biology</i> , 1998, 18, 7444-7454. | 1.1 | 23 |
| 3567 | Drug Pumping Mechanisms in <i>Canadida albicans</i> .. <i>Medical Mycology Journal</i> , 1998, 39, 73-78. | 0.9 | 18 |
| 3568 | Identification of a putative alpha-glucan synthase essential for cell wall construction and morphogenesis in fission yeast. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 1998, 95, 9161-9166. | 3.3 | 158 |
| 3569 | Cell Cycle Regulation of the <i>Saccharomyces cerevisiae</i> Polo-Like Kinase Cdc5p. <i>Molecular and Cellular Biology</i> , 1998, 18, 7360-7370. | 1.1 | 104 |
| 3570 | [28] Yeast secretory vesicle system for expression and functional characterization of P-glycoproteins. <i>Methods in Enzymology</i> , 1998, 292, 382-396. | 0.4 | 5 |
| 3571 | Isolation and characterization of the yeast <i>las21</i> mutants, which are sensitive to a local anesthetic, tetracaine.. <i>Genes and Genetic Systems</i> , 1998, 73, 365-375. | 0.2 | 10 |
| 3572 | Characterization of a calmodulin-binding transporter from the plasma membrane of barley aleurone. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 1998, 95, 1944-1949. | 3.3 | 175 |
| 3573 | Efficient Large-Scale Transformation of Yeast. <i>BioTechniques</i> , 1998, 24, 596-600. | 0.8 | 15 |
| 3574 | <i>Pichia stipitis</i> Genes for Alcohol Dehydrogenase with Fermentative and Respiratory Functions. <i>Applied and Environmental Microbiology</i> , 1998, 64, 1350-1358. | 1.4 | 67 |
| 3575 | The RNA recognition motif of yeast translation initiation factor Tif3/eIF4B is required but not sufficient for RNA strand-exchange and translational activity. <i>Rna</i> , 1998, 4, 1259-1267. | 1.6 | 23 |
| 3576 | Region specificity of chromosome III on gene expression in the yeast <i>Saccharomyces cerevisiae</i> .. <i>Journal of General and Applied Microbiology</i> , 1998, 44, 275-281. | 0.4 | 8 |
| 3577 | A Novel Asn344 Deletion in the Core Domain of Coagulation Factor XIII A Subunit: Its Effects on Protein Structure and Function. <i>Blood</i> , 1998, 92, 481-487. | 0.6 | 17 |
| 3578 | A role for U2/U6 helix Ib in 5' splice site selection. <i>Rna</i> , 1998, 4, 915-927. | 1.6 | 25 |
| 3579 | Chimeric rRNAs containing the GTPase centers of the developmentally regulated ribosomal rRNAs of <i>Plasmodium falciparum</i> are functionally distinct. <i>Rna</i> , 1998, 4, 594-602. | 1.6 | 42 |
| 3580 | Breeding of flocculent industrial alcohol yeast strains by self-cloning of the flocculation gene FLO1 and repeated-batch fermentation by transformants.. <i>Journal of General and Applied Microbiology</i> , 1998, 44, 347-353. | 0.4 | 26 |
| 3581 | The TOR (target of rapamycin) signal transduction pathway regulates the stability of translation initiation factor eIF4G in the yeast <i>Saccharomyces cerevisiae</i> . <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 1998, 95, 4264-4269. | 3.3 | 134 |
| 3582 | A yeast model for the study of Batten disease. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 1998, 95, 6915-6918. | 3.3 | 82 |

| # | ARTICLE | IF | CITATIONS |
|------|--|-----|-----------|
| 3583 | A translational repression assay procedure (TRAP) for RNA-protein interactions in vivo. Proceedings of the National Academy of Sciences of the United States of America, 1998, 95, 951-956. | 3.3 | 39 |
| 3584 | Translation Elongation Factor 2 Is Part of the Target for a New Family of Antifungals. Antimicrobial Agents and Chemotherapy, 1998, 42, 2694-2699. | 1.4 | 69 |
| 3585 | In vivo functional discrimination between plant thioredoxins by heterologous expression in the yeast <i>Saccharomyces cerevisiae</i> . Proceedings of the National Academy of Sciences of the United States of America, 1998, 95, 3312-3317. | 3.3 | 118 |
| 3586 | Mutations in VPS16 and MRT1 Stabilize mRNAs by Activating an Inhibitor of the Decapping Enzyme. Molecular and Cellular Biology, 1999, 19, 7568-7576. | 1.1 | 31 |
| 3587 | Functional analysis of aquaporin-2 mutants associated with nephrogenic diabetes insipidus by yeast expression. American Journal of Physiology - Renal Physiology, 1999, 277, F734-F741. | 1.3 | 20 |
| 3588 | The Yeast Nucleoporin Nup2p Is Involved in Nuclear Export of Importin β /Srp1p. Journal of Biological Chemistry, 1999, 274, 32360-32367. | 1.6 | 55 |
| 3589 | Transient interaction of BBP/ScSF1 and Mud2 with the splicing machinery affects the kinetics of spliceosome assembly. Rna, 1999, 5, 819-831. | 1.6 | 79 |
| 3590 | Isolation of centromeric DNA from <i>Saccharomyces bayanus</i> .. Journal of General and Applied Microbiology, 1999, 45, 89-92. | 0.4 | 2 |
| 3591 | Isolation and characterization of triacylglycerol-secreting mutant strain from yeast, <i>Saccharomyces cerevisiae</i> .. Journal of General and Applied Microbiology, 1999, 45, 1-6. | 0.4 | 17 |
| 3592 | The Nuclease Activity of Mre11 Is Required for Meiosis but Not for Mating Type Switching, End Joining, or Telomere Maintenance. Molecular and Cellular Biology, 1999, 19, 556-566. | 1.1 | 410 |
| 3593 | Ssy1p and Ptr3p Are Plasma Membrane Components of a Yeast System That Senses Extracellular Amino Acids. Molecular and Cellular Biology, 1999, 19, 5405-5416. | 1.1 | 186 |
| 3594 | Yap1p Activates Gene Transcription in an Oxidant-Specific Fashion. Molecular and Cellular Biology, 1999, 19, 8302-8313. | 1.1 | 139 |
| 3595 | Shr3p Mediates Specific COPII Coatomerâ€‘Cargo Interactions Required for the Packaging of Amino Acid Permeases Into ER-derived Transport Vesicles. Molecular Biology of the Cell, 1999, 10, 3549-3565. | 0.9 | 59 |
| 3596 | Pathways of As(III) detoxification in <i>Saccharomyces cerevisiae</i> . Proceedings of the National Academy of Sciences of the United States of America, 1999, 96, 5001-5006. | 3.3 | 391 |
| 3597 | Ctf19p: A Novel Kinetochore Protein in <i>Saccharomyces cerevisiae</i> and a Potential Link between the Kinetochore and Mitotic Spindle. Journal of Cell Biology, 1999, 145, 15-28. | 2.3 | 109 |
| 3598 | The Two Major Types of Plant Plasma Membrane H ⁺ -ATPases Show Different Enzymatic Properties and Confer Differential pH Sensitivity of Yeast Growth1. Plant Physiology, 1999, 119, 627-634. | 2.3 | 52 |
| 3599 | The <i>Saccharomyces cerevisiae</i> MER3 gene, encoding a novel helicase-like protein, is required for crossover control in meiosis. EMBO Journal, 1999, 18, 5714-5723. | 3.5 | 128 |
| 3600 | RNA Polymerase-specific Nucleosome Disruption by Transcription in Vivo. Journal of Biological Chemistry, 1999, 274, 16431-16436. | 1.6 | 22 |

| # | ARTICLE | IF | CITATIONS |
|------|--|-----|-----------|
| 3601 | A Novel Family of Yeast Chaperons Involved in the Distribution of V-ATPase and Other Membrane Proteins. <i>Journal of Biological Chemistry</i> , 1999, 274, 26885-26893. | 1.6 | 83 |
| 3602 | A Role for Tlg1p in the Transport of Proteins within the Golgi Apparatus of <i>Saccharomyces cerevisiae</i> . <i>Molecular Biology of the Cell</i> , 1999, 10, 2407-2423. | 0.9 | 50 |
| 3603 | Using a Biochemical Approach to Identify the Primary Dimerization Regions in Human DNA Topoisomerase III α . <i>Journal of Biological Chemistry</i> , 1999, 274, 26529-26536. | 1.6 | 19 |
| 3604 | The LPP1 and DPP1 Gene Products Account for Most of the Isoprenoid Phosphate Phosphatase Activities in <i>Saccharomyces cerevisiae</i> . <i>Journal of Biological Chemistry</i> , 1999, 274, 14831-14837. | 1.6 | 126 |
| 3605 | A Modulatory Role for Clathrin Light Chain Phosphorylation in Golgi Membrane Protein Localization during Vegetative Growth and during the Mating Response of <i>Saccharomyces cerevisiae</i> . <i>Molecular Biology of the Cell</i> , 1999, 10, 713-726. | 0.9 | 14 |
| 3606 | The Rho GTPase Rho3 Has a Direct Role in Exocytosis That Is Distinct from Its Role in Actin Polarity. <i>Molecular Biology of the Cell</i> , 1999, 10, 4121-4133. | 0.9 | 192 |
| 3607 | Phosphorylation and Regulation of Choline Kinase from <i>Saccharomyces cerevisiae</i> by Protein Kinase A. <i>Journal of Biological Chemistry</i> , 1999, 274, 9531-9538. | 1.6 | 36 |
| 3608 | Genetic Interactions between <i>KAR7/SEC71</i> , <i>KAR8/JEM1</i> , <i>KAR5</i> , and <i>KAR2</i> during Nuclear Fusion in <i>Saccharomyces cerevisiae</i> . <i>Molecular Biology of the Cell</i> , 1999, 10, 609-626. | 0.9 | 44 |
| 3609 | Functional Domain Analysis of the Yeast ABC Transporter Ycf1p by Site-directed Mutagenesis. <i>Journal of Biological Chemistry</i> , 1999, 274, 23584-23590. | 1.6 | 27 |
| 3610 | Identification of a Gene for a Polyamine Transport Protein in Yeast. <i>Journal of Biological Chemistry</i> , 1999, 274, 3265-3267. | 1.6 | 93 |
| 3611 | Characterization and Chromosomal Localization of USP3, a Novel Human Ubiquitin-specific Protease. <i>Journal of Biological Chemistry</i> , 1999, 274, 26878-26884. | 1.6 | 25 |
| 3612 | Disruption of YHC8, a Member of the TSR1 Gene Family, Reveals Its Direct Involvement in Yeast Protein Translocation. <i>Journal of Biological Chemistry</i> , 1999, 274, 11296-11302. | 1.6 | 10 |
| 3613 | Differential Regulation of the Kar3p Kinesin-related Protein by Two Associated Proteins, Cik1p and Vik1p. <i>Journal of Cell Biology</i> , 1999, 144, 1219-1233. | 2.3 | 100 |
| 3614 | Early Steps in Assembly of the Yeast Vacuolar H ⁺ -ATPase. <i>Journal of Biological Chemistry</i> , 1999, 274, 17275-17283. | 1.6 | 41 |
| 3615 | Molecular Dissection of Guanine Nucleotide Dissociation Inhibitor Function in Vivo. <i>Journal of Biological Chemistry</i> , 1999, 274, 14806-14817. | 1.6 | 52 |
| 3616 | Pokeweed Antiviral Protein Accesses Ribosomes by Binding to L3. <i>Journal of Biological Chemistry</i> , 1999, 274, 3859-3864. | 1.6 | 101 |
| 3617 | Helical Interactions and Membrane Disposition of the 16-kDa Proteolipid Subunit of the Vacuolar H ⁺ -ATPase Analyzed by Cysteine Replacement Mutagenesis. <i>Journal of Biological Chemistry</i> , 1999, 274, 25461-25470. | 1.6 | 22 |
| 3618 | High-Copy Suppressor Analysis Reveals a Physical Interaction between Sec34p and Sec35p, a Protein Implicated in Vesicle Docking. <i>Molecular Biology of the Cell</i> , 1999, 10, 3317-3329. | 0.9 | 78 |

| # | ARTICLE | IF | CITATIONS |
|------|--|-----|-----------|
| 3619 | L-Asparaginase II of <i>Saccharomyces cerevisiae</i> . , 1999, , 311-316. | | 1 |
| 3620 | Two Endoplasmic Reticulum (ER) Membrane Proteins That Facilitate ER-to-Golgi Transport of Glycosylphosphatidylinositol-anchored Proteins. <i>Molecular Biology of the Cell</i> , 1999, 10, 1043-1059. | 0.9 | 98 |
| 3621 | Nucleotide excision repair in a constitutive and inducible gene of a yeast minichromosome in intact cells. <i>Nucleic Acids Research</i> , 1999, 27, 3610-3620. | 6.5 | 13 |
| 3622 | Characterization of Fus3 Localization: Active Fus3 Localizes in Complexes of Varying Size and Specific Activity. <i>Molecular Biology of the Cell</i> , 1999, 10, 1553-1568. | 0.9 | 37 |
| 3623 | ATP hydrolysis activity of the DEAD box protein Rok1p is required for in vivo ROK1 function. <i>Nucleic Acids Research</i> , 1999, 27, 2753-2759. | 6.5 | 24 |
| 3624 | The GTP-binding Protein Rho1p Is Required for Cell Cycle Progression and Polarization of the Yeast Cell. <i>Journal of Cell Biology</i> , 1999, 146, 373-387. | 2.3 | 54 |
| 3625 | Cell Wall Stress Depolarizes Cell Growth via Hyperactivation of Rho1. <i>Journal of Cell Biology</i> , 1999, 147, 163-174. | 2.3 | 270 |
| 3626 | Structural and Functional Analysis of a Novel Coiled-Coil Protein Involved in Ypt6 GTPase-regulated Protein Transport in Yeast. <i>Molecular Biology of the Cell</i> , 1999, 10, 63-75. | 0.9 | 81 |
| 3627 | The Yck2 Yeast Casein Kinase 1 Isoform Shows Cell Cycle-specific Localization to Sites of Polarized Growth and Is Required for Proper Septin Organization. <i>Molecular Biology of the Cell</i> , 1999, 10, 1077-1092. | 0.9 | 63 |
| 3628 | Isolation of Supernumerary Yeast ATP Synthase Subunits e and i. <i>Journal of Biological Chemistry</i> , 1999, 274, 543-548. | 1.6 | 47 |
| 3629 | Protein Kinase C Enables the Regulatory Circuit That Connects Membrane Synthesis to Ribosome Synthesis in <i>Saccharomyces cerevisiae</i> . <i>Journal of Biological Chemistry</i> , 1999, 274, 13235-13241. | 1.6 | 86 |
| 3630 | Isolation and Characterization of the <i>Saccharomyces cerevisiae</i> EK11 Gene Encoding Ethanolamine Kinase. <i>Journal of Biological Chemistry</i> , 1999, 274, 14857-14866. | 1.6 | 79 |
| 3631 | Functional Analysis of a Homopolymeric (dA-dT) Element That Provides Nucleosomal Access to Yeast and Mammalian Transcription Factors. <i>Journal of Biological Chemistry</i> , 1999, 274, 23752-23760. | 1.6 | 30 |
| 3632 | Characterization and Function in Vivo of Two Novel Phospholipases B/Lysophospholipases from <i>Saccharomyces cerevisiae</i> . <i>Journal of Biological Chemistry</i> , 1999, 274, 28121-28127. | 1.6 | 99 |
| 3633 | Characterization of the Gene Encoding Serine Acetyltransferase, a Regulated Enzyme of Cysteine Biosynthesis from the Protist Parasites <i>Entamoeba histolytica</i> and <i>Entamoeba dispar</i> . <i>Journal of Biological Chemistry</i> , 1999, 274, 32445-32452. | 1.6 | 124 |
| 3634 | Characterization of a Novel ADP-ribosylation Factor-like Protein (γ ARL3) in <i>Saccharomyces cerevisiae</i> . <i>Journal of Biological Chemistry</i> , 1999, 274, 3819-3827. | 1.6 | 41 |
| 3635 | An Arabidopsis GSK3/shaggy-Like Gene That Complements Yeast Salt Stress-Sensitive Mutants Is Induced by NaCl and Abscisic Acid. <i>Plant Physiology</i> , 1999, 119, 1527-1534. | 2.3 | 91 |
| 3636 | A Transmembrane Hybrid-Type Histidine Kinase in Arabidopsis Functions as an Osmosensor. <i>Plant Cell</i> , 1999, 11, 1743-1754. | 3.1 | 501 |

| # | ARTICLE | IF | CITATIONS |
|------|---|-----|-----------|
| 3637 | A conditional U5 snRNA mutation affecting pre-mRNA splicing and nuclear pre-mRNA retention identifies SSD1/SRK1 as a general splicing mutant suppressor. <i>Nucleic Acids Research</i> , 1999, 27, 3455-3465. | 6.5 | 30 |
| 3638 | Yeast Transcriptional Regulator Leu3p. <i>Journal of Biological Chemistry</i> , 1999, 274, 19017-19024. | 1.6 | 38 |
| 3639 | Interaction of Translation Initiation Factor eIF4G with eIF4A in the Yeast <i>Saccharomyces cerevisiae</i> . <i>Journal of Biological Chemistry</i> , 1999, 274, 26720-26726. | 1.6 | 49 |
| 3640 | Pleiotropic Alterations in Lipid Metabolism in Yeast <i>sac1</i> Mutants: Relationship to <i>Bye</i> Sec14p and Inositol Auxotrophy. <i>Molecular Biology of the Cell</i> , 1999, 10, 2235-2250. | 0.9 | 138 |
| 3641 | Adaptor Complex-independent Clathrin Function in Yeast. <i>Molecular Biology of the Cell</i> , 1999, 10, 3643-3659. | 0.9 | 151 |
| 3642 | A Highly Conserved Signal Controls Degradation of 3-Hydroxy-3-methylglutaryl-coenzyme A (HMG-CoA) Reductase in Eukaryotes. <i>Journal of Biological Chemistry</i> , 1999, 274, 31671-31678. | 1.6 | 128 |
| 3643 | Construction of an Effective Host-Vector System for the Yeast <i>Saccharomyces exiguus</i> Yp74L-3. <i>Bioscience, Biotechnology and Biochemistry</i> , 1999, 63, 847-850. | 0.6 | 5 |
| 3644 | Human Calcitonin Tetrameric Gene: Comparative Expression in Yeast and Transgenic Potato Plants. <i>Biotechnology and Biotechnological Equipment</i> , 1999, 13, 20-24. | 0.5 | 2 |
| 3645 | Functional Analysis of the Zn ²⁺ -Cys ⁶ Transcription Factors Oaf1p and Pip2p. <i>Journal of Biological Chemistry</i> , 1999, 274, 22208-22216. | 1.6 | 64 |
| 3646 | Identification of SSF1, CNS1, and HCH1 as multicopy suppressors of a <i>Saccharomyces cerevisiae</i> Hsp90 loss-of-function mutation. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 1999, 96, 1409-1414. | 3.3 | 91 |
| 3647 | The polo-box-dependent induction of ectopic septal structures by a mammalian polo kinase, Plk, in <i>Saccharomyces cerevisiae</i> . <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 1999, 96, 14360-14365. | 3.3 | 42 |
| 3648 | Insect Immunity. <i>Journal of Biological Chemistry</i> , 1999, 274, 9320-9326. | 1.6 | 148 |
| 3649 | Cloning of Murine Translation Initiation Factor 6 and Functional Analysis of the Homologous Sequence YPR016c in <i>Saccharomyces cerevisiae</i> . <i>Journal of Biological Chemistry</i> , 1999, 274, 11653-11659. | 1.6 | 33 |
| 3650 | Identification of a Novel Region Critical for Calcineurin Function in Vivo and in Vitro. <i>Journal of Biological Chemistry</i> , 1999, 274, 18543-18551. | 1.6 | 23 |
| 3651 | Differential transmission of G1 cell cycle arrest and mating signals by <i>Saccharomyces cerevisiae</i> Ste5 mutants in the pheromone pathway. <i>Biochemistry and Cell Biology</i> , 1999, 77, 459-468. | 0.9 | 1 |
| 3652 | Participation of Bir1p, a member of the inhibitor of apoptosis family, in yeast chromosome segregation events. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 1999, 96, 13208-13213. | 3.3 | 113 |
| 3653 | Hepatitis C Virus Core Protein Binds to a DEAD Box RNA Helicase. <i>Journal of Biological Chemistry</i> , 1999, 274, 15751-15756. | 1.6 | 147 |
| 3654 | Post-translational Regulation of Adr1 Activity Is Mediated by Its DNA Binding Domain. <i>Journal of Biological Chemistry</i> , 1999, 274, 37575-37582. | 1.6 | 29 |

| # | ARTICLE | IF | CITATIONS |
|------|--|-----|-----------|
| 3655 | Leptomycin B inactivates CRM1/exportin 1 by covalent modification at a cysteine residue in the central conserved region. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 1999, 96, 9112-9117. | 3.3 | 953 |
| 3656 | In Vivo Characterization of a Thioredoxin h Target Protein Defines a New Peroxiredoxin Family. <i>Journal of Biological Chemistry</i> , 1999, 274, 19714-19722. | 1.6 | 213 |
| 3657 | Characterization of DP103, a Novel DEAD Box Protein That Binds to the Epstein-Barr Virus Nuclear Proteins EBNA2 and EBNA3C. <i>Journal of Biological Chemistry</i> , 1999, 274, 19136-19144. | 1.6 | 93 |
| 3658 | Overexpression of Arabidopsis Hexokinase in Tomato Plants Inhibits Growth, Reduces Photosynthesis, and Induces Rapid Senescence. <i>Plant Cell</i> , 1999, 11, 1253-1266. | 3.1 | 339 |
| 3659 | Functional Characterization of Human Methylenetetrahydrofolate Reductase in <i>Saccharomyces cerevisiae</i> . <i>Journal of Biological Chemistry</i> , 1999, 274, 32613-32618. | 1.6 | 41 |
| 3660 | Yeast flavin-containing monooxygenase generates oxidizing equivalents that control protein folding in the endoplasmic reticulum. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 1999, 96, 2687-2691. | 3.3 | 79 |
| 3661 | Starvation Induces Vacuolar Targeting and Degradation of the Tryptophan Permease in Yeast. <i>Journal of Cell Biology</i> , 1999, 146, 1227-1238. | 2.3 | 279 |
| 3662 | Yeast SMF1 Mediates H ⁺ -coupled Iron Uptake with Concomitant Uncoupled Cation Currents. <i>Journal of Biological Chemistry</i> , 1999, 274, 35089-35094. | 1.6 | 137 |
| 3663 | High-level secretory production of recombinant human lipocortin-I by <i>Saccharomyces cerevisiae</i> . <i>Process Biochemistry</i> , 1999, 35, 97-101. | 1.8 | 8 |
| 3664 | Overproduction of human parathyroid hormone by fed-batch culture of a <i>Saccharomyces cerevisiae</i> mutant lacking yeast aspartic protease 3. <i>Process Biochemistry</i> , 1999, 35, 503-508. | 1.8 | 12 |
| 3665 | Î-Integration of endo/exo-glucanase and Î ² -glucosidase genes into the yeast chromosomes for direct conversion of cellulose to ethanol. <i>Enzyme and Microbial Technology</i> , 1999, 25, 23-30. | 1.6 | 95 |
| 3666 | Identification and characterization of two missense mutations causing factor XIIIa deficiency. <i>British Journal of Haematology</i> , 1999, 104, 37-43. | 1.2 | 18 |
| 3667 | A male gametophyte-specific monosaccharide transporter in <i>Arabidopsis</i> . <i>Plant Journal</i> , 1999, 17, 191-201. | 2.8 | 143 |
| 3668 | Cloning and characterisation of PGA1 and PGA2: two G protein alpha-subunits from pea that promote growth in the yeast <i>Saccharomyces cerevisiae</i> . <i>Plant Journal</i> , 1999, 19, 237-247. | 2.8 | 23 |
| 3669 | Isolation and characterization of a flavonoid 3'-hydroxylase cDNA clone corresponding to the Ht1 locus of <i>Petunia hybrida</i> . <i>Plant Journal</i> , 1999, 19, 441-451. | 2.8 | 194 |
| 3670 | <i>Arabidopsis thaliana</i> AtHAL3: a flavoprotein related to salt and osmotic tolerance and plant growth. <i>Plant Journal</i> , 1999, 20, 529-539. | 2.8 | 104 |
| 3671 | Association between the endoplasmic reticulum and mitochondria of yeast facilitates interorganelle transport of phospholipids through membrane contact. <i>FEBS Journal</i> , 1999, 264, 545-553. | 0.2 | 260 |
| 3672 | Identification of Delta5-fatty acid desaturase from the cellular slime mold <i>Dictyostelium discoideum</i> . <i>FEBS Journal</i> , 1999, 265, 809-814. | 0.2 | 18 |

| # | ARTICLE | IF | CITATIONS |
|------|--|-----|-----------|
| 3673 | The presence of the alternatively spliced A2 cassette in the vacuolar H ⁺ -ATPase subunit A prevents assembly of the V1 catalytic domain. <i>FEBS Journal</i> , 1999, 266, 293-301. | 0.2 | 4 |
| 3674 | Physiological effects and adjuvanticity of recombinant brushtail possum TNF- $\hat{\pm}$. <i>Immunology and Cell Biology</i> , 1999, 77, 28-33. | 1.0 | 20 |
| 3675 | High copy number suppression of the meiotic arrest caused by <i>adm1</i> mutation: <i>REC114</i> imposes an early recombination block and <i>RAD54</i> promotes a <i>DMC1</i> -independent DSB repair pathway. <i>Genes To Cells</i> , 1999, 4, 425-444. | 0.5 | 89 |
| 3676 | Fission yeast APC/cyclosome subunits, <i>Cut20/Apc4</i> and <i>Cut23/Apc8</i> , in regulating metaphase-anaphase progression and cellular stress responses. <i>Genes To Cells</i> , 1999, 4, 445-463. | 0.5 | 37 |
| 3677 | <i>MCM21</i> and <i>MCM22</i> , two novel genes of the yeast <i>Saccharomyces cerevisiae</i> are required for chromosome transmission. <i>Molecular Microbiology</i> , 1999, 31, 349-360. | 1.2 | 52 |
| 3678 | A co-activator of nitrogen-regulated transcription in <i>Saccharomyces cerevisiae</i> . <i>Molecular Microbiology</i> , 1999, 31, 753-762. | 1.2 | 17 |
| 3679 | The WD protein <i>Cpc2p</i> is required for repression of <i>Gcn4</i> protein activity in yeast in the absence of amino-acid starvation. <i>Molecular Microbiology</i> , 1999, 31, 807-822. | 1.2 | 43 |
| 3680 | Multiple independent regulatory pathways control <i>UBI4</i> expression after heat shock in <i>Saccharomyces cerevisiae</i> . <i>Molecular Microbiology</i> , 1999, 31, 823-832. | 1.2 | 44 |
| 3681 | <i>Fps1p</i> controls the accumulation and release of the compatible solute glycerol in yeast osmoregulation. <i>Molecular Microbiology</i> , 1999, 31, 1087-1104. | 1.2 | 357 |
| 3682 | Genome-wide analysis of gene expression regulated by the yeast cell wall integrity signalling pathway. <i>Molecular Microbiology</i> , 1999, 34, 1049-1057. | 1.2 | 384 |
| 3683 | Genetic screens in yeast to identify mammalian nonreceptor modulators of G-protein signaling. <i>Nature Biotechnology</i> , 1999, 17, 878-883. | 9.4 | 196 |
| 3684 | Formation of AP-3 transport intermediates requires <i>Vps41</i> function. <i>Nature Cell Biology</i> , 1999, 1, 346-353. | 4.6 | 122 |
| 3685 | Involvement of mitochondrial aldehyde dehydrogenase <i>ALD5</i> in maintenance of the mitochondrial electron transport chain in <i>Saccharomyces cerevisiae</i> . <i>FEMS Microbiology Letters</i> , 1999, 181, 281-287. | 0.7 | 34 |
| 3686 | Expression and secretion of <i>Bacillus polymyxa</i> <i>neopullulanase</i> in <i>Saccharomyces cerevisiae</i> . <i>FEMS Microbiology Letters</i> , 1999, 170, 41-49. | 0.7 | 17 |
| 3687 | Subunit f of the yeast mitochondrial ATP synthase: topological and functional studies. <i>Journal of Bioenergetics and Biomembranes</i> , 1999, 31, 85-94. | 1.0 | 15 |
| 3688 | Title is missing!. <i>Biotechnology Letters</i> , 1999, 21, 225-229. | 1.1 | 5 |
| 3689 | Expression of the <i>INU2</i> gene for an endoinulinase of <i>Aspergillus ficuum</i> in <i>Saccharomyces cerevisiae</i> . <i>Biotechnology Letters</i> , 1999, 21, 621-623. | 1.1 | 17 |
| 3690 | A new transformation system of <i>Saccharomyces cerevisiae</i> with blasticidin S deaminase gene. <i>Biotechnology Letters</i> , 1999, 21, 969-971. | 1.1 | 15 |

| # | ARTICLE | IF | CITATIONS |
|------|--|-----|-----------|
| 3691 | BTF3 is a potential new substrate of protein kinase CK2. , 1999, 191, 121-128. | | 15 |
| 3692 | Title is missing!. Biotechnology Letters, 1999, 13, 605-608. | 0.5 | 3 |
| 3693 | Nuclear import of RPA in <i>Xenopus</i> egg extracts requires a novel protein XRIPalpha but not importin alpha. EMBO Journal, 1999, 18, 4348-4358. | 3.5 | 65 |
| 3694 | Elements essential for accumulation and function of small nucleolar RNAs directing site-specific pseudouridylation of ribosomal RNAs. EMBO Journal, 1999, 18, 457-469. | 3.5 | 106 |
| 3695 | The yeast non-Mendelian factor [ETA+] is a variant of [PSI+], a prion-like form of release factor eRF3. EMBO Journal, 1999, 18, 1182-1191. | 3.5 | 113 |
| 3696 | L-Asparaginase II of <i>Saccharomyces cerevisiae</i> : Activity Profile During Growth Using an ure2 Mutant P40-3C and a P40-3C+URE2 p Strain. Applied Biochemistry and Biotechnology, 1999, 77, 311-316. | 1.4 | 11 |
| 3697 | Effect of gene disruptions of the TCA cycle on production of succinic acid in <i>Saccharomyces cerevisiae</i> . Journal of Bioscience and Bioengineering, 1999, 87, 28-36. | 1.1 | 95 |
| 3698 | N-glycosylation is involved in the sensitivity of <i>Saccharomyces cerevisiae</i> to HM-1 killer toxin secreted from <i>Hansenula mrakii</i> IFO 0895. Applied Microbiology and Biotechnology, 1999, 51, 176-184. | 1.7 | 20 |
| 3699 | Construction of a flocculent <i>Saccharomyces cerevisiae</i> fermenting lactose. Applied Microbiology and Biotechnology, 1999, 51, 621-626. | 1.7 | 39 |
| 3700 | Independent production of two molecular forms of a recombinant <i>Rhizopus oryzae</i> lipase by KEX2-engineered strains of <i>Saccharomyces cerevisiae</i> . Applied Microbiology and Biotechnology, 1999, 52, 534-540. | 1.7 | 29 |
| 3701 | The Autonomously Replicating Sequence (ARS) of the Yeast <i>Saccharomyces exiguus</i> Yp74L-3. Current Microbiology, 1999, 38, 122-125. | 1.0 | 5 |
| 3702 | News & Notes: Stochastic Nucleosome Positioning in a Yeast Chromatin Region Is Not Dependent on Histone H1. Current Microbiology, 1999, 39, 168-172. | 1.0 | 16 |
| 3703 | The influence of mutation rad57-1 on the fidelity of DNA double-strand gap repair in <i>Saccharomyces cerevisiae</i> . Current Genetics, 1999, 34, 430-437. | 0.8 | 3 |
| 3704 | Transformation of <i>Candida oleophila</i> and survival of a transformant on orange fruit under field conditions. Current Genetics, 1999, 35, 51-57. | 0.8 | 17 |
| 3705 | Hyperosmotic stress represses the transcription of HXT2 and HXT4 genes in <i>Saccharomyces cerevisiae</i> . Folia Microbiologica, 1999, 44, 372-376. | 1.1 | 9 |
| 3706 | The yeast pheromone-responsive G1 \pm protein stimulates recovery from chronic pheromone treatment by two mechanisms that are activated at distinct levels of stimulus. Cell Biochemistry and Biophysics, 1999, 30, 193-212. | 0.9 | 12 |
| 3707 | Reconstitution of fibroblast growth factor receptor interactions in the yeast two hybrid system. Molecular Biotechnology, 1999, 11, 213-220. | 1.3 | 3 |
| 3708 | Functional implications of genetic interactions between genes encoding small GTPases involved in vesicular transport in yeast. Molecular Genetics and Genomics, 1999, 261, 80-91. | 2.4 | 10 |

| # | ARTICLE | IF | CITATIONS |
|------|---|-----|-----------|
| 3709 | YAP1 confers resistance to the fatty acid synthase inhibitor cerulenin through the transporter Flr1p in <i>Saccharomyces cerevisiae</i> . <i>Molecular Genetics and Genomics</i> , 1999, 261, 346-353. | 2.4 | 25 |
| 3710 | Dbf2 is implicated in a Cbt1-dependent pathway following a shift from glucose to galactose or non-fermentable carbon sources in <i>Saccharomyces cerevisiae</i> . <i>Molecular Genetics and Genomics</i> , 1999, 261, 402-407. | 2.4 | 5 |
| 3711 | Genetic evidence for interaction between components of the yeast 26S proteasome: combination of a mutation in RPN12 (a lid component gene) with mutations in RPT1 (an ATPase gene) causes synthetic lethality. <i>Molecular Genetics and Genomics</i> , 1999, 262, 145-153. | 2.4 | 8 |
| 3712 | Regulation of the <i>Saccharomyces cerevisiae</i> DLD1 gene encoding the mitochondrial protein D-lactate ferricytochrome c oxidoreductase by HAP1 and HAP2/3/4/5. <i>Molecular Genetics and Genomics</i> , 1999, 262, 623-632. | 2.4 | 30 |
| 3713 | Biological projectiles (phage, yeast, bacteria) for genetic transformation of plants. <i>In Vitro Cellular and Developmental Biology - Plant</i> , 1999, 35, 43-50. | 0.9 | 11 |
| 3714 | Cloning and characterization of a gene encoding phosphatidyl inositol-specific phospholipase C from <i>Trypanosoma cruzi</i> . <i>Molecular and Biochemical Parasitology</i> , 1999, 102, 283-295. | 0.5 | 16 |
| 3715 | Transcriptional regulation of the squalene synthase gene (ERG9) in the yeast <i>Saccharomyces cerevisiae</i> . <i>Biochimica Et Biophysica Acta Gene Regulatory Mechanisms</i> , 1999, 1445, 110-122. | 2.4 | 74 |
| 3716 | Isolation of a novel gene, <i>moc2</i> , encoding a putative RNA helicase as a suppressor of sterile strains in <i>Schizosaccharomyces pombe</i> . <i>Biochimica Et Biophysica Acta Gene Regulatory Mechanisms</i> , 1999, 1446, 93-101. | 2.4 | 23 |
| 3717 | A second-site mutation at glutamate-257 that restores the function of the mutant yeast ribosomal protein L5 containing lysine-270,271→arginine. <i>Biochimica Et Biophysica Acta Gene Regulatory Mechanisms</i> , 1999, 1489, 223-232. | 2.4 | 0 |
| 3718 | REV3 is required for spontaneous but not methylation damage-induced mutagenesis of <i>Saccharomyces cerevisiae</i> cells lacking O6-methylguanine DNA methyltransferase. <i>Mutation Research - Fundamental and Molecular Mechanisms of Mutagenesis</i> , 1999, 431, 155-165. | 0.4 | 9 |
| 3719 | Unfolded protein response-induced BiP/Kar2p production protects cell growth against accumulation of misfolded protein aggregates in the yeast endoplasmic reticulum. <i>European Journal of Cell Biology</i> , 1999, 78, 726-738. | 1.6 | 19 |
| 3720 | The processing of α 1,2-mannosidase of <i>Saccharomyces cerevisiae</i> depends on Rer1p for its localization in the endoplasmic reticulum. <i>European Journal of Cell Biology</i> , 1999, 78, 435-440. | 1.6 | 20 |
| 3721 | The <i>Saccharomyces cerevisiae</i> Cdc14 phosphatase is implicated in the structural organization of the nucleolus. <i>Biology of the Cell</i> , 1999, 91, 649-663. | 0.7 | 3 |
| 3722 | Effect of NAD ⁺ -dependent isocitrate dehydrogenase gene (IDH1, IDH2) disruption of sake yeast on organic acid composition in sake mash. <i>Journal of Bioscience and Bioengineering</i> , 1999, 88, 258-263. | 1.1 | 32 |
| 3723 | Mass mating method in combination with G418- and aureobasidin A-resistance markers for efficient selection of hybrids from homothallic strains in <i>Saccharomyces cerevisiae</i> . <i>Journal of Bioscience and Bioengineering</i> , 1999, 88, 468-471. | 1.1 | 7 |
| 3724 | DNA sequence of the β -isopropylmalate dehydrogenase gene and phylogenetic analysis of the yeast <i>Saccharomyces exiguus</i> Yp74L-3. <i>Journal of Bioscience and Bioengineering</i> , 1999, 87, 15-18. | 1.1 | 3 |
| 3725 | Construction of recombinant sake yeast containing a dominant FAS2 mutation without extraneous sequences by a two-step gene replacement protocol. <i>Journal of Bioscience and Bioengineering</i> , 1999, 87, 43-48. | 1.1 | 50 |
| 3726 | Characterization of a PDR1 mutant allele from a clotrimazole-resistant sake yeast mutant with improved fermentative activity. <i>Journal of Bioscience and Bioengineering</i> , 1999, 88, 20-25. | 1.1 | 12 |

| # | ARTICLE | IF | CITATIONS |
|------|--|-----|-----------|
| 3727 | In <i>Saccharomyces cerevisiae</i> , feedback inhibition of homocitrate synthase isoenzymes by lysine modulates the activation of LYS gene expression by Lys14p. <i>FEBS Journal</i> , 1999, 261, 163-170. | 0.2 | 74 |
| 3728 | Identification of Δ^{12} -fatty acid desaturase from arachidonic acid-producing <i>Mortierella</i> fungus by heterologous expression in the yeast <i>Saccharomyces cerevisiae</i> and the fungus <i>Aspergillus oryzae</i> . <i>FEBS Journal</i> , 1999, 261, 812-820. | 0.2 | 112 |
| 3729 | A role for the immunoglobulin-like domain of the human IL-6 receptor. Intracellular protein transport and shedding. <i>FEBS Journal</i> , 1999, 263, 438-446. | 0.2 | 47 |
| 3730 | Carbon source-dependent regulation of cell growth by murine protein kinase C epsilon expression in <i>Saccharomyces cerevisiae</i> . , 1999, 178, 216-226. | | 7 |
| 3731 | Genetically controlled cell lysis in the yeast <i>Saccharomyces cerevisiae</i> . , 1999, 64, 607-615. | | 19 |
| 3732 | A positive selection for plasmid loss in <i>Saccharomyces cerevisiae</i> using galactose-inducible growth inhibitory sequences. , 1999, 15, 1-10. | | 21 |
| 3733 | Disruption of six novel <i>Saccharomyces cerevisiae</i> genes reveals that YGL129c is necessary for growth in non-fermentable carbon sources, YGL128c for growth at low or high temperatures and YGL125w is implicated in the biosynthesis of methionine. , 1999, 15, 145-154. | | 24 |
| 3734 | Cloning and characterization of 1,3- β -glucanase-encoding genes from non-conventional yeasts. <i>Yeast</i> , 1999, 15, 91-109. | 0.8 | 21 |
| 3735 | Isolation and sequence of the GCR3 homologue from <i>Candida albicans</i> by complementation of Δ^{1} gcr3 strain of <i>Saccharomyces cerevisiae</i> . <i>Yeast</i> , 1999, 15, 323-327. | 0.8 | 0 |
| 3736 | Basic phenotypic analysis of six novel yeast genes reveals two essential genes and one which affects the growth rate. , 1999, 15, 351-360. | | 7 |
| 3737 | A single nucleotide change in the MNR1 (VCX1/HUM1) gene determines resistance to manganese in <i>Saccharomyces cerevisiae</i> . <i>Yeast</i> , 1999, 15, 371-375. | 0.8 | 19 |
| 3738 | The open reading frame YAL048c affects the secretion of proteinase A in <i>S. cerevisiae</i> . <i>Yeast</i> , 1999, 15, 427-434. | 0.8 | 12 |
| 3739 | Two-Dimensional analysis of proteins secreted by <i>Saccharomyces cerevisiae</i> regenerating protoplasts: a novel approach to study the cell wall. , 1999, 15, 459-472. | | 82 |
| 3740 | In vivo construction of cDNA libraries for use in the yeast two-hybrid system. , 1999, 15, 715-720. | | 20 |
| 3741 | A proposal for nomenclature of aldehyde dehydrogenases in <i>Saccharomyces cerevisiae</i> and characterization of the stress-inducible ALD2 and ALD3 genes. , 1999, 15, 829-842. | | 127 |
| 3742 | Three genes whose expression is induced by stress in <i>Saccharomyces cerevisiae</i> . <i>Yeast</i> , 1999, 15, 879-892. | 0.8 | 136 |
| 3743 | Transcription of the HXT4 gene is regulated by Gcr1p and Gcr2p in the yeast <i>S. cerevisiae</i> . , 1999, 15, 1045-1057. | | 14 |
| 3744 | Construction of a lactose-assimilating strain of baker's yeast. <i>Yeast</i> , 1999, 15, 1299-1305. | 0.8 | 21 |

| # | ARTICLE | IF | CITATIONS |
|------|---|-----|-----------|
| 3745 | ade9 is an allele ofSER1 and plays an indirect role in purine biosynthesis. Yeast, 1999, 15, 1347-1355. | 0.8 | 4 |
| 3746 | Mouse CAF1, a mouse homologue of the yeastPOP2 gene, complements the yeastpop2 null mutation. Yeast, 1999, 15, 1357-1364. | 0.8 | 10 |
| 3747 | Cysteine biosynthesis inSaccharomyces cerevisiae : a new outlook on pathway and regulation. Yeast, 1999, 15, 1365-1375. | 0.8 | 68 |
| 3748 | Molecular cloning and characterization of the geneH XK1 encoding the hexokinase fromYarrowia lipolytica. Yeast, 1999, 15, 1573-1584. | 0.8 | 35 |
| 3749 | Transformation ofCandida albicans by electroporation. Yeast, 1999, 15, 1609-1618. | 0.8 | 75 |
| 3750 | Cloning and characterization of theEXG1 gene from the yeastYarrowia lipolytica. Yeast, 1999, 15, 1631-1644. | 0.8 | 17 |
| 3751 | Selection of genes repressed by cAMP that are induced by nutritional limitation inSaccharomyces cerevisiae. Yeast, 1999, 15, 1733-1745. | 0.8 | 30 |
| 3752 | Transformation of a flocculatingSaccharomyces cerevisiae using lithium acetate and pYAC4. Journal of Basic Microbiology, 1999, 39, 37-41. | 1.8 | 6 |
| 3753 | Expression of a mammalian Na ⁺ /H ⁺ antiporter in Saccharomyces cerevisiae. Biochemistry and Cell Biology, 1999, 77, 25-31. | 0.9 | 10 |
| 3754 | Cell extensions in <i>pkc1</i> mutants of <i>Saccharomyces cerevisiae</i> . Canadian Journal of Microbiology, 1999, 45, 38-44. | 0.8 | 2 |
| 3755 | Functional Analysis of the Promoter of the YeastSNQ2Gene Encoding a Multidrug Resistance Transporter that Confers the Resistance to 4-NitroquinolineN-Oxide. Bioscience, Biotechnology and Biochemistry, 1999, 63, 162-167. | 0.6 | 15 |
| 3756 | Elements regulating the transcription of human interstitial retinoid-binding protein (IRBP) gene in cultured retinoblastoma cells. Current Eye Research, 1999, 18, 283-291. | 0.7 | 36 |
| 3757 | Involvement of Thioredoxin Peroxidase Type II (Ahp1p) ofSaccharomyces cerevisiaein Mn ²⁺ Homeostasis. Bioscience, Biotechnology and Biochemistry, 1999, 63, 1871-1881. | 0.6 | 15 |
| 3758 | Role of Carbohydrate Moiety in Carboxypeptidase Y: Structural Study of Mutant Enzyme Lacking Carbohydrate Moiety. Bioscience, Biotechnology and Biochemistry, 1999, 63, 1045-1050. | 0.6 | 12 |
| 3759 | Generation of the Amyloid- β Peptide N Terminus inSaccharomyces cerevisiae Expressing Human Alzheimer's Amyloid- β Precursor Protein. Journal of Biological Chemistry, 1999, 274, 33843-33846. | 1.6 | 18 |
| 3760 | Cloning and functional expression of a cDNA encoding a metabolic acyl-CoA Δ^9 -desaturase of the cabbage looper moth, <i>Trichoplusia ni</i> . Insect Biochemistry and Molecular Biology, 1999, 29, 435-443. | 1.2 | 54 |
| 3761 | Identification of the essential EPE1 gene involved in retention of secreted proteins on the cell surface of Saccharomyces cerevisiae cells. International Journal of Biochemistry and Cell Biology, 1999, 31, 903-914. | 1.2 | 4 |
| 3762 | SGT1 Encodes an Essential Component of the Yeast Kinetochore Assembly Pathway and a Novel Subunit of the SCF Ubiquitin Ligase Complex. Molecular Cell, 1999, 4, 21-33. | 4.5 | 357 |

| # | ARTICLE | IF | CITATIONS |
|------|--|------|-----------|
| 3763 | Near-Cognate Peptidyl-tRNAs Promote +1 Programmed Translational Frameshifting in Yeast. <i>Molecular Cell</i> , 1999, 4, 1005-1015. | 4.5 | 69 |
| 3764 | Yeast Sec14p Deficient in Phosphatidylinositol Transfer Activity Is Functional In Vivo. <i>Molecular Cell</i> , 1999, 4, 187-197. | 4.5 | 131 |
| 3765 | The Pachytene Checkpoint in <i>S. cerevisiae</i> Depends on Swe1-Mediated Phosphorylation of the Cyclin-Dependent Kinase Cdc28. <i>Molecular Cell</i> , 1999, 4, 805-814. | 4.5 | 80 |
| 3766 | Genetic interactions between error-prone and error-free postreplication repair pathways in <i>Saccharomyces cerevisiae</i> . <i>Mutation Research DNA Repair</i> , 1999, 435, 1-11. | 3.8 | 72 |
| 3767 | Identification of a gene, SOO1, which complements osmo-sensitivity and defect in in vitro β -1,3-glucan synthase activity in <i>Saccharomyces cerevisiae</i> . <i>Biochimica Et Biophysica Acta - Molecular Cell Research</i> , 1999, 1450, 145-154. | 1.9 | 13 |
| 3768 | Studies on the structures and antigenic properties of rabies virus glycoprotein analogues produced in yeast cells. <i>Vaccine</i> , 1999, 17, 205-218. | 1.7 | 23 |
| 3769 | Two distinct regions of the BPV1 E1 replication protein interact with the activation domain of E2. <i>Virus Research</i> , 1999, 65, 141-154. | 1.1 | 10 |
| 3770 | Introduction of the hiC6 Gene, which Encodes a Homologue of a Late Embryogenesis Abundant (LEA) Protein, Enhances Freezing Tolerance of Yeast. <i>Journal of Plant Physiology</i> , 1999, 155, 509-512. | 1.6 | 39 |
| 3771 | Glucose Signaling in Yeast Is Partially Mimicked by Galactose and Does Not Require the Tps1 Protein. <i>Molecular Cell Biology Research Communications: MCBRC: Part B of Biochemical and Biophysical Research Communications</i> , 1999, 1, 52-58. | 1.7 | 13 |
| 3772 | cAMP Is Involved in Transcriptional Regulation of Δ^9 -Desaturase during <i>Histoplasma capsulatum</i> Morphogenesis. <i>Molecular Cell Biology Research Communications: MCBRC: Part B of Biochemical and Biophysical Research Communications</i> , 1999, 2, 172-177. | 1.7 | 8 |
| 3773 | High-level expression of uniformly ^{15}N -labeled hen lysozyme in <i>Pichia pastoris</i> and identification of the site in hen lysozyme where phosphate ion binds using NMR measurements. <i>FEBS Letters</i> , 1999, 448, 33-37. | 1.3 | 48 |
| 3774 | Rpn4p acts as a transcription factor by binding to PACE, a nonamer box found upstream of 26S proteasomal and other genes in yeast. <i>FEBS Letters</i> , 1999, 450, 27-34. | 1.3 | 301 |
| 3775 | Enlargement of the endoplasmic reticulum membrane in <i>Saccharomyces cerevisiae</i> is not necessarily linked to the unfolded protein response via Ire1p. <i>FEBS Letters</i> , 1999, 453, 210-214. | 1.3 | 24 |
| 3776 | Induction of vacuolar Ca^{2+} -ATPase and $\text{H}^{+}/\text{Ca}^{2+}$ -exchange activity in yeast mutants lacking Pmrl, the Golgi Ca^{2+} -ATPase. <i>FEBS Letters</i> , 1999, 454, 181-186. | 1.3 | 60 |
| 3777 | Dominant-negative alleles of 14-3-3 proteins cause defects in actin organization and vesicle targeting in the yeast <i>Saccharomyces cerevisiae</i> . <i>FEBS Letters</i> , 1999, 460, 411-416. | 1.3 | 54 |
| 3778 | The E-box DNA binding protein Sgc1p suppresses the gcr2 mutation, which is involved in transcriptional activation of glycolytic genes in <i>Saccharomyces cerevisiae</i> . <i>FEBS Letters</i> , 1999, 463, 307-311. | 1.3 | 33 |
| 3779 | Monitoring of estrogen mimics by a recombinant yeast assay: synergy between natural and synthetic compounds?. <i>Science of the Total Environment</i> , 1999, 225, 69-79. | 3.9 | 53 |
| 3780 | RESPONSIVE-TO-ANTAGONIST1, a Menkes/Wilson Disease-Related Copper Transporter, Is Required for Ethylene Signaling in <i>Arabidopsis</i> . <i>Cell</i> , 1999, 97, 383-393. | 13.5 | 385 |

| # | ARTICLE | IF | CITATIONS |
|------|---|-----|-----------|
| 3781 | Aromatic amino acids in the Rieske iron-sulfur protein do not form an obligatory conduit for electron transfer from the iron-sulfur cluster to the heme of cytochrome c1 in the cytochrome bc1 complex. <i>Biochimica Et Biophysica Acta - Bioenergetics</i> , 1999, 1410, 237-247. | 0.5 | 5 |
| 3782 | Characterization of non-dominant lethal mutations in the yeast plasma membrane H ⁺ -ATPase gene. <i>Biochimica Et Biophysica Acta - Biomembranes</i> , 1999, 1417, 32-36. | 1.4 | 2 |
| 3783 | [29] Functional analyses of aquaporin water channel proteins. <i>Methods in Enzymology</i> , 1999, 294, 550-572. | 0.4 | 42 |
| 3784 | A 'distributed degron' allows regulated entry into the ER degradation pathway. <i>EMBO Journal</i> , 1999, 18, 5994-6004. | 3.5 | 81 |
| 3785 | Expression of the 180-kD Ribosome Receptor Induces Membrane Proliferation and Increased Secretory Activity in Yeast. <i>Journal of Cell Biology</i> , 1999, 146, 273-284. | 2.3 | 49 |
| 3786 | Novel Protein Kinases Associated with Calcineurin-like Calcium Sensors in Arabidopsis. <i>Plant Cell</i> , 1999, 11, 2393-2405. | 3.1 | 288 |
| 3787 | Mutations in a GTP-binding Motif of Eukaryotic Elongation Factor 1A Reduce Both Translational Fidelity and the Requirement for Nucleotide Exchange. <i>Journal of Biological Chemistry</i> , 1999, 274, 30297-30302. | 1.6 | 76 |
| 3788 | Integrity of Human YACs during Propagation in Recombination-Deficient Yeast Strains. <i>Genomics</i> , 1999, 56, 262-273. | 1.3 | 14 |
| 3789 | Identification of the INO1 gene of Mycobacterium tuberculosis H37Rv reveals a novel class of inositol-1-phosphate synthase enzyme 1 Edited by M. Yaniv. <i>Journal of Molecular Biology</i> , 1999, 291, 531-536. | 2.0 | 63 |
| 3790 | Methods of Assaying Bcl-2 and Bax Family Proteins in Yeast. <i>Methods</i> , 1999, 17, 292-304. | 1.9 | 18 |
| 3791 | Genetic Approaches to the Study of Protein-Protein Interactions. <i>Methods</i> , 1999, 19, 338-349. | 1.9 | 17 |
| 3792 | The Significance of Amino Acid Residue Asp446 for Enzymatic Stability of Rat UDP-Glucuronosyltransferase UGT1A6. <i>Archives of Biochemistry and Biophysics</i> , 1999, 363, 116-120. | 1.4 | 14 |
| 3793 | Cloning of anl-3-Hydroxyacyl-CoA Dehydrogenase That Interacts with the GLUT4 C-Terminus. <i>Archives of Biochemistry and Biophysics</i> , 1999, 363, 323-332. | 1.4 | 16 |
| 3794 | Probing the Affinity and Specificity of Yeast Alcohol Dehydrogenase I for Coenzymes. <i>Archives of Biochemistry and Biophysics</i> , 1999, 367, 240-249. | 1.4 | 25 |
| 3795 | Saccharomyces cerevisiae Expresses Two Genes Encoding Isozymes of Methylene tetrahydrofolate Reductase. <i>Archives of Biochemistry and Biophysics</i> , 1999, 372, 300-308. | 1.4 | 36 |
| 3796 | SHC1, a High pH Inducible Gene Required for Growth at Alkaline pH in Saccharomyces cerevisiae. <i>Biochemical and Biophysical Research Communications</i> , 1999, 255, 116-122. | 1.0 | 12 |
| 3797 | Cloned Prokaryotic Iron Superoxide Dismutase Protects Yeast Cells against Oxidative Stress Depending on Mitochondrial Location. <i>Biochemical and Biophysical Research Communications</i> , 1999, 256, 63-67. | 1.0 | 20 |
| 3798 | Difference in Strength of Autonomously Replicating Sequences among Repeats in the rDNA Region of Saccharomyces cerevisiae. <i>Biochemical and Biophysical Research Communications</i> , 1999, 266, 190-195. | 1.0 | 1 |

| # | ARTICLE | IF | CITATIONS |
|------|--|-----|-----------|
| 3799 | Saccharomyces cerevisiae Cells Harboring the Gene Encoding Sarcotoxin IA Secrete a Peptide That Is Toxic to Plant Pathogenic Bacteria. Protein Expression and Purification, 1999, 16, 120-124. | 0.6 | 19 |
| 3800 | Expression and Purification of the BmK M1 Neurotoxin from the Scorpion Buthus martensii Karsch. Protein Expression and Purification, 1999, 17, 358-365. | 0.6 | 29 |
| 3801 | Las21 participates in extracellular/cell surface phenomena in Saccharomyces cerevisiae.. Genes and Genetic Systems, 1999, 74, 241-256. | 0.2 | 16 |
| 3802 | Functional production and reconstitution of the human equilibrative nucleoside transporter (hENT1) in Saccharomyces cerevisiae. Biochemical Journal, 1999, 339, 21-32. | 1.7 | 97 |
| 3803 | Molecular and enzymic properties of recombinant 1,2-Î±-mannosidase from Aspergillus saitoi overexpressed in Aspergillus oryzae cells. Biochemical Journal, 1999, 339, 589-597. | 1.7 | 37 |
| 3804 | CLN3 expression is sufficient to restore G1-to-S-phase progression in Saccharomyces cerevisiae mutants defective in translation initiation factor eIF4E. Biochemical Journal, 1999, 340, 135-141. | 1.7 | 49 |
| 3805 | Breeding of sake yeast that produce enhanced levels of flavor by transformation with a mutated FAS 2 gene and their utilization in brewing rice flour shochu.. Journal of the Brewing Society of Japan, 1999, 94, 63-71. | 0.1 | 6 |
| 3806 | Transcriptional Induction by Aromatic Amino Acids in <i>Saccharomyces cerevisiae</i> . Molecular and Cellular Biology, 1999, 19, 3360-3371. | 1.1 | 102 |
| 3807 | The CCR4 and CAF1 Proteins of the CCR4-NOT Complex Are Physically and Functionally Separated from NOT2, NOT4, and NOT5. Molecular and Cellular Biology, 1999, 19, 6642-6651. | 1.1 | 138 |
| 3808 | Splicing of the Meiosis-Specific <i>HOP2</i> Transcript Utilizes a Unique 5â€² Splice Site. Molecular and Cellular Biology, 1999, 19, 7933-7943. | 1.1 | 18 |
| 3809 | Nrg1 Is a Transcriptional Repressor for Glucose Repression of <i>STA1</i> Gene Expression in <i>Saccharomyces cerevisiae</i> . Molecular and Cellular Biology, 1999, 19, 2044-2050. | 1.1 | 103 |
| 3810 | Mutational Disruption of Plasma Membrane Trafficking of <i>Saccharomyces cerevisiae</i> Yor1p, a Homologue of Mammalian Multidrug Resistance Protein. Molecular and Cellular Biology, 1999, 19, 2998-3009. | 1.1 | 89 |
| 3811 | Rec8p, a Meiotic Recombination and Sister Chromatid Cohesion Phosphoprotein of the Rad21p Family Conserved from Fission Yeast to Humans. Molecular and Cellular Biology, 1999, 19, 3515-3528. | 1.1 | 235 |
| 3812 | Accumulation of Single-Stranded DNA and Destabilization of Telomeric Repeats in Yeast Mutant Strains Carrying a Deletion of <i>RAD27</i> . Molecular and Cellular Biology, 1999, 19, 4143-4152. | 1.1 | 123 |
| 3813 | GCD14p, a Repressor of <i>GCN4</i> Translation, Cooperates with Gcd10p and Lhp1p in the Maturation of Initiator Methionyl-tRNA in <i>Saccharomyces cerevisiae</i> . Molecular and Cellular Biology, 1999, 19, 4167-4181. | 1.1 | 60 |
| 3814 | Rpn9 Is Required for Efficient Assembly of the Yeast 26S Proteasome. Molecular and Cellular Biology, 1999, 19, 6575-6584. | 1.1 | 63 |
| 3815 | Nucleolar Factors Direct the 2â€²-O-Ribose Methylation and Pseudouridylation of U6 Spliceosomal RNA. Molecular and Cellular Biology, 1999, 19, 6906-6917. | 1.1 | 149 |
| 3816 | A Novel Mechanism of Ion Homeostasis and Salt Tolerance in Yeast: the Hal4 and Hal5 Protein Kinases Modulate the Trk1-Trk2 Potassium Transporter. Molecular and Cellular Biology, 1999, 19, 3328-3337. | 1.1 | 179 |

| # | ARTICLE | IF | CITATIONS |
|------|---|-----|-----------|
| 3817 | Mid2 Is a Putative Sensor for Cell Integrity Signaling in <i>Saccharomyces cerevisiae</i> . <i>Molecular and Cellular Biology</i> , 1999, 19, 3969-3976. | 1.1 | 191 |
| 3818 | A G ¹ Cyclin Is Necessary for Maintenance of Filamentous Growth in <i>Candida albicans</i> . <i>Molecular and Cellular Biology</i> , 1999, 19, 4019-4027. | 1.1 | 101 |
| 3819 | The Yeast Trimeric Guanine Nucleotide-Binding Protein $\hat{\pm}$ Subunit, Gpa2p, Controls the Meiosis-Specific Kinase Ime2p Activity in Response to Nutrients. <i>Molecular and Cellular Biology</i> , 1999, 19, 6110-6119. | 1.1 | 50 |
| 3820 | Antagonistic Interactions between Yeast Chaperones Hsp104 and Hsp70 in Prion Curing. <i>Molecular and Cellular Biology</i> , 1999, 19, 1325-1333. | 1.1 | 258 |
| 3821 | The <i>Saccharomyces cerevisiae</i> CWH8 gene is required for full levels of dolichol-linked oligosaccharides in the endoplasmic reticulum and for efficient N-glycosylation. <i>Glycobiology</i> , 1999, 9, 243-253. | 1.3 | 32 |
| 3822 | Molecular Cloning and Functional Expression of the Human Golgi UDP-N-Acetylglucosamine Transporter. <i>Journal of Biochemistry</i> , 1999, 126, 68-77. | 0.9 | 66 |
| 3823 | Amino Acid Signaling in <i>Saccharomyces cerevisiae</i> : a Permease-Like Sensor of External Amino Acids and F-Box Protein Grr1p Are Required for Transcriptional Induction of the <i>AGP1</i> Gene, Which Encodes a Broad-Specificity Amino Acid Permease. <i>Molecular and Cellular Biology</i> , 1999, 19, 989-1001. | 1.1 | 256 |
| 3824 | The <i>msh2</i> Gene of <i>Schizosaccharomyces pombe</i> Is Involved in Mismatch Repair, Mating-Type Switching, and Meiotic Chromosome Organization. <i>Molecular and Cellular Biology</i> , 1999, 19, 241-250. | 1.1 | 54 |
| 3825 | Erf2, a Novel Gene Product That Affects the Localization and Palmitoylation of Ras2 in <i>Saccharomyces cerevisiae</i> . <i>Molecular and Cellular Biology</i> , 1999, 19, 6775-6787. | 1.1 | 164 |
| 3826 | Mutations in Elongation Factor $\hat{1}^2$, a Guanine Nucleotide Exchange Factor, Enhance Translational Fidelity. <i>Molecular and Cellular Biology</i> , 1999, 19, 5257-5266. | 1.1 | 69 |
| 3827 | Genetic Evidence for Pak1 Autoinhibition and Its Release by Cdc42. <i>Molecular and Cellular Biology</i> , 1999, 19, 602-611. | 1.1 | 92 |
| 3828 | Genetic and Physical Interactions Involving the Yeast Nuclear Cap-Binding Complex. <i>Molecular and Cellular Biology</i> , 1999, 19, 6543-6553. | 1.1 | 78 |
| 3829 | Specificity Determinants of Proteolytic Processing of <i>Aspergillus</i> PacC Transcription Factor Are Remote from the Processing Site, and Processing Occurs in Yeast If pH Signalling Is Bypassed. <i>Molecular and Cellular Biology</i> , 1999, 19, 1390-1400. | 1.1 | 68 |
| 3830 | Receptor Inhibition of Pheromone Signaling Is Mediated by the Ste4p G ¹ Subunit. <i>Molecular and Cellular Biology</i> , 1999, 19, 441-449. | 1.1 | 11 |
| 3831 | Binding of Gal4p and Bicoid to Nucleosomal Sites in Yeast in the Absence of Replication. <i>Molecular and Cellular Biology</i> , 1999, 19, 2977-2985. | 1.1 | 22 |
| 3832 | The Yeast C-Type Cyclin Ctk2p Is Phosphorylated and Rapidly Degraded by the Ubiquitin-Proteasome Pathway. <i>Molecular and Cellular Biology</i> , 1999, 19, 2527-2534. | 1.1 | 14 |
| 3833 | Cell Cycle Regulation of DNA Replication Initiator Factor Dbf4p. <i>Molecular and Cellular Biology</i> , 1999, 19, 4270-4278. | 1.1 | 90 |
| 3834 | Two Prion-Inducing Regions of Ure2p Are Nonoverlapping. <i>Molecular and Cellular Biology</i> , 1999, 19, 4516-4524. | 1.1 | 104 |

| # | ARTICLE | IF | CITATIONS |
|------|---|-----|-----------|
| 3835 | [23] RNA polymerase III-based two-hybrid system. <i>Methods in Enzymology</i> , 1999, 303, 411-422. | 0.4 | 9 |
| 3836 | An improved integration replacement/disruption method for mutagenesis of yeast essential genes.. <i>Genes and Genetic Systems</i> , 2000, 75, 33-39. | 0.2 | 11 |
| 3837 | Human and Mouse Fas (APO-1/CD95) Death Receptor Genes Each Contain a p53-responsive Element That Is Activated by p53 Mutants Unable to Induce Apoptosis. <i>Journal of Biological Chemistry</i> , 2000, 275, 3867-3872. | 1.6 | 104 |
| 3838 | [25] Translational repression assay procedure: A method to study RNA-protein interactions in yeast. <i>Methods in Enzymology</i> , 2000, 318, 374-384. | 0.4 | 4 |
| 3839 | Assays for Studying Bax-Induced Lethality in the Yeast <i>Saccharomyces cerevisiae</i> . <i>Methods in Enzymology</i> , 2000, 322, 283-296. | 0.4 | 13 |
| 3840 | Visualizing protein dynamics in yeast with green fluorescent protein. <i>Methods in Enzymology</i> , 2000, 327, 61-69. | 0.4 | 2 |
| 3841 | TAN-1813, a Novel Ras-Farnesyltransferase Inhibitor Produced by <i>Phoma</i> sp. Taxonomy, Fermentation, Isolation and Biological Activities In Vitro and In Vivo.. <i>Journal of Antibiotics</i> , 2000, 53, 765-778. | 1.0 | 23 |
| 3842 | The catabolic function of the ϵ -amino adipic acid pathway in plants is associated with unidirectional activity of lysine α -oxoglutarate reductase, but not saccharopine dehydrogenase. <i>Biochemical Journal</i> , 2000, 351, 215. | 1.7 | 17 |
| 3843 | Definition of a minimal munc18c domain that interacts with syntaxin 4. <i>Biochemical Journal</i> , 2000, 350, 741-746. | 1.7 | 10 |
| 3844 | Identification of copper ligands in <i>Aspergillus oryzae</i> tyrosinase by site-directed mutagenesis. <i>Biochemical Journal</i> , 2000, 350, 537-545. | 1.7 | 40 |
| 3845 | Two Distinct Nucleosome Alterations Characterize Chromatin Remodeling at the <i>Saccharomyces cerevisiae</i> ADH2 Promoter. <i>Journal of Biological Chemistry</i> , 2000, 275, 7612-7618. | 1.6 | 19 |
| 3846 | G418 Selection and stability of cloned genes integrated at chromosomal ψ sequences of <i>Saccharomyces cerevisiae</i> . <i>Biotechnology and Bioengineering</i> , 2000, 49, 45-51. | 1.7 | 43 |
| 3847 | Highly efficient secretion of heterologous proteins from <i>Saccharomyces cerevisiae</i> using inulinase signal peptides. , 2000, 49, 473-479. | | 19 |
| 3848 | Effect of gene amplification on threonine production by yeast. , 2000, 49, 667-674. | | 13 |
| 3849 | Site-specific integration of heterologous genes in yeast via Ty3 retrotransposition. , 2000, 51, 703-712. | | 2 |
| 3850 | Development of a LAC4 promoter-based gratuitous induction system in <i>Kluyveromyces lactis</i> . , 2000, 67, 408-416. | | 17 |
| 3851 | Evolutionary complementation for polymerase II CTD function. <i>Yeast</i> , 2000, 16, 57-64. | 0.8 | 37 |
| 3852 | Identification of a putative DEAD-box RNA helicase and a zinc-finger protein in <i>Candida albicans</i> by functional complementation of the <i>S. cerevisiae</i> rok1 mutation. <i>Yeast</i> , 2000, 16, 401-409. | 0.8 | 8 |

| # | ARTICLE | IF | CITATIONS |
|------|---|-----|-----------|
| 3853 | YHP1 encodes a new homeoprotein that binds to theIME1 promoter inSaccharomyces cerevisiae. Yeast, 2000, 16, 439-449. | 0.8 | 22 |
| 3854 | Molecular cloning of theCRM1 gene fromCandida albicans. Yeast, 2000, 16, 531-538. | 0.8 | 3 |
| 3855 | Disruption of six novel ORFs fromSaccharomyces cerevisiae chromosome VII and phenotypic analysis of the deletants. Yeast, 2000, 16, 621-630. | 0.8 | 6 |
| 3856 | Secretory production of recombinant human chymase as an active form inPichia pastoris. Yeast, 2000, 16, 315-323. | 0.8 | 8 |
| 3857 | Cloning and sequence analysis of theCandida boidiniiADE2 gene. Yeast, 2000, 16, 953-957. | 0.8 | 4 |
| 3858 | Co-deletion of theMSB3 andMSB4 coding regions affects bipolar budding and perturbs the organization of the actin cytoskeleton. Yeast, 2000, 16, 1015-1023. | 0.8 | 16 |
| 3859 | Identification of aCandida albicans homologue of thePHO85 gene, a negative regulator of the PHO system inSaccharomyces cerevisiae. Yeast, 2000, 16, 1045-1051. | 0.8 | 21 |
| 3860 | Cells of the yeastSaccharomyces cerevisiae are transformable by DNA under non-artificial conditions. Yeast, 2000, 16, 1107-1110. | 0.8 | 40 |
| 3861 | Inheritance of suppressors of the drug sensitivity of aNSR1 deleted yeast strain. Yeast, 2000, 16, 1147-1159. | 0.8 | 11 |
| 3862 | MET15 as a visual selection marker forCandida albicans. Yeast, 2000, 16, 1205-1215. | 0.8 | 29 |
| 3863 | Molecular cloning and characterization of theCandida albicansUBI3 gene coding for a ubiquitin-hybrid protein. Yeast, 2000, 16, 1413-1419. | 0.8 | 15 |
| 3864 | A Modular Set of Prokaryotic and Eukaryotic Expression Vectors. Analytical Biochemistry, 2000, 277, 109-120. | 1.1 | 70 |
| 3865 | Mis3 with a conserved RNA binding motif is essential for ribosome biogenesis and implicated in the start of cell growth and S phase checkpoint. Genes To Cells, 2000, 5, 525-541. | 0.5 | 14 |
| 3866 | Ebp2p, yeast homologue of a human protein that interacts with Epstein-Barr virus Nuclear Antigen 1, is required for pre-rRNA processing and ribosomal subunit assembly. Genes To Cells, 2000, 5, 543-553. | 0.5 | 59 |
| 3867 | The absence of ribonuclease H1 or H2 alters the sensitivity of Saccharomyces cerevisiae to hydroxyurea, caffeine and ethyl methanesulphonate: implications for roles of RNases H in DNA replication and repair. Genes To Cells, 2000, 5, 789-802. | 0.5 | 116 |
| 3868 | Recruitment of the yeast MADS-box proteins, ArgR1 and Mcm1 by the pleiotropic factor ArgR3 is required for their stability. Molecular Microbiology, 2000, 35, 15-31. | 1.2 | 53 |
| 3869 | Proper ascospore maturation requires the chs1+ chitin synthase gene in Schizosaccharomyces pombe. Molecular Microbiology, 2000, 35, 79-89. | 1.2 | 68 |
| 3870 | Mth1 receives the signal given by the glucose sensors Snf3 and Rgt2 in Saccharomyces cerevisiae. Molecular Microbiology, 2000, 35, 161-172. | 1.2 | 103 |

| # | ARTICLE | IF | CITATIONS |
|------|--|------|-----------|
| 3871 | FAP1, a homologue of human transcription factor NF-X1, competes with rapamycin for binding to FKBP12 in yeast. <i>Molecular Microbiology</i> , 2000, 37, 1480-1493. | 1.2 | 20 |
| 3872 | In vivo N-glycosylation of the Mep2 high-affinity ammonium transporter of <i>Saccharomyces cerevisiae</i> reveals an extracytosolic N-terminus. <i>Molecular Microbiology</i> , 2000, 38, 552-564. | 1.2 | 68 |
| 3873 | AtSTP3, a green leaf-specific, low affinity monosaccharide-H ⁺ symporter of <i>Arabidopsis thaliana</i> . <i>Plant, Cell and Environment</i> , 2000, 23, 175-184. | 2.8 | 68 |
| 3874 | Degradation of Aux/IAA proteins is essential for normal auxin signalling. <i>Plant Journal</i> , 2000, 21, 553-562. | 2.8 | 254 |
| 3875 | Cloning of the guanylate kinase homologues AGK-1 and AGK-2 from <i>Arabidopsis thaliana</i> and characterization of AGK-1. <i>FEBS Journal</i> , 2000, 267, 606-615. | 0.2 | 15 |
| 3876 | Contribution of Are1p and Are2p to steryl ester synthesis in the yeast <i>Saccharomyces cerevisiae</i> . <i>FEBS Journal</i> , 2000, 267, 1075-1082. | 0.2 | 158 |
| 3877 | A second functional Δ^5 fatty acid desaturase in the cellular slime mould <i>Dictyostelium discoideum</i> . <i>FEBS Journal</i> , 2000, 267, 1813-1818. | 0.2 | 21 |
| 3878 | Diazaborine resistance in yeast involves the efflux pumps Ycf1p and Flr1p and is enhanced by a gain-of-function allele of gene YAP1. <i>FEBS Journal</i> , 2000, 267, 4809-4816. | 0.2 | 43 |
| 3879 | An interlaboratory comparison of physiological and genetic properties of four <i>Saccharomyces cerevisiae</i> strains. <i>Enzyme and Microbial Technology</i> , 2000, 26, 706-714. | 1.6 | 488 |
| 3880 | Mutations in GAL2 or GAL4 alleviate catabolite repression produced by galactose in <i>Saccharomyces cerevisiae</i> . <i>Enzyme and Microbial Technology</i> , 2000, 26, 748-755. | 1.6 | 8 |
| 3881 | Preparation of high activity whole cell biocatalyst by permeabilization of recombinant flocculent yeast with alcohol. <i>Enzyme and Microbial Technology</i> , 2000, 27, 806-811. | 1.6 | 48 |
| 3882 | Mammalian Tumor Susceptibility Gene 101 (TSG101) and the Yeast Homologue, Vps23p, Both Function in Late Endosomal Trafficking. <i>Traffic</i> , 2000, 1, 248-258. | 1.3 | 371 |
| 3883 | A New Functional Domain of Guanine Nucleotide Dissociation Inhibitor ($\hat{\pm}$ -GDI) Involved in Rab Recycling. <i>Traffic</i> , 2000, 1, 270-281. | 1.3 | 37 |
| 3884 | Time-resolved fluorescence of tryptophans in yeast hexokinase-PI: effect of subunit dimerization and ligand binding. <i>Journal of Photochemistry and Photobiology B: Biology</i> , 2000, 55, 20-26. | 1.7 | 18 |
| 3885 | The human Rhesus-associated RhAG protein and a kidney homologue promote ammonium transport in yeast. <i>Nature Genetics</i> , 2000, 26, 341-344. | 9.4 | 325 |
| 3886 | The Src/Csk regulatory circuit arose early in metazoan evolution. <i>Oncogene</i> , 2000, 19, 3925-3930. | 2.6 | 27 |
| 3887 | A comprehensive analysis of protein-protein interactions in <i>Saccharomyces cerevisiae</i> . <i>Nature</i> , 2000, 403, 623-627. | 13.7 | 4,490 |
| 3888 | Functional dissection of in vivo interchromosome association in <i>Saccharomyces cerevisiae</i> . <i>Nature Cell Biology</i> , 2000, 2, 812-818. | 4.6 | 48 |

| # | ARTICLE | IF | CITATIONS |
|------|---|-----|-----------|
| 3889 | Cloning and sequencing analysis of TRP1 gene of <i>Flammulina velutipes</i> . <i>FEMS Microbiology Letters</i> , 2000, 190, 51-56. | 0.7 | 6 |
| 3890 | The immunosuppressive drug leflunomide affects mating-pheromone response and sporulation by different mechanisms in <i>Saccharomyces cerevisiae</i> . <i>FEMS Microbiology Letters</i> , 2000, 191, 57-60. | 0.7 | 2 |
| 3891 | Disruption of the ABC transporter genes PDR5, YOR1, and SNQ2, and their participation in improved fermentative activity of a sake yeast mutant showing pleiotropic drug resistance. <i>Journal of Bioscience and Bioengineering</i> , 2000, 89, 569-576. | 1.1 | 12 |
| 3892 | Effect of the FAA1 gene disruption of sake yeast on the accumulation of ethyl caproate in sake mash. <i>Journal of Bioscience and Bioengineering</i> , 2000, 89, 609-611. | 1.1 | 13 |
| 3893 | High-level expression of recombinant human serum albumin from the methylotrophic yeast <i>Pichia pastoris</i> with minimal protease production and activation. <i>Journal of Bioscience and Bioengineering</i> , 2000, 89, 55-61. | 1.1 | 174 |
| 3894 | A permease exhibiting a dual role for lysine and biotin uptake in <i>Saccharomyces cerevisiae</i> . <i>Journal of Bioscience and Bioengineering</i> , 2000, 89, 90-93. | 1.1 | 4 |
| 3895 | Amino acid residues contributing to stabilization of <i>Fusarium heterosporum</i> lipase. <i>Journal of Bioscience and Bioengineering</i> , 2000, 89, 446-450. | 1.1 | 5 |
| 3896 | Addition of oleic acid increases expression of recombinant human serum albumin by the AOX2 promoter in <i>Pichia pastoris</i> . <i>Journal of Bioscience and Bioengineering</i> , 2000, 89, 479-484. | 1.1 | 25 |
| 3897 | Cloning and functional expression of Rpn1, a regulatory-particle non-ATPase subunit 1, of proteasome from <i>Trypanosoma cruzi</i> . <i>Molecular and Biochemical Parasitology</i> , 2000, 110, 323-331. | 0.5 | 10 |
| 3898 | Functional cloning and mutational analysis of the human cDNA for phosphoacetylglucosamine mutase: identification of the amino acid residues essential for the catalysis. <i>Biochimica Et Biophysica Acta Gene Regulatory Mechanisms</i> , 2000, 1492, 369-376. | 2.4 | 35 |
| 3899 | Cloning, expression and regulation of <i>Schizosaccharomyces pombe</i> gene encoding thioltransferase. <i>Biochimica Et Biophysica Acta Gene Regulatory Mechanisms</i> , 2000, 1517, 171-175. | 2.4 | 7 |
| 3900 | Enthalpic destabilization of glycosylated lysozymes constructed by genetic modification. <i>BBA - Proteins and Proteomics</i> , 2000, 1481, 88-96. | 2.1 | 15 |
| 3901 | Characterization of the ubiquitin-specific protease activity of the mouse/human Unp/Unph oncoprotein. <i>BBA - Proteins and Proteomics</i> , 2000, 1481, 297-309. | 2.1 | 22 |
| 3902 | Agonistic and synergistic activity of tamoxifen in a yeast model system. <i>Biochemical Pharmacology</i> , 2000, 59, 177-185. | 2.0 | 22 |
| 3903 | Differential inhibition of human CYP3A4 and <i>Candida albicans</i> CYP51 with azole antifungal agents. <i>Chemico-Biological Interactions</i> , 2000, 125, 165-175. | 1.7 | 52 |
| 3904 | Molecular cloning of <i>Saccharomyces cerevisiae</i> MGC1/YDR473c gene which is essential for cell growth. , 2000, 77, 65-69. | | 0 |
| 3905 | Expression of <i>Bacillus</i> sp. β -xylosidase gene (xylB) in <i>Saccharomyces cerevisiae</i> . <i>Biotechnology Letters</i> , 2000, 22, 1025-1029. | 1.1 | 4 |
| 3906 | Differential regulation of testosterone vs. 5 α -dihydrotestosterone by selective androgen response elements. <i>Molecular and Cellular Biochemistry</i> , 2000, 206, 169-175. | 1.4 | 29 |

| # | ARTICLE | IF | CITATIONS |
|------|--|-----|-----------|
| 3907 | Title is missing!. Russian Journal of Genetics, 2000, 36, 1370-1374. | 0.2 | 1 |
| 3908 | Title is missing!. Russian Journal of Genetics, 2000, 36, 1375-1384. | 0.2 | 3 |
| 3909 | The Human T-Cell Leukemia Virus Type I (HTLV-I) X Region Encoded Protein p13II Interacts with Cellular Proteins. Virology, 2000, 277, 127-135. | 1.1 | 15 |
| 3910 | Studies on Insulin/IGF-1 Hybrid and IGF-1 Growth-Promoting Functional Region. IUBMB Life, 2000, 49, 321-325. | 1.5 | 7 |
| 3911 | An intact SH3 domain is required for myosin I-induced actin polymerization. EMBO Journal, 2000, 19, 4281-4291. | 3.5 | 107 |
| 3912 | A dual role for BBP/ScSF1 in nuclear pre-mRNA retention and splicing. EMBO Journal, 2000, 19, 1873-1886. | 3.5 | 89 |
| 3913 | Pseudohyphal growth is induced in <i>Saccharomyces cerevisiae</i> by a combination of stress and cAMP signalling. Antonie Van Leeuwenhoek, 2000, 78, 187-194. | 0.7 | 45 |
| 3914 | Nitrogen Regulation of <i>Saccharomyces cerevisiae</i> Invertase. Applied Biochemistry and Biotechnology, 2000, 84-86, 247-254. | 1.4 | 14 |
| 3915 | Construction of a flocculent brewer's yeast strain secreting <i>Aspergillus niger</i> Î²-galactosidase. Applied Microbiology and Biotechnology, 2000, 54, 97-103. | 1.7 | 21 |
| 3916 | Molecular Cloning and DNA Analysis of the Orotidine-5â€²-Phosphate Decarboxylase Gene from the Yeast <i>Saccharomyces exiguus</i> Yp74L-3. Current Microbiology, 2000, 41, 421-424. | 1.0 | 3 |
| 3917 | Mcchs1 , a Member of a Chitin Synthase Gene Family in <i>Mucor circinelloides</i> , Is Differentially Expressed During Dimorphism. Current Microbiology, 2000, 40, 169-175. | 1.0 | 15 |
| 3918 | The cadmium-resistant gene, CAD2 , which is a mutated putative copper-transporter gene (PCA1), controls the intracellular cadmium-level in the yeast <i>S. cerevisiae</i> . Current Genetics, 2000, 37, 79-86. | 0.8 | 57 |
| 3919 | UBC13 , a DNA-damage-inducible gene, is a member of the error-free postreplication repair pathway in <i>Saccharomyces cerevisiae</i> . Current Genetics, 2000, 37, 168-174. | 0.8 | 140 |
| 3920 | Characterization of novel rad6/ubc2 ubiquitin-conjugating enzyme mutants in yeast. Current Genetics, 2000, 37, 221-233. | 0.8 | 22 |
| 3921 | Multiple copies of MRG19 suppress transcription of the GAL1 promoter in a GAL80â€”dependent manner in <i>Saccharomyces cerevisiae</i> . Molecular Genetics and Genomics, 2000, 262, 1113-1122. | 2.4 | 16 |
| 3922 | Activation of basal transcription by a mutation in SIN4, a yeast global repressor, occurs through a mechanism different from activator-mediated transcriptional enhancement. Molecular Genetics and Genomics, 2000, 263, 48-59. | 2.4 | 12 |
| 3923 | Acquisition of a potential marker for insect transformation: isolation of a novel alcohol dehydrogenase gene from <i>Bactrocera oleae</i> by functional complementation in yeast. Molecular Genetics and Genomics, 2000, 263, 90-95. | 2.4 | 12 |
| 3924 | Protein overexport in a <i>Saccharomyces cerevisiae</i> mutant is not due to facilitated release of cell-surface proteins. Folia Microbiologica, 2000, 45, 251-254. | 1.1 | 1 |

| # | ARTICLE | IF | CITATIONS |
|------|--|-----|-----------|
| 3925 | Rapid purification of recombinant human lipocortin-I secreted from <i>Saccharomyces cerevisiae</i> . <i>Biotechnology and Bioprocess Engineering</i> , 2000, 5, 242-246. | 1.4 | 0 |
| 3926 | The tryptophan synthase-encoding <i>trpB</i> gene of <i>Aspergillus nidulans</i> is regulated by the cross-pathway control system. <i>Molecular Genetics and Genomics</i> , 2000, 263, 867-876. | 2.4 | 42 |
| 3927 | Intergenic complementation truncation mutants of cyclin-dependent kinase. <i>Molecular Genetics and Genomics</i> , 2000, 263, 222-231. | 2.4 | 0 |
| 3928 | Decreased meiotic reciprocal recombination in subtelomeric regions in <i>Saccharomyces cerevisiae</i> . <i>Chromosoma</i> , 2000, 109, 467-475. | 1.0 | 26 |
| 3929 | Mutations in helix 27 of the yeast <i>Saccharomyces cerevisiae</i> 18S rRNA affect the function of the decoding center of the ribosome. <i>Rna</i> , 2000, 6, 1174-1184. | 1.6 | 45 |
| 3930 | Elevated Temperature Greatly Improves Transformation of Fresh and Frozen Competent Cells in Yeast. <i>BioTechniques</i> , 2000, 28, 852-853. | 0.8 | 14 |
| 3931 | Global expression profiling of yeast treated with an inhibitor of amino acid biosynthesis, sulfometuron methyl. <i>Physiological Genomics</i> , 2000, 3, 83-92. | 1.0 | 105 |
| 3932 | Nonsense-mediated decay mutants do not affect programmed \hat{a}^1 frameshifting. <i>Rna</i> , 2000, 6, 952-961. | 1.6 | 70 |
| 3933 | Regulation of Yeast H ⁺ -ATPase by Protein Kinases Belonging to a Family Dedicated to Activation of Plasma Membrane Transporters. <i>Molecular and Cellular Biology</i> , 2000, 20, 7654-7661. | 1.1 | 167 |
| 3934 | Identifying a Core RNA Polymerase Surface Critical for Interactions with a Sigma-Like Specificity Factor. <i>Molecular and Cellular Biology</i> , 2000, 20, 7013-7023. | 1.1 | 19 |
| 3935 | Redirection of the Respiro-Fermentative Flux Distribution in <i>Saccharomyces cerevisiae</i> by Overexpression of the Transcription Factor Hap4p. <i>Applied and Environmental Microbiology</i> , 2000, 66, 1970-1973. | 1.4 | 85 |
| 3936 | Regulation of Hmg-Coa Reductase Degradation Requires the P-Type Atpase Cod1p/Spf1p. <i>Journal of Cell Biology</i> , 2000, 148, 915-924. | 2.3 | 103 |
| 3937 | Nucleoside Transporter Proteins of <i>Saccharomyces cerevisiae</i> . <i>Journal of Biological Chemistry</i> , 2000, 275, 25931-25938. | 1.6 | 70 |
| 3938 | Premature Structural Changes at Replication Origins in a Yeast Minichromosome Maintenance (MCM) Mutant. <i>Journal of Biological Chemistry</i> , 2000, 275, 18011-18021. | 1.6 | 58 |
| 3939 | Glucose-regulated Turnover of mRNA and the Influence of Poly(A) Tail Length on Half-life. <i>Journal of Biological Chemistry</i> , 2000, 275, 14155-14166. | 1.6 | 23 |
| 3940 | Biochemical and Functional Characterization of Inositol 1,3,4,5,6-Pentakisphosphate 2-Kinases. <i>Journal of Biological Chemistry</i> , 2000, 275, 36575-36583. | 1.6 | 83 |
| 3941 | The Family of SMF Metal Ion Transporters in Yeast Cells. <i>Journal of Biological Chemistry</i> , 2000, 275, 33388-33394. | 1.6 | 123 |
| 3942 | Aggregation of huntingtin in yeast varies with the length of the polyglutamine expansion and the expression of chaperone proteins. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 2000, 97, 1589-1594. | 3.3 | 524 |

| # | ARTICLE | IF | CITATIONS |
|------|--|-----|-----------|
| 3943 | d-Tyrosyl-tRNA ^{Tyr} Metabolism in <i>Saccharomyces cerevisiae</i> . <i>Journal of Biological Chemistry</i> , 2000, 275, 11626-11630. | 1.6 | 43 |
| 3944 | The Accumulation of Man ₆ GlcNAc ₂ -PP-dolichol in the <i>Saccharomyces cerevisiae</i> <i>alg9</i> Mutant Reveals a Regulatory Role for the Alg3p 1,3-Man Middle-arm Addition in Downstream Oligosaccharide-lipid and Glycoprotein Glycan Processing. <i>Journal of Biological Chemistry</i> , 2000, 275, 4267-4277. | 1.6 | 27 |
| 3945 | A Method for Determining the in Vivo Topology of Yeast Polytropic Membrane Proteins Demonstrates That Gap1p Fully Integrates into the Membrane Independently of Shr3p. <i>Journal of Biological Chemistry</i> , 2000, 275, 31488-31495. | 1.6 | 44 |
| 3946 | High-Level Rearrangement and Transcription of Yeast Artificial Chromosome-Based Mouse Ig ^h Transgenes Containing Distal Regions of the Contig. <i>Journal of Immunology</i> , 2000, 164, 812-824. | 0.4 | 8 |
| 3947 | Evidence for Contribution of Neutral Trehalase in Barotolerance of <i>Saccharomyces cerevisiae</i> . <i>Applied and Environmental Microbiology</i> , 2000, 66, 5182-5185. | 1.4 | 33 |
| 3948 | Functional Characterization of a Mammalian Sac1 and Mutants Exhibiting Substrate-specific Defects in Phosphoinositide Phosphatase Activity. <i>Journal of Biological Chemistry</i> , 2000, 275, 34293-34305. | 1.6 | 123 |
| 3949 | <i>Fusarium oxysporum</i> Fatty-acid Subterminal Hydroxylase (CYP505) Is a Membrane-bound Eukaryotic Counterpart of <i>Bacillus megaterium</i> Cytochrome P450BM3. <i>Journal of Biological Chemistry</i> , 2000, 275, 39734-39740. | 1.6 | 74 |
| 3950 | Regulation of Monovalent Ion Homeostasis and pH by the Ser-Thr Protein Phosphatase SIT4 in <i>Saccharomyces cerevisiae</i> . <i>Journal of Biological Chemistry</i> , 2000, 275, 30957-30961. | 1.6 | 38 |
| 3951 | Targeting of Aminopeptidase I to the Yeast Vacuole Is Mediated by Ssa1p, a Cytosolic Member of the 70-kDa Stress Protein Family. <i>Journal of Biological Chemistry</i> , 2000, 275, 34054-34059. | 1.6 | 10 |
| 3952 | TEP1, the yeast homolog of the human tumor suppressor gene PTEN/MMAC1/TEP1, is linked to the phosphatidylinositol pathway and plays a role in the developmental process of sporulation. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 2000, 97, 12672-12677. | 3.3 | 25 |
| 3953 | Sli2 (Ypk1), a Homologue of Mammalian Protein Kinase SGK, Is a Downstream Kinase in the Sphingolipid-Mediated Signaling Pathway of Yeast. <i>Molecular and Cellular Biology</i> , 2000, 20, 4411-4419. | 1.1 | 128 |
| 3954 | Analysis of the Role of the Hypervariable Region of Yeast Ras2p and Its Farnesylation in the Interaction with Exchange Factors and Adenylyl Cyclase. <i>Journal of Biological Chemistry</i> , 2000, 275, 17754-17761. | 1.6 | 12 |
| 3955 | Helicase and Capping Enzyme Active Site Mutations in Brome Mosaic Virus Protein 1a Cause Defects in Template Recruitment, Negative-Strand RNA Synthesis, and Viral RNA Capping. <i>Journal of Virology</i> , 2000, 74, 8803-8811. | 1.5 | 93 |
| 3956 | A missense mutation in the nuclear gene coding for the mitochondrial aspartyl-tRNA synthetase suppresses a mitochondrial tRNA ^{Asp} mutation. <i>Nucleic Acids Research</i> , 2000, 28, 1542-1547. | 6.5 | 2 |
| 3957 | The Kar3p Kinesin-related Protein Forms a Novel Heterodimeric Structure with Its Associated Protein Cik1p. <i>Molecular Biology of the Cell</i> , 2000, 11, 2373-2385. | 0.9 | 51 |
| 3958 | Characterization of Determinants for the Specificity of <i>Arabidopsis</i> Thioredoxins h in Yeast Complementation. <i>Journal of Biological Chemistry</i> , 2000, 275, 31641-31647. | 1.6 | 45 |
| 3959 | Gic2p May Link Activated Cdc42p to Components Involved in Actin Polarization, Including Bni1p and Bud6p (Aip3p). <i>Molecular and Cellular Biology</i> , 2000, 20, 6244-6258. | 1.1 | 72 |
| 3960 | mRNA splicing-mediated C-terminal replacement of transcription factor Hac1p is required for efficient activation of the unfolded protein response. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 2000, 97, 4660-4665. | 3.3 | 168 |

| # | ARTICLE | IF | CITATIONS |
|------|---|-----|-----------|
| 3961 | Peripheral Golgi protein GRASP65 is a target of mitotic polo-like kinase (Plk) and Cdc2. Proceedings of the National Academy of Sciences of the United States of America, 2000, 97, 12589-12594. | 3.3 | 116 |
| 3962 | Characterization of the Two Saccharopine Dehydrogenase Isozymes of Lysine Catabolism Encoded by the Single Composite AtLKR/SDHLocus of Arabidopsis. Plant Physiology, 2000, 124, 1363-1372. | 2.3 | 17 |
| 3963 | ARL4, an ARF-like Protein That Is Developmentally Regulated and Localized to Nuclei and Nucleoli. Journal of Biological Chemistry, 2000, 275, 37815-37823. | 1.6 | 46 |
| 3964 | Altered Distribution of the Yeast Plasma Membrane H ⁺ -ATPase as a Feature of Vacuolar H ⁺ -ATPase Null Mutants. Journal of Biological Chemistry, 2000, 275, 40088-40095. | 1.6 | 34 |
| 3965 | Characterization of Two 5-Aminoimidazole-4-carboxamide Ribonucleotide Transformylase/Inosine Monophosphate Cyclohydrolase Isozymes from <i>Saccharomyces cerevisiae</i> . Journal of Biological Chemistry, 2000, 275, 20920-20927. | 1.6 | 40 |
| 3966 | Suppression of sorbitol dependence in a strain bearing a mutation in the SRB1/PSA1/VIC9 gene encoding GDP-mannose pyrophosphorylase by PDE2 overexpression suggests a role for the Ras/cAMP signal-transduction pathway in the control of yeast cell-wall biogenesis. Microbiology (United Kingdom) 140:281-288. doi:10.1099/mic/0/01784314178431410000000000000000 | 0.7 | 28 |
| 3967 | Fruiting body development in <i>Coprinus cinereus</i> : regulated expression of two galectins secreted by a non-classical pathway The GenBank accession number for the sequence reported in this paper is AF130360. Microbiology (United Kingdom), 2000, 146, 1841-1853. | 0.7 | 104 |
| 3968 | Reduced virulence of <i>Candida albicans</i> mutants lacking the GNA1 gene encoding glucosamine-6-phosphate acetyltransferase. Microbiology (United Kingdom), 2000, 146, 1753-1758. | 0.7 | 38 |
| 3969 | Phosphatidylserine Synthesis Required for the Maximal Tryptophan Transport Activity in <i>Saccharomyces cerevisiae</i> . Bioscience, Biotechnology and Biochemistry, 2000, 64, 167-172. | 0.6 | 17 |
| 3970 | Requirement for Lysine-19 of the Yeast Mitochondrial ATPase Inhibitor for the Stability of the Inactivated Inhibitor-F1Fo Complex at Higher pH. Bioscience, Biotechnology and Biochemistry, 2000, 64, 89-95. | 0.6 | 5 |
| 3971 | Expression and Secretion of Scytalidopepsin B, an Acid Protease from <i>Scytalidium lignicolum</i> , in Yeast. Bioscience, Biotechnology and Biochemistry, 2000, 64, 1542-1546. | 0.6 | 11 |
| 3972 | Transport of Pyruvate in <i>Saccharomyces cerevisiae</i> and Cloning of the Gene Encoded Pyruvate Permease. Bioscience, Biotechnology and Biochemistry, 2000, 64, 980-984. | 0.6 | 38 |
| 3973 | Monitoring Activity of Caspases and Their Regulators in Yeast <i>Saccharomyces cerevisiae</i> . Methods in Enzymology, 2000, 322, 162-174. | 0.4 | 15 |
| 3974 | Cloning and Characterization of a Gene Complementing the Mutation of an Ethanol-sensitive Mutant of <i>SakeYeast</i> . Bioscience, Biotechnology and Biochemistry, 2000, 64, 229-236. | 0.6 | 58 |
| 3975 | Tetracycline-Regulatable System To Tightly Control Gene Expression in the Pathogenic Fungus <i>Candida albicans</i> . Infection and Immunity, 2000, 68, 6712-6719. | 1.0 | 152 |
| 3976 | Assembly and Preferential Localization of Nup116p on the Cytoplasmic Face of the Nuclear Pore Complex by Interaction with Nup82p. Molecular and Cellular Biology, 2000, 20, 5736-5748. | 1.1 | 47 |
| 3977 | Dual Lipid Modification Motifs in G α and G β Subunits Are Required for Full Activity of the Pheromone Response Pathway in <i>Saccharomyces cerevisiae</i> . Molecular Biology of the Cell, 2000, 11, 957-968. | 0.9 | 58 |
| 3978 | Association of Yeast Adenylyl Cyclase with Cyclase-Associated Protein CAP Forms a Second Ras-Binding Site Which Mediates Its Ras-Dependent Activation. Molecular and Cellular Biology, 2000, 20, 26-33. | 1.1 | 63 |

| # | ARTICLE | IF | CITATIONS |
|------|--|-----|-----------|
| 3979 | Cloning of the human cDNA which can complement the defect of the yeast mannosyltransferase I-deficient mutant alg 1. <i>Glycobiology</i> , 2000, 10, 321-327. | 1.3 | 19 |
| 3980 | The Dual-Specificity Protein Phosphatase Yvh1p Regulates Sporulation, Growth, and Glycogen Accumulation Independently of Catalytic Activity in <i>Saccharomyces cerevisiae</i> via the Cyclic AMP-Dependent Protein Kinase Cascade. <i>Journal of Bacteriology</i> , 2000, 182, 3517-3528. | 1.0 | 35 |
| 3981 | Batten disease: evaluation of CLN3 mutations on protein localization and function. <i>Human Molecular Genetics</i> , 2000, 9, 735-744. | 1.4 | 86 |
| 3982 | Molecular Cloning and Characterization of ATP-Phosphoribosyl Transferase from <i>Arabidopsis</i> , a Key Enzyme in the Histidine Biosynthetic Pathway. <i>Plant Physiology</i> , 2000, 122, 907-914. | 2.3 | 41 |
| 3983 | Identification of a Novel Family of Nonclassic Yeast Phosphatidylinositol Transfer Proteins Whose Function Modulates Phospholipase D Activity and Sec14p-independent Cell Growth. <i>Molecular Biology of the Cell</i> , 2000, 11, 1989-2005. | 0.9 | 140 |
| 3984 | In <i>Saccharomyces cerevisiae</i> , Expression of Arginine Catabolic Genes CAR1 and CAR2 in Response to Exogenous Nitrogen Availability Is Mediated by the Ume6 (CargR1)-Sin3 (CargR2)-Rpd3 (CargR3) Complex. <i>Journal of Bacteriology</i> , 2000, 182, 3158-3164. | 1.0 | 33 |
| 3985 | Vav3 Mediates Receptor Protein Tyrosine Kinase Signaling, Regulates GTPase Activity, Modulates Cell Morphology, and Induces Cell Transformation. <i>Molecular and Cellular Biology</i> , 2000, 20, 9212-9224. | 1.1 | 131 |
| 3986 | Multiple Signals from Dysfunctional Mitochondria Activate the Pleiotropic Drug Resistance Pathway in <i>Saccharomyces cerevisiae</i> . <i>Journal of Biological Chemistry</i> , 2000, 275, 37347-37356. | 1.6 | 175 |
| 3987 | Depletion of the Squalene Synthase (ERG9) Gene Does Not Impair Growth of <i>Candida glabrata</i> in Mice. <i>Antimicrobial Agents and Chemotherapy</i> , 2000, 44, 2411-2418. | 1.4 | 49 |
| 3988 | Molecular Interactions Involved in the Transactivation of the Human T-Cell Leukemia Virus Type 1 Promoter Mediated by Tax and CREB-2 (ATF-4). <i>Molecular and Cellular Biology</i> , 2000, 20, 3470-3481. | 1.1 | 64 |
| 3989 | Yeast flavin-containing monooxygenase is induced by the unfolded protein response. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 2000, 97, 121-126. | 3.3 | 35 |
| 3990 | Crk1, a Novel Cdc2-Related Protein Kinase, Is Required for Hyphal Development and Virulence in <i>Candida albicans</i> . <i>Molecular and Cellular Biology</i> , 2000, 20, 8696-8708. | 1.1 | 60 |
| 3991 | A new double-stranded RNA-binding protein that interacts with PKR. <i>Nucleic Acids Research</i> , 2000, 28, 1407-1417. | 6.5 | 29 |
| 3992 | Role of HSP90 in Salt Stress Tolerance via Stabilization and Regulation of Calcineurin. <i>Molecular and Cellular Biology</i> , 2000, 20, 9262-9270. | 1.1 | 122 |
| 3993 | Isolation of the MIG1 Gene from <i>Candida albicans</i> and Effects of Its Disruption on Catabolite Repression. <i>Journal of Bacteriology</i> , 2000, 182, 320-326. | 1.0 | 67 |
| 3994 | MTABC3, a Novel Mitochondrial ATP-binding Cassette Protein Involved in Iron Homeostasis. <i>Journal of Biological Chemistry</i> , 2000, 275, 17536-17540. | 1.6 | 118 |
| 3995 | <i>Schizosaccharomyces pombe</i> Rho2p GTPase Regulates Cell Wall β -Glucan Biosynthesis through the Protein Kinase Pck2p. <i>Molecular Biology of the Cell</i> , 2000, 11, 4393-4401. | 0.9 | 89 |
| 3996 | Molecular Basis for Rab Prenylation. <i>Journal of Cell Biology</i> , 2000, 150, 89-104. | 2.3 | 277 |

| # | ARTICLE | IF | CITATIONS |
|------|--|-----|-----------|
| 3997 | Regulatory Mechanisms for Modulation of Signaling through the Cell Integrity Slt2-mediated Pathway in <i>Saccharomyces cerevisiae</i> . <i>Journal of Biological Chemistry</i> , 2000, 275, 1511-1519. | 1.6 | 316 |
| 3998 | Interaction Specificity of Arabidopsis Calcineurin B-Like Calcium Sensors and Their Target Kinases. <i>Plant Physiology</i> , 2000, 124, 1844-1853. | 2.3 | 192 |
| 3999 | Cytoplasmic Sequestration of Rel Proteins by I κ B β Requires CRM1-Dependent Nuclear Export. <i>Molecular and Cellular Biology</i> , 2000, 20, 2269-2284. | 1.1 | 147 |
| 4000 | Lst1p and Sec24p Cooperate in Sorting of the Plasma Membrane Atpase into Copii Vesicles in <i>Saccharomyces cerevisiae</i> . <i>Journal of Cell Biology</i> , 2000, 151, 973-984. | 2.3 | 133 |
| 4001 | A Lecithin Cholesterol Acyltransferase-like Gene Mediates Diacylglycerol Esterification in Yeast. <i>Journal of Biological Chemistry</i> , 2000, 275, 15609-15612. | 1.6 | 236 |
| 4002 | Initiation of Protein Synthesis in <i>Saccharomyces cerevisiae</i> Mitochondria without Formylation of the Initiator tRNA. <i>Journal of Bacteriology</i> , 2000, 182, 2886-2892. | 1.0 | 62 |
| 4003 | Communication between the ATPase and Cleavage/Religation Domains of Human Topoisomerase II α . <i>Journal of Biological Chemistry</i> , 2000, 275, 13041-13048. | 1.6 | 41 |
| 4004 | Essential Function of the Polo Box of Cdc5 in Subcellular Localization and Induction of Cytokinetic Structures. <i>Molecular and Cellular Biology</i> , 2000, 20, 286-298. | 1.1 | 151 |
| 4005 | Sbe2p and Sbe22p, Two Homologous Golgi Proteins Involved in Yeast Cell Wall Formation. <i>Molecular Biology of the Cell</i> , 2000, 11, 435-452. | 0.9 | 40 |
| 4006 | Functions and Functional Domains of the GTPase Cdc42p. <i>Molecular Biology of the Cell</i> , 2000, 11, 339-354. | 0.9 | 79 |
| 4007 | Identification of <i>Cryptosporidium parvum</i> Dihydrofolate Reductase Inhibitors by Complementation in <i>Saccharomyces cerevisiae</i> . <i>Antimicrobial Agents and Chemotherapy</i> , 2000, 44, 1019-1028. | 1.4 | 41 |
| 4008 | Sec24p and Iss1p Function Interchangeably in Transport Vesicle Formation from the Endoplasmic Reticulum in <i>Saccharomyces cerevisiae</i> . <i>Molecular Biology of the Cell</i> , 2000, 11, 983-998. | 0.9 | 70 |
| 4009 | The Role of Nuclear Cap Binding Protein Cbc1p of Yeast in mRNA Termination and Degradation. <i>Molecular and Cellular Biology</i> , 2000, 20, 2827-2838. | 1.1 | 66 |
| 4010 | Distinct Roles for the Yeast Phosphatidylinositol 4-Kinases, Stt4p and Pik1p, in Secretion, Cell Growth, and Organelle Membrane Dynamics. <i>Molecular Biology of the Cell</i> , 2000, 11, 2673-2689. | 0.9 | 327 |
| 4011 | RRS1, a Conserved Essential Gene, Encodes a Novel Regulatory Protein Required for Ribosome Biogenesis in <i>Saccharomyces cerevisiae</i> . <i>Molecular and Cellular Biology</i> , 2000, 20, 2066-2074. | 1.1 | 80 |
| 4012 | CHL1 is a nuclear protein with an essential ATP binding site that exhibits a size-dependent effect on chromosome segregation. <i>Nucleic Acids Research</i> , 2000, 28, 3056-3064. | 6.5 | 34 |
| 4013 | Tryptophan Permease Gene TAT2 Confers High-Pressure Growth in <i>Saccharomyces cerevisiae</i> . <i>Molecular and Cellular Biology</i> , 2000, 20, 8093-8102. | 1.1 | 147 |
| 4014 | Identification of Domains and Residues within the ϵ Subunit of Eukaryotic Translation Initiation Factor 2B (eIF2B ϵ) Required for Guanine Nucleotide Exchange Reveals a Novel Activation Function Promoted by eIF2B Complex Formation. <i>Molecular and Cellular Biology</i> , 2000, 20, 3965-3976. | 1.1 | 99 |

| # | ARTICLE | IF | CITATIONS |
|------|---|-----|-----------|
| 4015 | Localization and Signaling of G β 2 Subunit Ste4p Are Controlled by a β -Factor Receptor and the α -Specific Protein Apg7p. <i>Molecular and Cellular Biology</i> , 2000, 20, 8826-8835. | 1.1 | 27 |
| 4016 | ArgR11, a Component of the ArgR-Mcm1 Complex Involved in the Control of Arginine Metabolism in <i>Saccharomyces cerevisiae</i> , Is the Sensor of Arginine. <i>Molecular and Cellular Biology</i> , 2000, 20, 2087-2097. | 1.1 | 45 |
| 4017 | The Gcd10p/Gcd14p complex is the essential two-subunit tRNA(1-methyladenosine) methyltransferase of <i>Saccharomyces cerevisiae</i> . <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 2000, 97, 5173-5178. | 3.3 | 131 |
| 4018 | Isolation of <i>Trypanosoma brucei</i> CYC2 and CYC3 Cyclin Genes by Rescue of a Yeast G1Cyclin Mutant. <i>Journal of Biological Chemistry</i> , 2000, 275, 8315-8323. | 1.6 | 47 |
| 4019 | Sweet Tooth, a Novel Receptor Protein-tyrosine Kinase with C-type Lectin-like Extracellular Domains. <i>Journal of Biological Chemistry</i> , 2000, 275, 10323-10330. | 1.6 | 37 |
| 4020 | Cloning, Expression, Characterization, and Nucleophile Identification of Family 3, <i>Aspergillus niger</i> β -Glucosidase. <i>Journal of Biological Chemistry</i> , 2000, 275, 4973-4980. | 1.6 | 115 |
| 4021 | Mutations in Yeast ARV1 Alter Intracellular Sterol Distribution and Are Complemented by Human ARV1. <i>Journal of Biological Chemistry</i> , 2000, 275, 40667-40670. | 1.6 | 95 |
| 4022 | Aut7p, a Soluble Autophagic Factor, Participates in Multiple Membrane Trafficking Processes. <i>Journal of Biological Chemistry</i> , 2000, 275, 32966-32973. | 1.6 | 62 |
| 4023 | Stalk Segment 4 of the Yeast Plasma Membrane H ⁺ -ATPase. <i>Journal of Biological Chemistry</i> , 2000, 275, 20545-20550. | 1.6 | 14 |
| 4024 | Characterization of a (2R,3R)-2,3-Butanediol Dehydrogenase as the <i>Saccharomyces cerevisiae</i> YAL060W Gene Product. <i>Journal of Biological Chemistry</i> , 2000, 275, 35876-35885. | 1.6 | 114 |
| 4025 | Kinectin-Kinesin Binding Domains and Their Effects on Organelle Motility. <i>Journal of Biological Chemistry</i> , 2000, 275, 32854-32860. | 1.6 | 50 |
| 4026 | The Aconitase Function of Iron Regulatory Protein 1. <i>Journal of Biological Chemistry</i> , 2000, 275, 16227-16234. | 1.6 | 58 |
| 4027 | A Family of ADP-Ribosylation Factor Effectors That Can Alter Membrane Transport through the trans-Golgi. <i>Molecular Biology of the Cell</i> , 2000, 11, 1241-1255. | 0.9 | 258 |
| 4028 | Functional Expression of Regiospecific Cytochrome P450 Limonene Hydroxylases from Mint (<i>Mentha</i>) Tj ETQq1 1 0.784314 rgBT /Overl 379, 127-136. | 1.4 | 102 |
| 4029 | Molecular Cloning of cDNA Encoding β -N-Acetylgalactosaminidase from <i>Acremonium</i> sp. and Its Expression in Yeast. <i>Archives of Biochemistry and Biophysics</i> , 2000, 384, 305-310. | 1.4 | 18 |
| 4030 | Interaction of the PDZ Domain of Human PICK1 with Class I ADP-Ribosylation Factors. <i>Biochemical and Biophysical Research Communications</i> , 2000, 267, 149-155. | 1.0 | 52 |
| 4031 | yam8+, a <i>Schizosaccharomyces pombe</i> Gene, Is a Potential Homologue of the <i>Saccharomyces cerevisiae</i> MID1 Gene Encoding a Stretch- Activated Ca ²⁺ -Permeable Channel. <i>Biochemical and Biophysical Research Communications</i> , 2000, 269, 265-269. | 1.0 | 14 |
| 4032 | Stimulation of Yeast 3-Phosphoglycerate Kinase Gene Promoter by Paraquat. <i>Biochemical and Biophysical Research Communications</i> , 2000, 270, 1036-1040. | 1.0 | 13 |

| # | ARTICLE | IF | CITATIONS |
|------|--|-----|-----------|
| 4033 | Functional Expression of the Pore Forming Subunit of the ATP-Sensitive Potassium Channel in <i>Saccharomyces cerevisiae</i> . <i>Biochemical and Biophysical Research Communications</i> , 2000, 272, 403-409. | 1.0 | 6 |
| 4034 | Hsl7p, the Yeast Homologue of Human JBP1, Is a Protein Methyltransferase. <i>Biochemical and Biophysical Research Communications</i> , 2000, 274, 105-111. | 1.0 | 57 |
| 4035 | The Cysteine- and Glycine-Rich LIM Domain Protein CRP2 Specifically Interacts with a Novel Human Protein (CRP2BP). <i>Biochemical and Biophysical Research Communications</i> , 2000, 274, 655-663. | 1.0 | 22 |
| 4036 | Expression of ZRC1 Coding for Suppressor of Zinc Toxicity Is Induced by Zinc-starvation Stress in Zap1-Dependent Fashion in <i>Saccharomyces cerevisiae</i> . <i>Biochemical and Biophysical Research Communications</i> , 2000, 276, 879-884. | 1.0 | 37 |
| 4037 | A Pivotal Role of Zn-Binding Residues in the Function of the Copper Chaperone for SOD1. <i>Biochemical and Biophysical Research Communications</i> , 2000, 276, 999-1004. | 1.0 | 20 |
| 4038 | Positive Regulation of Transcription of Homeoprotein-Encoding YHP1 by the Two-Component Regulator Sln1 in <i>Saccharomyces cerevisiae</i> . <i>Biochemical and Biophysical Research Communications</i> , 2000, 278, 344-348. | 1.0 | 4 |
| 4039 | NBP1 (Nap1 Binding Protein 1), an essential gene for G2/M transition of <i>Saccharomyces cerevisiae</i> , encodes a protein of distinct sub-nuclear localization. <i>Gene</i> , 2000, 246, 395-404. | 1.0 | 21 |
| 4040 | Complementation of <i>Saccharomyces cerevisiae</i> mutations in genes involved in translation and protein folding (EFB1 and SSB1) with <i>Candida albicans</i> cloned genes. <i>Research in Microbiology</i> , 2000, 151, 739-746. | 1.0 | 17 |
| 4041 | The Activation Loop of Phosphatidylinositol Phosphate Kinases Determines Signaling Specificity. <i>Molecular Cell</i> , 2000, 5, 1-11. | 4.5 | 214 |
| 4042 | Cloning and expression of cDNAs encoding plant V-ATPase subunits in the corresponding yeast null mutants. <i>Biochimica Et Biophysica Acta - Bioenergetics</i> , 2000, 1459, 489-498. | 0.5 | 13 |
| 4043 | Genetic characterization of the 534DPPR motif of the yeast plasma membrane H ⁺ -ATPase. <i>Biochimica Et Biophysica Acta - Biomembranes</i> , 2000, 1468, 99-106. | 1.4 | 2 |
| 4044 | Sequencing and heterologous expression in <i>Saccharomyces cerevisiae</i> of a <i>Cryptococcus neoformans</i> cDNA encoding a plasma membrane H ⁺ -ATPase. <i>Biochimica Et Biophysica Acta - Biomembranes</i> , 2000, 1509, 103-110. | 1.4 | 4 |
| 4045 | The cell wall integrity/remodeling MAPK cascade is involved in glucose activation of the yeast plasma membrane H ⁺ -ATPase. <i>Biochimica Et Biophysica Acta - Biomembranes</i> , 2000, 1509, 189-194. | 1.4 | 13 |
| 4046 | The Leu-132 of the Ste4(GÎ²) subunit is essential for proper coupling of the G protein with the Ste2 Î± factor receptor during the mating pheromone response in yeast. <i>FEBS Letters</i> , 2000, 467, 22-26. | 1.3 | 17 |
| 4047 | Biochemical characterization and subcellular localization of the sterol C-24(28) reductase, Erg4p, from the yeast <i>Saccharomyces cerevisiae</i> . <i>FEBS Letters</i> , 2000, 470, 83-87. | 1.3 | 75 |
| 4048 | Stimulation of the yeast high osmolarity glycerol (HOG) pathway: evidence for a signal generated by a change in turgor rather than by water stress. <i>FEBS Letters</i> , 2000, 472, 159-165. | 1.3 | 81 |
| 4049 | Cloning and characterization of a root specific high-affinity sulfate transporter from <i>Arabidopsis thaliana</i> . <i>FEBS Letters</i> , 2000, 475, 65-69. | 1.3 | 106 |
| 4050 | Possible His to Asp phosphorelay signaling in an <i>Arabidopsis</i> two-component system. <i>FEBS Letters</i> , 2000, 478, 227-232. | 1.3 | 147 |

| # | ARTICLE | IF | CITATIONS |
|------|--|-----|-----------|
| 4051 | Heterologous expression of a mammalian epithelial sodium channel in yeast. <i>FEBS Letters</i> , 2000, 481, 77-80. | 1.3 | 11 |
| 4052 | Recent Developments in Molecular Genetics of <i>Candida Albicans</i> . <i>Annual Review of Microbiology</i> , 2000, 54, 463-498. | 2.9 | 104 |
| 4053 | HRD Gene Dependence of Endoplasmic Reticulum-associated Degradation. <i>Molecular Biology of the Cell</i> , 2000, 11, 1697-1708. | 0.9 | 100 |
| 4054 | An NH ₂ -terminal deleted plasma membrane H ⁺ -ATPase is a dominant negative mutant and is sequestered in endoplasmic reticulum derived structures. <i>Biochemistry and Cell Biology</i> , 2000, 78, 51-58. | 0.9 | 12 |
| 4055 | Defect in Cell Wall Integrity of the Yeast <i>Saccharomyces cerevisiae</i> Caused by a Mutation of the GDP-mannose Pyrophosphorylase Gene <i>VIG9</i> . <i>Bioscience, Biotechnology and Biochemistry</i> , 2000, 64, 1937-1941. | 0.6 | 30 |
| 4056 | Improving synthetic lethal screens by regulating the yeast centromere sequence. <i>Genome</i> , 2000, 43, 910-917. | 0.9 | 8 |
| 4057 | Introduction of Kex2 cleavage sites in fusion proteins for monitoring localization and transport in yeast secretory pathway. <i>Methods in Enzymology</i> , 2000, 327, 107-118. | 0.4 | 4 |
| 4058 | Interaction of the Rabies Virus P Protein with the LC8 Dynein Light Chain. <i>Journal of Virology</i> , 2000, 74, 10212-10216. | 1.5 | 279 |
| 4059 | Expression of glucose oxidase by using recombinant yeast. <i>Journal of Biotechnology</i> , 2000, 81, 35-44. | 1.9 | 77 |
| 4060 | Lactose utilization by <i>Saccharomyces cerevisiae</i> strains expressing <i>Kluyveromyces lactis</i> LAC genes. <i>Journal of Biotechnology</i> , 2000, 84, 97-106. | 1.9 | 39 |
| 4061 | Deletion of the <i>MAG1</i> DNA glycosylase gene suppresses alkylation-induced killing and mutagenesis in yeast cells lacking AP endonucleases. <i>Mutation Research DNA Repair</i> , 2001, 487, 137-147. | 3.8 | 44 |
| 4062 | Expression and Characterization of Recombinant Human Antithrombin III in <i>Pichia pastoris</i> . <i>Protein Expression and Purification</i> , 2001, 23, 55-65. | 0.6 | 43 |
| 4063 | Demonstration That Menthofuran Synthase of Mint (<i>Mentha</i>) Is a Cytochrome P450 Monooxygenase: Cloning, Functional Expression, and Characterization of the Responsible Gene. <i>Archives of Biochemistry and Biophysics</i> , 2001, 390, 279-286. | 1.4 | 106 |
| 4064 | Cloning and Characterization of <i>Ginkgo biloba</i> Levopimaradiene Synthase, Which Catalyzes the First Committed Step in Ginkgolide Biosynthesis. <i>Archives of Biochemistry and Biophysics</i> , 2001, 392, 263-269. | 1.4 | 94 |
| 4065 | Functional Characterization of Two Ripening-related Sucrose Transporters from Grape Berries. <i>Annals of Botany</i> , 2001, 87, 125-129. | 1.4 | 55 |
| 4066 | Rpg1p/Tif32p, a Subunit of Translation Initiation Factor 3, Interacts with Actin-Associated Protein Sla2p. <i>Biochemical and Biophysical Research Communications</i> , 2001, 282, 1244-1250. | 1.0 | 14 |
| 4067 | Biochemical Analysis of Yeast $G1\pm$ Mutants That Enhance Adaptation to Pheromone. <i>Biochemical and Biophysical Research Communications</i> , 2001, 284, 247-254. | 1.0 | 9 |
| 4068 | Interaction of Myc-Associated Zinc Finger Protein with DCC, the Product of a Tumor-Suppressor Gene, during the Neural Differentiation of P19 EC Cells. <i>Biochemical and Biophysical Research Communications</i> , 2001, 286, 1087-1097. | 1.0 | 16 |

| # | ARTICLE | IF | CITATIONS |
|------|---|-----|-----------|
| 4069 | Identification of Structural Domains Affecting Transactivation Potential of Nm23. <i>Biochemical and Biophysical Research Communications</i> , 2001, 289, 738-743. | 1.0 | 15 |
| 4070 | A Secretion Signal Is Present in the <i>Collybia velutipes</i> Oxalate Decarboxylase Gene. <i>Biochemical and Biophysical Research Communications</i> , 2001, 289, 807-812. | 1.0 | 12 |
| 4071 | Cloning and Characterization of a Potential Transcriptional Activator of Human β -Globin Genes. <i>Blood Cells, Molecules, and Diseases</i> , 2001, 27, 1-15. | 0.6 | 6 |
| 4072 | Genetic interaction between yeast <i>Saccharomyces cerevisiae</i> release factors and the decoding region of 18 S rRNA. <i>Journal of Molecular Biology</i> , 2001, 305, 715-727. | 2.0 | 28 |
| 4073 | Denatured state thermodynamics: residual structure, chain stiffness and scaling factors ¹¹ Edited by C. R. Matthews. <i>Journal of Molecular Biology</i> , 2001, 311, 1091-1104. | 2.0 | 64 |
| 4074 | The Tandem Affinity Purification (TAP) Method: A General Procedure of Protein Complex Purification. <i>Methods</i> , 2001, 24, 218-229. | 1.9 | 1,550 |
| 4075 | Transformation Systems of non- <i>Saccharomyces</i> Yeasts. <i>Critical Reviews in Biotechnology</i> , 2001, 21, 177-218. | 5.1 | 35 |
| 4076 | Tryptophan Fluorescence of Yeast Actin Resolved via Conserved Mutations. <i>Biophysical Journal</i> , 2001, 80, 427-434. | 0.2 | 33 |
| 4077 | Genetic dissection of the yeast 26S proteasome: Cell cycle defects caused by the \hat{r}^{prn9} mutation. <i>Biochimie</i> , 2001, 83, 333-340. | 1.3 | 15 |
| 4078 | Cloning and characterization of a cDNA encoding hexokinase from tomato. <i>Plant Science</i> , 2001, 160, 209-218. | 1.7 | 27 |
| 4079 | The REC41 gene of <i>Saccharomyces cerevisiae</i> : isolation and genetic analysis. <i>Mutation Research DNA Repair</i> , 2001, 486, 41-52. | 3.8 | 2 |
| 4080 | Identification of glycogen phosphorylase and creatine kinase as calpain substrates in skeletal muscle. <i>International Journal of Biochemistry and Cell Biology</i> , 2001, 33, 531-540. | 1.2 | 30 |
| 4081 | Nhp6, an HMG1 Protein, Functions in SNR6 Transcription by RNA Polymerase III in <i>S. cerevisiae</i> . <i>Molecular Cell</i> , 2001, 7, 309-318. | 4.5 | 66 |
| 4082 | The cAMP response element binding protein-2 (CREB-2) can interact with the C/EBP-homologous protein (CHOP). <i>FEBS Letters</i> , 2001, 502, 57-62. | 1.3 | 34 |
| 4083 | Features of V-ATPases that distinguish them from F-ATPases. <i>FEBS Letters</i> , 2001, 504, 223-228. | 1.3 | 30 |
| 4084 | Molecular characterization of two ammonium transporters from the ectomycorrhizal fungus <i>Hebeloma cylindrosporum</i> . <i>FEBS Letters</i> , 2001, 505, 393-398. | 1.3 | 60 |
| 4085 | Genetic evidence that antibacterial activity of lysozyme is independent of its catalytic function. <i>FEBS Letters</i> , 2001, 506, 27-32. | 1.3 | 285 |
| 4086 | Elements from the cAMP signaling pathway are involved in the control of expression of the yeast gluconeogenic gene <i>FBP1</i> . <i>FEBS Letters</i> , 2001, 506, 262-266. | 1.3 | 13 |

| # | ARTICLE | IF | CITATIONS |
|------|--|-----|-----------|
| 4087 | Characterisation of a concentrative type of adenosine transporter from <i>Arabidopsis thaliana</i> (ENT1, At). <i>FEBS Letters</i> , 2001, 509, 370-374. | 1.3 | 53 |
| 4088 | Identification of functional regions of Cbp3p, an enzyme-specific chaperone required for the assembly of ubiquinol-cytochrome c reductase in yeast mitochondria The costs of publication of this article were defrayed in part by the payment of page charges. This article must therefore be hereby marked "advertisement" in accordance with 18 U.S.C. Section 1734 solely to indicate this fact. 1. <i>Biochimica Et Biophysica Acta - Bioenergetics</i> , 2001, 1506, 103-116. | 0.5 | 14 |
| 4089 | Multiple Yap1p-binding Sites Mediate Induction of the Yeast Major Facilitator FLR1 Gene in Response to Drugs, Oxidants, and Alkylating Agents. <i>Journal of Biological Chemistry</i> , 2001, 276, 1138-1145. | 1.6 | 93 |
| 4090 | Functional production of mammalian concentrative nucleoside transporters in <i>Saccharomyces cerevisiae</i> . <i>Molecular Membrane Biology</i> , 2001, 18, 73-79. | 2.0 | 7 |
| 4091 | Ribosomal protein L5 helps anchor peptidyl-tRNA to the P-site in <i>Saccharomyces cerevisiae</i> . <i>Rna</i> , 2001, 7, 1084-1096. | 1.6 | 54 |
| 4092 | Programmed +1 frameshifting stimulated by complementarity between a downstream mRNA sequence and an error-correcting region of rRNA. <i>Rna</i> , 2001, 7, 275-284. | 1.6 | 28 |
| 4093 | A Streamlined Process to Phenotypically Profile Heterologous cDNAs in Parallel Using Yeast Cell-Based Assays. <i>Genome Research</i> , 2001, 11, 1899-1912. | 2.4 | 30 |
| 4094 | Evidence that Fungal MEP Proteins Mediate Diffusion of the Uncharged Species NH ₃ across the Cytoplasmic Membrane. <i>Molecular and Cellular Biology</i> , 2001, 21, 5733-5741. | 1.1 | 77 |
| 4095 | Chapter 20 Studies of apoptosis proteins in yeast. <i>Methods in Cell Biology</i> , 2001, 66, 453-468. | 0.5 | 6 |
| 4096 | Nis1 encoded by YNL078W: a new neck protein of <i>Saccharomyces cerevisiae</i> . <i>Genes and Genetic Systems</i> , 2001, 76, 335-343. | 0.2 | 30 |
| 4097 | Identification of Zinc Finger Proteins Bound to a Silencer Region in the Rat Glutathione Transferase P Gene. <i>Biological and Pharmaceutical Bulletin</i> , 2001, 24, 144-150. | 0.6 | 2 |
| 4098 | Molecular transformation, gene cloning, and gene expression systems for filamentous fungi. <i>Applied Mycology and Biotechnology</i> , 2001, 1, 199-238. | 0.3 | 3 |
| 4099 | Multiple polyamine transport systems on the vacuolar membrane in yeast. <i>Biochemical Journal</i> , 2001, 353, 681. | 1.7 | 84 |
| 4100 | RGS18 is a myeloerythroid lineage-specific regulator of G-protein-signalling molecule highly expressed in megakaryocytes. <i>Biochemical Journal</i> , 2001, 359, 109. | 1.7 | 36 |
| 4101 | LIM-domain protein cysteine- and glycine-rich protein 2 (CRP2) is a novel marker of hepatic stellate cells and binding partner of the protein inhibitor of activated STAT1. <i>Biochemical Journal</i> , 2001, 359, 485. | 1.7 | 31 |
| 4102 | <i>Saccharomyces cerevisiae</i> Multidrug Resistance Gene Expression Inversely Correlates with the Status of the FO Component of the Mitochondrial ATPase. <i>Journal of Biological Chemistry</i> , 2001, 276, 47844-47852. | 1.6 | 76 |
| 4103 | Purification and Characterization of YACs Containing Large Inserts. , 2001, Chapter 5, Unit 5.7. | | 0 |
| 4104 | Breeding of sake yeasts that produce enhanced levels of flavor by transformation with a mutated LEU4 gene or leudh gene. <i>Journal of the Brewing Society of Japan</i> , 2001, 96, 129-136. | 0.1 | 3 |

| # | ARTICLE | IF | CITATIONS |
|------|---|-----|-----------|
| 4105 | Control of Higher Alcohol Production by Manipulation of the <i>BAP2</i> Gene in Brewing Yeast. <i>Journal of the American Society of Brewing Chemists</i> , 2001, 59, 157-162. | 0.8 | 41 |
| 4107 | Regulative fine-tuning of the two novel DAHP isoenzymes <i>aroFp</i> and <i>aroGp</i> of the filamentous fungus <i>Aspergillus nidulans</i> . <i>Archives of Microbiology</i> , 2001, 175, 112-121. | 1.0 | 11 |
| 4108 | Intelligent yeast strains with the ability to self-monitor the concentrations of intra- and extracellular phosphate or ammonium ion by emission of fluorescence from the cell surface. <i>Applied Microbiology and Biotechnology</i> , 2001, 57, 702-707. | 1.7 | 21 |
| 4109 | Function of the prosequence for in vivo folding and secretion of active <i>Rhizopus oryzae</i> lipase in <i>Saccharomyces cerevisiae</i> . <i>Applied Microbiology and Biotechnology</i> , 2001, 55, 454-462. | 1.7 | 45 |
| 4110 | Quantitative evaluation of the enhanced green fluorescent protein displayed on the cell surface of <i>Saccharomyces cerevisiae</i> by fluorometric and confocal laser scanning microscopic analyses. <i>Applied Microbiology and Biotechnology</i> , 2001, 55, 471-475. | 1.7 | 90 |
| 4111 | Biotransformation of tryptamine and secologanin into plant terpenoid indole alkaloids by transgenic yeast. <i>Applied Microbiology and Biotechnology</i> , 2001, 56, 420-424. | 1.7 | 81 |
| 4112 | Cell surface-engineered yeast displaying a histidine oligopeptide (hexa-His) has enhanced adsorption and tolerance to heavy metal ions. <i>Applied Microbiology and Biotechnology</i> , 2001, 57, 697-701. | 1.7 | 87 |
| 4113 | Subunit C of the Vacuolar-Type ATPase from the Vanadium-Rich Ascidian <i>Ascidia sydneiensis samea</i> Rescued the pH Sensitivity of Yeast <i>vma5</i> Mutants. <i>Marine Biotechnology</i> , 2001, 3, 316-321. | 1.1 | 19 |
| 4114 | Secretory expression of human growth hormone in <i>Saccharomyces cerevisiae</i> using three different leader sequences. <i>Biotechnology and Bioprocess Engineering</i> , 2001, 6, 306-309. | 1.4 | 13 |
| 4115 | Expression of nutritionally well-balanced protein, <i>AmA1</i> , in <i>Saccharomyces cerevisiae</i> . <i>Biotechnology and Bioprocess Engineering</i> , 2001, 6, 173-178. | 1.4 | 1 |
| 4116 | A 14-3-3 protein homologue is expressed in feline enteroepithelial-stages of <i>Toxoplasma gondii</i> . <i>Veterinary Parasitology</i> , 2001, 96, 65-74. | 0.7 | 18 |
| 4117 | The protein kinases <i>AtMAP3K1</i> and <i>BnMAP3K1</i> are functional homologues of <i>S. pombe cdc7p</i> and may be involved in cell division. <i>Plant Journal</i> , 2001, 26, 637-649. | 2.8 | 35 |
| 4118 | Existence of a tightly regulated water channel in <i>Saccharomyces cerevisiae</i> . <i>FEBS Journal</i> , 2001, 268, 334-343. | 0.2 | 45 |
| 4119 | Carboxyl group of residue Asp647 as possible proton donor in catalytic reaction of α -glucosidase from <i>Schizosaccharomyces pombe</i> . <i>FEBS Journal</i> , 2001, 268, 2270-2280. | 0.2 | 67 |
| 4120 | Identification and characterization of the genes for N-acetylglucosamine kinase and N-acetylglucosamine-phosphate deacetylase in the pathogenic fungus <i>Candida albicans</i> . <i>FEBS Journal</i> , 2001, 268, 2498-2505. | 0.2 | 80 |
| 4121 | The ABC transporter <i>Pdr5p</i> mediates the efflux of nonsteroidal ecdysone agonists in <i>Saccharomyces cerevisiae</i> . <i>FEBS Journal</i> , 2001, 268, 3416-3422. | 0.2 | 14 |
| 4122 | Probing the role of oligomerization in the high thermal stability of <i>Pyrococcus furiosus</i> ornithine carbamoyltransferase by site-specific mutants. <i>FEBS Journal</i> , 2001, 268, 3937-3942. | 0.2 | 30 |
| 4123 | Molecular and physiological characterization of the NAD-dependent glycerol 3-phosphate dehydrogenase in the filamentous fungus <i>Aspergillus nidulans</i> . <i>Molecular Microbiology</i> , 2001, 39, 145-157. | 1.2 | 58 |

| # | ARTICLE | IF | CITATIONS |
|------|---|-----|-----------|
| 4124 | The elimination of the yeast [PSI ⁺] prion by guanidine hydrochloride is the result of Hsp104 inactivation. <i>Molecular Microbiology</i> , 2001, 40, 1357-1369. | 1.2 | 235 |
| 4125 | Genetic analysis of the signalling pathway activated by external amino acids in <i>Saccharomyces cerevisiae</i> . <i>Molecular Microbiology</i> , 2001, 41, 489-502. | 1.2 | 54 |
| 4126 | Enhanced production of <i>Aspergillus ficuum</i> endoinulinase in <i>Saccharomyces cerevisiae</i> by using the SUC2-deletion mutation. <i>Enzyme and Microbial Technology</i> , 2001, 29, 107-110. | 1.6 | 15 |
| 4127 | The Class C Vps Complex Functions at Multiple Stages of the Vacuolar Transport Pathway. <i>Traffic</i> , 2001, 2, 476-486. | 1.3 | 142 |
| 4128 | Clathrin Interactions with C-Terminal Regions of the Yeast AP-1 $\hat{1}^2$ and $\hat{1}^3$ Subunits are Important for AP-1 Association with Clathrin Coats. <i>Traffic</i> , 2001, 2, 565-576. | 1.3 | 26 |
| 4129 | Neurotensin induces mating in <i>Saccharomyces cerevisiae</i> cells that express human neurotensin receptor type 1 in place of the endogenous pheromone receptor. <i>FEBS Journal</i> , 2001, 268, 4860-4867. | 0.2 | 14 |
| 4130 | Use of <i>Saccharomyces cerevisiae</i> in the identification of novel transcription factor DNA binding specificities. <i>Yeast</i> , 2001, 18, 445-454. | 0.8 | 2 |
| 4131 | Isolation and sequence of the MIG1 homologue from the yeast <i>Candida utilis</i> . <i>Yeast</i> , 2001, 18, 597-603. | 0.8 | 6 |
| 4132 | Identification of <i>Candida tropicalis</i> HSR1, a gene of the heat-shock factor-related family, which confers salt tolerance in <i>Saccharomyces cerevisiae</i> . <i>Yeast</i> , 2001, 18, 605-610. | 0.8 | 2 |
| 4133 | Isolation of GCR1, a major transcription factor of glycolytic genes in <i>Saccharomyces cerevisiae</i> , from <i>Kluyveromyces lactis</i> . <i>Yeast</i> , 2001, 18, 729-735. | 0.8 | 13 |
| 4134 | A single-copy suppressor of the <i>Saccharomyces cerevisiae</i> late-mitotic mutant <i>scdc15</i> and <i>dbf2</i> is encoded by the <i>Candida albicans</i> CDC14 gene. <i>Yeast</i> , 2001, 18, 849-858. | 0.8 | 3 |
| 4135 | Molecular cloning of the RPS0 gene from <i>Candida tropicalis</i> . <i>Yeast</i> , 2001, 18, 971-980. | 0.8 | 3 |
| 4136 | High efficiency transformation of <i>Schizosaccharomyces pombe</i> pretreated with thiol compounds by electroporation. <i>Yeast</i> , 2001, 18, 1015-1021. | 0.8 | 79 |
| 4137 | Meiotic crossing-over in the regions of homology between homeologous chromosomes V. <i>Yeast</i> , 2001, 18, 1173-1183. | 0.8 | 4 |
| 4138 | Functional isolation of the <i>Candida albicans</i> FCR3 gene encoding a bZip transcription factor homologous to <i>Saccharomyces cerevisiae</i> Yap3p. <i>Yeast</i> , 2001, 18, 1217-1225. | 0.8 | 26 |
| 4139 | Isolation of a <i>Candida glabrata</i> homologue of RAP1, a regulator of transcription and telomere function in <i>Saccharomyces cerevisiae</i> . <i>Yeast</i> , 2001, 18, 1277-1284. | 0.8 | 19 |
| 4140 | Mode of action of the ecdysone agonist tebufenozide (RH-5992), and an exclusion mechanism to explain resistance to it. <i>Pest Management Science</i> , 2001, 57, 951-957. | 1.7 | 52 |
| 4141 | Yeast GGA proteins interact with GTP-bound Arf and facilitate transport through the Golgi. <i>Yeast</i> , 2001, 18, 1-18. | 0.8 | 91 |

| # | ARTICLE | IF | CITATIONS |
|------|--|-----|-----------|
| 4142 | The <i>Schizosaccharomyces pombe</i> GPI8 gene complements a <i>Saccharomyces cerevisiae</i> GPI8 anchoring mutant. <i>Yeast</i> , 2001, 18, 33-39. | 0.8 | 8 |
| 4143 | Isolation of a <i>Candida albicans</i> gene, tightly linked to URA3, coding for a putative transcription factor that suppresses a <i>Saccharomyces cerevisiae</i> aft1 mutation. <i>Yeast</i> , 2001, 18, 301-311. | 0.8 | 39 |
| 4144 | Mutagenesis of γ -tubulin cysteine residues in <i>Saccharomyces cerevisiae</i> : Mutation of cysteine 354 results in cold-stable microtubules. <i>Cytoskeleton</i> , 2001, 49, 67-77. | 4.4 | 47 |
| 4145 | Expression of the small tyrosine phosphatase (Stp1) in <i>Saccharomyces cerevisiae</i> : A study on protein tyrosine phosphorylation. <i>Electrophoresis</i> , 2001, 22, 576-585. | 1.3 | 16 |
| 4146 | Functional studies of yeast actin mutants corresponding to human cardiomyopathy mutations. <i>Journal of Muscle Research and Cell Motility</i> , 2001, 22, 665-674. | 0.9 | 21 |
| 4147 | Yeast genes YOL002C and SUL1 are involved in neomycin resistance. <i>World Journal of Microbiology and Biotechnology</i> , 2001, 17, 399-402. | 1.7 | 3 |
| 4148 | Title is missing!. <i>Molecular Biology</i> , 2001, 35, 356-364. | 0.4 | 27 |
| 4149 | A putative protein inhibitor of activated STAT (PIASy) interacts with p53 and inhibits p53-mediated transactivation but not apoptosis. <i>Apoptosis: an International Journal on Programmed Cell Death</i> , 2001, 6, 221-234. | 2.2 | 61 |
| 4150 | Title is missing!. <i>Russian Journal of Genetics</i> , 2001, 37, 1014-1024. | 0.2 | 2 |
| 4151 | Autonomous Replication of a <i>Bombyx mori</i> Transgene Fragment in the Yeast <i>Saccharomyces cerevisiae</i> . <i>Russian Journal of Genetics</i> , 2001, 37, 1257-1265. | 0.2 | 0 |
| 4152 | Evidence for the attachment of Hsp150/Pir2 to the cell wall of <i>Saccharomyces cerevisiae</i> through disulfide bridges. <i>FEMS Yeast Research</i> , 2001, 1, 241-245. | 1.1 | 37 |
| 4153 | Impact of the six nucleotides downstream of the stop codon on translation termination. <i>EMBO Reports</i> , 2001, 2, 787-793. | 2.0 | 196 |
| 4154 | Ski7p G protein interacts with the exosome and the Ski complex for 3'-to-5' mRNA decay in yeast. <i>EMBO Journal</i> , 2001, 20, 4684-4693. | 3.5 | 183 |
| 4155 | An acidic sequence of a putative yeast Golgi membrane protein binds COPII and facilitates ER export. <i>EMBO Journal</i> , 2001, 20, 6742-6750. | 3.5 | 118 |
| 4156 | Cloning and sequence analysis of cnaA gene encoding the catalytic subunit of calcineurin from <i>Aspergillus oryzae</i> . <i>FEMS Microbiology Letters</i> , 2001, 204, 169-174. | 0.7 | 12 |
| 4157 | The <i>Candida albicans</i> ERG26 gene encoding the C-3 sterol dehydrogenase (C-4 decarboxylase) is essential for growth. <i>FEMS Yeast Research</i> , 2001, 1, 93-101. | 1.1 | 17 |
| 4158 | Modulating the distribution of fluxes among respiration and fermentation by overexpression of HAP4 in <i>Saccharomyces cerevisiae</i> . <i>FEMS Yeast Research</i> , 2001, 1, 139-149. | 1.1 | 41 |
| 4159 | Overexpression of a novel gene, Cms1, can rescue the growth arrest of a <i>Saccharomyces cerevisiae</i> mcm10 suppressor. <i>Cell Research</i> , 2001, 11, 285-291. | 5.7 | 1 |

| # | ARTICLE | IF | CITATIONS |
|------|--|-----|-----------|
| 4160 | Mutagenesis of dihydrofolate reductase from Plasmodium falciparum: analysis in Saccharomyces cerevisiae of triple mutant alleles resistant to pyrimethamine or WR99210. Molecular and Biochemical Parasitology, 2001, 113, 139-150. | 0.5 | 34 |
| 4161 | Novel alleles of the Plasmodium falciparum dhfr highly resistant to pyrimethamine and chlorocycloguanil, but not WR99210. Molecular and Biochemical Parasitology, 2001, 117, 91-102. | 0.5 | 44 |
| 4162 | Positive and negative regulation of squalene synthase (ERG9), an ergosterol biosynthetic gene, in Saccharomyces cerevisiae. Biochimica Et Biophysica Acta Gene Regulatory Mechanisms, 2001, 1517, 177-189. | 2.4 | 47 |
| 4163 | Characterization and regulation of Schizosaccharomyces pombe gene encoding thioredoxin. Biochimica Et Biophysica Acta Gene Regulatory Mechanisms, 2001, 1518, 194-199. | 2.4 | 13 |
| 4164 | Characterization and regulation of glutathione S-transferase gene from Schizosaccharomyces pombe. Biochimica Et Biophysica Acta Gene Regulatory Mechanisms, 2001, 1520, 179-185. | 2.4 | 22 |
| 4165 | Activation of Ty transposition by mutagens. Mutation Research - Fundamental and Molecular Mechanisms of Mutagenesis, 2001, 474, 93-103. | 0.4 | 45 |
| 4166 | Early immunodiagnosis of fasciolosis in ruminants using recombinant Fasciola hepatica cathepsin L-like protease. International Journal for Parasitology, 2001, 31, 728-737. | 1.3 | 106 |
| 4167 | Expression of a Glutamate Decarboxylase Homologue Is Required for Normal Oxidative Stress Tolerance in Saccharomyces cerevisiae. Journal of Biological Chemistry, 2001, 276, 244-250. | 1.6 | 190 |
| 4168 | Activation of the Cyclin-dependent Kinase CTDK-I Requires the Heterodimerization of Two Unstable Subunits. Journal of Biological Chemistry, 2001, 276, 8005-8013. | 1.6 | 22 |
| 4169 | The Yeast ALG11 Gene Specifies Addition of the Terminal $\hat{\pm}1,2$ -Man to the Man5GlcNAc2-PP-dolicholN-Glycosylation Intermediate Formed on the Cytosolic Side of the Endoplasmic Reticulum. Journal of Biological Chemistry, 2001, 276, 21828-21840. | 1.6 | 64 |
| 4170 | Functional Analysis of the Protein-interacting Domains of Chloroplast SRP43. Journal of Biological Chemistry, 2001, 276, 24654-24660. | 1.6 | 52 |
| 4171 | Functional Analysis of the Trypanosomal AAA ProteinTbVCP with trans-Dominant ATP Hydrolysis Mutants. Journal of Biological Chemistry, 2001, 276, 21512-21520. | 1.6 | 78 |
| 4172 | Complementing Yeast rho1 Mutation Groups with Distinct Functional Defects. Journal of Biological Chemistry, 2001, 276, 46165-46171. | 1.6 | 39 |
| 4173 | Analysis of the Pore of the Unusual Major Intrinsic Protein Channel, Yeast Fps1p. Journal of Biological Chemistry, 2001, 276, 36543-36549. | 1.6 | 27 |
| 4174 | Phosphorylation of the Yeast Phospholipid Synthesis Regulatory Protein Opi1p by Protein Kinase C. Journal of Biological Chemistry, 2001, 276, 29915-29923. | 1.6 | 50 |
| 4175 | Activation of the Ras-cAMP Signal Transduction Pathway Inhibits the Proteasome-independent Degradation of Misfolded Protein Aggregates in the Endoplasmic Reticulum Lumen. Journal of Biological Chemistry, 2001, 276, 41444-41454. | 1.6 | 15 |
| 4176 | Mediation of Elicitin Activity on Tobacco Is Assumed by Elicitin-Sterol Complexes. Molecular Biology of the Cell, 2001, 12, 2825-2834. | 0.9 | 103 |
| 4177 | TOR Modulates GCN4 -Dependent Expression of Genes Turned on by Nitrogen Limitation. Journal of Bacteriology, 2001, 183, 2331-2334. | 1.0 | 89 |

| # | ARTICLE | IF | CITATIONS |
|------|--|-----|-----------|
| 4178 | Interactions and Nuclear Import of the N and P Proteins of Sonchus Yellow Net Virus, a Plant Nucleorhabdovirus. <i>Journal of Virology</i> , 2001, 75, 9393-9406. | 1.5 | 69 |
| 4179 | The <i>Schizosaccharomyces pombe</i> spo20 ⁺ Gene Encoding a Homologue of <i>Saccharomyces cerevisiae</i> Sec14 Plays an Important Role in Forespore Membrane Formation. <i>Molecular Biology of the Cell</i> , 2001, 12, 901-917. | 0.9 | 74 |
| 4180 | Functional analysis of human MLH1 and MSH2 missense variants and hybrid human-yeast MLH1 proteins in <i>Saccharomyces cerevisiae</i> . <i>Human Molecular Genetics</i> , 2001, 10, 1889-1900. | 1.4 | 68 |
| 4181 | A Truncation Mutant of the 95-Kilodalton Subunit of Transcription Factor IIIc Reveals Asymmetry in Ty3 Integration. <i>Molecular and Cellular Biology</i> , 2001, 21, 7839-7851. | 1.1 | 32 |
| 4182 | Overproduction of PDR3 Suppresses Mitochondrial Import Defects Associated with a TOM70 Null Mutation by Increasing the Expression of TOM72 in <i>Saccharomyces cerevisiae</i> . <i>Molecular and Cellular Biology</i> , 2001, 21, 7576-7586. | 1.1 | 16 |
| 4183 | Sac1 Lipid Phosphatase and Stt4 Phosphatidylinositol 4-Kinase Regulate a Pool of Phosphatidylinositol 4-Phosphate That Functions in the Control of the Actin Cytoskeleton and Vacuole Morphology. <i>Molecular Biology of the Cell</i> , 2001, 12, 2396-2411. | 0.9 | 216 |
| 4184 | In Vitro and In Vivo Effects of 14 α -Demethylase (ERG11) Depletion in <i>Candida glabrata</i> . <i>Antimicrobial Agents and Chemotherapy</i> , 2001, 45, 3037-3045. | 1.4 | 32 |
| 4185 | Glucose-Induced Monoubiquitination of the <i>Saccharomyces cerevisiae</i> Galactose Transporter Is Sufficient To Signal Its Internalization. <i>Journal of Bacteriology</i> , 2001, 183, 3083-3088. | 1.0 | 66 |
| 4186 | Yol082p, a Novel CVT Protein Involved in the Selective Targeting of Aminopeptidase I to the Yeast Vacuole. <i>Journal of Biological Chemistry</i> , 2001, 276, 29210-29217. | 1.6 | 68 |
| 4187 | Phosphoglucomutase Is an in Vivo Lithium Target in Yeast. <i>Journal of Biological Chemistry</i> , 2001, 276, 37794-37801. | 1.6 | 73 |
| 4188 | Yeast Cdc42 functions at a late step in exocytosis, specifically during polarized growth of the emerging bud. <i>Journal of Cell Biology</i> , 2001, 155, 581-592. | 2.3 | 151 |
| 4189 | Wsc1 and Mid2 Are Cell Surface Sensors for Cell Wall Integrity Signaling That Act through Rom2, a Guanine Nucleotide Exchange Factor for Rho1. <i>Molecular and Cellular Biology</i> , 2001, 21, 271-280. | 1.1 | 288 |
| 4190 | Ras and rap 1 interaction with af-6 effector target. <i>Methods in Enzymology</i> , 2001, 332, 151-168. | 0.4 | 17 |
| 4192 | The GATA Transcription Factors GLN3 and GAT1 Link TOR to Salt Stress in <i>Saccharomyces cerevisiae</i> . <i>Journal of Biological Chemistry</i> , 2001, 276, 34441-34444. | 1.6 | 84 |
| 4193 | Circular Minichromosomes Become Highly Recombinogenic in Topoisomerase-deficient Yeast Cells. <i>Journal of Biological Chemistry</i> , 2001, 276, 2243-2248. | 1.6 | 14 |
| 4194 | Mutations in the TATA-binding Protein, Affecting Transcriptional Activation, Show Synthetic Lethality with the TAF145 Gene Lacking the TAF N-terminal Domain in <i>Saccharomyces cerevisiae</i> . <i>Journal of Biological Chemistry</i> , 2001, 276, 395-405. | 1.6 | 29 |
| 4195 | Cloning and Characterization of Liver-specific Isoform of Chk1 Gene from Rat. <i>Journal of Biological Chemistry</i> , 2001, 276, 48863-48870. | 1.6 | 13 |
| 4196 | Isolation and Characterization of a Gene Specific to Lager Brewing Yeast That Encodes a Branched-Chain Amino Acid Permease. <i>Applied and Environmental Microbiology</i> , 2001, 67, 3455-3462. | 1.4 | 43 |

| # | ARTICLE | IF | CITATIONS |
|------|---|-----|-----------|
| 4197 | GDP Dissociation Inhibitor Domain II Required for Rab GTPase Recycling. <i>Journal of Biological Chemistry</i> , 2001, 276, 8014-8020. | 1.6 | 19 |
| 4199 | Quantitation of the RNA Polymerase II Transcription Machinery in Yeast. <i>Journal of Biological Chemistry</i> , 2001, 276, 47150-47153. | 1.6 | 62 |
| 4200 | Molecular cloning of a cytochrome P450 taxane 10 \hat{A} -hydroxylase cDNA from <i>Taxus</i> and functional expression in yeast. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 2001, 98, 1501-1506. | 3.3 | 174 |
| 4201 | Human BIN3 Complements the F-actin Localization Defects Caused by Loss of Hob3p, the Fission Yeast Homolog of Rvs161p. <i>Journal of Biological Chemistry</i> , 2001, 276, 21670-21677. | 1.6 | 23 |
| 4202 | Characterization of Transsulfuration and Cysteine Biosynthetic Pathways in the Protozoan Hemoflagellate, <i>Trypanosoma cruzi</i> . <i>Journal of Biological Chemistry</i> , 2001, 276, 6516-6523. | 1.6 | 98 |
| 4203 | Diazaborine Treatment of Baker's Yeast Results in Stabilization of Aberrant mRNAs. <i>Journal of Biological Chemistry</i> , 2001, 276, 36419-36424. | 1.6 | 14 |
| 4204 | The Effect of the <i>erg26-1</i> Mutation on the Regulation of Lipid Metabolism in <i>Saccharomyces cerevisiae</i> . <i>Journal of Biological Chemistry</i> , 2001, 276, 12702-12711. | 1.6 | 53 |
| 4205 | Dimer Formation of Octaprenyl-diphosphate Synthase (IspB) Is Essential for Chain Length Determination of Ubiquinone. <i>Journal of Biological Chemistry</i> , 2001, 276, 7876-7883. | 1.6 | 52 |
| 4206 | Role of Arg-166 in Yeast Cytochrome c1. <i>Journal of Biological Chemistry</i> , 2001, 276, 18450-18456. | 1.6 | 5 |
| 4207 | Expression of V _{HH} Antibody Fragments in <i>Saccharomyces cerevisiae</i> . , 2002, 178, 359-366. | | 12 |
| 4208 | Saturation Mutagenesis of 5S rRNA in <i>Saccharomyces cerevisiae</i> . <i>Molecular and Cellular Biology</i> , 2001, 21, 8264-8275. | 1.1 | 47 |
| 4209 | An Oxysterol-derived Positive Signal for 3-Hydroxy-3-methylglutaryl-CoA Reductase Degradation in Yeast. <i>Journal of Biological Chemistry</i> , 2001, 276, 8681-8694. | 1.6 | 48 |
| 4210 | Ubiquitin Is Required for Sorting to the Vacuole of the Yeast General Amino Acid Permease, Gap1. <i>Journal of Biological Chemistry</i> , 2001, 276, 43949-43957. | 1.6 | 208 |
| 4211 | NADP-Glutamate Dehydrogenase Isoenzymes of <i>Saccharomyces cerevisiae</i> . <i>Journal of Biological Chemistry</i> , 2001, 276, 43775-43783. | 1.6 | 142 |
| 4212 | Stalk Segment 5 of the Yeast Plasma Membrane H ⁺ -ATPase. <i>Journal of Biological Chemistry</i> , 2001, 276, 22485-22490. | 1.6 | 16 |
| 4213 | A protein interaction map for cell polarity development. <i>Journal of Cell Biology</i> , 2001, 154, 549-576. | 2.3 | 294 |
| 4214 | Regulation of the <i>Saccharomyces cerevisiae</i> DPP1-encoded Diacylglycerol Pyrophosphate Phosphatase by Zinc. <i>Journal of Biological Chemistry</i> , 2001, 276, 10126-10133. | 1.6 | 57 |
| 4215 | The <i>Schizosaccharomyces pombe spo3⁺</i> Gene Is Required for Assembly of the Forespore Membrane and Genetically Interacts with <i>psy1⁺</i> -encoding Syntaxin-like Protein. <i>Molecular Biology of the Cell</i> , 2001, 12, 3955-3972. | 0.9 | 131 |

| # | ARTICLE | IF | CITATIONS |
|------|--|-----|-----------|
| 4216 | Screening for Modulators of Spermine Tolerance Identifies Sky1, the SR Protein Kinase of <i>Saccharomyces cerevisiae</i> , as a Regulator of Polyamine Transport and Ion Homeostasis. <i>Molecular and Cellular Biology</i> , 2001, 21, 175-184. | 1.1 | 47 |
| 4217 | Genetic and Biochemical Analysis of the Yeast Plasma Membrane Ssy1p-Ptr3p-Ssy5p Sensor of Extracellular Amino Acids. <i>Molecular and Cellular Biology</i> , 2001, 21, 814-826. | 1.1 | 141 |
| 4218 | O2R, a Novel Regulatory Element Mediating Rox1p-Independent O ₂ and Unsaturated Fatty Acid Repression of OLE1 in <i>Saccharomyces cerevisiae</i> . <i>Journal of Bacteriology</i> , 2001, 183, 745-751. | 1.0 | 33 |
| 4219 | Evidence for an Intrinsic Toxicity of Phosphatidylcholine to Sec14p-dependent Protein Transport from the Yeast Golgi Complex. <i>Molecular Biology of the Cell</i> , 2001, 12, 1117-1129. | 0.9 | 60 |
| 4220 | Mitochondrial DNA ligase function in <i>Saccharomyces cerevisiae</i> . <i>Nucleic Acids Research</i> , 2001, 29, 1582-1589. | 6.5 | 29 |
| 4221 | High-Efficiency Transformation of Plasmid DNA into Yeast. , 2001, 177, 085-097. | | 41 |
| 4222 | The Transcriptional Activation Domain of the Plant-Specific Dof1 Factor Functions in Plant, Animal, and Yeast Cells. <i>Plant and Cell Physiology</i> , 2001, 42, 813-822. | 1.5 | 56 |
| 4223 | Transcriptional Autoregulation and Inhibition of mRNA Translation of Amino Acid Regulator Gene <i>cpcA</i> of Filamentous Fungus <i>Aspergillus nidulans</i> . <i>Molecular Biology of the Cell</i> , 2001, 12, 2846-2857. | 0.9 | 116 |
| 4224 | Vps41p Function in the Alkaline Phosphatase Pathway Requires Homo-oligomerization and Interaction with AP-3 through Two Distinct Domains. <i>Molecular Biology of the Cell</i> , 2001, 12, 37-51. | 0.9 | 80 |
| 4225 | Domain Interactions in the Yeast ATP Binding Cassette Transporter Ycf1p: Intragenic Suppressor Analysis of Mutations in the Nucleotide Binding Domains. <i>Journal of Bacteriology</i> , 2001, 183, 4761-4770. | 1.0 | 23 |
| 4226 | Effects of Anoxia and the Mitochondrion on Expression of Aerobic Nuclear COX Genes in Yeast. <i>Journal of Biological Chemistry</i> , 2001, 276, 7593-7601. | 1.6 | 53 |
| 4227 | Early Cell Cycle Box-Mediated Transcription of CLN3 and SWI4 Contributes to the Proper Timing of the G ₁ -to-S Transition in Budding Yeast. <i>Molecular and Cellular Biology</i> , 2001, 21, 4140-4148. | 1.1 | 49 |
| 4228 | Yeast Mps1p Phosphorylates the Spindle Pole Component Spc110p in the N-terminal Domain. <i>Journal of Biological Chemistry</i> , 2001, 276, 17958-17967. | 1.6 | 28 |
| 4229 | The Basic Helix-Loop-Helix Transcription Factor Cph2 Regulates Hyphal Development in <i>Candida albicans</i> Partly via Tec1. <i>Molecular and Cellular Biology</i> , 2001, 21, 6418-6428. | 1.1 | 110 |
| 4230 | Repression of GCN4 mRNA Translation by Nitrogen Starvation in <i>Saccharomyces cerevisiae</i> . <i>Journal of Biological Chemistry</i> , 2001, 276, 25661-25671. | 1.6 | 42 |
| 4231 | The Npr1 Kinase Controls Biosynthetic and Endocytic Sorting of the Yeast Gap1 Permease. <i>Journal of Biological Chemistry</i> , 2001, 276, 43939-43948. | 1.6 | 147 |
| 4232 | Enhanced Iron Uptake of <i>Saccharomyces cerevisiae</i> by Heterologous Expression of a Tadpole Ferritin Gene. <i>Applied and Environmental Microbiology</i> , 2001, 67, 1280-1283. | 1.4 | 29 |
| 4233 | Topological and Mutational Analysis of <i>Saccharomyces cerevisiae</i> Ste14p, Founding Member of the Isoprenylcysteine Carboxyl Methyltransferase Family. <i>Molecular Biology of the Cell</i> , 2001, 12, 1957-1971. | 0.9 | 70 |

| # | ARTICLE | IF | CITATIONS |
|------|---|-----|-----------|
| 4234 | Repression of rRNA synthesis due to a secretory defect requires the C-terminal silencing domain of Rap1p in <i>Saccharomyces cerevisiae</i> . <i>Nucleic Acids Research</i> , 2001, 29, 3297-3303. | 6.5 | 19 |
| 4235 | Cross-talk between Transcriptional Regulators of Multidrug Resistance in <i>Saccharomyces cerevisiae</i> . <i>Journal of Biological Chemistry</i> , 2001, 276, 8812-8819. | 1.6 | 48 |
| 4236 | Yak1p, a DYRK family kinase, translocates to the nucleus and phosphorylates yeast Pop2p in response to a glucose signal. <i>Genes and Development</i> , 2001, 15, 1217-1228. | 2.7 | 138 |
| 4237 | Substrate Recognition by UDP-galactose and CMP-sialic Acid Transporters. <i>Journal of Biological Chemistry</i> , 2001, 276, 21555-21561. | 1.6 | 55 |
| 4238 | Upc2p and Ecm22p, Dual Regulators of Sterol Biosynthesis in <i>Saccharomyces cerevisiae</i> . <i>Molecular and Cellular Biology</i> , 2001, 21, 6395-6405. | 1.1 | 238 |
| 4239 | Hsp70 Molecular Chaperone Facilitates Endoplasmic Reticulum-associated Protein Degradation of Cystic Fibrosis Transmembrane Conductance Regulator in Yeast. <i>Molecular Biology of the Cell</i> , 2001, 12, 1303-1314. | 0.9 | 260 |
| 4240 | Cdc5 Interacts with the Wee1 Kinase in Budding Yeast. <i>Molecular and Cellular Biology</i> , 2001, 21, 4949-4959. | 1.1 | 48 |
| 4241 | Discrete Targeting Signals Direct Pmp47 to Oleate-induced Peroxisomes in <i>Saccharomyces cerevisiae</i> . <i>Journal of Biological Chemistry</i> , 2001, 276, 10897-10905. | 1.6 | 55 |
| 4242 | Translational control of human p53 expression in yeast mediated by 5'-UTR-ORF structural interaction. <i>Nucleic Acids Research</i> , 2001, 29, 1222-1227. | 6.5 | 23 |
| 4243 | Transcriptional Regulation of the Two Sterol Esterification Genes in the Yeast <i>Saccharomyces cerevisiae</i> . <i>Journal of Bacteriology</i> , 2001, 183, 4950-4957. | 1.0 | 53 |
| 4244 | The Phosphotyrosyl Phosphatase Activator, Ncs1p (Rrd1p), Functions with Cla4p To Regulate the G ₂ /M Transition in <i>Saccharomyces cerevisiae</i> . <i>Molecular and Cellular Biology</i> , 2001, 21, 488-500. | 1.1 | 41 |
| 4245 | In Vivo Action of the HRD Ubiquitin Ligase Complex: Mechanisms of Endoplasmic Reticulum Quality Control and Sterol Regulation. <i>Molecular and Cellular Biology</i> , 2001, 21, 4276-4291. | 1.1 | 113 |
| 4246 | The CWH8 Gene Encodes a Dolichyl Pyrophosphate Phosphatase with a Luminally Oriented Active Site in the Endoplasmic Reticulum of <i>Saccharomyces cerevisiae</i> . <i>Journal of Biological Chemistry</i> , 2001, 276, 41455-41464. | 1.6 | 65 |
| 4247 | New secretory strategies for <i>Kluyveromyces lactis</i> Î ² -galactosidase. <i>Protein Engineering, Design and Selection</i> , 2001, 14, 379-386. | 1.0 | 39 |
| 4248 | Interaction of Cellular Apoptosis Regulating Proteins with Adenovirus Anti-apoptosis Protein E1B-19K. <i>Journal of Biological Chemistry</i> , 2001, 276, 211-218. | | 3 |
| 4249 | Rapid Degradation of Auxin/Indoleacetic Acid Proteins Requires Conserved Amino Acids of Domain II and Is Proteasome Dependent. <i>Plant Cell</i> , 2001, 13, 2349-2360. | 3.1 | 260 |
| 4250 | A Novel Screening for Inhibitors of a Pleiotropic Drug Resistant Pump, Pdr5, in <i>Saccharomyces cerevisiae</i> . <i>Bioscience, Biotechnology and Biochemistry</i> , 2001, 65, 1589-1595. | 0.6 | 13 |
| 4251 | Insect Immunity. <i>Journal of Biological Chemistry</i> , 2001, 276, 4085-4092. | 1.6 | 210 |

| # | ARTICLE | IF | CITATIONS |
|------|---|-----|-----------|
| 4252 | Functional analysis of R4D4 gene required for nucleotide excision repair of UV-induced DNA damage in <i>Saccharomyces cerevisiae</i> . <i>Korean Journal of Biological Sciences</i> , 2002, 6, 311-315. | 0.1 | 1 |
| 4253 | <i>Saccharomyces cerevisiae</i> Mer3 Is a DNA Helicase Involved in Meiotic Crossing Over. <i>Molecular and Cellular Biology</i> , 2002, 22, 3281-3291. | 1.1 | 59 |
| 4254 | Nob1p is required for biogenesis of the 26S proteasome and degraded upon its maturation in <i>Saccharomyces cerevisiae</i> . <i>Genes and Development</i> , 2002, 16, 3142-3157. | 2.7 | 80 |
| 4255 | The SUMO Ubiquitin-Protein Isopeptide Ligase Family Member Miz1/PIASx1/Siz2 Is a Transcriptional Cofactor for TFII-I. <i>Journal of Biological Chemistry</i> , 2002, 277, 43185-43193. | 1.6 | 31 |
| 4256 | Functional and Molecular Characterization of Nucleobase Transport by Recombinant Human and Rat Equilibrative Nucleoside Transporters 1 and 2. <i>Journal of Biological Chemistry</i> , 2002, 277, 24938-24948. | 1.6 | 157 |
| 4257 | In Vivo and in Vitro Phosphorylation of Two Isoforms of Yeast Pyruvate Kinase by Protein Kinase A. <i>Journal of Biological Chemistry</i> , 2002, 277, 30477-30487. | 1.6 | 50 |
| 4258 | The Arabidopsis AtSTE24 Is a CAAX Protease with Broad Substrate Specificity. <i>Journal of Biological Chemistry</i> , 2002, 277, 29856-29864. | 1.6 | 62 |
| 4259 | Refined molecular hinge between allosteric and catalytic domain determines allosteric regulation and stability of fungal chorismate mutase. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 2002, 99, 6631-6636. | 3.3 | 12 |
| 4260 | Identification of Interdependent Signals Required for Anterograde Traffic of the ATP-binding Cassette Transporter Protein Yor1p. <i>Journal of Biological Chemistry</i> , 2002, 277, 34860-34869. | 1.6 | 23 |
| 4261 | Stalk Segment 5 of the Yeast Plasma Membrane H ⁺ -ATPase. <i>Journal of Biological Chemistry</i> , 2002, 277, 40981-40988. | 1.6 | 5 |
| 4262 | The Initiation Factor eIF4A Is Involved in the Response to Lithium Stress in <i>Saccharomyces cerevisiae</i> . <i>Journal of Biological Chemistry</i> , 2002, 277, 21542-21548. | 1.6 | 43 |
| 4263 | Critical Function of the Ras-associating Domain as a Primary Ras-binding Site for Regulation of <i>Saccharomyces cerevisiae</i> Adenylyl Cyclase. <i>Journal of Biological Chemistry</i> , 2002, 277, 3117-3123. | 1.6 | 26 |
| 4264 | Neither Reb1p nor Poly(dA-dT) Elements Are Responsible for the Highly Specific Chromatin Organization at the ILV1 Promoter. <i>Journal of Biological Chemistry</i> , 2002, 277, 3202-3209. | 1.6 | 12 |
| 4265 | Disease mechanism for retinitis pigmentosa (RP11) caused by mutations in the splicing factor gene PRPF31. <i>Human Molecular Genetics</i> , 2002, 11, 3209-3219. | 1.4 | 75 |
| 4266 | Overexpression, purification, and partial characterization of <i>Saccharomyces cerevisiae</i> processing alpha glucosidase I. <i>Glycobiology</i> , 2002, 12, 229-234. | 1.3 | 32 |
| 4267 | Identification of a Novel Mitogen-Activated Protein Kinase Kinase Activation Domain Recognized by the Inhibitor PD 184352. <i>Molecular and Cellular Biology</i> , 2002, 22, 7593-7602. | 1.1 | 64 |
| 4268 | Normal Assembly of 60 S Ribosomal Subunits Is Required for the Signaling in Response to a Secretory Defect in <i>Saccharomyces cerevisiae</i> . <i>Journal of Biological Chemistry</i> , 2002, 277, 18334-18339. | 1.6 | 46 |
| 4269 | Actin cable dynamics in budding yeast. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 2002, 99, 751-756. | 3.3 | 180 |

| # | ARTICLE | IF | CITATIONS |
|------|---|-----|-----------|
| 4270 | Isolation and Functional Analysis of a Gene, <i>tcsB</i> , Encoding a Transmembrane Hybrid-Type Histidine Kinase from <i>Aspergillus nidulans</i> . <i>Applied and Environmental Microbiology</i> , 2002, 68, 5304-5310. | 1.4 | 80 |
| 4271 | Lack of Pseudouridine 38/39 in the Anticodon Arm of Yeast Cytoplasmic tRNA Decreases in Vivo Recoding Efficiency. <i>Journal of Biological Chemistry</i> , 2002, 277, 30445-30453. | 1.6 | 66 |
| 4272 | Phosphorylation of <i>Saccharomyces cerevisiae</i> Choline Kinase on Ser30 and Ser85 by Protein Kinase A Regulates Phosphatidylcholine Synthesis by the CDP-choline Pathway. <i>Journal of Biological Chemistry</i> , 2002, 277, 34978-34986. | 1.6 | 39 |
| 4273 | Geranyl Diphosphate:4-Hydroxybenzoate Geranyltransferase from <i>Lithospermum erythrorhizon</i> . <i>Journal of Biological Chemistry</i> , 2002, 277, 6240-6246. | 1.6 | 133 |
| 4274 | Pyrimethamine and WR99210 exert opposing selection on dihydrofolate reductase from <i>Plasmodium vivax</i> . <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 2002, 99, 13137-13141. | 3.3 | 79 |
| 4275 | Overexpression, purification, and site-directed spin labeling of the Nrap metal transporter from <i>Mycobacterium leprae</i> . <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 2002, 99, 8608-8613. | 3.3 | 19 |
| 4276 | Novel G-Protein Complex Whose Requirement Is Linked to the Translational Status of the Cell. <i>Molecular and Cellular Biology</i> , 2002, 22, 2564-2574. | 1.1 | 67 |
| 4277 | The Retinoblastoma Family of Proteins Directly Represses Transcription in <i>Saccharomyces cerevisiae</i> . <i>Journal of Biological Chemistry</i> , 2002, 277, 8797-8801. | 1.6 | 5 |
| 4278 | GLFG and FxFG Nucleoporins Bind to Overlapping Sites on Importin- β . <i>Journal of Biological Chemistry</i> , 2002, 277, 50597-50606. | 1.6 | 203 |
| 4279 | Analysis of oxysterol binding protein homologue Kes1p function in regulation of Sec14p-dependent protein transport from the yeast Golgi complex. <i>Journal of Cell Biology</i> , 2002, 157, 63-78. | 2.3 | 219 |
| 4280 | [12] Role of yeast flavin-containing monooxygenase in maintenance of thiol-disulfide redox potential. <i>Methods in Enzymology</i> , 2002, 348, 113-121. | 0.4 | 8 |
| 4281 | Wine Yeast Strains Engineered for Glycogen Overproduction Display Enhanced Viability under Glucose Deprivation Conditions. <i>Applied and Environmental Microbiology</i> , 2002, 68, 3339-3344. | 1.4 | 43 |
| 4282 | Engineering of Polyploid <i>Saccharomyces cerevisiae</i> for Secretion of Large Amounts of Fungal Glucoamylase. <i>Applied and Environmental Microbiology</i> , 2002, 68, 5693-5697. | 1.4 | 13 |
| 4283 | Gene Expression Analysis of Cold and Freeze Stress in Baker's Yeast. <i>Applied and Environmental Microbiology</i> , 2002, 68, 3024-3030. | 1.4 | 59 |
| 4284 | Yeast Rpi1 Is a Putative Transcriptional Regulator That Contributes to Preparation for Stationary Phase. <i>Eukaryotic Cell</i> , 2002, 1, 56-65. | 3.4 | 21 |
| 4285 | Complex transcriptional circuitry at the G1/S transition in <i>Saccharomyces cerevisiae</i> . <i>Genes and Development</i> , 2002, 16, 3017-3033. | 2.7 | 236 |
| 4286 | Characterization of a Hyperthermophilic P-type ATPase from <i>Methanococcus jannaschii</i> Expressed in Yeast. <i>Journal of Biological Chemistry</i> , 2002, 277, 29608-29616. | 1.6 | 24 |
| 4287 | Yeast Cells Lacking the ARV1 Gene Harbor Defects in Sphingolipid Metabolism. <i>Journal of Biological Chemistry</i> , 2002, 277, 36152-36160. | 1.6 | 59 |

| # | ARTICLE | IF | CITATIONS |
|------|---|-----|-----------|
| 4288 | Molecular and Biochemical Characterization of Two Plant Inositol Polyphosphate 6-/3-/5-Kinases. <i>Journal of Biological Chemistry</i> , 2002, 277, 42711-42718. | 1.6 | 120 |
| 4289 | Myosin I Is Required for Hypha Formation in <i>Candida albicans</i> . <i>Eukaryotic Cell</i> , 2002, 1, 213-228. | 3.4 | 68 |
| 4290 | Novel <i>Saccharomyces cerevisiae</i> Screen Identifies WR99210 Analogues That Inhibit <i>Mycobacterium tuberculosis</i> Dihydrofolate Reductase. <i>Antimicrobial Agents and Chemotherapy</i> , 2002, 46, 3362-3369. | 1.4 | 54 |
| 4291 | Regulation of Fab1 Phosphatidylinositol 3-Phosphate 5-Kinase Pathway by Vac7 Protein and Fig4, a Polyphosphoinositide Phosphatase Family Member. <i>Molecular Biology of the Cell</i> , 2002, 13, 1238-1251. | 0.9 | 159 |
| 4292 | Identification of Putative Direct Effectors for $G_{i\pm o}$, Using Yeast Two-Hybrid Method. <i>Methods in Enzymology</i> , 2002, 345, 140-149. | 0.4 | 2 |
| 4293 | The Complementary Strand of the Human T-Cell Leukemia Virus Type 1 RNA Genome Encodes a bZIP Transcription Factor That Down-Regulates Viral Transcription. <i>Journal of Virology</i> , 2002, 76, 12813-12822. | 1.5 | 444 |
| 4294 | Retromer function in endosome-to-Golgi retrograde transport is regulated by the yeast Vps34 PtdIns 3-kinase. <i>Journal of Cell Science</i> , 2002, 115, 3889-3900. | 1.2 | 201 |
| 4295 | A developmentally regulated ARF-like 5 protein (ARL5), localized to nuclei and nucleoli, interacts with heterochromatin protein 1. <i>Journal of Cell Science</i> , 2002, 115, 4433-4445. | 1.2 | 41 |
| 4296 | Gene characterized for membrane desaturase that produces (E)-11 isomers of mono- and diunsaturated fatty acids. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 2002, 99, 620-624. | 3.3 | 63 |
| 4297 | A Chinese Cabbage cDNA with High Sequence Identity to Phospholipid Hydroperoxide Glutathione Peroxidases Encodes a Novel Isoform of Thioredoxin-dependent Peroxidase. <i>Journal of Biological Chemistry</i> , 2002, 277, 12572-12578. | 1.6 | 106 |
| 4298 | In the Absence of the First Membrane-spanning Segment of Subunit4(b), the Yeast ATP Synthase Is Functional but Does Not Dimerize or Oligomerize. <i>Journal of Biological Chemistry</i> , 2002, 277, 10739-10745. | 1.6 | 76 |
| 4299 | Quality Control in the Yeast Secretory Pathway. <i>Journal of Biological Chemistry</i> , 2002, 277, 21027-21040. | 1.6 | 38 |
| 4300 | Differential Role of the Proline-rich Domain of Nuclear Factor 1-C Splice Variants in DNA Binding and Transactivation. <i>Journal of Biological Chemistry</i> , 2002, 277, 16383-16390. | 1.6 | 7 |
| 4301 | Cloning and Characterization of Methenyltetrahydrofolate Synthetase from <i>Saccharomyces cerevisiae</i> . <i>Journal of Biological Chemistry</i> , 2002, 277, 20205-20213. | 1.6 | 37 |
| 4302 | Rpf2p, an Evolutionarily Conserved Protein, Interacts with Ribosomal Protein L11 and Is Essential for the Processing of 27 SB Pre-rRNA to 25 S rRNA and the 60 S Ribosomal Subunit Assembly in <i>Saccharomyces cerevisiae</i> . <i>Journal of Biological Chemistry</i> , 2002, 277, 28780-28786. | 1.6 | 46 |
| 4303 | BLOC-1, a Novel Complex Containing the Pallidin and Muted Proteins Involved in the Biogenesis of Melanosomes and Platelet-dense Granules. <i>Journal of Biological Chemistry</i> , 2002, 277, 28191-28199. | 1.6 | 154 |
| 4304 | A Novel Membrane Protein, Ros3p, Is Required for Phospholipid Translocation across the Plasma Membrane in <i>Saccharomyces cerevisiae</i> . <i>Journal of Biological Chemistry</i> , 2002, 277, 37855-37862. | 1.6 | 117 |
| 4305 | The Synthesis of Inositol Hexakisphosphate. <i>Journal of Biological Chemistry</i> , 2002, 277, 31857-31862. | 1.6 | 84 |

| # | ARTICLE | IF | CITATIONS |
|------|--|-----|-----------|
| 4306 | Failure to Relax Negative Supercoiling of DNA Is a Primary Cause of Mitotic Hyper-recombination in Topoisomerase-deficient Yeast Cells. <i>Journal of Biological Chemistry</i> , 2002, 277, 37207-37211. | 1.6 | 28 |
| 4307 | Characterization of Mammalian eIF2A and Identification of the Yeast Homolog. <i>Journal of Biological Chemistry</i> , 2002, 277, 37079-37087. | 1.6 | 64 |
| 4308 | The <i>Saccharomyces cerevisiae</i> LSB6 Gene Encodes Phosphatidylinositol 4-Kinase Activity. <i>Journal of Biological Chemistry</i> , 2002, 277, 47709-47718. | 1.6 | 75 |
| 4309 | A Homologue of CROC-1 in a Ciliated Protist (<i>Sterkiella histriomuscorum</i>) Testifies to the Ancient Origin of the Ubiquitin-conjugating Enzyme Variant Family. <i>Molecular Biology and Evolution</i> , 2002, 19, 39-48. | 3.5 | 19 |
| 4310 | Recruitment of Gcn5-containing Complexes during c-Myc-dependent Gene Activation. <i>Journal of Biological Chemistry</i> , 2002, 277, 23399-23406. | 1.6 | 27 |
| 4311 | Production in yeast of β -galactosidase A, a lysosomal enzyme applicable to enzyme replacement therapy for Fabry disease. <i>Glycobiology</i> , 2002, 12, 821-828. | 1.3 | 56 |
| 4312 | Oligomerization and Cooperative RNA Synthesis Activity of Hepatitis C Virus RNA-Dependent RNA Polymerase. <i>Journal of Virology</i> , 2002, 76, 3865-3872. | 1.5 | 132 |
| 4313 | ADP-Ribosylation Factor (ARF) Interaction Is Not Sufficient for Yeast GGA Protein Function or Localization. <i>Molecular Biology of the Cell</i> , 2002, 13, 3078-3095. | 0.9 | 36 |
| 4314 | Sterol-dependent Regulation of Sphingolipid Metabolism in <i>Saccharomyces cerevisiae</i> . <i>Journal of Biological Chemistry</i> , 2002, 277, 26177-26184. | 1.6 | 53 |
| 4315 | The Human Homolog of the Rat Inositol Phosphate Multikinase Is an Inositol 1,3,4,6-Tetrakisphosphate 5-Kinase. <i>Journal of Biological Chemistry</i> , 2002, 277, 43836-43843. | 1.6 | 65 |
| 4316 | Chromatin Structural Analyses of the Mouse <i>Igf1^{re}</i> Gene Locus Reveal New Hypersensitive Sites Specifying a Transcriptional Silencer and Enhancer. <i>Journal of Biological Chemistry</i> , 2002, 277, 32640-32649. | 1.6 | 61 |
| 4317 | The Core Protein of Hepatitis C Virus Is Imported into the Nucleus by Transport Receptor Kap123p but Inhibits Kap121p-dependent Nuclear Import of Yeast API-like Transcription Factor in Yeast Cells. <i>Journal of Biological Chemistry</i> , 2002, 277, 39634-39641. | 1.6 | 26 |
| 4318 | Cofilin, But Not Profilin, Is Required for Myosin-I-Induced Actin Polymerization and the Endocytic Uptake in Yeast. <i>Molecular Biology of the Cell</i> , 2002, 13, 4074-4087. | 0.9 | 32 |
| 4319 | The Eukaryotic Two-Component Histidine Kinase Sln1p Regulates OCH1 via the Transcription Factor, Skn7p. <i>Molecular Biology of the Cell</i> , 2002, 13, 412-424. | 0.9 | 93 |
| 4320 | Function of a Plant Stress-Induced Gene, HVA22. Synthetic Enhancement Screen with Its Yeast Homolog Reveals Its Role in Vesicular Traffic. <i>Plant Physiology</i> , 2002, 130, 1121-1131. | 2.3 | 64 |
| 4321 | HTL1 Encodes a Novel Factor That Interacts with the RSC Chromatin Remodeling Complex in <i>Saccharomyces cerevisiae</i> . <i>Molecular and Cellular Biology</i> , 2002, 22, 8165-8174. | 1.1 | 27 |
| 4322 | Intragenic Promoter Adaptation and Facilitated RNA Polymerase III Recycling in the Transcription of SCR1, the 7SL RNA Gene of <i>Saccharomyces cerevisiae</i> . <i>Journal of Biological Chemistry</i> , 2002, 277, 6903-6914. | 1.6 | 43 |
| 4323 | Yeast Protein Kinases and the RHO1 Exchange Factor TUS1 Are Novel Components of the Cell Integrity Pathway in Yeast. <i>Molecular and Cellular Biology</i> , 2002, 22, 1329-1339. | 1.1 | 127 |

| # | ARTICLE | IF | CITATIONS |
|------|---|------|-----------|
| 4324 | Transcriptional Profiling Identifies Two Members of the ATP-binding Cassette Transporter Superfamily Required for Sterol Uptake in Yeast. <i>Journal of Biological Chemistry</i> , 2002, 277, 32466-32472. | 1.6 | 225 |
| 4325 | Large-Scale Identification of Putative Exported Proteins in <i>Candida albicans</i> by Genetic Selection. <i>Eukaryotic Cell</i> , 2002, 1, 514-525. | 3.4 | 25 |
| 4326 | Removal of a Single α -Tubulin Gene Intron Suppresses Cell Cycle Arrest Phenotypes of Splicing Factor Mutations in <i>Saccharomyces cerevisiae</i> . <i>Molecular and Cellular Biology</i> , 2002, 22, 801-815. | 1.1 | 69 |
| 4327 | The DGA1 Gene Determines a Second Triglyceride Synthetic Pathway in Yeast. <i>Journal of Biological Chemistry</i> , 2002, 277, 8877-8881. | 1.6 | 285 |
| 4328 | A Novel Connection between the Yeast Cdc42 GTPase and the Slt2-mediated Cell Integrity Pathway Identified through the Effect of Secreted <i>Salmonella</i> GTPase Modulators. <i>Journal of Biological Chemistry</i> , 2002, 277, 27094-27102. | 1.6 | 26 |
| 4329 | Pheromone induction promotes Ste11 degradation through a MAPK feedback and ubiquitin-dependent mechanism. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 2002, 99, 9160-9165. | 3.3 | 49 |
| 4330 | Novel PtdIns(3)P-binding protein Etf1 functions as an effector of the Vps34 PtdIns 3-kinase in autophagy. <i>Journal of Cell Biology</i> , 2002, 158, 761-772. | 2.3 | 85 |
| 4331 | Characterization of the <i>Saccharomyces cerevisiae</i> YMR318C (ADH6) gene product as a broad specificity NADPH-dependent alcohol dehydrogenase: relevance in aldehyde reduction. <i>Biochemical Journal</i> , 2002, 361, 163. | 1.7 | 100 |
| 4332 | Cloning and characterization of the 5' flanking region of the oxalate decarboxylase gene from <i>Flammulina velutipes</i> . <i>Biochemical Journal</i> , 2002, 367, 67-75. | 1.7 | 24 |
| 4333 | Isolation of Temperature-sensitive <i>Saccharomyces cerevisiae</i> with a Mutation in <i>erg25</i> for C-4 Sterol Methyl Oxidase.. <i>Journal of Antibiotics</i> , 2002, 55, 962-968. | 1.0 | 1 |
| 4334 | The GCR2 Gene Is Required for the Transcriptional Activation of Retrotransposon Ty2-917 in <i>Saccharomyces cerevisiae</i> .. <i>Biological and Pharmaceutical Bulletin</i> , 2002, 25, 1212-1213. | 0.6 | 4 |
| 4335 | Genetic characterization of genes encoding enzymes catalyzing addition of phospho-ethanolamine to the glycosylphosphatidylinositol anchor in <i>Saccharomyces cerevisiae</i> .. <i>Genes and Genetic Systems</i> , 2002, 77, 309-322. | 0.2 | 15 |
| 4336 | Problems with Methods used to Screen Estrogenic Chemicals by Yeast Two-Hybrid Assays.. <i>Journal of Health Science</i> , 2002, 48, 83-88. | 0.9 | 4 |
| 4337 | Functional Assays for Mammalian G-Protein-Coupled Receptors in Yeast. <i>Methods in Enzymology</i> , 2002, 344, 92-111. | 0.4 | 16 |
| 4338 | Subcellular localization of the yeast proteome. <i>Genes and Development</i> , 2002, 16, 707-719. | 2.7 | 667 |
| 4339 | Mga2p Is a Putative Sensor for Low Temperature and Oxygen to Induce OLE1 Transcription in <i>Saccharomyces cerevisiae</i> . <i>Biochemical and Biophysical Research Communications</i> , 2002, 291, 707-713. | 1.0 | 98 |
| 4340 | Comparison of the interaction of uridine, cytidine, and other pyrimidine nucleoside analogues with recombinant human equilibrative nucleoside transporter 2 (hENT2) produced in <i>Saccharomyces cerevisiae</i> . <i>Biochemistry and Cell Biology</i> , 2002, 80, 639-644. | 0.9 | 41 |
| 4341 | Kre1p, the Plasma Membrane Receptor for the Yeast K1 Viral Toxin. <i>Cell</i> , 2002, 108, 395-405. | 13.5 | 117 |

| # | ARTICLE | IF | CITATIONS |
|------|---|------|-----------|
| 4342 | Cid13 Is a Cytoplasmic Poly(A) Polymerase that Regulates Ribonucleotide Reductase mRNA. <i>Cell</i> , 2002, 109, 563-573. | 13.5 | 130 |
| 4343 | GDH1 expression is regulated by GLN3, GCN4, and HAP4 under respiratory growth. <i>Biochemical and Biophysical Research Communications</i> , 2002, 293, 79-85. | 1.0 | 43 |
| 4344 | Molecular cloning, characterisation, and expression of a neutral trehalase from the insect pathogenic fungus <i>Metarhizium anisopliae</i> . <i>Journal of Invertebrate Pathology</i> , 2002, 80, 127-137. | 1.5 | 31 |
| 4345 | Centrosomal Anchoring of the Protein Kinase CK1 $\hat{\imath}$ Mediated by Attachment to the Large, Coiled-coil Scaffolding Protein CG-NAP/AKAP450. <i>Journal of Molecular Biology</i> , 2002, 322, 785-797. | 2.0 | 102 |
| 4346 | Association of Cu,Zn-Type Superoxide Dismutase with Mitochondria and Peroxisomes. <i>Archives of Biochemistry and Biophysics</i> , 2002, 399, 96-102. | 1.4 | 38 |
| 4347 | GRACILE Syndrome, a Lethal Metabolic Disorder with Iron Overload, Is Caused by a Point Mutation in BCS1L. <i>American Journal of Human Genetics</i> , 2002, 71, 863-876. | 2.6 | 263 |
| 4348 | Endosome-Associated Complex, ESCRT-II, Recruits Transport Machinery for Protein Sorting at the Multivesicular Body. <i>Developmental Cell</i> , 2002, 3, 283-289. | 3.1 | 589 |
| 4349 | Escrt-III. <i>Developmental Cell</i> , 2002, 3, 271-282. | 3.1 | 799 |
| 4350 | The gene encoding a hybrid ubiquitin fusion protein involved in ribosome biogenesis is essential for growth. <i>FEMS Yeast Research</i> , 2002, 2, 25-30. | 1.1 | 0 |
| 4351 | HSM2 (HMO1) gene participates in mutagenesis control in yeast <i>Saccharomyces cerevisiae</i> . <i>DNA Repair</i> , 2002, 1, 287-297. | 1.3 | 22 |
| 4352 | Structural and functional conservation of error-free DNA postreplication repair in <i>Schizosaccharomyces pombe</i> . <i>DNA Repair</i> , 2002, 1, 869-880. | 1.3 | 20 |
| 4353 | Acyl-CoA Z9- and Z10-desaturase genes from a New Zealand leafroller moth species, <i>Planotortrix octo</i> . <i>Insect Biochemistry and Molecular Biology</i> , 2002, 32, 961-966. | 1.2 | 57 |
| 4354 | Moth desaturase characterized that produces both Z and E isomers of $\hat{\imath}^{11}$ -tetradecenoic acids. <i>Insect Biochemistry and Molecular Biology</i> , 2002, 32, 1489-1495. | 1.2 | 46 |
| 4355 | The tomato hexokinase LeHXK1 cloning, mapping, expression pattern and phylogenetic relationships. <i>Plant Science</i> , 2002, 163, 581-590. | 1.7 | 46 |
| 4356 | Cloning and characterization of glucokinase from a methylotrophic yeast <i>Hansenula polymorpha</i> : different effects on glucose repression in <i>H. polymorpha</i> and <i>Saccharomyces cerevisiae</i> . <i>Gene</i> , 2002, 296, 195-203. | 1.0 | 22 |
| 4357 | Analysis of fungal P450foxy (CYP505), a fused protein of cytochrome P450 and its reductase. <i>International Congress Series</i> , 2002, 1233, 143-149. | 0.2 | 1 |
| 4358 | Arabidopsis CAP Regulates the Actin Cytoskeleton Necessary for Plant Cell Elongation and Division. <i>Plant Cell</i> , 2002, 14, 149-163. | 3.1 | 70 |
| 4359 | <i>Saccharomyces cerevisiae</i> nucleolar protein Nop7p is necessary for biogenesis of 60S ribosomal subunits. <i>Rna</i> , 2002, 8, 150-165. | 1.6 | 79 |

| # | ARTICLE | IF | CITATIONS |
|------|---|-----|-----------|
| 4360 | Isolation and characterization of new <i>Saccharomyces cerevisiae</i> mutants perturbed in nuclear pore complex assembly. <i>BMC Genetics</i> , 2002, 3, 17. | 2.7 | 40 |
| 4361 | RPL29 codes for a non-essential protein of the 60S ribosomal subunit in <i>Saccharomyces cerevisiae</i> and exhibits synthetic lethality with mutations in genes for proteins required for subunit coupling. <i>Biochimica Et Biophysica Acta Gene Regulatory Mechanisms</i> , 2002, 1574, 255-261. | 2.4 | 34 |
| 4362 | Construction of engineered yeast with the ability of binding to cellulose. <i>Journal of Molecular Catalysis B: Enzymatic</i> , 2002, 17, 197-202. | 1.8 | 23 |
| 4363 | A role of <i>Saccharomyces cerevisiae</i> fatty Acid activation protein 4 in palmitoyl-CoA pool for growth in the presence of ethanol. <i>Journal of Bioscience and Bioengineering</i> , 2002, 93, 288-295. | 1.1 | 10 |
| 4364 | Extremely simple, rapid and highly efficient transformation method for the yeast <i>Saccharomyces cerevisiae</i> using glutathione and early log phase cells. <i>Journal of Bioscience and Bioengineering</i> , 2002, 94, 166-171. | 1.1 | 2 |
| 4365 | High-level production of prourokinase-annexin V chimeras in the methylotrophic yeast <i>Pichia pastoris</i> . <i>Journal of Bioscience and Bioengineering</i> , 2002, 94, 467-473. | 1.1 | 15 |
| 4366 | <i>C. elegans</i> PAT-4/ILK Functions as an Adaptor Protein within Integrin Adhesion Complexes. <i>Current Biology</i> , 2002, 12, 787-797. | 1.8 | 286 |
| 4367 | Engineering of baker's yeasts, <i>E. coli</i> and <i>Bacillus</i> hosts for the production of <i>Bacillus subtilis</i> Lipase A. <i>Biotechnology and Bioengineering</i> , 2002, 78, 339-345. | 1.7 | 29 |
| 4368 | Integrated bioprocessing in <i>Saccharomyces cerevisiae</i> using green fluorescent protein as a fusion partner. <i>Biotechnology and Bioengineering</i> , 2002, 79, 682-693. | 1.7 | 15 |
| 4369 | A Rapid Screening Method to Monitor Expression of Recombinant Proteins from Various Prokaryotic and Eukaryotic Expression Systems Using Matrix-Assisted Laser Desorption Ionization "Time-of-Flight Mass Spectrometry. <i>Analytical Biochemistry</i> , 2002, 305, 242-250. | 1.1 | 11 |
| 4370 | Yeast methionine aminopeptidase type 1 is ribosome-associated and requires its N-terminal zinc finger domain for normal function in vivo. <i>Journal of Cellular Biochemistry</i> , 2002, 85, 678-688. | 1.2 | 56 |
| 4371 | Cloning and molecular characterization of the Δ^6 -desaturase from two <i>Echium</i> plant species: Production of GLA by heterologous expression in yeast and tobacco. <i>Lipids</i> , 2002, 37, 417-426. | 0.7 | 50 |
| 4372 | Spacer-elongated cell wall fusion proteins improve cell surface expression in the yeast <i>Saccharomyces cerevisiae</i> . <i>Applied Microbiology and Biotechnology</i> , 2002, 58, 637-644. | 1.7 | 37 |
| 4373 | Production of active bovine cathepsin C (dipeptidyl aminopeptidase I) in the methylotrophic yeast <i>Candida boidinii</i> . <i>Applied Microbiology and Biotechnology</i> , 2002, 59, 252-258. | 1.7 | 12 |
| 4374 | Cell surface-engineered yeast with ability to bind, and self-aggregate in response to, copper ion. <i>Applied Microbiology and Biotechnology</i> , 2002, 59, 259-264. | 1.7 | 58 |
| 4375 | Sequence-based approach for identification of cell wall proteins in <i>Saccharomyces cerevisiae</i> . <i>Current Genetics</i> , 2002, 40, 311-316. | 0.8 | 21 |
| 4376 | Genome-wide expression analysis of genes affected by amino acid sensor Ssy1p in <i>Saccharomyces cerevisiae</i> . <i>Current Genetics</i> , 2002, 41, 63-72. | 0.8 | 31 |
| 4377 | Role of RNA surveillance proteins Upf1/CpaR, Upf2 and Upf3 in the translational regulation of yeast CPA1 gene. <i>Current Genetics</i> , 2002, 41, 224-231. | 0.8 | 13 |

| # | ARTICLE | IF | CITATIONS |
|------|---|-----|-----------|
| 4378 | Functional dissection of Pdr1p, a regulator of multidrug resistance in <i>Saccharomyces cerevisiae</i> . <i>Molecular Genetics and Genomics</i> , 2002, 267, 96-106. | 1.0 | 41 |
| 4379 | Roles of disulfide bridges in scorpion toxin BmK ϵ M1 analyzed by mutagenesis. <i>Chemical Biology and Drug Design</i> , 2002, 60, 247-256. | 1.2 | 27 |
| 4380 | Human and <i>Drosophila</i> UDP-galactose transporters transport UDP-N -acetylgalactosamine in addition to UDP-galactose. <i>FEBS Journal</i> , 2002, 269, 128-138. | 0.2 | 80 |
| 4381 | Synthesis and in vivo modulatory activity of protein kinase C of xanthone derivatives. <i>Bioorganic and Medicinal Chemistry</i> , 2002, 10, 3219-3227. | 1.4 | 37 |
| 4382 | A GyrB-GyrA fusion protein expressed in yeast cells is able to remove DNA supercoils but cannot substitute eukaryotic topoisomerase II. <i>Genes To Cells</i> , 2002, 7, 249-257. | 0.5 | 17 |
| 4383 | MBLR, a new RING finger protein resembling mammalian Polycomb gene products, is regulated by cell cycle-dependent phosphorylation. <i>Genes To Cells</i> , 2002, 7, 835-850. | 0.5 | 44 |
| 4384 | Analysis of the oxidative stress regulation of the <i>Candida albicans</i> transcription factor, Cap1p. <i>Molecular Microbiology</i> , 2002, 36, 618-629. | 1.2 | 131 |
| 4385 | The yeast transcriptome in aerobic and hypoxic conditions: effects of hap1, rox1, rox3 and rbi1 deletions. <i>Molecular Microbiology</i> , 2002, 43, 545-555. | 1.2 | 77 |
| 4386 | A cytoplasmic coiled-coil domain is required for histidine kinase activity of the yeast osmosensor, SLN1. <i>Molecular Microbiology</i> , 2002, 43, 459-473. | 1.2 | 48 |
| 4387 | Translational readthrough of the PDE2 stop codon modulates cAMP levels in <i>Saccharomyces cerevisiae</i> . <i>Molecular Microbiology</i> , 2002, 43, 641-652. | 1.2 | 107 |
| 4388 | The proper folding of a long C-terminal segment of the yeast Lys14p regulator is required for activation of LYS genes in response to the metabolic effector. <i>Molecular Microbiology</i> , 2002, 43, 1629-1639. | 1.2 | 12 |
| 4389 | Rho1p mutations specific for regulation of β (1 \rightarrow 3)glucan synthesis and the order of assembly of the yeast cell wall. <i>Molecular Microbiology</i> , 2002, 44, 1167-1183. | 1.2 | 36 |
| 4390 | The RHO1-GAPs SAC7, BEM2 and BAG7 control distinct RHO1 functions in <i>Saccharomyces cerevisiae</i> . <i>Molecular Microbiology</i> , 2002, 45, 1433-1441. | 1.2 | 55 |
| 4391 | Regulation of the yeast Rlm1 transcription factor by the Mpk1 cell wall integrity MAP kinase. <i>Molecular Microbiology</i> , 2002, 46, 781-789. | 1.2 | 209 |
| 4392 | Characterization of a <i>Saccharomyces cerevisiae</i> NADP(H)-dependent alcohol dehydrogenase (ADHVII), a member of the cinnamyl alcohol dehydrogenase family. <i>FEBS Journal</i> , 2002, 269, 5738-5745. | 0.2 | 102 |
| 4393 | Molecular characterization of MRC19 of <i>Saccharomyces cerevisiae</i> . <i>FEBS Journal</i> , 2002, 269, 5840-5850. | 0.2 | 10 |
| 4394 | Phylogenetic footprinting reveals multiple regulatory elements involved in control of the meiotic recombination gene, REC102. <i>Yeast</i> , 2002, 19, 99-114. | 0.8 | 11 |
| 4395 | Molecular and functional analysis of a MIG1 homologue from the yeast <i>Schwanniomyces occidentalis</i> . <i>Yeast</i> , 2002, 19, 459-465. | 0.8 | 5 |

| # | ARTICLE | IF | CITATIONS |
|------|---|-----|-----------|
| 4396 | TheSCR1 gene fromSchwanniomycetes occidentalisencodes a highly hydrophobic polypeptide, which confers ribosomal resistance to cycloheximide. <i>Yeast</i> , 2002, 19, 735-743. | 0.8 | 1 |
| 4397 | The role of the 5' untranslated region (UTR) in glucose-dependent mRNA decay. <i>Yeast</i> , 2002, 19, 887-902. | 0.8 | 26 |
| 4398 | A singleFKShomologue inYarrowia lipolyticaais essential for viability. <i>Yeast</i> , 2002, 19, 1003-1014. | 0.8 | 16 |
| 4399 | ARL1 and membrane traffic inSaccharomyces cerevisiae. <i>Yeast</i> , 2002, 19, 1039-1056. | 0.8 | 54 |
| 4400 | Genetic interactions linkARF1,YPT31/32 andTRS130. <i>Yeast</i> , 2002, 19, 1075-1086. | 0.8 | 28 |
| 4401 | Cloning and analysis ofCoEXG1, a secreted 1,3-?-glucanase of the yeast biocontrol agentCandida oleophila. <i>Yeast</i> , 2002, 19, 1171-1182. | 0.8 | 21 |
| 4402 | SHAM-sensitive alternative respiration in the xylose-metabolizing yeastPichia stipitis. <i>Yeast</i> , 2002, 19, 1203-1220. | 0.8 | 45 |
| 4403 | Isolation and characterization of the geneURA3 encoding the orotidine-5?-phosphate decarboxylase fromTorulaspora delbrueckii. <i>Yeast</i> , 2002, 19, 1431-1435. | 0.8 | 11 |
| 4404 | Expression of a functional human interleukin-18 in yeast. <i>Enzyme and Microbial Technology</i> , 2002, 30, 703-709. | 1.6 | 8 |
| 4405 | Overexpression of B-type cyclins alters chromosomal segregation. <i>Oncogene</i> , 2002, 21, 2051-2057. | 2.6 | 67 |
| 4406 | An integrated approach for finding overlooked genes in yeast. <i>Nature Biotechnology</i> , 2002, 20, 58-63. | 9.4 | 112 |
| 4407 | Dual cell wall/mitochondria localization of the SUN family proteins. <i>FEMS Microbiology Letters</i> , 2002, 207, 165-172. | 0.7 | 45 |
| 4408 | Cloning and characterization of vmaA, the gene encoding a 69-kDa catalytic subunit of the vacuolar H ⁺ -ATPase during alkaline pH mediated growth ofAspergillus oryzae. <i>FEMS Microbiology Letters</i> , 2002, 209, 277-282. | 0.7 | 10 |
| 4409 | Characterization of theCaNAG3,CaNAG4, andCaNAG6genes of the pathogenic fungusCandida albicans: possible involvement of these genes in the susceptibilities of cytotoxic agents. <i>FEMS Microbiology Letters</i> , 2002, 212, 15-21. | 0.7 | 41 |
| 4410 | TheCandida albicans UBI3gene encoding a hybrid ubiquitin fusion protein involved in ribosome biogenesis is essential for growth. <i>FEMS Yeast Research</i> , 2002, 2, 25-30. | 1.1 | 8 |
| 4411 | Amphipathic property of free thiol group contributes to an increase inthe catalytic efficiency of carboxypeptidase-?Y. <i>FEBS Journal</i> , 2002, 269, 3220-3225. | 0.2 | 4 |
| 4412 | New Targets for Antivirals: The Ribosomal A-Site and the Factors That Interact with It. <i>Virology</i> , 2002, 300, 60-70. | 1.1 | 49 |
| 4413 | Identification of SnIP1, a novel protein that interacts with SNF1-related protein kinase (SnRK1). <i>Plant Molecular Biology</i> , 2002, 49, 31-44. | 2.0 | 22 |

| # | ARTICLE | IF | CITATIONS |
|------|--|-----|-----------|
| 4414 | Coregulation of Alternative Transcripts of the STA2 Gene in the Yeast <i>Saccharomyces cerevisiae</i> . <i>Russian Journal of Genetics</i> , 2002, 38, 1121-1126. | 0.2 | 1 |
| 4415 | Mutations in yeast Rad51 that partially bypass the requirement for Rad55 and Rad57 in DNA repair by increasing the stability of Rad51-DNA complexes. <i>EMBO Journal</i> , 2002, 21, 3160-3170. | 3.5 | 104 |
| 4416 | Recombination-dependent mtDNA partitioning: in vivo role of Mhr1p to promote pairing of homologous DNA. <i>EMBO Journal</i> , 2002, 21, 4730-4740. | 3.5 | 85 |
| 4417 | Trp17 and Glu20 residues in conserved WMN(D/E)PN motif are essential for <i>Aspergillus ficuum</i> endoinulinase (EC 3.2.1.7) activity. <i>Biochemistry (Moscow)</i> , 2003, 68, 658-661. | 0.7 | 5 |
| 4418 | Monitoring of the tissue distribution of fibroblast growth factor containing a high mannose-type sugar chain produced in mutant yeast. <i>Glycoconjugate Journal</i> , 2003, 20, 385-397. | 1.4 | 11 |
| 4419 | Role of acetylated human AP-endonuclease (APE1/Ref-1) in regulation of the parathyroid hormone gene. <i>EMBO Journal</i> , 2003, 22, 6299-6309. | 3.5 | 158 |
| 4420 | Domains of eIF1A that mediate binding to eIF2, eIF3 and eIF5B and promote ternary complex recruitment in vivo. <i>EMBO Journal</i> , 2003, 22, 193-204. | 3.5 | 120 |
| 4421 | Internal initiation drives the synthesis of Ure2 protein lacking the prion domain and affects [URE3] propagation in yeast cells. <i>EMBO Journal</i> , 2003, 22, 1199-1209. | 3.5 | 68 |
| 4422 | A novel eukaryotic factor for cytosolic Fe-S cluster assembly. <i>EMBO Journal</i> , 2003, 22, 4826-4835. | 3.5 | 150 |
| 4423 | An interaction between an <i>Arabidopsis</i> poly(A) polymerase and a homologue of the 100 kDa subunit of CPSF. <i>Plant Molecular Biology</i> , 2003, 51, 373-384. | 2.0 | 16 |
| 4424 | The p1 Protein of the Yeast Transposon Ty1 Can Be Used for the Construction of Bi-Functional Virus-Like Particles. <i>Journal of Molecular Microbiology and Biotechnology</i> , 2003, 5, 97-104. | 1.0 | 10 |
| 4425 | An improved transformation protocol for the human fungal pathogen <i>Candida albicans</i> . <i>Current Genetics</i> , 2003, 42, 339-343. | 0.8 | 193 |
| 4426 | High-efficiency electroporation by freezing intact yeast cells with addition of calcium. <i>Current Genetics</i> , 2003, 43, 206-211. | 0.8 | 40 |
| 4427 | Rsf1p, a protein required for respiratory growth of <i>Saccharomyces cerevisiae</i> . <i>Current Genetics</i> , 2003, 43, 263-272. | 0.8 | 24 |
| 4428 | Cloning and biochemical characterization of hexokinase from the methylotrophic yeast <i>Hansenula polymorpha</i> . <i>Current Genetics</i> , 2003, 44, 268-276. | 0.8 | 16 |
| 4429 | Display of a functional hetero-oligomeric catalytic antibody on the yeast cell surface. <i>Applied Microbiology and Biotechnology</i> , 2003, 62, 226-232. | 1.7 | 53 |
| 4430 | Sequential cloned gene integration in the yeast <i>Kluyveromyces lactis</i> . <i>Applied Microbiology and Biotechnology</i> , 2003, 62, 523-527. | 1.7 | 29 |
| 4431 | Enhanced expression and functional characterization of the human ferritin H- and L-chain genes in <i>Saccharomyces cerevisiae</i> . <i>Applied Microbiology and Biotechnology</i> , 2003, 63, 57-63. | 1.7 | 16 |

| # | ARTICLE | IF | CITATIONS |
|------|--|-----|-----------|
| 4432 | Bioadsorption of cadmium ion by cell surface-engineered yeasts displaying metallothionein and hexa-His. <i>Applied Microbiology and Biotechnology</i> , 2003, 63, 182-186. | 1.7 | 61 |
| 4433 | Analysis of the distribution of the kinetochore protein Ndc10p in <i>Saccharomyces cerevisiae</i> using 3-D modeling of mitotic spindles. <i>Chromosoma</i> , 2003, 111, 417-428. | 1.0 | 29 |
| 4434 | A signal peptide secretion screen in <i>Fucus distichus</i> embryos reveals expression of glucanase, EGF domain-containing, and LRR receptor kinase-like polypeptides during asymmetric cell growth. <i>Planta</i> , 2003, 217, 931-950. | 1.6 | 26 |
| 4435 | Mutations in GCR1 affect SUC2 gene expression in <i>Saccharomyces cerevisiae</i> . <i>Molecular Genetics and Genomics</i> , 2003, 268, 825-831. | 1.0 | 19 |
| 4436 | The <i>Aspergillus parasiticus</i> protein AFLJ interacts with the aflatoxin pathway-specific regulator AFLR. <i>Molecular Genetics and Genomics</i> , 2003, 268, 711-719. | 1.0 | 148 |
| 4437 | Gal11 is a general activator of basal transcription, whose activity is regulated by the general repressor Sin4 in yeast. <i>Molecular Genetics and Genomics</i> , 2003, 269, 68-77. | 1.0 | 11 |
| 4438 | Merging of multiple signals regulating Δ^9 fatty acid desaturase gene transcription in <i>Saccharomyces cerevisiae</i> . <i>Molecular Genetics and Genomics</i> , 2003, 269, 370-380. | 1.0 | 17 |
| 4439 | Mutation of a conserved CDK site converts a metazoan Elongation Factor 1B β subunit into a replacement for yeast eEF1B β . <i>Molecular Genetics and Genomics</i> , 2003, 269, 776-788. | 1.0 | 8 |
| 4440 | Cis-acting elements sufficient for induction of FDH1 expression by formate in the methylotrophic yeast <i>Candida boidinii</i> . <i>Molecular Genetics and Genomics</i> , 2003, 270, 273-280. | 1.0 | 15 |
| 4441 | Microarray analyses of the metabolic responses of <i>Saccharomyces cerevisiae</i> to organic solvent dimethyl sulfoxide. <i>Journal of Industrial Microbiology and Biotechnology</i> , 2003, 30, 57-69. | 1.4 | 40 |
| 4442 | Influence of Ras function on ethanol stress response of sake yeast. <i>Journal of Bioscience and Bioengineering</i> , 2003, 96, 474-480. | 1.1 | 9 |
| 4443 | Characterization of an Δ -ketoglutarate-resistant sake yeast mutant with high organic acid productivity. <i>Journal of Bioscience and Bioengineering</i> , 2003, 96, 332-336. | 1.1 | 12 |
| 4444 | Repeated chromosome splitting targeted to Δ sequences in <i>Saccharomyces cerevisiae</i> . <i>Journal of Bioscience and Bioengineering</i> , 2003, 96, 397-400. | 1.1 | 7 |
| 4445 | Ura Δ host strains for genetic manipulation and heterologous expression of <i>Torulaspora delbrueckii</i> . <i>International Journal of Food Microbiology</i> , 2003, 86, 79-86. | 2.1 | 6 |
| 4446 | Novel tumorigenic rearrangement, Δ rfp/ret, in a papillary thyroid carcinoma from externally irradiated patient. <i>Mutation Research - Fundamental and Molecular Mechanisms of Mutagenesis</i> , 2003, 527, 81-90. | 0.4 | 51 |
| 4447 | β -cardiac actin (ACTC) binds to the band 3 (AE1) cardiac isoform. <i>Journal of Cellular Biochemistry</i> , 2003, 89, 1215-1221. | 1.2 | 9 |
| 4448 | N-terminal methionine removal and methionine metabolism in <i>Saccharomyces cerevisiae</i> . <i>Journal of Cellular Biochemistry</i> , 2003, 89, 964-974. | 1.2 | 25 |
| 4449 | Ecdysone agonists: Mechanism and importance in controlling insect pests of agriculture and forestry. <i>Archives of Insect Biochemistry and Physiology</i> , 2003, 54, 187-199. | 0.6 | 88 |

| # | ARTICLE | IF | CITATIONS |
|------|---|-----|-----------|
| 4450 | Kinetic analysis of enhanced thermal stability of an alkaline protease with engineered twin disulfide bridges and calcium-dependent stability. <i>Biotechnology and Bioengineering</i> , 2003, 81, 187-192. | 1.7 | 10 |
| 4451 | Activation of <i>chsC</i> transcription by <i>AbaA</i> during asexual development of <i>Aspergillus nidulans</i> . <i>FEMS Microbiology Letters</i> , 2003, 220, 241-246. | 0.7 | 25 |
| 4452 | Characterization of <i>gdp1+</i> as encoding a GDPase in the fission yeast <i>Schizosaccharomyces pombe</i> . <i>FEMS Microbiology Letters</i> , 2003, 228, 33-38. | 0.7 | 7 |
| 4453 | Inhibition of protein kinase C by synthetic xanthone derivatives. <i>Bioorganic and Medicinal Chemistry</i> , 2003, 11, 1215-1225. | 1.4 | 34 |
| 4454 | Isolation and characterization of the gene from the yeast. <i>FEMS Yeast Research</i> , 2003, 4, 51-57. | 1.1 | 4 |
| 4455 | Development of combinatorial bioengineering using yeast cell surface display – order-made design of cell and protein for bio-monitoring. <i>Biosensors and Bioelectronics</i> , 2003, 19, 123-130. | 5.3 | 27 |
| 4456 | Cell cycle regulator <i>Cdc14</i> is expressed during sporulation but not hyphal growth in the fungus-like oomycete <i>Phytophthora infestans</i> . <i>Molecular Microbiology</i> , 2003, 50, 487-494. | 1.2 | 90 |
| 4457 | Identification of a mitochondrial transporter for basic amino acids in <i>Arabidopsis thaliana</i> by functional reconstitution into liposomes and complementation in yeast. <i>Plant Journal</i> , 2003, 33, 1027-1035. | 2.8 | 85 |
| 4458 | Three <i>SAC1</i> -like genes show overlapping patterns of expression in <i>Arabidopsis</i> but are remarkably silent during embryo development. <i>Plant Journal</i> , 2003, 34, 293-306. | 2.8 | 39 |
| 4459 | Functional analyses of mammalian protein kinase C isozymes in budding yeast and mammalian fibroblasts. <i>Genes To Cells</i> , 2003, 2, 601-614. | 0.5 | 19 |
| 4460 | A brute force postgenome approach to identify temperature-sensitive mutations that negatively interact with <i>separase</i> and <i>securin</i> plasmids. <i>Genes To Cells</i> , 2003, 8, 341-355. | 0.5 | 9 |
| 4461 | Molecular characterization, function and regulation of ammonium transporters (<i>Amt</i>) and ammonium-metabolizing enzymes (<i>GS</i> , <i>NADP-GDH</i>) in the ectomycorrhizal fungus <i>Hebeloma cylindrosporum</i> . <i>Molecular Microbiology</i> , 2003, 47, 411-430. | 1.2 | 113 |
| 4462 | <i>Rga5p</i> is a specific <i>Rho1p</i> GTPase-activating protein that regulates cell integrity in <i>Schizosaccharomyces pombe</i> . <i>Molecular Microbiology</i> , 2003, 47, 507-518. | 1.2 | 51 |
| 4463 | <i>Ty3</i> requires yeast <i>La</i> homologous protein for wild-type frequencies of transposition. <i>Molecular Microbiology</i> , 2003, 49, 501-515. | 1.2 | 7 |
| 4464 | <i>NET1</i> and <i>HFI1</i> genes of yeast mediate both chromosome maintenance and mitochondrial rho? mutagenesis. <i>Yeast</i> , 2003, 20, 955-971. | 0.8 | 6 |
| 4465 | The <i>CL1</i> gene of <i>Pichia pastoris</i> , transcriptional regulation and use of its promoter. <i>Yeast</i> , 2003, 20, 1097-1108. | 0.8 | 52 |
| 4466 | <i>LYS2</i> gene and its mutation in <i>Kluyveromyces lactis</i> . <i>Yeast</i> , 2003, 20, 1171-1175. | 0.8 | 4 |
| 4467 | A DNA region of <i>Torulaspora delbrueckii</i> containing the <i>HIS3</i> gene: sequence, gene order and evolution. <i>Yeast</i> , 2003, 20, 1359-1368. | 0.8 | 3 |

| # | ARTICLE | IF | CITATIONS |
|------|--|-----|-----------|
| 4468 | The role of the GATA factors Gln3p, Nil1p, Dal80p and the Ure2p on ASP3 regulation in <i>Saccharomyces cerevisiae</i> . <i>Yeast</i> , 2003, 20, 31-37. | 0.8 | 20 |
| 4469 | RHO1 (YIRHO1) is a non-essential gene in <i>Yarrowia lipolytica</i> and complements rho1 ^Δ lethality in <i>Saccharomyces cerevisiae</i> . <i>Yeast</i> , 2003, 20, 343-350. | 0.8 | 5 |
| 4470 | Characterization of a disulphide-bound Pir-cell wall protein (Pir-CWP) of <i>Yarrowia lipolytica</i> . <i>Yeast</i> , 2003, 20, 417-426. | 0.8 | 33 |
| 4471 | Isolation of the MNN9 gene of <i>Yarrowia lipolytica</i> (YIMNN9) and phenotype analysis of a mutant ylmnn9 ^Δ strain. <i>Yeast</i> , 2003, 20, 633-644. | 0.8 | 10 |
| 4472 | New mutations of <i>Saccharomyces cerevisiae</i> that partially relieve both glucose and galactose repression activate the protein kinase Snf1. <i>FEMS Yeast Research</i> , 2003, 3, 77-84. | 1.1 | 2 |
| 4473 | Expression of recombinant GFP-actin fusion protein in the methylotrophic yeast <i>Pichia pastoris</i> . <i>FEMS Yeast Research</i> , 2003, 3, 105-111. | 1.1 | 4 |
| 4474 | Activation state of protein kinase A as measured in permeabilised <i>Saccharomyces cerevisiae</i> cells correlates with PKA-controlled phenotypes in vivo. <i>FEMS Yeast Research</i> , 2003, 3, 119-126. | 1.1 | 5 |
| 4475 | The localization change of Ybr078w/Ecm33, a yeast GPI-associated protein, from the plasma membrane to the cell wall, affecting the cellular function. <i>FEMS Microbiology Letters</i> , 2003, 218, 175-180. | 0.7 | 38 |
| 4476 | Engineering tolerance and accumulation of lead and cadmium in transgenic plants. <i>Nature Biotechnology</i> , 2003, 21, 914-919. | 9.4 | 381 |
| 4477 | Regulation of the maltase gene promoter in and. <i>FEMS Yeast Research</i> , 2003, 4, 165-173. | 1.1 | 21 |
| 4478 | Application of the reusable, selectable marker to industrial yeast: construction and evaluation of heterothallic wine strains of <i>S. cerevisiae</i> , possessing minimal foreign DNA sequences. <i>FEMS Yeast Research</i> , 2003, 4, 339-347. | 1.1 | 65 |
| 4479 | Construction of a Trp commercial baker's yeast strain by using food-safe-grade dominant drug resistance cassettes. <i>FEMS Yeast Research</i> , 2003, 4, 329-338. | 1.1 | 10 |
| 4480 | The delivery of ADP/ATP carrier protein to mitochondria probed by fusions with green fluorescent protein and β -galactosidase. <i>FEMS Yeast Research</i> , 2003, 4, 315-321. | 1.1 | 6 |
| 4481 | The mechanism of action of the <i>Pseudomonas aeruginosa</i> -encoded type III cytotoxin, ExoU. <i>EMBO Journal</i> , 2003, 22, 2959-2969. | 3.5 | 321 |
| 4482 | Gef1p, a New Guanine Nucleotide Exchange Factor for Cdc42p, Regulates Polarity in <i>Schizosaccharomyces pombe</i> . <i>Molecular Biology of the Cell</i> , 2003, 14, 313-323. | 0.9 | 81 |
| 4483 | Characterization of HRD3, a <i>Schizosaccharomyces pombe</i> gene involved in DNA repair and cell viability. <i>Korean Journal of Biological Sciences</i> , 2003, 7, 159-164. | 0.1 | 1 |
| 4484 | Cytoskeletal Activation of a Checkpoint Kinase. <i>Molecular Cell</i> , 2003, 12, 663-673. | 4.5 | 48 |
| 4485 | Glucose and Type 2A Protein Phosphatase Regulate the Interaction Between Catalytic and Regulatory Subunits of AMP-activated Protein Kinase. <i>Journal of Molecular Biology</i> , 2003, 333, 201-209. | 2.0 | 44 |

| # | ARTICLE | IF | CITATIONS |
|------|--|-----|-----------|
| 4486 | Sub-families of β -Barrel Enzymes: A New Adenine Deaminase Family. <i>Journal of Molecular Biology</i> , 2003, 334, 1117-1131. | 2.0 | 32 |
| 4487 | Ethanol Tolerance in the Yeast <i>Saccharomyces cerevisiae</i> Is Dependent on Cellular Oleic Acid Content. <i>Applied and Environmental Microbiology</i> , 2003, 69, 1499-1503. | 1.4 | 282 |
| 4488 | Aspartate-186 in the head group of the yeast iron-sulfur protein of the cytochrome bc1 complex contributes to the protein conformation required for efficient electron transfer. <i>Biochimica Et Biophysica Acta - Bioenergetics</i> , 2003, 1607, 65-78. | 0.5 | 5 |
| 4489 | Functional Expression of a Bacterial Heavy Metal Transporter in <i>Arabidopsis</i> Enhances Resistance to and Decreases Uptake of Heavy Metals. <i>Plant Physiology</i> , 2003, 133, 589-596. | 2.3 | 136 |
| 4490 | Mutational analysis and localization of the inositol transporters of <i>Saccharomyces cerevisiae</i> . <i>Journal of Bioscience and Bioengineering</i> , 2003, 96, 291-297. | 1.1 | 10 |
| 4491 | Metalloid tolerance based on phytochelatins is not functionally equivalent to the arsenite transporter Acr3p. <i>Biochemical and Biophysical Research Communications</i> , 2003, 304, 293-300. | 1.0 | 51 |
| 4492 | Association of hepatitis B virus polymerase with promyelocytic leukemia nuclear bodies mediated by the S100 family protein p11. <i>Biochemical and Biophysical Research Communications</i> , 2003, 305, 1049-1056. | 1.0 | 29 |
| 4493 | Transcriptional regulation of phosphate-responsive genes in low-affinity phosphate-transporter-defective mutants in <i>Saccharomyces cerevisiae</i> . <i>Biochemical and Biophysical Research Communications</i> , 2003, 306, 843-850. | 1.0 | 55 |
| 4494 | Growth temperature downshift induces antioxidant response in <i>Saccharomyces cerevisiae</i> . <i>Biochemical and Biophysical Research Communications</i> , 2003, 307, 308-314. | 1.0 | 56 |
| 4495 | The anti-neoplastic and novel topoisomerase II-mediated cytotoxicity of neoamphimedine, a marine pyridoacridine. <i>Biochemical Pharmacology</i> , 2003, 66, 447-458. | 2.0 | 53 |
| 4496 | Nucleo-mitochondrial interactions in <i>Saccharomyces cerevisiae</i> : characterization of a nuclear gene suppressing a defect in mitochondrial tRNA ^{Asp} processing. <i>Gene</i> , 2003, 303, 63-68. | 1.0 | 8 |
| 4497 | Glucose dependant negative translational control of the heterologous expression of the preS2 HBV antigen in yeast. <i>Gene</i> , 2003, 311, 165-170. | 1.0 | 8 |
| 4498 | A single amino acid substitution in MSH5 results in DNA alkylation tolerance. <i>Gene</i> , 2003, 315, 177-182. | 1.0 | 14 |
| 4499 | Cloning of CnLYSO1, a novel extracellular lysophospholipase of the pathogenic fungus <i>Cryptococcus neoformans</i> . <i>Gene</i> , 2003, 316, 67-78. | 1.0 | 12 |
| 4500 | Oxidative stresses elevate the expression of cytochrome c peroxidase in <i>Saccharomyces cerevisiae</i> . <i>Biochimica Et Biophysica Acta - General Subjects</i> , 2003, 1623, 1-5. | 1.1 | 38 |
| 4501 | Generation of disruption cassettes in vivo using a PCR product and <i>Saccharomyces cerevisiae</i> . <i>Journal of Microbiological Methods</i> , 2003, 52, 141-145. | 0.7 | 12 |
| 4502 | Efficient transformation of the osmotolerant yeast <i>Zygosaccharomyces rouxii</i> by electroporation. <i>Journal of Microbiological Methods</i> , 2003, 55, 481-484. | 0.7 | 31 |
| 4503 | Conditions affecting production of functional muscle recombinant β -tropomyosin in <i>Saccharomyces cerevisiae</i> . <i>Protein Expression and Purification</i> , 2003, 30, 105-111. | 0.6 | 4 |

| # | ARTICLE | IF | CITATIONS |
|------|--|-----|-----------|
| 4504 | RNA Polymerase II/TFIIIF Structure and Conserved Organization of the Initiation Complex. <i>Molecular Cell</i> , 2003, 12, 1003-1013. | 4.5 | 86 |
| 4505 | Potential of human estrogen receptor β -mediated gene expression by steroid receptor coactivator-1 (SRC-1) in <i>Saccharomyces cerevisiae</i> . <i>Journal of Steroid Biochemistry and Molecular Biology</i> , 2003, 86, 15-26. | 1.2 | 8 |
| 4506 | Yeast as a Host to Screen for Modulators and Regulatory Regions of Mammalian Protein Kinase C Isoforms. , 2003, 233, 491-516. | | 3 |
| 4507 | Importance of the Conserved Aromatic Residues in the Scorpion β -Like Toxin BmK M1. <i>Journal of Biological Chemistry</i> , 2003, 278, 24125-24131. | 1.6 | 52 |
| 4508 | Functional Dissection of the Eukaryotic-specific tRNA-interacting Factor of Lysyl-tRNA Synthetase. <i>Journal of Biological Chemistry</i> , 2003, 278, 1472-1479. | 1.6 | 39 |
| 4509 | Isoform-selectivity of PKC Inhibitors Acting at the Regulatory and Catalytic Domain of Mammalian PKC- β , - δ , - ϵ and - ζ . <i>Journal of Enzyme Inhibition and Medicinal Chemistry</i> , 2003, 18, 475-483. | 2.5 | 21 |
| 4510 | Sphingosine Phosphate Lyase Expression Is Essential for Normal Development in <i>Caenorhabditis elegans</i> . <i>Journal of Biological Chemistry</i> , 2003, 278, 22341-22349. | 1.6 | 64 |
| 4511 | Human EFO1p exhibits acetyltransferase activity and is a unique combination of linker histone and Ctf7p/Eco1p chromatid cohesion establishment domains. <i>Nucleic Acids Research</i> , 2003, 31, 6334-6343. | 6.5 | 47 |
| 4512 | A Protein Complex Containing Mdm10p, Mdm12p, and Mmm1p Links Mitochondrial Membranes and DNA to the Cytoskeleton-based Segregation Machinery. <i>Molecular Biology of the Cell</i> , 2003, 14, 4618-4627. | 0.9 | 242 |
| 4513 | A novel human protein of the maternal centriole is required for the final stages of cytokinesis and entry into S phase. <i>Journal of Cell Biology</i> , 2003, 161, 535-545. | 2.3 | 244 |
| 4514 | Uridine Binding Motifs of Human Concentrative Nucleoside Transporters 1 and 3 Produced in <i>Saccharomyces cerevisiae</i> . <i>Molecular Pharmacology</i> , 2003, 64, 1512-1520. | 1.0 | 59 |
| 4515 | A Dispensable Yeast Ribosomal Protein Optimizes Peptidyltransferase Activity and Affects Translocation. <i>Journal of Biological Chemistry</i> , 2003, 278, 3314-3322. | 1.6 | 34 |
| 4516 | Role for Arf3p in Development of Polarity, but Not Endocytosis, in <i>Saccharomyces cerevisiae</i> . <i>Molecular Biology of the Cell</i> , 2003, 14, 3834-3847. | 0.9 | 34 |
| 4517 | Genetic Improvement of Baker's Yeasts. <i>Applied Mycology and Biotechnology</i> , 2003, 3, 213-240. | 0.3 | 1 |
| 4518 | Phosphorylation of the Yeast Phospholipid Synthesis Regulatory Protein Opi1p by Protein Kinase A. <i>Journal of Biological Chemistry</i> , 2003, 278, 20673-20680. | 1.6 | 42 |
| 4519 | G α 3 Recruits Rho1 to the Site of Polarized Growth during Mating in Budding Yeast. <i>Journal of Biological Chemistry</i> , 2003, 278, 21798-21804. | 1.6 | 30 |
| 4520 | Constriction and Dnm1p Recruitment Are Distinct Processes in Mitochondrial Fission. <i>Molecular Biology of the Cell</i> , 2003, 14, 1953-1963. | 0.9 | 168 |
| 4521 | Expression of Heteropolymeric Ferritin Improves Iron Storage in <i>Saccharomyces cerevisiae</i> . <i>Applied and Environmental Microbiology</i> , 2003, 69, 1999-2005. | 1.4 | 32 |

| # | ARTICLE | IF | CITATIONS |
|------|---|-----|-----------|
| 4522 | Molecular Cloning of Endo- β -1,4-Glucanase Genes, <i>rce1</i> , <i>rce2</i> , and <i>rce3</i> , from <i>Rhizopus oryzae</i> . <i>Journal of Bacteriology</i> , 2003, 185, 1749-1756. | 1.0 | 22 |
| 4523 | Effect of the Pheromone-Responsive G β and Phosphatase Proteins of <i>Saccharomyces cerevisiae</i> on the Subcellular Localization of the Fus3 Mitogen-Activated Protein Kinase. <i>Molecular and Cellular Biology</i> , 2003, 23, 1135-1150. | 1.1 | 46 |
| 4524 | Sir2p suppresses recombination of replication forks stalled at the replication fork barrier of ribosomal DNA in <i>Saccharomyces cerevisiae</i> . <i>Nucleic Acids Research</i> , 2003, 31, 893-898. | 6.5 | 35 |
| 4525 | The endo- β -1,3-glucanase <i>eng1p</i> is required for dissolution of the primary septum during cell separation in <i>Schizosaccharomyces pombe</i> . <i>Journal of Cell Science</i> , 2003, 116, 1689-1698. | 1.2 | 163 |
| 4526 | Identification of stop codon readthrough genes in <i>Saccharomyces cerevisiae</i> . <i>Nucleic Acids Research</i> , 2003, 31, 2289-2296. | 6.5 | 100 |
| 4527 | Amino Acid Starvation and <i>Gcn4p</i> Regulate Adhesive Growth and <i>FLO11</i> Gene Expression in <i>Saccharomyces cerevisiae</i> . <i>Molecular Biology of the Cell</i> , 2003, 14, 4272-4284. | 0.9 | 93 |
| 4528 | <i>Saccharomyces cerevisiae</i> Histidine Phosphotransferase <i>Ypd1p</i> Shuttles between the Nucleus and Cytoplasm for <i>SLN1</i> -Dependent Phosphorylation of <i>Ssk1p</i> and <i>Skn7p</i> . <i>Eukaryotic Cell</i> , 2003, 2, 1304-1314. | 3.4 | 78 |
| 4529 | Functional Similarities and Differences between <i>Candida albicans</i> <i>Cdr1p</i> and <i>Cdr2p</i> Transporters. <i>Antimicrobial Agents and Chemotherapy</i> , 2003, 47, 1543-1554. | 1.4 | 50 |
| 4530 | RNA-protein interactions promote asymmetric sorting of the <i>ASH1</i> mRNA ribonucleoprotein complex. <i>Rna</i> , 2003, 9, 1383-1399. | 1.6 | 34 |
| 4531 | An in vivo dual-luciferase assay system for studying translational recoding in the yeast <i>Saccharomyces cerevisiae</i> . <i>Rna</i> , 2003, 9, 1019-1024. | 1.6 | 141 |
| 4532 | Involvement of <i>CjMDR1</i> , a plant multidrug-resistance-type ATP-binding cassette protein, in alkaloid transport in <i>Coptis japonica</i> . <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 2003, 100, 751-756. | 3.3 | 256 |
| 4533 | Replication of Carnation Italian Ringspot Virus Defective Interfering RNA in <i>Saccharomyces cerevisiae</i> . <i>Journal of Virology</i> , 2003, 77, 2116-2123. | 1.5 | 57 |
| 4534 | Cell-type-dependent repression of yeast α -specific genes requires <i>Itc1p</i> , a subunit of the <i>Isw2p</i> α - <i>Itc1p</i> chromatin remodelling complex. <i>Microbiology (United Kingdom)</i> , 2003, 149, 341-351. | 0.7 | 39 |
| 4535 | Molecular Cloning and Characterization of <i>CYP719</i> , a Methyleneedioxy Bridge-forming Enzyme That Belongs to a Novel P450 Family, from cultured <i>Coptis japonica</i> Cells. <i>Journal of Biological Chemistry</i> , 2003, 278, 38557-38565. | 1.6 | 170 |
| 4536 | Functional Interactions between <i>Hsp90</i> and the Co-chaperones <i>Cns1</i> and <i>Cpr7</i> in <i>Saccharomyces cerevisiae</i> . <i>Journal of Biological Chemistry</i> , 2003, 278, 32692-32701. | 1.6 | 45 |
| 4537 | The Ran GTPase cycle is required for yeast nuclear pore complex assembly. <i>Journal of Cell Biology</i> , 2003, 160, 1041-1053. | 2.3 | 104 |
| 4538 | The Yeast hnRNP-like Protein <i>Hrp1/Nab4</i> Accumulates in the Cytoplasm after Hyperosmotic Stress: A Novel <i>Fps1</i> -dependent Response. <i>Molecular Biology of the Cell</i> , 2003, 14, 3929-3941. | 0.9 | 18 |
| 4539 | Mechanical Link between Cohesion Establishment and DNA Replication: <i>Ctf7p/Eco1p</i> , a Cohesion Establishment Factor, Associates with Three Different Replication Factor C Complexes. <i>Molecular and Cellular Biology</i> , 2003, 23, 2999-3007. | 1.1 | 101 |

| # | ARTICLE | IF | CITATIONS |
|------|---|-----|-----------|
| 4540 | Immobilization of <i>Saccharomyces cerevisiae</i> Cells to Protein G-Sepharose by Cell Wall Engineering. <i>Journal of Molecular Microbiology and Biotechnology</i> , 2003, 5, 161-166. | 1.0 | 8 |
| 4541 | Dual Subcellular Distribution of Cytochrome b5 in Plant, Cauliflower, Cells. <i>Journal of Biochemistry</i> , 2003, 133, 115-121. | 0.9 | 43 |
| 4542 | Collaborative Competition Mechanism for Gene Activation In Vivo. <i>Molecular and Cellular Biology</i> , 2003, 23, 1623-1632. | 1.1 | 165 |
| 4543 | Septins, under Cla4p Regulation, and the Chitin Ring Are Required for Neck Integrity in Budding Yeast. <i>Molecular Biology of the Cell</i> , 2003, 14, 2128-2141. | 0.9 | 82 |
| 4544 | Delayed rRNA Processing Results in Significant Ribosome Biogenesis and Functional Defects. <i>Molecular and Cellular Biology</i> , 2003, 23, 1602-1613. | 1.1 | 29 |
| 4545 | Biochemical support for the V-ATPase rotary mechanism: antibody against HA-tagged Vma7p or Vma16p but not Vma10p inhibits activity. <i>Journal of Experimental Biology</i> , 2003, 206, 3227-3237. | 0.8 | 16 |
| 4546 | Substrate Recognition by Nucleotide Sugar Transporters. <i>Journal of Biological Chemistry</i> , 2003, 278, 22887-22893. | 1.6 | 49 |
| 4547 | Molecular Dissection of the Hydrophobic Segments H3 and H4 of the Yeast Ca ²⁺ Channel Component Mid1. <i>Journal of Biological Chemistry</i> , 2003, 278, 9647-9654. | 1.6 | 24 |
| 4548 | Transcription of genes encoding trans-acting factors required for rRNA maturation/ribosomal subunit assembly is coordinately regulated with ribosomal protein genes and involves Rap1 in <i>Saccharomyces cerevisiae</i> . <i>Nucleic Acids Research</i> , 2003, 31, 1969-1973. | 6.5 | 20 |
| 4549 | Functional Interactions between Yeast Translation Eukaryotic Elongation Factor (eEF) 1A and eEF3. <i>Journal of Biological Chemistry</i> , 2003, 278, 6985-6991. | 1.6 | 66 |
| 4550 | Calcineurin Is Essential for <i>Candida albicans</i> Survival in Serum and Virulence. <i>Eukaryotic Cell</i> , 2003, 2, 422-430. | 3.4 | 177 |
| 4551 | Cloning, functional analysis and post-transcriptional regulation of a type II DNA topoisomerase from <i>Leishmania infantum</i> . A new potential target for anti-parasite drugs. <i>Nucleic Acids Research</i> , 2003, 31, 4917-4928. | 6.5 | 26 |
| 4552 | Toxicity of Metal Ions Used in Dental Alloys: A Study in the Yeast <i>Saccharomyces cerevisiae</i> . <i>Drug and Chemical Toxicology</i> , 2003, 26, 75-85. | 1.2 | 28 |
| 4553 | Co-translational, Intraribosomal Cleavage of Polypeptides by the Foot-and-mouth Disease Virus 2A Peptide. <i>Journal of Biological Chemistry</i> , 2003, 278, 11441-11448. | 1.6 | 149 |
| 4554 | Molecular Cloning and Identification of 3'-Phosphoadenosine 5'-Phosphosulfate Transporter. <i>Journal of Biological Chemistry</i> , 2003, 278, 25958-25963. | 1.6 | 123 |
| 4555 | Yeast Epiarginase Regulation, an Enzyme-Enzyme Activity Control. <i>Journal of Biological Chemistry</i> , 2003, 278, 21550-21558. | 1.6 | 7 |
| 4556 | Human Mitochondrial C1-Tetrahydrofolate Synthase. <i>Journal of Biological Chemistry</i> , 2003, 278, 43178-43187. | 1.6 | 51 |
| 4557 | Ser3p (Yer081wp) and Ser33p (Yil074cp) Are Phosphoglycerate Dehydrogenases in <i>Saccharomyces cerevisiae</i> . <i>Journal of Biological Chemistry</i> , 2003, 278, 10264-10272. | 1.6 | 63 |

| # | ARTICLE | IF | CITATIONS |
|------|--|-----|-----------|
| 4558 | Phosphorylation of CTP Synthetase on Ser36, Ser330, Ser354, and Ser454 Regulates the Levels of CTP and Phosphatidylcholine Synthesis in <i>Saccharomyces cerevisiae</i> . <i>Journal of Biological Chemistry</i> , 2003, 278, 20785-20794. | 1.6 | 44 |
| 4559 | Involvement of the VDE Homing Endonuclease and Rapamycin in Regulation of the <i>Saccharomyces cerevisiae</i> GSH11 Gene Encoding the High Affinity Glutathione Transporter. <i>Journal of Biological Chemistry</i> , 2003, 278, 39632-39636. | 1.6 | 16 |
| 4560 | Kinectin Anchors the Translation Elongation Factor-1 β to the Endoplasmic Reticulum. <i>Journal of Biological Chemistry</i> , 2003, 278, 32115-32123. | 1.6 | 29 |
| 4561 | Functional Characterization and Localization of Acetyl-CoA Hydrolase, Ach1p, in <i>Saccharomyces cerevisiae</i> . <i>Journal of Biological Chemistry</i> , 2003, 278, 17203-17209. | 1.6 | 45 |
| 4562 | Degradation of Normal mRNA in the Nucleus of <i>Saccharomyces cerevisiae</i> . <i>Molecular and Cellular Biology</i> , 2003, 23, 5502-5515. | 1.1 | 108 |
| 4563 | A yeast-based functional assay for the detection of the mutant androgen receptor in prostate cancer. <i>European Journal of Endocrinology</i> , 2003, 148, 99-110. | 1.9 | 20 |
| 4564 | Novel Rho GTPase Involved in Cytokinesis and Cell Wall Integrity in the Fission Yeast <i>Schizosaccharomyces pombe</i> . <i>Eukaryotic Cell</i> , 2003, 2, 521-533. | 3.4 | 48 |
| 4565 | The Central Region of Human T-Cell Leukemia Virus Type 1 Tax Protein Contains Distinct Domains Involved in Subunit Dimerization. <i>Journal of Virology</i> , 2003, 77, 13028-13035. | 1.5 | 33 |
| 4566 | The Proton Transfer Step Catalyzed by Yeast Pyruvate Kinase. <i>Journal of Biological Chemistry</i> , 2003, 278, 12660-12671. | 1.6 | 22 |
| 4567 | The Unfolded Protein Response Is Required for Haploid Tolerance in Yeast. <i>Journal of Biological Chemistry</i> , 2003, 278, 11818-11827. | 1.6 | 17 |
| 4568 | Molecular Evolution of the Rab-Escort-Protein/Guanine-Nucleotide-Dissociation-Inhibitor Superfamily. <i>Molecular Biology of the Cell</i> , 2003, 14, 3857-3867. | 0.9 | 31 |
| 4569 | Detection of Protein Kinase-Binding Partners by the Yeast Two-Hybrid Analysis. , 2003, 233, 327-344. | | 0 |
| 4570 | Analysis of Transcriptional Repression by Mig1 in <i>Saccharomyces cerevisiae</i> Using a Reporter Assay. <i>Methods in Enzymology</i> , 2003, 371, 602-614. | 0.4 | 3 |
| 4571 | Topology Models for 37 <i>Saccharomyces cerevisiae</i> Membrane Proteins Based on C-terminal Reporter Fusions and Predictions. <i>Journal of Biological Chemistry</i> , 2003, 278, 10208-10213. | 1.6 | 58 |
| 4572 | Dissociation of Steroid Receptor Coactivator 1 and Nuclear Receptor Corepressor Recruitment to the Human Glucocorticoid Receptor by Modification of the Ligand-Receptor Interface: The Role of Tyrosine 735. <i>Molecular Endocrinology</i> , 2003, 17, 845-859. | 3.7 | 56 |
| 4573 | Inhibition of $\hat{\pm}$, $\hat{2}$ I, $\hat{1}$, $\hat{1}$ and $\hat{1}$ η Protein Kinase C Isoforms by Xanthonolignoids. <i>Journal of Enzyme Inhibition and Medicinal Chemistry</i> , 2003, 18, 357-370. | 2.5 | 18 |
| 4574 | Effect of expression of the <i>Escherichia coli</i> nth gene in <i>Saccharomyces cerevisiae</i> on the toxicity of ionizing radiation and hydrogen peroxide. <i>International Journal of Radiation Biology</i> , 2003, 79, 747-755. | 1.0 | 3 |
| 4575 | Functional characterization in yeast of genetic variants in the human equilibrative nucleoside transporter, ENT1. <i>Pharmacogenetics and Genomics</i> , 2003, 13, 297-301. | 5.7 | 71 |

| # | ARTICLE | IF | CITATIONS |
|------|---|------|-----------|
| 4576 | Splyregulation of sphingolipid signaling molecules is essential for Drosophila development. <i>Development (Cambridge)</i> , 2003, 130, 2443-2453. | 1.2 | 106 |
| 4577 | A neuronal-specific differentiation protein that directly modulates retinoid receptor transcriptional activation. <i>Nuclear Receptor</i> , 2003, 1, 7. | 10.0 | 6 |
| 4578 | Nitrogen-source regulation of yeast gamma-glutamyl transpeptidase synthesis involves the regulatory network including the GATA zinc-finger factors Gln3, Nil1/Gat1 and Gzf3. <i>Biochemical Journal</i> , 2003, 371, 589-595. | 1.7 | 64 |
| 4579 | Inhibition of protein kinase C catalytic activity by additional regions within the human protein kinase Calpha-regulatory domain lying outside of the pseudosubstrate sequence. <i>Biochemical Journal</i> , 2003, 373, 571-581. | 1.7 | 17 |
| 4580 | Molecular Evolution of the Rab-Escort-Protein/Guanine-Nucleotide-Dissociation-Inhibitor Superfamily. <i>Molecular Biology of the Cell</i> , 2003, 14, 3857-3867. | 0.9 | 35 |
| 4581 | Meiotic and Mitotic Phenotypes Conferred by the blm1-1 Mutation in <i>Saccharomyces cerevisiae</i> and MSH4 Suppression of the Bleomycin Hypersusceptibility. <i>International Journal of Molecular Sciences</i> , 2003, 4, 1-12. | 1.8 | 4 |
| 4582 | Screening for suppressors of temperature sensitivity in a yeast mutant defective in vacuolar protein degradation. <i>Genetics and Molecular Biology</i> , 2003, 26, 89-98. | 0.6 | 1 |
| 4583 | Transformation of yeast using calcium alginate microbeads with surface-immobilized chromosomal DNA. <i>BioTechniques</i> , 2003, 35, 734-740. | 0.8 | 22 |
| 4584 | Expression of the bacterial recA gene impairs genetic recombination and sporulation in a <i>Saccharomyces cerevisiae</i> diploid strain. <i>Genetics and Molecular Biology</i> , 2003, 26, 213-220. | 0.6 | 1 |
| 4585 | Large-scale yeast transformation in low-percentage agarose medium. <i>BioTechniques</i> , 2004, 36, 40-44. | 0.8 | 10 |
| 4586 | Ybr267w is a New Cytoplasmic Protein Belonging to the Mitotic Signaling Network of <i>Saccharomyces cerevisiae</i> . <i>Cell Structure and Function</i> , 2004, 29, 1-15. | 0.5 | 30 |
| 4587 | Molecular Cloning of a Gene Encoding Endo-Î²-D-1,4-Glucanase PCE1 from <i>Phycomyces nitens</i> . <i>Bioscience, Biotechnology and Biochemistry</i> , 2004, 68, 2299-2305. | 0.6 | 12 |
| 4588 | The Phox Homology (PX) Domain Protein Interaction Network in Yeast. <i>Molecular and Cellular Proteomics</i> , 2004, 3, 1053-1064. | 2.5 | 53 |
| 4589 | Three Arabidopsis MBF1 Homologs with Distinct Expression Profiles Play Roles as Transcriptional Co-activators. <i>Plant and Cell Physiology</i> , 2004, 45, 225-231. | 1.5 | 65 |
| 4590 | A Novel Salt-tolerant l-myo-Inositol-1-phosphate Synthase from <i>Porteresia coarctata</i> (Roxb.) Tateoka, a Halophytic Wild Rice. <i>Journal of Biological Chemistry</i> , 2004, 279, 28539-28552. | 1.6 | 169 |
| 4591 | Cloning and Expression of clt Genes Encoding Milk-Clotting Proteases from <i>Myxococcus xanthus</i> 422. <i>Applied and Environmental Microbiology</i> , 2004, 70, 6337-6341. | 1.4 | 18 |
| 4592 | Vacuole Size Control: Regulation of PtdIns(3,5)P2 Levels by the Vacuole-associated Vac14-Fig4 Complex, a PtdIns(3,5)P2-specific Phosphatase. <i>Molecular Biology of the Cell</i> , 2004, 15, 24-36. | 0.9 | 191 |
| 4593 | A Novel Arabidopsis Gene Required for Ethanol Tolerance is Conserved Among Plants and Archaea. <i>Plant and Cell Physiology</i> , 2004, 45, 659-666. | 1.5 | 13 |

| # | ARTICLE | IF | CITATIONS |
|------|--|-----|-----------|
| 4594 | A method for cloning and sequencing long palindromic DNA junctions. <i>Nucleic Acids Research</i> , 2004, 32, e155-e155. | 6.5 | 20 |
| 4595 | Bipartite 3' cis-acting Signal for Replication in Yeast 23 S RNA Virus and Its Repair. <i>Journal of Biological Chemistry</i> , 2004, 279, 13215-13223. | 1.6 | 17 |
| 4596 | Synthesis of Novel Lipids in <i>Saccharomyces cerevisiae</i> by Heterologous Expression of an Unspecific Bacterial Acyltransferase. <i>Applied and Environmental Microbiology</i> , 2004, 70, 7119-7125. | 1.4 | 119 |
| 4597 | DNA Fragment Transplacement in <i>Saccharomyces cerevisiae</i> : Some Genetic Considerations. , 2004, 262, 157-172. | | 5 |
| 4598 | Pdr3 is required for DNA damage induction of MAG1 and DDI1 via a bi-directional promoter element. <i>Nucleic Acids Research</i> , 2004, 32, 5066-5075. | 6.5 | 27 |
| 4599 | Comparison of Two Alternative Dominant Selectable Markers for Wine Yeast Transformation. <i>Applied and Environmental Microbiology</i> , 2004, 70, 7018-7023. | 1.4 | 23 |
| 4600 | <i>Arabidopsis</i> HMA2, a Divalent Heavy Metal-Transporting PIB-Type ATPase, Is Involved in Cytoplasmic Zn ²⁺ Homeostasis. <i>Plant Physiology</i> , 2004, 136, 3712-3723. | 2.3 | 206 |
| 4601 | The External Amino Acid Signaling Pathway Promotes Activation of Stp1 and Uga35/Dal81 Transcription Factors for Induction of the AGP1 Gene in <i>Saccharomyces cerevisiae</i> . <i>Genetics</i> , 2004, 166, 1727-1739. | 1.2 | 45 |
| 4602 | Genetic Evidence of a Role for Membrane Lipid Composition in the Regulation of Soluble NEM-Sensitive Factor Receptor Function in <i>Saccharomyces cerevisiae</i> . <i>Genetics</i> , 2004, 166, 89-97. | 1.2 | 26 |
| 4603 | Copper and Iron Are the Limiting Factors for Growth of the Yeast <i>Saccharomyces cerevisiae</i> in an Alkaline Environment. <i>Journal of Biological Chemistry</i> , 2004, 279, 19698-19704. | 1.6 | 118 |
| 4604 | Functional Characterization of Myosin I Tail Regions in <i>Candida albicans</i> . <i>Eukaryotic Cell</i> , 2004, 3, 1272-1286. | 3.4 | 24 |
| 4605 | Multiple Targeting Modules on Peroxisomal Proteins Are Not Redundant: Discrete Functions of Targeting Signals within Pmp47 and Pex8p. <i>Molecular Biology of the Cell</i> , 2004, 15, 1702-1710. | 0.9 | 22 |
| 4606 | U17/snrR30 Is a Ubiquitous snoRNA with Two Conserved Sequence Motifs Essential for 18S rRNA Production. <i>Molecular and Cellular Biology</i> , 2004, 24, 1769-1778. | 1.1 | 84 |
| 4607 | Degradation of Mutated Bovine Pancreatic Trypsin Inhibitor in the Yeast Vacuole Suggests Post-endoplasmic Reticulum Protein Quality Control. <i>Journal of Biological Chemistry</i> , 2004, 279, 15289-15297. | 1.6 | 64 |
| 4608 | Human MutL homolog (MLH1) function in DNA mismatch repair: a prospective screen for missense mutations in the ATPase domain. <i>Nucleic Acids Research</i> , 2004, 32, 5321-5338. | 6.5 | 38 |
| 4609 | Potent inhibition of huntingtin aggregation and cytotoxicity by a disulfide bond-free single-domain intracellular antibody. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 2004, 101, 17616-17621. | 3.3 | 173 |
| 4610 | A Short-range Gradient of Histone H3 Acetylation and Tup1p Redistribution at the Promoter of the <i>Saccharomyces cerevisiae</i> SUC2 Gene. <i>Journal of Biological Chemistry</i> , 2004, 279, 7678-7684. | 1.6 | 18 |
| 4611 | Nuclear surveillance and degradation of hypomodified initiator tRNA ^{Met} in <i>S. cerevisiae</i> . <i>Genes and Development</i> , 2004, 18, 1227-1240. | 2.7 | 426 |

| # | ARTICLE | IF | CITATIONS |
|------|---|-----|-----------|
| 4612 | Simultaneous yet Independent Regulation of Actin Cytoskeletal Organization and Translation Initiation by Glucose in <i>Saccharomyces cerevisiae</i> . <i>Molecular Biology of the Cell</i> , 2004, 15, 1544-1556. | 0.9 | 43 |
| 4613 | Rare, Highly Pyrimethamine-Resistant Alleles of the <i>Plasmodium falciparum</i> Dihydrofolate Reductase Gene from 5 African Sites. <i>Journal of Infectious Diseases</i> , 2004, 190, 1783-1792. | 1.9 | 19 |
| 4614 | Characterization of a glycosylphosphatidylinositol-bound cell-wall protein (GPI-CWP) in <i>Yarrowia lipolytica</i> . <i>Microbiology (United Kingdom)</i> , 2004, 150, 53-60. | 0.7 | 28 |
| 4615 | Yeast Kre1p is GPI-anchored and involved in both cell wall assembly and architecture. <i>Microbiology (United Kingdom)</i> , 2004, 150, 3209-3218. | 0.7 | 33 |
| 4616 | Ebp2p, the Yeast Homolog of Epstein-Barr Virus Nuclear Antigen 1-binding Protein 2, Interacts with Factors of Both the 60 S and the 40 S Ribosomal Subunit Assembly. <i>Journal of Biological Chemistry</i> , 2004, 279, 25353-25358. | 1.6 | 28 |
| 4617 | Genetic Interactions of DST1 in <i>Saccharomyces cerevisiae</i> Suggest a Role of TFIIIS in the Initiation-Elongation Transition. <i>Genetics</i> , 2004, 166, 1215-1227. | 1.2 | 48 |
| 4618 | Position-dependent function of a B block promoter element implies a specialized chromatin structure on the <i>S.cerevisiae</i> U6 RNA gene, SNR6. <i>Nucleic Acids Research</i> , 2004, 32, 4297-4305. | 6.5 | 8 |
| 4619 | Roles of Phosphoinositides and of Spo14p (phospholipase D)-generated Phosphatidic Acid during Yeast Sporulation. <i>Molecular Biology of the Cell</i> , 2004, 15, 207-218. | 0.9 | 63 |
| 4620 | Mutational analysis of the cytoplasmic domain of the Wsc1 cell wall stress sensor. <i>Microbiology (United Kingdom)</i> , 2004, 150, 3281-3288. | 0.7 | 32 |
| 4621 | Nuclear Export of hnRNP Hrp1p and Nuclear Export of hnRNP Npl3p Are Linked and Influenced by the Methylation State of Npl3p. <i>Molecular and Cellular Biology</i> , 2004, 24, 10742-10756. | 1.1 | 48 |
| 4622 | Evidence against a direct role for the Upf proteins in frameshifting or nonsense codon readthrough. <i>Rna</i> , 2004, 10, 1721-1729. | 1.6 | 32 |
| 4623 | Assembly of Turnip Yellow Mosaic Virus Replication Complexes: Interaction between the Proteinase and Polymerase Domains of the Replication Proteins. <i>Journal of Virology</i> , 2004, 78, 7945-7957. | 1.5 | 50 |
| 4624 | Receptor (CD155)-Dependent Endocytosis of Poliovirus and Retrograde Axonal Transport of the Endosome. <i>Journal of Virology</i> , 2004, 78, 7186-7198. | 1.5 | 114 |
| 4625 | Functional Unit of the RNA Polymerase II C-Terminal Domain Lies within Heptapeptide Pairs. <i>Eukaryotic Cell</i> , 2004, 3, 735-740. | 3.4 | 62 |
| 4626 | Biochemical evidence of translational infidelity and decreased peptidyltransferase activity by a sarcin/ricin domain mutation of yeast 25S rRNA. <i>Nucleic Acids Research</i> , 2004, 32, 5398-5408. | 6.5 | 10 |
| 4627 | Chl1p, a DNA Helicase-Like Protein in Budding Yeast, Functions in Sister-Chromatid Cohesion. <i>Genetics</i> , 2004, 166, 33-42. | 1.2 | 139 |
| 4628 | RAD51 -Dependent Break-Induced Replication in Yeast. <i>Molecular and Cellular Biology</i> , 2004, 24, 2344-2351. | 1.1 | 172 |
| 4629 | Dihydrofolate Reductase Mutations in <i>Plasmodium vivax</i> from Indonesia and Therapeutic Response to Sulfadoxine plus Pyrimethamine. <i>Journal of Infectious Diseases</i> , 2004, 189, 744-750. | 1.9 | 95 |

| # | ARTICLE | IF | CITATIONS |
|------|--|-----|-----------|
| 4630 | Cloning of S-Adenosyl-L-Methionine:C-24- β -Sterol-Methyltransferase (ERG6) from <i>Leishmania donovani</i> and Characterization of mRNAs in Wild-Type and Amphotericin B-Resistant Promastigotes. <i>Antimicrobial Agents and Chemotherapy</i> , 2004, 48, 2409-2414. | 1.4 | 65 |
| 4631 | The p36 and p95 replicase proteins of Carnation Italian ringspot virus cooperate in stabilizing defective interfering RNA. <i>Journal of General Virology</i> , 2004, 85, 2429-2433. | 1.3 | 15 |
| 4632 | Pyridoxine biosynthesis in yeast: participation of ribose 5-phosphate ketol-isomerase. <i>Biochemical Journal</i> , 2004, 379, 65-70. | 1.7 | 24 |
| 4633 | Molecular Cloning and Characterization of a Human Multisubstrate Specific Nucleotide-sugar Transporter Homologous to <i>Drosophila</i> fringe connection. <i>Journal of Biological Chemistry</i> , 2004, 279, 26469-26474. | 1.6 | 61 |
| 4634 | Domain Organization of Phytochelatin Synthase. <i>Journal of Biological Chemistry</i> , 2004, 279, 14686-14693. | 1.6 | 72 |
| 4635 | Regulation of Phospholipid Synthesis in <i>Saccharomyces cerevisiae</i> by Zinc. <i>Journal of Biological Chemistry</i> , 2004, 279, 21976-21983. | 1.6 | 75 |
| 4636 | Dissecting the Cell-killing Mechanism of the Topoisomerase II-targeting Drug ICRF-193. <i>Journal of Biological Chemistry</i> , 2004, 279, 28100-28105. | 1.6 | 19 |
| 4637 | Regulation of the Yeast EK11-encoded Ethanolamine Kinase by Inositol and Choline. <i>Journal of Biological Chemistry</i> , 2004, 279, 35353-35359. | 1.6 | 15 |
| 4638 | Cytoplasmic Inositol Hexakisphosphate Production Is Sufficient for Mediating the Gle1-mRNA Export Pathway. <i>Journal of Biological Chemistry</i> , 2004, 279, 51022-51032. | 1.6 | 45 |
| 4639 | DNA Stimulates Mec1-mediated Phosphorylation of Replication Protein A. <i>Journal of Biological Chemistry</i> , 2004, 279, 26762-26767. | 1.6 | 30 |
| 4640 | Role for the Ran Binding Protein, Mog1p, in <i>Saccharomyces cerevisiae</i> SLN1-SKN7 Signal Transduction. <i>Eukaryotic Cell</i> , 2004, 3, 1544-1556. | 3.4 | 24 |
| 4641 | Sgt1 Associates with Hsp90: an Initial Step of Assembly of the Core Kinetochores Complex. <i>Molecular and Cellular Biology</i> , 2004, 24, 8069-8079. | 1.1 | 77 |
| 4642 | Host Factors That Affect Ty3 Retrotransposition in <i>Saccharomyces cerevisiae</i> . <i>Genetics</i> , 2004, 168, 1159-1176. | 1.2 | 47 |
| 4643 | Swi5 Acts in Meiotic DNA Joint Molecule Formation in <i>Schizosaccharomyces pombe</i> . <i>Genetics</i> , 2004, 168, 1891-1898. | 1.2 | 50 |
| 4644 | CDC7/DBF4 Functions in the Translesion Synthesis Branch of the RAD6 Epistasis Group in <i>Saccharomyces cerevisiae</i> . <i>Genetics</i> , 2004, 167, 1597-1610. | 1.2 | 71 |
| 4645 | The major 5' determinant in stop codon read-through involves two adjacent adenines. <i>Nucleic Acids Research</i> , 2004, 32, 415-421. | 6.5 | 74 |
| 4646 | The Yeast RNA-binding Protein Rbp1p Modifies the Stability of Mitochondrial Porin mRNA. <i>Journal of Biological Chemistry</i> , 2004, 279, 453-462. | 1.6 | 14 |
| 4647 | The ER-Golgi v-SNARE Bet1p is required for cross-linking β -agglutinin to the cell wall in yeast. <i>Microbiology (United Kingdom)</i> , 2004, 150, 3219-3228. | 0.7 | 6 |

| # | ARTICLE | IF | CITATIONS |
|------|---|-----|-----------|
| 4648 | Hindering the Strand Passage Reaction of Human Topoisomerase III α without Disturbing DNA Cleavage, ATP Hydrolysis, or the Operation of the N-terminal Clamp. <i>Journal of Biological Chemistry</i> , 2004, 279, 28093-28099. | 1.6 | 7 |
| 4649 | Activation of the RAS/Cyclic AMP Pathway Suppresses a TOR Deficiency in Yeast. <i>Molecular and Cellular Biology</i> , 2004, 24, 338-351. | 1.1 | 239 |
| 4650 | Applicability of Tandem Affinity Purification MudPIT to Pathway Proteomics in Yeast. <i>Molecular and Cellular Proteomics</i> , 2004, 3, 226-237. | 2.5 | 130 |
| 4651 | The Meiosis-specific Protein Kinase Ime2 Directs Phosphorylation of Replication Protein A. <i>Journal of Biological Chemistry</i> , 2004, 279, 6163-6170. | 1.6 | 32 |
| 4652 | ASH1 mRNA Anchoring Requires Reorganization of the Myo4p-She3p-She2p Transport Complex. <i>Journal of Biological Chemistry</i> , 2004, 279, 46286-46294. | 1.6 | 21 |
| 4653 | Regulation of Phospholipid Synthesis in the Yeast <i>cki1Δ eki1Δ</i> Mutant Defective in the Kennedy Pathway. <i>Journal of Biological Chemistry</i> , 2004, 279, 12081-12087. | 1.6 | 18 |
| 4654 | In Vivo Analysis of Synaptonemal Complex Formation During Yeast Meiosis. <i>Genetics</i> , 2004, 167, 51-63. | 1.2 | 38 |
| 4655 | Yeast ARL1 encodes a regulator of K ⁺ influx. <i>Journal of Cell Science</i> , 2004, 117, 2309-2320. | 1.2 | 37 |
| 4656 | The Transducer Domain Is Important for Clamp Operation in Human DNA Topoisomerase III α . <i>Journal of Biological Chemistry</i> , 2004, 279, 1684-1691. | 1.6 | 22 |
| 4657 | A Mutation in the Yeast Mitochondrial Core RNA Polymerase, Rpo41, Confers Defects in Both Specificity Factor Interaction and Promoter Utilization. <i>Journal of Biological Chemistry</i> , 2004, 279, 2012-2019. | 1.6 | 18 |
| 4658 | Localization of a promoter in the putative internal ribosome entry site of the <i>Saccharomyces cerevisiae</i> TIF4631 gene. <i>Rna</i> , 2004, 10, 277-286. | 1.6 | 23 |
| 4659 | Negative Regulation of Phosphatidylinositol 4,5-Bisphosphate Levels by the INP51-associated Proteins TAX4 and IRS4. <i>Journal of Biological Chemistry</i> , 2004, 279, 39604-39610. | 1.6 | 20 |
| 4660 | Elucidation of the Interactions between C99, Presenilin, and Nicastrin by the Split-Ubiquitin Assay. <i>DNA and Cell Biology</i> , 2004, 23, 59-65. | 0.9 | 11 |
| 4661 | Gain and Loss of Fruit Flavor Compounds Produced by Wild and Cultivated Strawberry Species. <i>Plant Cell</i> , 2004, 16, 3110-3131. | 3.1 | 427 |
| 4662 | The Cotton GhNHX1 Gene Encoding a Novel Putative Tonoplast Na ⁺ /H ⁺ Antiporter Plays an Important Role in Salt Stress. <i>Plant and Cell Physiology</i> , 2004, 45, 600-607. | 1.5 | 240 |
| 4663 | The Translation Elongation Factor eEF1B Plays a Role in the Oxidative Stress Response Pathway. <i>RNA Biology</i> , 2004, 1, 89-94. | 1.5 | 46 |
| 4664 | Mutations in AAC2, equivalent to human adPEO-associated ANT1 mutations, lead to defective oxidative phosphorylation in <i>Saccharomyces cerevisiae</i> and affect mitochondrial DNA stability. <i>Human Molecular Genetics</i> , 2004, 13, 923-934. | 1.4 | 71 |
| 4665 | The mRNA Export Factor Human Gle1 Interacts with the Nuclear Pore Complex Protein Nup155. <i>Molecular and Cellular Proteomics</i> , 2004, 3, 145-155. | 2.5 | 52 |

| # | ARTICLE | IF | CITATIONS |
|------|--|-----|-----------|
| 4666 | Characterization of the products of the genes SNO1 and SNZ1 involved in pyridoxine synthesis in <i>Saccharomyces cerevisiae</i> . <i>FEBS Journal</i> , 2004, 271, 745-752. | 0.2 | 65 |
| 4667 | Further insights into the assembly of the yeast cytochrome <i>b</i> ₁ complex based on analysis of single and double deletion mutants lacking supernumerary subunits and cytochrome <i>b</i> . <i>FEBS Journal</i> , 2004, 271, 1209-1218. | 0.2 | 44 |
| 4668 | Two functionally redundant isoforms of <i>Drosophila melanogaster</i> eukaryotic initiation factor 4B are involved in cap-dependent translation, cell survival, and proliferation. <i>FEBS Journal</i> , 2004, 271, 2923-2936. | 0.2 | 24 |
| 4669 | The highly conserved extracellular peptide, DSYG(893-896), is a critical structure for sodium pump function. <i>FEBS Journal</i> , 2004, 271, 3821-3831. | 0.2 | 8 |
| 4670 | Autoregulation of the HAC1 gene is required for sustained activation of the yeast unfolded protein response. <i>Genes To Cells</i> , 2004, 9, 95-104. | 0.5 | 53 |
| 4671 | Phosphorylation of glucosamine-6-phosphate synthase is important but not essential for germination and mycelial growth of <i>Candida albicans</i> . <i>FEMS Microbiology Letters</i> , 2004, 235, 73-80. | 0.7 | 17 |
| 4672 | FLO11 is essential for flor formation caused by the C-terminal deletion of NRG1 in <i>Saccharomyces cerevisiae</i> . <i>FEMS Microbiology Letters</i> , 2004, 237, 425-430. | 0.7 | 34 |
| 4673 | Cloning and characterization of the gene encoding a high-affinity maltose transporter from. <i>FEMS Yeast Research</i> , 2004, 4, 467-476. | 1.1 | 16 |
| 4674 | Inactivation of the RRB1-Pescadillo pathway involved in ribosome biogenesis induces chromosomal instability. <i>Oncogene</i> , 2004, 23, 8597-8602. | 2.6 | 61 |
| 4675 | Identification of novel dominant INO2 ^Δ mutants with an Opi- phenotype. <i>Molecular Microbiology</i> , 2004, 52, 1271-1280. | 1.2 | 14 |
| 4676 | Isolation of novel catalytic antibody clones from combinatorial library displayed on yeast-cell surface. <i>Journal of Molecular Catalysis B: Enzymatic</i> , 2004, 28, 247-251. | 1.8 | 8 |
| 4677 | Functional analysis of chemically-induced mutations at the flounder TP53 locus, the FACIM assay. <i>Mutation Research - Fundamental and Molecular Mechanisms of Mutagenesis</i> , 2004, 552, 51-60. | 0.4 | 8 |
| 4678 | A hap1 mutation in a laboratory strain of <i>Saccharomyces cerevisiae</i> results in decreased expression of ergosterol-related genes and cellular ergosterol content compared to sake yeast. <i>Journal of Bioscience and Bioengineering</i> , 2004, 98, 159-166. | 1.1 | 45 |
| 4679 | Efficient selection of hybrids by protoplast fusion using drug resistance markers and reporter genes in <i>Saccharomyces cerevisiae</i> . <i>Journal of Bioscience and Bioengineering</i> , 2004, 98, 353-358. | 1.1 | 14 |
| 4680 | The promoter of the yeast OPI1 regulatory gene. <i>Journal of Bioscience and Bioengineering</i> , 2004, 97, 369-373. | 1.1 | 6 |
| 4681 | Non-histone proteins Nhp6A and Nhp6B are required for the regulated expression of SUC2 gene of <i>Saccharomyces cerevisiae</i> . <i>Journal of Bioscience and Bioengineering</i> , 2004, 98, 9-13. | 1.1 | 3 |
| 4682 | Characterization of phorbol esters activity on individual mammalian protein kinase C isoforms, using the yeast phenotypic assay. <i>European Journal of Pharmacology</i> , 2004, 491, 101-110. | 1.7 | 27 |
| 4683 | Suppressor analysis points to the subtle role of the LAG1 ceramide synthase gene in determining yeast longevity. <i>Experimental Gerontology</i> , 2004, 39, 999-1009. | 1.2 | 45 |

| # | ARTICLE | IF | CITATIONS |
|------|---|-----|-----------|
| 4684 | Isolation and characterization of Arabidopsis thaliana ISU1 gene. <i>Biochimica Et Biophysica Acta Gene Regulatory Mechanisms</i> , 2004, 1680, 171-175. | 2.4 | 18 |
| 4685 | Lead optimization of antifungal peptides with 3D NMR structures analysis. <i>Protein Science</i> , 2004, 13, 703-713. | 3.1 | 37 |
| 4686 | Functional expression in yeast of an N-deleted form of At-ACA8, a plasma membrane Ca ²⁺ -ATPase of Arabidopsis thaliana, and characterization of a hyperactive mutant. <i>Planta</i> , 2004, 218, 814-823. | 1.6 | 33 |
| 4687 | sll1722, an unassigned open reading frame of Synechocystis PCC7942/6803, codes for l- myo -inositol 1-phosphate synthase. <i>Planta</i> , 2004, 218, 989-998. | 1.6 | 14 |
| 4688 | Analyses of the effects of Rck2p mutants on Pbs2pDD-induced toxicity in Saccharomyces cerevisiae identify a MAP kinase docking motif, and unexpected functional inactivation due to acidic substitution of T379. <i>Molecular Genetics and Genomics</i> , 2004, 271, 208-219. | 1.0 | 44 |
| 4689 | TamA interacts with LeuB, the homologue of Saccharomyces cerevisiae Leu3p, to regulate gdhA expression in Aspergillus nidulans. <i>Molecular Genetics and Genomics</i> , 2004, 272, 452-459. | 1.0 | 20 |
| 4690 | Expression of a functional human tumor necrosis factor- α (hTNF- α) in yeast Saccharomyces cerevisiae. <i>Biotechnology and Bioprocess Engineering</i> , 2004, 9, 292-296. | 1.4 | 10 |
| 4691 | Genetic and cytological characterization of the RecA-homologous proteins Rad51 and Dmc1 of Schizosaccharomyces pombe. <i>Current Genetics</i> , 2004, 44, 317-328. | 0.8 | 22 |
| 4692 | Regulation of glycolysis in Kluyveromyces lactis : role of KIGCR1 and KIGCR2 in glucose uptake and catabolism. <i>Current Genetics</i> , 2004, 45, 129-139. | 0.8 | 19 |
| 4693 | Genetic analysis of apomictic wine yeasts. <i>Current Genetics</i> , 2004, 45, 187-196. | 0.8 | 9 |
| 4694 | Functional characterisation and transcriptional regulation of the KIHEM12 gene from Kluyveromyces lactis. <i>Current Genetics</i> , 2004, 46, 147-57. | 0.8 | 6 |
| 4695 | Isolation and characterization of Candida albicans homologue of RAP1, a repressor and activator protein gene in Saccharomyces cerevisiae.. <i>Yeast</i> , 2004, 21, 1-10. | 0.8 | 11 |
| 4696 | A simple and efficient procedure for transformation of Schizosaccharomyces pombe. <i>Yeast</i> , 2004, 21, 613-617. | 0.8 | 73 |
| 4697 | Role of the N-terminal region of Rap1p in the transcriptional activation of glycolytic genes in Saccharomyces cerevisiae. <i>Yeast</i> , 2004, 21, 851-866. | 0.8 | 20 |
| 4698 | Characterization of triacylglycerols from overwintering prepupae of the alfalfa pollinator Megachile rotundata (Hymenoptera: Megachilidae). <i>Archives of Insect Biochemistry and Physiology</i> , 2004, 57, 1-14. | 0.6 | 28 |
| 4699 | Synergistic operation of four -acting elements mediate high level transcription in. <i>FEMS Yeast Research</i> , 2004, 5, 29-41. | 1.1 | 10 |
| 4700 | Catalytic Reaction of Basidiomycete Lentinula edodes Cytochrome P450, Le.CYP1 Enzyme Produced in Yeast. <i>Bioscience, Biotechnology and Biochemistry</i> , 2004, 68, 79-84. | 0.6 | 4 |
| 4701 | Uridine Recognition Motifs of Human Equilibrative Nucleoside Transporters 1 and 2 Produced in Saccharomyces cerevisiae. <i>Nucleosides, Nucleotides and Nucleic Acids</i> , 2004, 23, 361-373. | 0.4 | 45 |

| # | ARTICLE | IF | CITATIONS |
|------|--|------|-----------|
| 4702 | Role of Pir1 in the construction of the <i>Candida albicans</i> cell wall. <i>Microbiology (United Kingdom)</i> , 2004, 150, 3151-3161. | 0.7 | 57 |
| 4703 | Role of the yeast Rrp1 protein in the dynamics of pre-ribosome maturation. <i>Rna</i> , 2004, 10, 813-827. | 1.6 | 89 |
| 4704 | Three Decades of Fungal Transformation: Key Concepts and Applications. , 2004, 267, 297-314. | | 27 |
| 4705 | Systematic analysis of bicistronic reporter assay data. <i>Nucleic Acids Research</i> , 2004, 32, e160-e160. | 6.5 | 90 |
| 4706 | Mis16 and Mis18 Are Required for CENP-A Loading and Histone Deacetylation at Centromeres. <i>Cell</i> , 2004, 118, 715-729. | 13.5 | 391 |
| 4707 | Involvement of two endonuclease III homologs in the base excision repair pathway for the processing of DNA alkylation damage in <i>Saccharomyces cerevisiae</i> . <i>DNA Repair</i> , 2004, 3, 51-59. | 1.3 | 39 |
| 4708 | Cloning and characterization in <i>Pichia pastoris</i> of PNO1 gene required for phosphomannosylation of N-linked oligosaccharides. <i>Gene</i> , 2004, 324, 129-137. | 1.0 | 20 |
| 4709 | The European flounder (<i>Platichthys flesus</i>) TP53 functions as a temperature-sensitive transcription factor which inhibits cell growth in yeast. <i>Gene</i> , 2004, 324, 97-104. | 1.0 | 6 |
| 4710 | Functional co-expression of xenobiotic metabolizing enzymes, rat cytochrome P450 1A1 and UDP-glucuronosyltransferase 1A6, in yeast microsomes. <i>Biochimica Et Biophysica Acta - General Subjects</i> , 2004, 1672, 86-92. | 1.1 | 23 |
| 4711 | ARL1 participates with ATC1/LIC4 to regulate responses of yeast cells to ions. <i>Biochemical and Biophysical Research Communications</i> , 2004, 315, 617-623. | 1.0 | 13 |
| 4712 | Uptake of GABA and putrescine by UGA4 on the vacuolar membrane in <i>Saccharomyces cerevisiae</i> . <i>Biochemical and Biophysical Research Communications</i> , 2004, 315, 1082-1087. | 1.0 | 49 |
| 4713 | Molecular insights on DNA delivery into <i>Saccharomyces cerevisiae</i> . <i>Biochemical and Biophysical Research Communications</i> , 2004, 317, 100-107. | 1.0 | 28 |
| 4714 | A novel induction mechanism of the rat CYP1A2 gene mediated by Ah receptorâ€“Arnt heterodimer. <i>Biochemical and Biophysical Research Communications</i> , 2004, 318, 746-755. | 1.0 | 93 |
| 4715 | The yeast genes, ARL1 and CCZ1, interact to control membrane traffic and ion homeostasis. <i>Biochemical and Biophysical Research Communications</i> , 2004, 319, 840-846. | 1.0 | 9 |
| 4716 | Molecular cloning and biochemical characterization of <i>Candida albicans</i> acyl-CoA:sterol acyltransferase, a potential target of antifungal agents. <i>Biochemical and Biophysical Research Communications</i> , 2004, 319, 911-919. | 1.0 | 13 |
| 4717 | Cloning and characterization of the lanosterol 14 α -demethylase (ERG11) gene in <i>Cryptococcus neoformans</i> . <i>Biochemical and Biophysical Research Communications</i> , 2004, 324, 719-728. | 1.0 | 36 |
| 4718 | Molecular cloning, overexpression, and purification of <i>Micrococcus luteus</i> K-3-type glutaminase from <i>Aspergillus oryzae</i> RIB40. <i>Protein Expression and Purification</i> , 2004, 38, 272-278. | 0.6 | 21 |
| 4719 | Isolation and functional analysis of the glycerol permease activity of two new nodulin-like intrinsic proteins from salt stressed roots of the halophyte <i>Atriplex nummularia</i> . <i>Plant Science</i> , 2004, 166, 633-640. | 1.7 | 29 |

| # | ARTICLE | IF | CITATIONS |
|------|---|-----|-----------|
| 4720 | Identification of a nuclear targeting signal in YopM from <i>Yersinia</i> spp.. <i>Microbial Pathogenesis</i> , 2004, 36, 247-261. | 1.3 | 66 |
| 4721 | Expression and evolution of Δ^9 and Δ^{11} desaturase genes in the moth <i>Spodoptera littoralis</i> . <i>Insect Biochemistry and Molecular Biology</i> , 2004, 34, 1315-1328. | 1.2 | 36 |
| 4722 | Substrate and Metal Complexes of 3-Deoxy-d-arabino-heptulosonate-7-phosphate Synthase from <i>Saccharomyces cerevisiae</i> Provide New Insights into the Catalytic Mechanism. <i>Journal of Molecular Biology</i> , 2004, 337, 675-690. | 2.0 | 45 |
| 4723 | Structural Analysis of the Human Golgi-associated Plant Pathogenesis Related Protein GAPR-1 Implicates Dimerization as a Regulatory Mechanism. <i>Journal of Molecular Biology</i> , 2004, 339, 173-183. | 2.0 | 66 |
| 4724 | Complementation of Yeast Arc1p by the p43 Component of the Human Multisynthetase Complex does not Require its Association with Yeast MetRS and GluRS. <i>Journal of Molecular Biology</i> , 2004, 340, 15-27. | 2.0 | 13 |
| 4725 | A Conserved Element in the Yeast RNase MRP RNA Subunit can Participate in a Long-range Base-pairing Interaction. <i>Journal of Molecular Biology</i> , 2004, 341, 375-388. | 2.0 | 24 |
| 4726 | Apo and Holo Structures of an NADP(H)-dependent Cinnamyl Alcohol Dehydrogenase from <i>Saccharomyces cerevisiae</i> . <i>Journal of Molecular Biology</i> , 2004, 341, 1049-1062. | 2.0 | 49 |
| 4727 | <i>Neurospora crassa</i> CyPBP37: A Cytosolic Stress Protein that is able to Replace Yeast Thi4p Function in the Synthesis of Vitamin B1. <i>Journal of Molecular Biology</i> , 2004, 344, 1147-1157. | 2.0 | 31 |
| 4728 | Influence of compartmental localization on the function of yeast NADP ⁺ -specific isocitrate dehydrogenases. <i>Archives of Biochemistry and Biophysics</i> , 2004, 423, 235-246. | 1.4 | 18 |
| 4729 | The interaction between E-tropomodulin and thymosin β^2 -10 rescues tumor cells from thymosin β^2 -10 mediated apoptosis by restoring actin architecture. <i>FEBS Letters</i> , 2004, 557, 57-63. | 1.3 | 21 |
| 4730 | Expression of the Cymbidium Ringspot Virus 33-Kilodalton Protein in <i>Saccharomyces cerevisiae</i> and Molecular Dissection of the Peroxisomal Targeting Signal. <i>Journal of Virology</i> , 2004, 78, 4744-4752. | 1.5 | 59 |
| 4731 | Yeast Two-Hybrid Assay for Studying Protein-Protein Interactions. , 2004, 270, 403-422. | | 28 |
| 4732 | Three Decades of Fungal Transformation: Novel Technologies. , 2004, 267, 315-326. | | 18 |
| 4733 | Cytochrome P450 Taxadiene 5 β -Hydroxylase, a Mechanistically Unusual Monooxygenase Catalyzing the First Oxygenation Step of Taxol Biosynthesis. <i>Chemistry and Biology</i> , 2004, 11, 379-387. | 6.2 | 155 |
| 4734 | Lack of interaction between AFLR and AFLJ contributes to nonaflatoxicity of <i>Aspergillus sojae</i> . <i>Journal of Biotechnology</i> , 2004, 107, 245-253. | 1.9 | 59 |
| 4735 | Engineered autolytic yeast strains secreting <i>Kluyveromyces lactis</i> β -galactosidase for production of heterologous proteins in lactose media. <i>Journal of Biotechnology</i> , 2004, 109, 131-137. | 1.9 | 27 |
| 4736 | Dual targeting of yeast catalase A to peroxisomes and mitochondria. <i>Biochemical Journal</i> , 2004, 380, 393-400. | 1.7 | 112 |
| 4737 | Characterization of three novel members of the <i>Arabidopsis thaliana</i> equilibrative nucleoside transporter (ENT) family. <i>Biochemical Journal</i> , 2004, 383, 19-26. | 1.7 | 69 |

| # | ARTICLE | IF | CITATIONS |
|------|--|-----|-----------|
| 4738 | Saccharomyces cerevisiae ubiquitin-like protein Rub1 conjugates to cullin proteins Rtt101 and Cul3 in vivo. <i>Biochemical Journal</i> , 2004, 377, 459-467. | 1.7 | 25 |
| 4739 | Forchlorfenuron, a phenylurea cytokinin, disturbs septin organization in <i>Saccharomyces cerevisiae</i> . <i>Genes and Genetic Systems</i> , 2004, 79, 199-206. | 0.2 | 39 |
| 4740 | Genomics of <i>Candida albicans</i> . <i>Applied Mycology and Biotechnology</i> , 2004, 4, 99-135. | 0.3 | 0 |
| 4741 | Identification and functional characterization of variants in human concentrative nucleoside transporter 3, hCNT3 (SLC28A3), arising from single nucleotide polymorphisms in coding regions of the hCNT3 gene. <i>Pharmacogenetics and Genomics</i> , 2005, 15, 173-182. | 0.7 | 28 |
| 4742 | Manipulation of morphological transition in <i>Saccharomyces cerevisiae</i> mutant J19. <i>Biochemical Engineering Journal</i> , 2005, 27, 72-76. | 1.8 | 1 |
| 4743 | Overexpression of <i>scs1-1</i> . A plausible strategy to obtain wine yeast strains undergoing accelerated autolysis. <i>FEMS Microbiology Letters</i> , 2005, 246, 1-9. | 0.7 | 22 |
| 4744 | Clustering of genes in : Cloning of the maltose permease gene and expression from the divergent intergenic region between the maltose permease and maltase genes. <i>FEMS Yeast Research</i> , 2005, 5, 1019-1028. | 1.1 | 18 |
| 4745 | In vitro assessment of biocompatibility of biomaterials by using fluorescent yeast biosensor. <i>Current Applied Physics</i> , 2005, 5, 444-448. | 1.1 | 13 |
| 4746 | Induction of antigen-specific immune responses by oral vaccination with <i>Saccharomyces cerevisiae</i> expressing <i>Actinobacillus pleuropneumoniae</i> ApxIIA. <i>FEMS Immunology and Medical Microbiology</i> , 2005, 43, 155-164. | 2.7 | 39 |
| 4747 | Screening and identification of the levoglucosan kinase gene (<i>lgk</i>) from <i>Aspergillus niger</i> by LC-ESI-MS/MS and RT-PCR. <i>FEMS Microbiology Letters</i> , 2005, 251, 313-319. | 0.7 | 13 |
| 4748 | Multiple genetic and biochemical interactions of <i>Brr2</i> , <i>Prp8</i> , <i>Prp31</i> , <i>Prp1</i> and <i>Prp4</i> kinase suggest a function in the control of the activation of spliceosomes in <i>Schizosaccharomyces pombe</i> . <i>Current Genetics</i> , 2005, 48, 151-161. | 0.8 | 32 |
| 4749 | A Family of Basic Amino Acid Transporters of the Vacuolar Membrane from <i>Saccharomyces cerevisiae</i> . <i>Journal of Biological Chemistry</i> , 2005, 280, 4851-4857. | 1.6 | 81 |
| 4750 | The amino-terminal non-catalytic region of <i>Salmonella typhimurium</i> SigD affects actin organization in yeast and mammalian cells. <i>Cellular Microbiology</i> , 2005, 7, 1432-1446. | 1.1 | 27 |
| 4751 | The HOG MAP kinase pathway is required for the induction of methylglyoxal-responsive genes and determines methylglyoxal resistance in <i>Saccharomyces cerevisiae</i> . <i>Molecular Microbiology</i> , 2005, 56, 228-239. | 1.2 | 61 |
| 4752 | Convergent evolution of hydroxylation mechanisms in the fungal kingdom: molybdenum cofactor-independent hydroxylation of xanthine via $\hat{1}\pm$ -ketoglutarate-dependent dioxygenases. <i>Molecular Microbiology</i> , 2005, 57, 276-290. | 1.2 | 40 |
| 4753 | <i>Swi/SNF-GCN5</i> -dependent chromatin remodelling determines induced expression of <i>GDH3</i> , one of the paralogous genes responsible for ammonium assimilation and glutamate biosynthesis in <i>Saccharomyces cerevisiae</i> . <i>Molecular Microbiology</i> , 2005, 57, 291-305. | 1.2 | 33 |
| 4754 | A novel isoform of glucan, water dikinase phosphorylates pre-phosphorylated $\hat{1}\pm$ -glucans and is involved in starch degradation in <i>Arabidopsis</i> . <i>Plant Journal</i> , 2005, 41, 595-605. | 2.8 | 180 |
| 4755 | Mutational analysis of the yeast multidrug resistance ABC transporter <i>Pdr5p</i> with altered drug specificity. <i>Genes To Cells</i> , 2005, 10, 409-420. | 0.5 | 39 |

| # | ARTICLE | IF | CITATIONS |
|------|---|-----|-----------|
| 4756 | Translation elongation factor 1A is essential for regulation of the actin cytoskeleton and cell morphology. <i>Nature Structural and Molecular Biology</i> , 2005, 12, 772-778. | 3.6 | 247 |
| 4757 | ERS1 encodes a functional homologue of the human lysosomal cystine transporter. <i>FEBS Journal</i> , 2005, 272, 2497-2511. | 2.2 | 45 |
| 4758 | Nonclassical PITPs Activate PLD via the Stt4p PtdIns-4-kinase and Modulate Function of Late Stages of Exocytosis in Vegetative Yeast. <i>Traffic</i> , 2005, 6, 1157-1172. | 1.3 | 62 |
| 4759 | Two New Group 3 LEA Genes of Wheat and Their Functional Analysis in Yeast. <i>Journal of Integrative Plant Biology</i> , 2005, 47, 1372-1381. | 4.1 | 21 |
| 4760 | Production of recombinant human antithrombin by <i>Pichia pastoris</i> . <i>Journal of Bioscience and Bioengineering</i> , 2005, 99, 264-271. | 1.1 | 17 |
| 4761 | Evidence of a dominant negative mutant of yeast methionine aminopeptidase type 2 in <i>Saccharomyces cerevisiae</i> . <i>Journal of Cellular Biochemistry</i> , 2005, 94, 656-668. | 1.2 | 3 |
| 4762 | Use of the cell wall protein Pir4 as a fusion partner for the expression of <i>Bacillus</i> sp. BP-7 xylanase A in <i>Saccharomyces cerevisiae</i> . <i>Biotechnology and Bioengineering</i> , 2005, 89, 690-697. | 1.7 | 39 |
| 4763 | Enzymatic Polymerization Behavior Using Cellulose-Binding Domain Deficient Endoglucanase II. <i>Macromolecular Bioscience</i> , 2005, 5, 623-628. | 2.1 | 23 |
| 4764 | Construction of a selective cleavage system for a protein displayed on the cell surface of yeast. <i>Applied Microbiology and Biotechnology</i> , 2005, 69, 423-427. | 1.7 | 13 |
| 4765 | Salt-dependent expression of ammonium assimilation genes in the halotolerant yeast, <i>Debaryomyces hansenii</i> . <i>Current Genetics</i> , 2005, 47, 163-171. | 0.8 | 10 |
| 4766 | The YJR127C/ZMS1 gene product is involved in glycerol-based respiratory growth of the yeast <i>Saccharomyces cerevisiae</i> . <i>Current Genetics</i> , 2005, 48, 235-246. | 0.8 | 25 |
| 4767 | Characterization of <i>rec15</i> , an early meiotic recombination gene in <i>Schizosaccharomyces pombe</i> . <i>Current Genetics</i> , 2005, 48, 323-333. | 0.8 | 12 |
| 4768 | An Os-1 family histidine kinase from a filamentous fungus confers fungicide-sensitivity to yeast. <i>Current Genetics</i> , 2005, 47, 298-306. | 0.8 | 72 |
| 4769 | Novel interaction of the Hsp90 chaperone machine with Ssl2, an essential DNA helicase in <i>Saccharomyces cerevisiae</i> . <i>Current Genetics</i> , 2005, 47, 368-380. | 0.8 | 28 |
| 4770 | Mechanism of polyamine tolerance in yeast: novel regulators and insights. <i>Cellular and Molecular Life Sciences</i> , 2005, 62, 3106-3116. | 2.4 | 30 |
| 4771 | Gln3p and Nil1p regulation of invertase activity and expression in. <i>FEMS Yeast Research</i> , 2005, 5, 605-609. | 1.1 | 7 |
| 4772 | Deficiency of Pkc1 activity affects glycerol metabolism in. <i>FEMS Yeast Research</i> , 2005, 5, 767-776. | 1.1 | 11 |
| 4773 | GABAA receptor-associated protein (GABARAP) induces apoptosis by interacting with DEAD (Asp-Glu-Ala-Asp/His) box polypeptide 47 (DDX 47). <i>Biotechnology Letters</i> , 2005, 27, 623-628. | 1.1 | 29 |

| # | ARTICLE | IF | CITATIONS |
|------|--|-----|-----------|
| 4774 | Programmed Cell Death 6 (PDCD6) Protein Interacts with Death-Associated Protein Kinase 1 (DAPk1): Additive Effect on Apoptosis via Caspase-3 Dependent Pathway. <i>Biotechnology Letters</i> , 2005, 27, 1011-1015. | 1.1 | 46 |
| 4775 | Non-reciprocal chromosomal bridge-induced translocation (BIT) by targeted DNA integration in yeast. <i>Chromosoma</i> , 2005, 114, 15-27. | 1.0 | 19 |
| 4776 | Structural and functional analysis of 5S rRNA in <i>Saccharomyces cerevisiae</i> . <i>Molecular Genetics and Genomics</i> , 2005, 274, 235-247. | 1.0 | 40 |
| 4777 | Sir-dependent downregulation of various aging processes. <i>Molecular Genetics and Genomics</i> , 2005, 274, 539-547. | 1.0 | 9 |
| 4778 | Transcriptional profiling of <i>Saccharomyces cerevisiae</i> cells under adhesion-inducing conditions. <i>Molecular Genetics and Genomics</i> , 2005, 273, 382-393. | 1.0 | 22 |
| 4779 | A novel expression system for genomic DNA loci using a human artificial chromosome vector with transformation-associated recombination cloning. <i>Journal of Human Genetics</i> , 2005, 50, 592-599. | 1.1 | 24 |
| 4780 | Expression of the Apx toxins of <i>Actinobacillus pleuropneumoniae</i> in <i>Saccharomyces cerevisiae</i> and its induction of immune response in mice. <i>Biotechnology and Bioprocess Engineering</i> , 2005, 10, 362-366. | 1.4 | 4 |
| 4781 | Expression of fungal phytase on the cell surface of <i>Saccharomyces cerevisiae</i> . <i>Biotechnology and Bioprocess Engineering</i> , 2005, 10, 576. | 1.4 | 16 |
| 4783 | Influence of low glycolytic activities in <i>gcr1</i> and <i>gcr2</i> mutants on the expression of other metabolic pathway genes in <i>Saccharomyces cerevisiae</i> . <i>Yeast</i> , 2005, 22, 111-127. | 0.8 | 29 |
| 4784 | Physiological effects of unassembled chaperonin Cct subunits in the yeast <i>Saccharomyces cerevisiae</i> . <i>Yeast</i> , 2005, 22, 219-239. | 0.8 | 40 |
| 4785 | Expression of GCR1, the transcriptional activator of glycolytic enzyme genes in the yeast <i>Saccharomyces cerevisiae</i> , is positively autoregulated by Gcr1p. <i>Yeast</i> , 2005, 22, 305-319. | 0.8 | 14 |
| 4786 | Inverted repeat of a large segment unveiled on the right arm of <i>Saccharomyces cerevisiae</i> chromosome II. <i>Yeast</i> , 2005, 22, 321-336. | 0.8 | 4 |
| 4787 | New drug-resistant cassettes for gene disruption and epitope tagging in <i>Schizosaccharomyces pombe</i> . <i>Yeast</i> , 2005, 22, 583-591. | 0.8 | 252 |
| 4788 | A novel yeast genomic DNA library on a geneticin-resistance vector. <i>Yeast</i> , 2005, 22, 653-657. | 0.8 | 18 |
| 4789 | The <i>Saccharomyces cerevisiae</i> ESU1 gene, which is responsible for enhancement of termination suppression, corresponds to the 3' terminal half of GAL11. <i>Yeast</i> , 2005, 22, 895-906. | 0.8 | 5 |
| 4790 | Condensed protocol for competent cell preparation and transformation of the methylotrophic yeast <i>Pichia pastoris</i> . <i>BioTechniques</i> , 2005, 38, 44-48. | 0.8 | 249 |
| 4791 | DNA-Transformation und Charakterisierung transgener Organismen. , 2005, , 141-220. | | 0 |
| 4792 | Two-hybrid reporter vectors for gap repair cloning. <i>BioTechniques</i> , 2005, 38, 927-934. | 0.8 | 13 |

| # | ARTICLE | IF | CITATIONS |
|------|--|-----|-----------|
| 4793 | The Pleckstrin Homology Domain Proteins Slm1 and Slm2 Are Required for Actin Cytoskeleton Organization in Yeast and Bind Phosphatidylinositol-4,5-Bisphosphate and TORC2. <i>Molecular Biology of the Cell</i> , 2005, 16, 1883-1900. | 0.9 | 120 |
| 4794 | The Ess1 prolyl isomerase is dispensable for growth but required for virulence in <i>Cryptococcus neoformans</i> . <i>Microbiology (United Kingdom)</i> , 2005, 151, 1593-1605. | 0.7 | 19 |
| 4795 | Yeast Glutamine-fructose-6-phosphate Aminotransferase (Gfa1) Requires Methionine Aminopeptidase Activity for Proper Function. <i>Journal of Biological Chemistry</i> , 2005, 280, 14356-14360. | 1.6 | 13 |
| 4796 | Harnessing Natural Diversity to Probe Metabolic Pathways. <i>PLoS Genetics</i> , 2005, 1, e80. | 1.5 | 58 |
| 4797 | Genome-Wide Screen for Inner Nuclear Membrane Protein Targeting in <i>Saccharomyces cerevisiae</i> . <i>Genetics</i> , 2005, 170, 1553-1560. | 1.2 | 59 |
| 4798 | A Three-Stemmed mRNA Pseudoknot in the SARS Coronavirus Frameshift Signal. <i>PLoS Biology</i> , 2005, 3, e172. | 2.6 | 158 |
| 4799 | Functional expression of the CMP-sialic acid transporter in <i>Escherichia coli</i> and its identification as a simple mobile carrier. <i>Glycobiology</i> , 2005, 16, 73-81. | 1.3 | 25 |
| 4800 | Ability of Sit4p To Promote K ⁺ Efflux via Nha1p Is Modulated by Sap155p and Sap185p. <i>Eukaryotic Cell</i> , 2005, 4, 1041-1049. | 3.4 | 18 |
| 4801 | A Role for the Protease-sensitive Loop Region of Shiga-like Toxin 1 in the Retrotranslocation of Its A1 Domain from the Endoplasmic Reticulum Lumen. <i>Journal of Biological Chemistry</i> , 2005, 280, 23310-23318. | 1.6 | 55 |
| 4802 | A Gibberellin-Regulated Calcineurin B in Rice Localizes to the Tonoplast and Is Implicated in Vacuole Function. <i>Plant Physiology</i> , 2005, 138, 1347-1358. | 2.3 | 69 |
| 4803 | The Spatial Organization of Lipid Synthesis in the Yeast <i>Saccharomyces cerevisiae</i> Derived from Large Scale Green Fluorescent Protein Tagging and High Resolution Microscopy. <i>Molecular and Cellular Proteomics</i> , 2005, 4, 662-672. | 2.5 | 150 |
| 4804 | Identification of programmed translational -1 frameshifting sites in the genome of <i>Saccharomyces cerevisiae</i> . <i>Genome Research</i> , 2005, 15, 1411-1420. | 2.4 | 6 |
| 4805 | The dimorphic yeast <i>Yarrowia lipolytica</i> possesses an atypical phosphofructokinase: characterization of the enzyme and its encoding gene. <i>Microbiology (United Kingdom)</i> , 2005, 151, 1465-1474. | 0.7 | 16 |
| 4806 | Novel Characteristics of the Biological Properties of the Yeast <i>Saccharomyces cerevisiae</i> Eukaryotic Initiation Factor 2A. <i>Journal of Biological Chemistry</i> , 2005, 280, 15601-15611. | 1.6 | 49 |
| 4807 | End3p-Mediated Endocytosis Is Required for Spore Wall Formation in <i>Saccharomyces cerevisiae</i> . <i>Genetics</i> , 2005, 170, 1561-1574. | 1.2 | 22 |
| 4808 | Identification of Functionally Interacting SNAREs by Using Complementary Substitutions in the Conserved 'O' Layer. <i>Molecular Biology of the Cell</i> , 2005, 16, 2263-2274. | 0.9 | 26 |
| 4809 | The bipartite structure of the tRNA m1A58 methyltransferase from <i>S. cerevisiae</i> is conserved in humans. <i>Rna</i> , 2005, 11, 1281-1290. | 1.6 | 118 |
| 4810 | Ace2p Controls the Expression of Genes Required for Cell Separation in <i>Schizosaccharomyces pombe</i> . <i>Molecular Biology of the Cell</i> , 2005, 16, 2003-2017. | 0.9 | 78 |

| # | ARTICLE | IF | CITATIONS |
|------|--|-----|-----------|
| 4811 | Saccharomyces cerevisiae Sps1p Regulates Trafficking of Enzymes Required for Spore Wall Synthesis. Eukaryotic Cell, 2005, 4, 536-544. | 3.4 | 31 |
| 4812 | Cell Polarity Protein Spa2P Associates With Proteins Involved In Actin Function In Saccharomyces Cerevisiae. Molecular Biology of the Cell, 2005, 16, 4595-4608. | 0.9 | 33 |
| 4813 | Hepatitis C Virus Internal Ribosome Entry Site-Dependent Translation in Saccharomyces cerevisiae Is Independent of Polypyrimidine Tract-Binding Protein, Poly(rC)-Binding Protein 2, and La Protein. Journal of Virology, 2005, 79, 10126-10137. | 1.5 | 30 |
| 4814 | Cloning and characterization of the phosphoglucomutase of Trypanosoma cruzi and functional complementation of a Saccharomyces cerevisiae PGM null mutant. Glycobiology, 2005, 15, 1359-1367. | 1.3 | 11 |
| 4815 | Loss of SOD1 and LYS7 Sensitizes Saccharomyces cerevisiae to Hydroxyurea and DNA Damage Agents and Downregulates MEC1 Pathway Effectors. Molecular and Cellular Biology, 2005, 25, 10273-10285. | 1.1 | 36 |
| 4816 | Abundant Expression in Vascular Tissue of Plant TAF10, an Orthologous Gene for TATA Box-binding Protein-associated Factor 10, in Flaveria trinervia and Abnormal Morphology of Arabidopsis thaliana Transformants on its Overexpression. Plant and Cell Physiology, 2005, 46, 108-117. | 1.5 | 21 |
| 4817 | Gpi19, the Saccharomyces cerevisiae Homologue of Mammalian PIG-P, Is a Subunit of the Initial Enzyme for Glycosylphosphatidylinositol Anchor Biosynthesis. Eukaryotic Cell, 2005, 4, 1801-1807. | 3.4 | 28 |
| 4818 | Synthetic Lethal Screen. , 2006, 313, 161-170. | | 8 |
| 4819 | Effects of N-Glycosylation and Inositol on the ER Stress Response in Yeast Saccharomyces cerevisiae. Bioscience, Biotechnology and Biochemistry, 2005, 69, 1274-1280. | 0.6 | 6 |
| 4820 | Knocking out Ubiquitin Proteasome System Function In Vivo and In Vitro with Genetically Encodable Tandem Ubiquitin. Methods in Enzymology, 2005, 399, 64-74. | 0.4 | 5 |
| 4821 | Interplay between GCN2 and GCN4 expression, translation elongation factor 1 mutations and translational fidelity in yeast. Nucleic Acids Research, 2005, 33, 4584-4592. | 6.5 | 5 |
| 4822 | Isolation and Characterization of Anthocyanin 5-O-Glucosyltransferase in Perilla frutescens var. crispa by Differential Display. , 2006, 317, 255-266. | | 5 |
| 4823 | The Saccharomyces cerevisiae YLL012/YEH1, YLR020/YEH2, and TGL1 Genes Encode a Novel Family of Membrane-Anchored Lipases That Are Required for Steryl Ester Hydrolysis. Molecular and Cellular Biology, 2005, 25, 1655-1668. | 1.1 | 124 |
| 4824 | Efficient Production of L-Lactic Acid by Metabolically Engineered Saccharomyces cerevisiae with a Genome-Integrated L-Lactate Dehydrogenase Gene. Applied and Environmental Microbiology, 2005, 71, 1964-1970. | 1.4 | 118 |
| 4825 | Excretion of Putrescine and Spermidine by the Protein Encoded by YKL174c (TPO5) in Saccharomyces cerevisiae. Journal of Biological Chemistry, 2005, 280, 12637-12642. | 1.6 | 44 |
| 4826 | Mutual Interference between Genomic RNA Replication and Subgenomic mRNA Transcription in Brome Mosaic Virus. Journal of Virology, 2005, 79, 1438-1451. | 1.5 | 27 |
| 4827 | The highly conserved glutamic acid 791 of Rpb2 is involved in the binding of NTP and Mg(B) in the active center of human RNA polymerase II. Nucleic Acids Research, 2005, 33, 2629-2639. | 6.5 | 13 |
| 4828 | Characteristics of the Polyamine Transporter TPO1 and Regulation of Its Activity and Cellular Localization by Phosphorylation. Journal of Biological Chemistry, 2005, 280, 9646-9652. | 1.6 | 69 |

| # | ARTICLE | IF | CITATIONS |
|------|--|-----|-----------|
| 4829 | Characterizing the Sphingolipid Signaling Pathway That Remediate Defects Associated with Loss of the Yeast Amphiphysin-like Orthologs, Rvs161p and Rvs167p. <i>Journal of Biological Chemistry</i> , 2005, 280, 4270-4278. | 1.6 | 24 |
| 4830 | Expression of Human CTP Synthetase in <i>Saccharomyces cerevisiae</i> Reveals Phosphorylation by Protein Kinase A. <i>Journal of Biological Chemistry</i> , 2005, 280, 38328-38336. | 1.6 | 39 |
| 4831 | Molecular Organization of Target of Rapamycin Complex 2. <i>Journal of Biological Chemistry</i> , 2005, 280, 30697-30704. | 1.6 | 197 |
| 4832 | Transcriptional Regulation by Lge1p Requires a Function Independent of Its Role in Histone H2B Ubiquitination*. <i>Journal of Biological Chemistry</i> , 2005, 280, 2759-2770. | 1.6 | 30 |
| 4833 | Phosphorylation of the Yeast Choline Kinase by Protein Kinase C. <i>Journal of Biological Chemistry</i> , 2005, 280, 26105-26112. | 1.6 | 27 |
| 4834 | Oxidant-specific Folding of Yap1p Regulates Both Transcriptional Activation and Nuclear Localization. <i>Journal of Biological Chemistry</i> , 2005, 280, 40524-40533. | 1.6 | 65 |
| 4835 | Sequence Context Effects on Oligo(dT) Termination Signal Recognition by <i>Saccharomyces cerevisiae</i> RNA Polymerase III. <i>Journal of Biological Chemistry</i> , 2005, 280, 19551-19562. | 1.6 | 97 |
| 4836 | Regulation of the PIS1-encoded Phosphatidylinositol Synthase in <i>Saccharomyces cerevisiae</i> by Zinc. <i>Journal of Biological Chemistry</i> , 2005, 280, 29017-29024. | 1.6 | 37 |
| 4837 | Regulation of Sphingosine-1-phosphate Lyase Gene Expression by Members of the GATA Family of Transcription Factors. <i>Journal of Biological Chemistry</i> , 2005, 280, 18403-18410. | 1.6 | 32 |
| 4838 | Genetic Interactions Between Mediator and the Late G1-Specific Transcription Factor Swi6 in <i>Saccharomyces cerevisiae</i> . <i>Genetics</i> , 2005, 171, 477-488. | 1.2 | 13 |
| 4839 | <i>Yarrowia lipolytica</i> Mutants Devoid of Pyruvate Carboxylase Activity Show an Unusual Growth Phenotype. <i>Eukaryotic Cell</i> , 2005, 4, 356-364. | 3.4 | 21 |
| 4840 | Molecular basis of the mammalian potency of the scorpion Î±-toxin, BmK M1. <i>FASEB Journal</i> , 2005, 19, 1-24. | 0.2 | 35 |
| 4841 | Identification of Two Novel Human Acyl-CoA Wax Alcohol Acyltransferases. <i>Journal of Biological Chemistry</i> , 2005, 280, 14755-14764. | 1.6 | 99 |
| 4842 | Characterizing Sterol Defect Suppressors Uncovers a Novel Transcriptional Signaling Pathway Regulating Zymosterol Biosynthesis. <i>Journal of Biological Chemistry</i> , 2005, 280, 35904-35913. | 1.6 | 29 |
| 4843 | Novel CIPK1-Associated Proteins in <i>Arabidopsis</i> Contain an Evolutionarily Conserved C-Terminal Region That Mediates Nuclear Localization. <i>Plant Physiology</i> , 2005, 139, 138-150. | 2.3 | 82 |
| 4844 | The Smk1p MAP kinase negatively regulates Gsc2p, a 1,3-Î±-glucan synthase, during spore wall morphogenesis in <i>Saccharomyces cerevisiae</i> . <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 2005, 102, 12431-12436. | 3.3 | 35 |
| 4845 | A Chaperone Pathway in Protein Disaggregation. <i>Journal of Biological Chemistry</i> , 2005, 280, 23869-23875. | 1.6 | 257 |
| 4846 | Rabies Virus P Protein Interacts with STAT1 and Inhibits Interferon Signal Transduction Pathways. <i>Journal of Virology</i> , 2005, 79, 14411-14420. | 1.5 | 188 |

| # | ARTICLE | IF | CITATIONS |
|------|---|-----|-----------|
| 4847 | A Genomic Screen for Yeast Vacuolar Membrane ATPase Mutants. <i>Genetics</i> , 2005, 170, 1539-1551. | 1.2 | 86 |
| 4848 | Novel <i>Plasmodium vivax</i> dhfr Alleles from the Indonesian Archipelago and Papua New Guinea: Association with Pyrimethamine Resistance Determined by a <i>Saccharomyces cerevisiae</i> Expression System. <i>Antimicrobial Agents and Chemotherapy</i> , 2005, 49, 733-740. | 1.4 | 60 |
| 4849 | Receptor for Retrograde Transport in the Apicomplexan Parasite <i>Toxoplasma gondii</i> . <i>Eukaryotic Cell</i> , 2005, 4, 432-442. | 3.4 | 33 |
| 4850 | Loss of Meiotic Rereplication Block in <i>Saccharomyces cerevisiae</i> Cells Defective in Cdc28p Regulation. <i>Eukaryotic Cell</i> , 2005, 4, 55-62. | 3.4 | 15 |
| 4851 | Rho4 GTPase Is Involved in Secretion of Glucanases during Fission Yeast Cytokinesis. <i>Eukaryotic Cell</i> , 2005, 4, 1639-1645. | 3.4 | 40 |
| 4852 | Evolution of 3-deoxy-D-arabino-heptulosonate-7-phosphate synthase-encoding genes in the yeast <i>Saccharomyces cerevisiae</i> . <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 2005, 102, 9784-9789. | 3.3 | 43 |
| 4853 | A Single Mms2 Residue Insertion into a Ubc13 Pocket Determines the Interface Specificity of a Human Lys63 Ubiquitin Conjugation Complex*. <i>Journal of Biological Chemistry</i> , 2005, 280, 17891-17900. | 1.6 | 44 |
| 4854 | Binding and Functions of ADP-ribosylation Factor on Mammalian and Yeast Peroxisomes. <i>Journal of Biological Chemistry</i> , 2005, 280, 34489-34499. | 1.6 | 30 |
| 4855 | Distinct regulation of Ubc13 functions by the two ubiquitin-conjugating enzyme variants Mms2 and Uev1A. <i>Journal of Cell Biology</i> , 2005, 170, 745-755. | 2.3 | 151 |
| 4856 | Synthetic Genetic Array Analysis of the PtdIns 4-kinase Pik1p Identifies Components in a Golgi-specific Ypt31/rab-GTPase Signaling Pathway. <i>Molecular Biology of the Cell</i> , 2005, 16, 776-793. | 0.9 | 112 |
| 4857 | Role for Gcs1p in Regulation of Arl1p at Trans-Golgi Compartments. <i>Molecular Biology of the Cell</i> , 2005, 16, 4024-4033. | 0.9 | 39 |
| 4858 | Disruption of MRC19 results in altered nitrogen metabolic status and defective pseudohyphal development in <i>Saccharomyces cerevisiae</i> . <i>Microbiology (United Kingdom)</i> , 2005, 151, 91-98. | 0.7 | 5 |
| 4859 | Interaction between the Shuttling mRNA Export Factor Gle1 and the Nucleoporin hCG1: A Conserved Mechanism in the Export of Hsp70 mRNA. <i>Molecular Biology of the Cell</i> , 2005, 16, 4304-4315. | 0.9 | 70 |
| 4860 | Dominant-negative mutant phenotypes and the regulation of translation elongation factor 2 levels in yeast. <i>Nucleic Acids Research</i> , 2005, 33, 5740-5748. | 6.5 | 40 |
| 4861 | Genetic Manipulation of Yeast to Identify Genes Involved in Regulation of Chemosensitivity. , 2005, 111, 241-256. | | 0 |
| 4862 | Mutations in Mre11 Phosphoesterase Motif I That Impair <i>Saccharomyces cerevisiae</i> Mre11-Rad50-Xrs2 Complex Stability in Addition to Nuclease Activity. <i>Genetics</i> , 2005, 171, 1561-1570. | 1.2 | 76 |
| 4863 | Involvement of actin and polarisome in morphological change during spore germination of <i>Saccharomyces cerevisiae</i> . <i>Yeast</i> , 2005, 22, 129-139. | 0.8 | 26 |
| 4864 | HABP1/p32/gC1qR induces aberrant growth and morphology in through its N-terminal $\hat{\pm}$ helix. <i>Experimental Cell Research</i> , 2005, 309, 250-263. | 1.2 | 14 |

| # | ARTICLE | IF | CITATIONS |
|------|--|-----|-----------|
| 4865 | Molecular Cloning, Bacterial Overexpression and Characterization of L-myo- inositol 1- Phosphate Synthase from a Monocotyledonous Resurrection Plant, <i>Xerophyta viscosa</i> Baker. <i>Journal of Plant Biochemistry and Biotechnology</i> , 2005, 14, 95-99. | 0.9 | 9 |
| 4866 | Anchorage of <i>Candida albicans</i> Ssr1 to the cell wall, and transcript profiling of the null mutant. <i>Research in Microbiology</i> , 2005, 156, 911-920. | 1.0 | 13 |
| 4867 | Expression of the cry11A gene of <i>Bacillus thuringiensis</i> ssp. <i>israelensis</i> in <i>Saccharomyces cerevisiae</i> . <i>Canadian Journal of Microbiology</i> , 2005, 51, 165-170. | 0.8 | 4 |
| 4868 | Genetically Engineered Wine Yeast Produces a High Concentration of L-Lactic Acid of Extremely High Optical Purity. <i>Applied and Environmental Microbiology</i> , 2005, 71, 2789-2792. | 1.4 | 120 |
| 4869 | Uridine Binding and Transportability Determinants of Human Concentrative Nucleoside Transporters. <i>Molecular Pharmacology</i> , 2005, 68, 830-839. | 1.0 | 38 |
| 4870 | The CCAAT-binding Complex of Eukaryotes: Evolution of a Second NLS in the HapB Subunit of the Filamentous Fungus <i>Aspergillus nidulans</i> Despite Functional Conservation at the Molecular Level between Yeast, <i>A. nidulans</i> and Human. <i>Journal of Molecular Biology</i> , 2005, 352, 517-533. | 2.0 | 28 |
| 4871 | The Ty1 transposition assay: a new short-term test for detection of carcinogens. <i>Journal of Microbiological Methods</i> , 2005, 61, 1-8. | 0.7 | 26 |
| 4872 | An Extended Signal Involved in Eukaryotic α -1 Frameshifting Operates through Modification of the E Site tRNA. <i>Molecular Cell</i> , 2005, 17, 61-68. | 4.5 | 58 |
| 4873 | Isolation of trehalose-6-phosphate phosphatase gene from tobacco and its functional analysis in yeast cells. <i>Journal of Plant Physiology</i> , 2005, 162, 215-223. | 1.6 | 32 |
| 4874 | Functional expression of the rat organic anion transporter 1 (rOAT1) in <i>Saccharomyces cerevisiae</i> . <i>Biochimica Et Biophysica Acta - Biomembranes</i> , 2005, 1720, 44-51. | 1.4 | 4 |
| 4875 | Uptake of putrescine and spermidine by Gap1p on the plasma membrane in <i>Saccharomyces cerevisiae</i> . <i>Biochemical and Biophysical Research Communications</i> , 2005, 328, 1028-1033. | 1.0 | 41 |
| 4876 | Enniatin has a new function as an inhibitor of Pdr5p, one of the ABC transporters in <i>Saccharomyces cerevisiae</i> . <i>Biochemical and Biophysical Research Communications</i> , 2005, 328, 1119-1125. | 1.0 | 106 |
| 4877 | A new function of isonitrile as an inhibitor of the Pdr5p multidrug ABC transporter in <i>Saccharomyces cerevisiae</i> . <i>Biochemical and Biophysical Research Communications</i> , 2005, 330, 622-628. | 1.0 | 34 |
| 4878 | The yeast osmosensitive mutant <i>tps1^Δ</i> transformed by the cauliflower <i>BobTIP1;1</i> aquaporin withstand a hypo-osmotic shock. <i>FEBS Letters</i> , 2005, 579, 3872-3880. | 1.3 | 9 |
| 4879 | The polycomb group gene product Mel-18 interacts with cyclin D2 and modulates its activity. <i>FEBS Letters</i> , 2005, 579, 5275-5280. | 1.3 | 11 |
| 4880 | Identification and characterization of human Golgi nucleotide sugar transporter SLC35D2, a novel member of the SLC35 nucleotide sugar transporter family. <i>Genomics</i> , 2005, 85, 106-116. | 1.3 | 55 |
| 4881 | Metabolic Engineering of <i>Saccharomyces cerevisiae</i> for Efficient Production of Pure L-(+)-Lactic Acid. , 2006, , 795-807. | | 2 |
| 4882 | Identification of a second-site suppressor mutation of the GTPase defect associated with McCune-Albright syndrome: A model using the yeast heterotrimeric G protein, GPA1. <i>Archives of Physiology and Biochemistry</i> , 2006, 112, 166-173. | 1.0 | 2 |

| # | ARTICLE | IF | CITATIONS |
|------|--|-----|-----------|
| 4883 | The <i>Saccharomyces cerevisiae</i> Lipin Homolog Is a Mg ²⁺ -dependent Phosphatidate Phosphatase Enzyme*. <i>Journal of Biological Chemistry</i> , 2006, 281, 9210-9218. | 1.6 | 481 |
| 4884 | A Downshift in Temperature Activates the High Osmolarity Glycerol (HOG) Pathway, Which Determines Freeze Tolerance in <i>Saccharomyces cerevisiae</i> . <i>Journal of Biological Chemistry</i> , 2006, 281, 4638-4645. | 1.6 | 164 |
| 4885 | HOXA5 Regulates hMLH1 Expression in Breast Cancer Cells. <i>Neoplasia</i> , 2006, 8, 250-258. | 2.3 | 26 |
| 4886 | Secretion and properties of a hybrid <i>Kluyveromyces lactis</i> - <i>Aspergillus niger</i> beta-galactosidase. <i>Microbial Cell Factories</i> , 2006, 5, 41. | 1.9 | 33 |
| 4887 | d-Lactic acid production by metabolically engineered <i>Saccharomyces cerevisiae</i> . <i>Journal of Bioscience and Bioengineering</i> , 2006, 101, 172-177. | 1.1 | 102 |
| 4888 | Purification and Assays of <i>Saccharomyces cerevisiae</i> Homologous Recombination Proteins. <i>Methods in Enzymology</i> , 2006, 408, 445-463. | 0.4 | 41 |
| 4889 | Identification of eukaryotic secreted and cell surface proteins using the yeast secretion trap screen. <i>Nature Protocols</i> , 2006, 1, 2439-2447. | 5.5 | 30 |
| 4890 | Characterization of the second external alternative dehydrogenase from mitochondria of the respiratory yeast <i>Kluyveromyces lactis</i> . <i>Biochimica Et Biophysica Acta - Bioenergetics</i> , 2006, 1757, 1476-1484. | 0.5 | 24 |
| 4891 | Ternary Complex Formation of Ino2p-Ino4p Transcription Factors and Apl2p Adaptin \hat{I}^2 Subunit in Yeast. <i>Bioscience, Biotechnology and Biochemistry</i> , 2006, 70, 2604-2612. | 0.6 | 6 |
| 4892 | Pokeweed antiviral protein depurinates the sarcin/ricin loop of the rRNA prior to binding of aminoacyl-tRNA to the ribosomal A-site. <i>Rna</i> , 2006, 12, 1683-1692. | 1.6 | 15 |
| 4893 | A Ca ²⁺ signaling pathway regulates a K ⁺ channel for low-K response in <i>Arabidopsis</i> . <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 2006, 103, 12625-12630. | 3.3 | 412 |
| 4894 | Identification of Genes Affecting Lipid Content Using Transposon Mutagenesis in <i>Saccharomyces cerevisiae</i> . <i>Bioscience, Biotechnology and Biochemistry</i> , 2006, 70, 646-653. | 0.6 | 41 |
| 4895 | FLO11Is the Primary Factor in Flor Formation Caused by Cell Surface Hydrophobicity in Wild-Type Flor Yeast. <i>Bioscience, Biotechnology and Biochemistry</i> , 2006, 70, 660-666. | 0.6 | 54 |
| 4896 | Novel allosteric properties produced by residue substitutions in the subunit interface of yeast NAD ⁺ -specific isocitrate dehydrogenase. <i>Archives of Biochemistry and Biophysics</i> , 2006, 453, 207-216. | 1.4 | 6 |
| 4897 | A novel cervical cancer suppressor 3 (CCS-3) interacts with the BTB domain of PLZF and inhibits the cell growth by inducing apoptosis. <i>FEBS Letters</i> , 2006, 580, 4073-4080. | 1.3 | 25 |
| 4898 | Cloning and characterization of the <i>erg1</i> gene of <i>Trichoderma harzianum</i> : Effect of the <i>erg1</i> silencing on ergosterol biosynthesis and resistance to terbinafine. <i>Fungal Genetics and Biology</i> , 2006, 43, 164-178. | 0.9 | 77 |
| 4899 | Mutation D104G in ANT1 gene: Complementation study in <i>Saccharomyces cerevisiae</i> as a model system. <i>Biochemical and Biophysical Research Communications</i> , 2006, 341, 810-815. | 1.0 | 17 |
| 4900 | Calcium signaling and sugar-induced activation of plasma membrane H ⁺ -ATPase in <i>Saccharomyces cerevisiae</i> cells. <i>Biochemical and Biophysical Research Communications</i> , 2006, 343, 1234-1243. | 1.0 | 24 |

| # | ARTICLE | IF | CITATIONS |
|------|---|-----|-----------|
| 4901 | Gcn5p contributes to the bidirectional character of the UGA3â€“GLT1 yeast promoter. <i>Biochemical and Biophysical Research Communications</i> , 2006, 348, 989-996. | 1.0 | 2 |
| 4902 | Scanning Mutagenesis Studies Reveal Multiple Distinct Regions within the Human Protein Kinase C Alpha Regulatory Domain Important for Phorbol Ester-dependent Activation of the Enzyme. <i>Journal of Molecular Biology</i> , 2006, 357, 820-832. | 2.0 | 4 |
| 4903 | A novel system of genetic transformation allows multiple integrations of a desired gene in <i>Saccharomyces cerevisiae</i> chromosomes. <i>Journal of Microbiological Methods</i> , 2006, 67, 437-445. | 0.7 | 10 |
| 4904 | Cdc48 is required for the stability of Cut1/separase in mitotic anaphase. <i>Journal of Structural Biology</i> , 2006, 156, 50-61. | 1.3 | 25 |
| 4905 | Efficient expression of the 15-kDa form of infectious pancreatic necrosis virus VP5 by suppression of a UGA codon. <i>Virus Research</i> , 2006, 122, 61-68. | 1.1 | 2 |
| 4906 | High-Resolution Mutation Mapping Reveals Parallel Experimental Evolution in Yeast. <i>PLoS Biology</i> , 2006, 4, e256. | 2.6 | 73 |
| 4907 | Characterization of an Arabidopsis inositol 1,3,4,5,6-pentakisphosphate 2-kinase (AtIPK1). <i>Biochemical Journal</i> , 2006, 394, 95-103. | 1.7 | 57 |
| 4908 | Interactions between Hairy/Enhancer of Split-related proteins and the pancreatic transcription factor Ptf1-p48 modulate function of the PTF1 transcriptional complex. <i>Biochemical Journal</i> , 2006, 393, 679-685. | 1.7 | 33 |
| 4909 | Characterization of yeast pyruvate kinase 1 as a protein kinase A substrate, and specificity of the phosphorylation site sequence in the whole protein. <i>Biochemical Journal</i> , 2006, 396, 117-126. | 1.7 | 35 |
| 4910 | Mg ²⁺ and a Key Lysine Modulate Exchange Activity of Eukaryotic Translation Elongation Factor 1B \pm . <i>Journal of Biological Chemistry</i> , 2006, 281, 19457-19468. | 1.6 | 34 |
| 4911 | Genetic and proteomic evidences support the localization of yeast enolase in the cell surface. <i>Proteomics</i> , 2006, 6, S107-S118. | 1.3 | 68 |
| 4912 | Spsgt1, a new essential gene of <i>Schizosaccharomyces pombe</i> , is involved in carbohydrate metabolism. <i>Yeast</i> , 2006, 23, 35-53. | 0.8 | 18 |
| 4913 | Isolation of Δ 12 and Δ 3-fatty acid desaturase genes from the yeast <i>Kluyveromyces lactis</i> and their heterologous expression to produce linoleic and Δ -linolenic acids in <i>Saccharomyces cerevisiae</i> . <i>Yeast</i> , 2006, 23, 605-612. | 0.8 | 60 |
| 4914 | Molecular characterization of the Tim9 homologue from the methylotrophic yeast <i>Pichia methanolica</i> . <i>Yeast</i> , 2006, 23, 701-706. | 0.8 | 1 |
| 4915 | A two-step method for the introduction of single or multiple defined point mutations into the genome of <i>Saccharomyces cerevisiae</i> . <i>Yeast</i> , 2006, 23, 825-831. | 0.8 | 46 |
| 4916 | Yeast Derlin Dfm1 interacts with Cdc48 and functions in ER homeostasis. <i>Yeast</i> , 2006, 23, 1053-1064. | 0.8 | 71 |
| 4917 | High frequency gene transfer by microprojectile bombardment of intact conidia from the entomopathogenic fungus <i>Paecilomyces fumosoroseus</i> . <i>FEMS Microbiology Letters</i> , 2006, 156, 95-99. | 0.7 | 16 |
| 4918 | STE11 disruption reveals the central role of a MAPK pathway in dimorphism and mating in <i>Yarrowia lipolytica</i> . <i>FEMS Yeast Research</i> , 2006, 6, 801-815. | 1.1 | 35 |

| # | ARTICLE | IF | CITATIONS |
|------|---|-----|-----------|
| 4919 | CaMac1, a <i>Candida albicans</i> Copper Ion-sensing Transcription Factor, Promotes Filamentous and Invasive Growth in <i>Saccharomyces cerevisiae</i> . <i>Acta Biochimica Et Biophysica Sinica</i> , 2006, 38, 213-217. | 0.9 | 15 |
| 4920 | Sel1p/Ubx2p Participates in a Distinct Cdc48p-Dependent Endoplasmic Reticulum-Associated Degradation Pathway. <i>Traffic</i> , 2006, 7, 1213-1223. | 1.3 | 20 |
| 4921 | Mating type regulation of cellular tolerance to DNA damage is specific to the DNA post-replication repair and mutagenesis pathway. <i>Molecular Microbiology</i> , 2006, 59, 637-650. | 1.2 | 11 |
| 4922 | The UGA3-GLT1 intergenic region constitutes a promoter whose bidirectional nature is determined by chromatin organization in <i>Saccharomyces cerevisiae</i> . <i>Molecular Microbiology</i> , 2006, 59, 1790-1806. | 1.2 | 18 |
| 4923 | PDR16-mediated azole resistance in <i>Candida albicans</i> . <i>Molecular Microbiology</i> , 2006, 60, 1546-1562. | 1.2 | 62 |
| 4924 | The phosphatase Ppt1 is a dedicated regulator of the molecular chaperone Hsp90. <i>EMBO Journal</i> , 2006, 25, 367-376. | 3.5 | 153 |
| 4925 | Application of sequential integration for metabolic engineering of 1,2-propanediol production in yeast. <i>Metabolic Engineering</i> , 2006, 8, 58-65. | 3.6 | 41 |
| 4926 | Metabolic Engineering of <i>Saccharomyces cerevisiae</i> for Efficient Production of Pure L-(+)-Lactic Acid. <i>Applied Biochemistry and Biotechnology</i> , 2006, 131, 795-807. | 1.4 | 47 |
| 4927 | <i>Arabidopsis thaliana</i> UBC13: Implication of Error-free DNA Damage Tolerance and Lys63-linked Polyubiquitylation in Plants. <i>Plant Molecular Biology</i> , 2006, 61, 241-253. | 2.0 | 105 |
| 4928 | Molecular Cloning of the Phytase Gene from <i>Citrobacter Æbraakii</i> and its Expression in <i>Saccharomyces Æcerevisiae</i> . <i>Biotechnology Letters</i> , 2006, 28, 33-38. | 1.1 | 33 |
| 4929 | FLO11 mediated filamentous growth of the yeast <i>Saccharomyces cerevisiae</i> depends on the expression of the ribosomal RPS26 genes. <i>Molecular Genetics and Genomics</i> , 2006, 276, 113-125. | 1.0 | 22 |
| 4930 | Identification of a bHLH-type G-box binding factor and its regulation activity with G-box and Box I elements of the PsCHS1 promoter. <i>Plant Cell Reports</i> , 2006, 26, 85-93. | 2.8 | 22 |
| 4931 | Effective display of metallothionein tandem repeats on the bioadsorption of cadmium ion. <i>Applied Microbiology and Biotechnology</i> , 2006, 70, 458-463. | 1.7 | 68 |
| 4932 | Enhanced Protein Export in <i>Saccharomyces cerevisiae nud1</i> Mutants Is an Active Process. <i>Current Microbiology</i> , 2006, 53, 496-501. | 1.0 | 2 |
| 4933 | The yeast CPC2/ASC1 gene is regulated by the transcription factors Fhl1p and Ifh1p. <i>Current Genetics</i> , 2006, 49, 218-228. | 0.8 | 11 |
| 4934 | A branched-chain amino acid aminotransferase gene isolated from <i>Hordeum vulgare</i> is differentially regulated by drought stress. <i>Theoretical and Applied Genetics</i> , 2006, 113, 965-976. | 1.8 | 46 |
| 4935 | A <i>Phytophthora sojae</i> gene of glyceraldehyde-3-phosphate dehydrogenase (GAPDH) induced in host infection and its anti-oxidative function in yeast. <i>Science Bulletin</i> , 2006, 51, 1316-1323. | 1.7 | 5 |
| 4936 | Protection against oxidative stress through SUA7/TFIIB regulation in <i>Saccharomyces cerevisiae</i> . <i>Free Radical Biology and Medicine</i> , 2006, 41, 1684-1693. | 1.3 | 4 |

| # | ARTICLE | IF | CITATIONS |
|------|---|-----|-----------|
| 4937 | TRIP6 transcriptional co-activator is a novel substrate of AMP-activated protein kinase. Cellular Signalling, 2006, 18, 1702-1712. | 1.7 | 26 |
| 4938 | Experimental Evolution of Mating Discrimination in Budding Yeast. Current Biology, 2006, 16, 280-286. | 1.8 | 60 |
| 4939 | Cellular localization and phosphorylation of Hrb1p is independent of Sky1p. Biochimica Et Biophysica Acta - Molecular Cell Research, 2006, 1763, 207-213. | 1.9 | 4 |
| 4940 | Hsp78 chaperone functions in restoration of mitochondrial network following heat stress. Biochimica Et Biophysica Acta - Molecular Cell Research, 2006, 1763, 141-151. | 1.9 | 42 |
| 4941 | Identification and characterization of triosephosphate isomerase that specifically interacts with the integrin α 5 β 1b cytoplasmic domain. Biochemical Pharmacology, 2006, 72, 551-557. | 2.0 | 17 |
| 4942 | A Small-Molecule-Protein Interaction System with Split-Ubiquitin as Sensor. ChemBioChem, 2006, 7, 936-942. | 1.3 | 18 |
| 4943 | Yeast expression of the VP8* fragment of the rotavirus spike protein and its use as immunogen in mice. Biotechnology and Bioengineering, 2006, 93, 89-98. | 1.7 | 21 |
| 4944 | New pFA-cassettes for PCR-based gene manipulation in <i>Candida albicans</i> . Journal of Basic Microbiology, 2006, 46, 416-429. | 1.8 | 75 |
| 4945 | The [PSI ⁺] Prion of <i>Saccharomyces cerevisiae</i> Can Be Propagated by an Hsp104 Orthologue from <i>Candida albicans</i> . Eukaryotic Cell, 2006, 5, 217-225. | 3.4 | 27 |
| 4946 | Genomic Reconstruction by Serial Mitotic Recombination of Yeast Artificial Chromosomes. , 2006, 349, 117-126. | | 0 |
| 4947 | The Arf-GTPase-Activating Protein Gcs1p Is Essential for Sporulation and Regulates the Phospholipase D Spo14p. Eukaryotic Cell, 2006, 5, 112-124. | 3.4 | 18 |
| 4948 | Regulation of Salt Tolerance by <i>Torulospira delbrueckii</i> Calcineurin Target Crz1p. Eukaryotic Cell, 2006, 5, 469-479. | 3.4 | 31 |
| 4949 | Targeted Gene Silencing in the Model Mushroom <i>Coprinopsis cinerea</i> (<i>Coprinus cinereus</i>) by Expression of Homologous Hairpin RNAs. Eukaryotic Cell, 2006, 5, 732-744. | 3.4 | 73 |
| 4950 | A Genomewide Screen Reveals a Role of Mitochondria in Anaerobic Uptake of Sterols in Yeast. Molecular Biology of the Cell, 2006, 17, 90-103. | 0.9 | 75 |
| 4951 | Cytological analysis of <i>Saccharomyces cerevisiae</i> cells supporting cymbidium ringspot virus defective interfering RNA replication. Journal of General Virology, 2006, 87, 705-714. | 1.3 | 30 |
| 4952 | Yeh1 Constitutes the Major Steryl Ester Hydrolase under Heme-Deficient Conditions in <i>Saccharomyces cerevisiae</i> . Eukaryotic Cell, 2006, 5, 1018-1025. | 3.4 | 31 |
| 4953 | The rad51-K191R ATPase-Defective Mutant Is Impaired for Presynaptic Filament Formation. Molecular and Cellular Biology, 2006, 26, 9544-9554. | 1.1 | 40 |
| 4954 | Yeast Gcn4p Stabilization Is Initiated by the Dissociation of the Nuclear Pho85p/Pcl5p Complex. Molecular Biology of the Cell, 2006, 17, 2952-2962. | 0.9 | 22 |

| # | ARTICLE | IF | CITATIONS |
|------|---|-----|-----------|
| 4955 | Fission Yeast Rho5p GTPase Is a Functional Parologue of Rho1p That Plays a Role in Survival of Spores and Stationary-Phase Cells. <i>Eukaryotic Cell</i> , 2006, 5, 435-446. | 3.4 | 18 |
| 4956 | Cross-talk among Rho GTPases Acting Downstream of PI 3-Kinase Induces Mesenchymal Transformation of Corneal Endothelial Cells Mediated by FGF-2. , 2006, 47, 2358. | | 36 |
| 4957 | Analyses of SUM1-1-Mediated Long-Range Repression. <i>Genetics</i> , 2006, 172, 99-112. | 1.2 | 10 |
| 4958 | Molecular Cloning and Characterization of a Novel 3â€²-Phosphoadenosine 5â€²-Phosphosulfate Transporter, PAPST2. <i>Journal of Biological Chemistry</i> , 2006, 281, 10945-10953. | 1.6 | 67 |
| 4959 | Production of Xylitol from d-Xylose by a Xylitol Dehydrogenase Gene-Disrupted Mutant of <i>Candida tropicalis</i> . <i>Applied and Environmental Microbiology</i> , 2006, 72, 4207-4213. | 1.4 | 98 |
| 4960 | The yeast Igl family member Sro7p is an effector of the secretory Rab GTPase Sec4p. <i>Journal of Cell Biology</i> , 2006, 172, 55-66. | 2.3 | 81 |
| 4961 | The Phosphatidylinositol 4,5-Biphosphate and TORC2 Binding Proteins Slm1 and Slm2 Function in Sphingolipid Regulation. <i>Molecular and Cellular Biology</i> , 2006, 26, 5861-5875. | 1.1 | 125 |
| 4962 | The RAD6/BRE1 Histone Modification Pathway in <i>Saccharomyces</i> Confers Radiation Resistance Through a RAD51-Dependent Process That Is Independent of RAD18. <i>Genetics</i> , 2006, 173, 1951-1968. | 1.2 | 54 |
| 4963 | Arl1p is involved in transport of the GPI-anchored protein Gas1p from the late Golgi to the plasma membrane. <i>Journal of Cell Science</i> , 2006, 119, 3845-3855. | 1.2 | 36 |
| 4964 | Specific and Nonspecific Membrane-binding Determinants Cooperate in Targeting Phosphatidylinositol Transfer Protein Î²-Isoform to the Mammalian Trans-Golgi Network. <i>Molecular Biology of the Cell</i> , 2006, 17, 2498-2512. | 0.9 | 38 |
| 4965 | The Effect of Pyruvate Decarboxylase Gene Knockout in <i>Saccharomyces cerevisiae</i> on L-Lactic Acid Production. <i>Bioscience, Biotechnology and Biochemistry</i> , 2006, 70, 1148-1153. | 0.6 | 80 |
| 4966 | DNA Polymerase 4 of <i>Saccharomyces cerevisiae</i> Is Important for Accurate Repair of Methyl-Methanesulfonate-Induced DNA Damage. <i>Genetics</i> , 2006, 172, 89-98. | 1.2 | 8 |
| 4967 | Plasma Membrane Localization of Ras Requires Class C Vps Proteins and Functional Mitochondria in <i>Saccharomyces cerevisiae</i> . <i>Molecular and Cellular Biology</i> , 2006, 26, 3243-3255. | 1.1 | 52 |
| 4968 | Mutational and Functional Analysis of the Cryptic N-Terminal Targeting Signal for Both Mitochondria and Peroxisomes in Yeast Peroxisomal Citrate Synthase Cit2p. <i>Journal of Biochemistry</i> , 2006, 140, 121-133. | 0.9 | 8 |
| 4969 | Deletion of BCY1 from the <i>Saccharomyces cerevisiae</i> Genome Is Semidominant and Induces Autolytic Phenotypes Suitable for Improvement of Sparkling Wines. <i>Applied and Environmental Microbiology</i> , 2006, 72, 2351-2358. | 1.4 | 45 |
| 4970 | The Integral Membrane Protein Pom34p Functionally Links Nucleoporin Subcomplexes. <i>Genetics</i> , 2006, 172, 1441-1457. | 1.2 | 51 |
| 4971 | The E and G Subunits of the Yeast V-ATPase Interact Tightly and Are Both Present at More Than One Copy per V1 Complex. <i>Journal of Biological Chemistry</i> , 2006, 281, 22752-22760. | 1.6 | 65 |
| 4972 | Unique Classes of Mutations in the <i>Saccharomyces cerevisiae</i> G-Protein Translation Elongation Factor 1A Suppress the Requirement for Guanine Nucleotide Exchange. <i>Genetics</i> , 2006, 174, 651-663. | 1.2 | 13 |

| # | ARTICLE | IF | CITATIONS |
|------|--|-----|-----------|
| 4973 | Hog1 Mitogen-Activated Protein Kinase Plays Conserved and Distinct Roles in the Osmotolerant Yeast <i>Torulasporea delbrueckii</i> . <i>Eukaryotic Cell</i> , 2006, 5, 1410-1419. | 3.4 | 15 |
| 4974 | Inducible Yeast System for Viral RNA Recombination Reveals Requirement for an RNA Replication Signal on Both Parental RNAs. <i>Journal of Virology</i> , 2006, 80, 8316-8328. | 1.5 | 17 |
| 4975 | Knockdown of an Arbuscular Mycorrhiza-inducible Phosphate Transporter Gene of <i>Lotus japonicus</i> Suppresses Mutualistic Symbiosis. <i>Plant and Cell Physiology</i> , 2006, 47, 807-817. | 1.5 | 193 |
| 4976 | Cell Surface Expression of Bacterial Esterase A by <i>Saccharomyces cerevisiae</i> and Its Enhancement by Constitutive Activation of the Cellular Unfolded Protein Response. <i>Applied and Environmental Microbiology</i> , 2006, 72, 7140-7147. | 1.4 | 32 |
| 4977 | Comparative study of the effects of heptameric slippery site composition on -1 frameshifting among different eukaryotic systems. <i>Rna</i> , 2006, 12, 666-673. | 1.6 | 45 |
| 4978 | Casein Kinase II Phosphorylation of the Yeast Phospholipid Synthesis Transcription Factor Opi1p. <i>Journal of Biological Chemistry</i> , 2006, 281, 4754-4761. | 1.6 | 26 |
| 4979 | Specific Effects of Ribosome-Tethered Molecular Chaperones on Programmed +1 Ribosomal Frameshifting. <i>Eukaryotic Cell</i> , 2006, 5, 762-770. | 3.4 | 28 |
| 4980 | An mRNA sequence derived from a programmed frameshifting signal decreases codon discrimination during translation initiation. <i>Rna</i> , 2006, 12, 1154-1160. | 1.6 | 2 |
| 4981 | Chromatin-Modifying Enzymes Are Essential When the <i>Saccharomyces cerevisiae</i> Morphogenesis Checkpoint Is Constitutively Activated. <i>Genetics</i> , 2006, 174, 1135-1149. | 1.2 | 16 |
| 4982 | The Ribosomal Shunt Translation Strategy of Cauliflower Mosaic Virus Has Evolved from Ancient Long Terminal Repeats. <i>Journal of Virology</i> , 2006, 80, 3811-3822. | 1.5 | 8 |
| 4983 | An Arc of Unpaired "Hinge Bases" Facilitates Information Exchange among Functional Centers of the Ribosome. <i>Molecular and Cellular Biology</i> , 2006, 26, 8992-9002. | 1.1 | 40 |
| 4984 | An intimate collaboration between peroxisomes and lipid bodies. <i>Journal of Cell Biology</i> , 2006, 173, 719-731. | 2.3 | 329 |
| 4985 | <i>Caenorhabditis elegans</i> UNC-96 Is a New Component of M-Lines That Interacts with UNC-98 and Paramyosin and Is Required in Adult Muscle for Assembly and/or Maintenance of Thick Filaments. <i>Molecular Biology of the Cell</i> , 2006, 17, 3832-3847. | 0.9 | 40 |
| 4986 | Determinants of Rbp1p Localization in Specific Cytoplasmic mRNA-processing Foci, P-bodies. <i>Journal of Biological Chemistry</i> , 2006, 281, 29379-29390. | 1.6 | 12 |
| 4987 | Signaling Alkaline pH Stress in the Yeast <i>Saccharomyces cerevisiae</i> through the Wsc1 Cell Surface Sensor and the Slit2 MAPK Pathway. <i>Journal of Biological Chemistry</i> , 2006, 281, 39785-39795. | 1.6 | 107 |
| 4988 | Kinectin-dependent Assembly of Translation Elongation Factor-1 Complex on Endoplasmic Reticulum Regulates Protein Synthesis. <i>Journal of Biological Chemistry</i> , 2006, 281, 33621-33634. | 1.6 | 27 |
| 4989 | Pheromone-Induced Degradation of Ste12 Contributes to Signal Attenuation and the Specificity of Developmental Fate. <i>Eukaryotic Cell</i> , 2006, 5, 2147-2160. | 3.4 | 17 |
| 4990 | In Vivo Targeting of the Yeast Pop2 Deadenylase Subunit to Reporter Transcripts Induces Their Rapid Degradation and Generates New Decay Intermediates. <i>Journal of Biological Chemistry</i> , 2006, 281, 25940-25947. | 1.6 | 31 |

| # | ARTICLE | IF | CITATIONS |
|------|---|-----|-----------|
| 4991 | Metabolic Engineering of <i>Saccharomyces cerevisiae</i> Producing Nicotianamine: Potential for Industrial Biosynthesis of a Novel Antihypertensive Substrate. <i>Bioscience, Biotechnology and Biochemistry</i> , 2006, 70, 1408-1415. | 0.6 | 18 |
| 4992 | Dissecting toxin immunity in virus-infected killer yeast uncovers an intrinsic strategy of self-protection. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 2006, 103, 3810-3815. | 3.3 | 66 |
| 4993 | Translation Elongation Factor 2 Anticodon Mimicry Domain Mutants Affect Fidelity and Diphtheria Toxin Resistance. <i>Journal of Biological Chemistry</i> , 2006, 281, 32639-32648. | 1.6 | 99 |
| 4994 | Characterization of the Transport Mechanism and Permeant Binding Profile of the Uridine Permease Fui1p of <i>Saccharomyces cerevisiae</i> . <i>Journal of Biological Chemistry</i> , 2006, 281, 28210-28221. | 1.6 | 13 |
| 4995 | PKR1 Encodes an Assembly Factor for the Yeast V-Type ATPase. <i>Journal of Biological Chemistry</i> , 2006, 281, 32025-32035. | 1.6 | 34 |
| 4996 | The Transcriptional Response of the Yeast Na ⁺ -ATPase ENA1 Gene to Alkaline Stress Involves Three Main Signaling Pathways*. <i>Journal of Biological Chemistry</i> , 2006, 281, 36632-36642. | 1.6 | 80 |
| 4997 | Effects of C-terminal Truncations on Trafficking of the Yeast Plasma Membrane H ⁺ -ATPase. <i>Journal of Biological Chemistry</i> , 2006, 281, 23887-23898. | 1.6 | 15 |
| 4998 | Domain and Nucleotide Dependence of the Interaction between <i>Saccharomyces cerevisiae</i> Translation Elongation Factors 3 and 1A. <i>Journal of Biological Chemistry</i> , 2006, 281, 32318-32326. | 1.6 | 34 |
| 4999 | Specific modulation of apoptosis and Bcl-xL phosphorylation in yeast by distinct mammalian protein kinase C isoforms. <i>Journal of Cell Science</i> , 2006, 119, 3171-3181. | 1.2 | 41 |
| 5000 | DNA Damage Checkpoints Are Involved in Postreplication Repair. <i>Genetics</i> , 2006, 174, 1789-1800. | 1.2 | 34 |
| 5001 | Regulation of the <i>Saccharomyces cerevisiae</i> EKI1-encoded Ethanolamine Kinase by Zinc Depletion. <i>Journal of Biological Chemistry</i> , 2006, 281, 13110-13116. | 1.6 | 22 |
| 5002 | TEDS Site Phosphorylation of the Yeast Myosins I Is Required for Ligand-induced but Not for Constitutive Endocytosis of the G Protein-coupled Receptor Ste2p. <i>Journal of Biological Chemistry</i> , 2006, 281, 11104-11114. | 1.6 | 28 |
| 5003 | Long Chain Base Tolerance in <i>Saccharomyces cerevisiae</i> Is Induced by Retrograde Signals from the Mitochondria. <i>Journal of Biological Chemistry</i> , 2006, 281, 6376-6384. | 1.6 | 47 |
| 5004 | Characterization of Mmp37p, a <i>Saccharomyces cerevisiae</i> Mitochondrial Matrix Protein with a Role in Mitochondrial Protein Import. <i>Molecular Biology of the Cell</i> , 2006, 17, 4051-4062. | 0.9 | 56 |
| 5005 | Identification and Characterization of a Novel <i>Drosophila</i> 3'-Phosphoadenosine 5'-Phosphosulfate Transporter. <i>Journal of Biological Chemistry</i> , 2006, 281, 28508-28517. | 1.6 | 30 |
| 5006 | Vma9p (Subunit e) Is an Integral Membrane V0 Subunit of the Yeast V-ATPase. <i>Journal of Biological Chemistry</i> , 2006, 281, 15312-15319. | 1.6 | 30 |
| 5007 | <i>DGA1</i> (diacylglycerol acyltransferase gene) overexpression and leucine biosynthesis significantly increase lipid accumulation in the <i>snf2</i> disruptant of <i>Saccharomyces cerevisiae</i> . <i>Biochemical Journal</i> , 2007, 408, 61-68. | 1.7 | 109 |
| 5008 | MpkA-Dependent and -Independent Cell Wall Integrity Signaling in <i>Aspergillus nidulans</i> . <i>Eukaryotic Cell</i> , 2007, 6, 1497-1510. | 3.4 | 157 |

| # | ARTICLE | IF | CITATIONS |
|------|--|-----|-----------|
| 5009 | Chromosome mobility during meiotic prophase in <i>Saccharomyces cerevisiae</i> . Proceedings of the National Academy of Sciences of the United States of America, 2007, 104, 16934-16939. | 3.3 | 144 |
| 5010 | Phosphorylation of Human CTP Synthetase 1 by Protein Kinase C. Journal of Biological Chemistry, 2007, 282, 17613-17622. | 1.6 | 33 |
| 5011 | Role of the Synthase Domain of Ags1p in Cell Wall β -Glucan Biosynthesis in Fission Yeast. Journal of Biological Chemistry, 2007, 282, 18969-18979. | 1.6 | 27 |
| 5012 | DNA Sequences from Arabidopsis, Which Encode Protein Kinases and Function as Upstream Regulators of Snf1 in Yeast. Journal of Biological Chemistry, 2007, 282, 10472-10479. | 1.6 | 45 |
| 5013 | The Cellular Functions of the Yeast Lipin Homolog Pah1p Are Dependent on Its Phosphatidate Phosphatase Activity. Journal of Biological Chemistry, 2007, 282, 37026-37035. | 1.6 | 150 |
| 5014 | Identification of a Novel Lysophospholipid Acyltransferase in <i>Saccharomyces cerevisiae</i> . Journal of Biological Chemistry, 2007, 282, 30562-30569. | 1.6 | 103 |
| 5015 | Phosphorylation of Human CTP Synthetase 1 by Protein Kinase A. Journal of Biological Chemistry, 2007, 282, 5367-5377. | 1.6 | 20 |
| 5016 | Deletion of the nuclear exosome component RRP6 leads to continued accumulation of the histone mRNA HTB1 in S-phase of the cell cycle in <i>Saccharomyces cerevisiae</i> . Nucleic Acids Research, 2007, 35, 6268-6279. | 6.5 | 32 |
| 5017 | Cymbidium ringspot virus defective interfering RNA replication in yeast cells occurs on endoplasmic reticulum-derived membranes in the absence of peroxisomes. Journal of General Virology, 2007, 88, 1634-1642. | 1.3 | 18 |
| 5018 | The Alternative Pathway of Glutathione Degradation Is Mediated by a Novel Protein Complex Involving Three New Genes in <i>Saccharomyces cerevisiae</i> . Genetics, 2007, 175, 1137-1151. | 1.2 | 67 |
| 5019 | Respiratory Deficiency Mediates the Regulation of CHO1-encoded Phosphatidylserine Synthase by mRNA Stability in <i>Saccharomyces cerevisiae</i> . Journal of Biological Chemistry, 2007, 282, 31217-31227. | 1.6 | 8 |
| 5020 | An mRNA sequence derived from the yeast EST3 gene stimulates programmed +1 translational frameshifting. Rna, 2007, 13, 606-613. | 1.6 | 18 |
| 5021 | Proteomic analysis of the U1 snRNP of <i>Schizosaccharomyces pombe</i> reveals three essential organism-specific proteins. Nucleic Acids Research, 2007, 35, 1391-1401. | 6.5 | 13 |
| 5022 | Conserved amino acids in each subunit of the heterologomeric tRNA methylase from <i>Saccharomyces cerevisiae</i> contribute to tRNA binding. Nucleic Acids Research, 2007, 35, 6808-6819. | 6.5 | 32 |
| 5023 | Development of a Highly Efficient Gene Targeting System Induced by Transient Repression of YKU80 Expression in <i>Candida glabrata</i> . Eukaryotic Cell, 2007, 6, 1239-1247. | 3.4 | 85 |
| 5024 | Absence of Btn1p in the yeast model for juvenile Batten disease may cause arginine to become toxic to yeast cells. Human Molecular Genetics, 2007, 16, 1007-1016. | 1.4 | 18 |
| 5025 | Roles of the negatively charged N-terminal extension of <i>Saccharomyces cerevisiae</i> ribosomal protein S5 revealed by characterization of a yeast strain containing human ribosomal protein S5. Rna, 2007, 13, 2116-2128. | 1.6 | 28 |
| 5026 | Function-Altering SNPs in the Human Multidrug Transporter Gene ABCB1 Identified Using a <i>Saccharomyces</i> -Based Assay. PLoS Genetics, 2007, 3, e39. | 1.5 | 53 |

| # | ARTICLE | IF | CITATIONS |
|------|---|-----|-----------|
| 5027 | GCP6 Binds to Intermediate Filaments: A Novel Function of Keratins in the Organization of Microtubules in Epithelial Cells. <i>Molecular Biology of the Cell</i> , 2007, 18, 781-794. | 0.9 | 44 |
| 5028 | Identification of functional, endogenous programmed +1 ribosomal frameshift signals in the genome of <i>Saccharomyces cerevisiae</i> . <i>Nucleic Acids Research</i> , 2007, 35, 165-174. | 6.5 | 145 |
| 5029 | Overexpression of Autophagy-Related Genes Inhibits Yeast Filamentous Growth. <i>Autophagy</i> , 2007, 3, 604-609. | 4.3 | 12 |
| 5030 | Yeast Phenotypic Assays on Translational Control. <i>Methods in Enzymology</i> , 2007, 429, 105-137. | 0.4 | 20 |
| 5031 | Shuffling of Promoters for Multiple Genes To Optimize Xylose Fermentation in an Engineered <i>Saccharomyces cerevisiae</i> Strain. <i>Applied and Environmental Microbiology</i> , 2007, 73, 6072-6077. | 1.4 | 90 |
| 5032 | The Hsp40 Molecular Chaperone Ydj1p, Along With the Protein Kinase C Pathway, Affects Cell-Wall Integrity in the Yeast <i>Saccharomyces cerevisiae</i> . <i>Genetics</i> , 2007, 175, 1649-1664. | 1.2 | 26 |
| 5033 | A Large-scale Protein-Protein Interaction Analysis in <i>Synechocystis</i> sp. PCC6803. <i>DNA Research</i> , 2007, 14, 207-216. | 1.5 | 170 |
| 5034 | Improper Organization of the Actin Cytoskeleton Affects Protein Synthesis at Initiation. <i>Molecular and Cellular Biology</i> , 2007, 27, 1974-1989. | 1.1 | 77 |
| 5035 | Genome-Wide Expression and Location Analyses of the <i>Candida albicans</i> Tac1p Regulon. <i>Eukaryotic Cell</i> , 2007, 6, 2122-2138. | 3.4 | 118 |
| 5036 | Shs1 Plays Separable Roles in Septin Organization and Cytokinesis in <i>Saccharomyces cerevisiae</i> . <i>Genetics</i> , 2007, 177, 215-229. | 1.2 | 48 |
| 5037 | Negative Transcriptional Regulation of Multidrug Resistance Gene Expression by an Hsp70 Protein. <i>Journal of Biological Chemistry</i> , 2007, 282, 26822-26831. | 1.6 | 29 |
| 5038 | Atg18 Regulates Organelle Morphology and Fab1 Kinase Activity Independent of Its Membrane Recruitment by Phosphatidylinositol 3,5-Bisphosphate. <i>Molecular Biology of the Cell</i> , 2007, 18, 4232-4244. | 0.9 | 112 |
| 5039 | The Yeast PH Domain Proteins Slm1 and Slm2 Are Targets of Sphingolipid Signaling during the Response to Heat Stress. <i>Molecular and Cellular Biology</i> , 2007, 27, 633-650. | 1.1 | 76 |
| 5040 | Heterologous Production of Dihomo- δ^3 -Linolenic Acid in <i>Saccharomyces cerevisiae</i> . <i>Applied and Environmental Microbiology</i> , 2007, 73, 6965-6971. | 1.4 | 41 |
| 5041 | A specific role for the C-terminal region of the Poly(A)-binding protein in mRNA decay. <i>Nucleic Acids Research</i> , 2007, 35, 6017-6028. | 6.5 | 45 |
| 5042 | Yeast Cells Lacking the <i>CIT1</i> -encoded Mitochondrial Citrate Synthase Are Hypersusceptible to Heat- or Aging-induced Apoptosis. <i>Molecular Biology of the Cell</i> , 2007, 18, 3556-3567. | 0.9 | 24 |
| 5043 | Novel <i>sfi1</i> Alleles Uncover Additional Functions for Sfi1p in Bipolar Spindle Assembly and Function. <i>Molecular Biology of the Cell</i> , 2007, 18, 2047-2056. | 0.9 | 27 |
| 5044 | The <i>Saccharomyces</i> Homolog of Mammalian RACK1, Cpc2/Asc1p, Is Required for FLO11-dependent Adhesive Growth and Dimorphism. <i>Molecular and Cellular Proteomics</i> , 2007, 6, 1968-1979. | 2.5 | 53 |

| # | ARTICLE | IF | CITATIONS |
|------|---|-----|-----------|
| 5045 | One- plus Two-hybrid System, a Novel Yeast Genetic Selection for Specific Missense Mutations Disrupting Protein/Protein Interactions. <i>Molecular and Cellular Proteomics</i> , 2007, 6, 1727-1740. | 2.5 | 23 |
| 5046 | PsTRXh1 and PsTRXh2 Are Both Pea h-Type Thioredoxins with Antagonistic Behavior in Redox Imbalances. <i>Plant Physiology</i> , 2007, 143, 300-311. | 2.3 | 35 |
| 5047 | Conformational Dynamics of the Major Yeast Phosphatidylinositol Transfer Protein Sec14p: Insight into the Mechanisms of Phospholipid Exchange and Diseases of Sec14p-Like Protein Deficiencies. <i>Molecular Biology of the Cell</i> , 2007, 18, 1928-1942. | 0.9 | 55 |
| 5048 | <i>Saccharomyces cerevisiae</i> Ebs1p is a putative ortholog of human Smg7 and promotes nonsense-mediated mRNA decay. <i>Nucleic Acids Research</i> , 2007, 35, 7688-7697. | 6.5 | 63 |
| 5049 | DNA Recombination-Initiation Plays a Role in the Extremely Biased Inheritance of Yeast [rho â'] Mitochondrial DNA That Contains the Replication Origin ori5. <i>Molecular and Cellular Biology</i> , 2007, 27, 1133-1145. | 1.1 | 49 |
| 5050 | Saturation mutagenesis of a +1 programmed frameshift-inducing mRNA sequence derived from a yeast retrotransposon. <i>Rna</i> , 2007, 13, 1940-1947. | 1.6 | 15 |
| 5051 | Degradation of hypomodified tRNA_i^{Met} in vivo involves RNA-dependent ATPase activity of the DExH helicase Mtr4p. <i>Rna</i> , 2008, 14, 107-116. | 1.6 | 84 |
| 5052 | A multifunctional desaturase involved in the biosynthesis of the processionary moth sex pheromone. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 2007, 104, 16444-16449. | 3.3 | 46 |
| 5053 | A Conserved Sequence Immediately N-terminal to the Bateman Domains in AMP-activated Protein Kinase Î³ Subunits Is Required for the Interaction with the Î² Subunits. <i>Journal of Biological Chemistry</i> , 2007, 282, 16117-16125. | 1.6 | 25 |
| 5054 | RAVE Is Essential for the Efficient Assembly of the C Subunit with the Vacuolar H ⁺ -ATPase. <i>Journal of Biological Chemistry</i> , 2007, 282, 26185-26194. | 1.6 | 73 |
| 5055 | Evidence for the Direct Involvement of the Proteasome in the Proteolytic Processing of the <i>Aspergillus nidulans</i> Zinc Finger Transcription Factor PacC. <i>Journal of Biological Chemistry</i> , 2007, 282, 34735-34747. | 1.6 | 78 |
| 5056 | Efficient Cargo Sorting by ESCRT-I and the Subsequent Release of ESCRT-I from Multivesicular Bodies Requires the Subunit Mvb12. <i>Molecular Biology of the Cell</i> , 2007, 18, 636-645. | 0.9 | 70 |
| 5057 | NPFXD-mediated Endocytosis Is Required for Polarity and Function of a Yeast Cell Wall Stress Sensor. <i>Molecular Biology of the Cell</i> , 2007, 18, 57-65. | 0.9 | 85 |
| 5058 | Overexpression of the Calcineurin Target CRZ1 Provides Freeze Tolerance and Enhances the Fermentative Capacity of Baker's Yeast. <i>Applied and Environmental Microbiology</i> , 2007, 73, 4824-4831. | 1.4 | 29 |
| 5059 | U2 toggles iteratively between the stem Ila and stem Ilc conformations to promote pre-mRNA splicing. <i>Genes and Development</i> , 2007, 21, 821-834. | 2.7 | 100 |
| 5060 | Molecular cloning and characterization of a cDNA for low-temperature inducible cytosolic glucose 6-phosphate dehydrogenase gene from <i>Chlorella vulgaris</i> and expression of the gene in <i>Saccharomyces cerevisiae</i> . <i>Plant Science</i> , 2007, 172, 649-658. | 1.7 | 14 |
| 5061 | Mre11 and Ku regulation of double-strand break repair by gene conversion and break-induced replication. <i>DNA Repair</i> , 2007, 6, 797-808. | 1.3 | 25 |
| 5062 | Expression pattern of inositol phosphate-related enzymes in rice (<i>Oryza sativa</i> L.): Implications for the phytic acid biosynthetic pathway. <i>Gene</i> , 2007, 405, 55-64. | 1.0 | 109 |

| # | ARTICLE | IF | CITATIONS |
|------|---|-----|-----------|
| 5063 | Functional characterization of KIHAP1: A model to foresee different mechanisms of transcriptional regulation by Hap1p in yeasts. <i>Gene</i> , 2007, 405, 96-107. | 1.0 | 18 |
| 5064 | Regulation of human estrogen receptor $\hat{\pm}$ -mediated gene transactivation in <i>Saccharomyces cerevisiae</i> by human coactivator and corepressor proteins. <i>Journal of Steroid Biochemistry and Molecular Biology</i> , 2007, 103, 189-195. | 1.2 | 4 |
| 5065 | A new method for the construction of a mutant library with a predictable occurrence rate using Poisson distribution. <i>Journal of Microbiological Methods</i> , 2007, 69, 442-450. | 0.7 | 5 |
| 5066 | Development of a transformation system for gene knock-out in the flavinogenic yeast <i>Pichia guilliermondii</i> . <i>Journal of Microbiological Methods</i> , 2007, 70, 13-19. | 0.7 | 26 |
| 5067 | Ras-pathway has a dual role in yeast galactose metabolism. <i>FEBS Letters</i> , 2007, 581, 2009-2016. | 1.3 | 5 |
| 5068 | Mitochondrial Phosphate Carrier Deficiency: A Novel Disorder of Oxidative Phosphorylation. <i>American Journal of Human Genetics</i> , 2007, 80, 478-484. | 2.6 | 142 |
| 5069 | Fluidization of Membrane Lipids Enhances the Tolerance of <i>Saccharomyces cerevisiae</i> to Freezing and Salt Stress. <i>Applied and Environmental Microbiology</i> , 2007, 73, 110-116. | 1.4 | 181 |
| 5070 | Expression of a Novel 90-kDa Protein, Lsd90, Involved in the Metabolism of Very Long-chain Fatty Acid-containing Phospholipids in a Mitosis-defective Fission Yeast Mutant. <i>Journal of Biochemistry</i> , 2007, 143, 369-375. | 0.9 | 7 |
| 5071 | Bacterial Regulatory Networks Include Direct Contact of Response Regulator Proteins: Interaction of RegA and NtrX in <i>Rhodobacter capsulatus</i> . <i>Journal of Molecular Microbiology and Biotechnology</i> , 2007, 13, 126-139. | 1.0 | 25 |
| 5072 | Fine-Tuning of Translation Termination Efficiency in <i>Saccharomyces cerevisiae</i> Involves Two Factors in Close Proximity to the Exit Tunnel of the Ribosome. <i>Genetics</i> , 2007, 177, 1527-1537. | 1.2 | 18 |
| 5073 | DNA-Mediated Transformation. <i>Methods in Molecular Biology</i> , 2007, 389, 27-42. | 0.4 | 19 |
| 5074 | TIMP-1 regulates cell proliferation by interacting with the ninth zinc finger domain of PLZF. <i>Journal of Cellular Biochemistry</i> , 2007, 101, 57-67. | 1.2 | 23 |
| 5075 | Functional motifs outside the kinase domain of yeast Srb10p. Their role in transcriptional regulation and protein-interactions with Tup1p and Srb11p. <i>Biochimica Et Biophysica Acta - Proteins and Proteomics</i> , 2007, 1774, 1227-1235. | 1.1 | 7 |
| 5076 | Proteomic analysis of the amyloid precursor protein fragment C99: expression in yeast. <i>Analytical Biochemistry</i> , 2007, 370, 162-170. | 1.1 | 9 |
| 5077 | The pheromone-induced nuclear accumulation of the Fus3 MAPK in yeast depends on its phosphorylation state and on Dig1 and Dig2. <i>BMC Cell Biology</i> , 2007, 8, 44. | 3.0 | 21 |
| 5078 | Dosage rescue by UBC4 restores cell wall integrity in <i>Saccharomyces cerevisiae</i> lacking the myosin type II gene MYO1. <i>Yeast</i> , 2007, 24, 343-355. | 0.8 | 8 |
| 5079 | Thetxl1+gene from <i>Schizosaccharomyces pombe</i> encodes a new thioredoxin-like 1 protein that participates in the antioxidant defence against tert-butyl hydroperoxide. <i>Yeast</i> , 2007, 24, 481-490. | 0.8 | 16 |
| 5080 | Direct in vivo access to potential gene targets of the RPD3 histone deacetylase using fitness-based interferential genetics. <i>Yeast</i> , 2007, 24, 575-587. | 0.8 | 5 |

| # | ARTICLE | IF | CITATIONS |
|------|--|-----|-----------|
| 5081 | Yeast vectors for the integration/expression of any sequence at theTYR1 locus. <i>Yeast</i> , 2007, 24, 761-766. | 0.8 | 3 |
| 5082 | High-efficiency yeast transformation using the LiAc/SS carrier DNA/PEG method. <i>Nature Protocols</i> , 2007, 2, 31-34. | 5.5 | 2,082 |
| 5083 | Quick and easy yeast transformation using the LiAc/SS carrier DNA/PEG method. <i>Nature Protocols</i> , 2007, 2, 35-37. | 5.5 | 368 |
| 5084 | Large-scale high-efficiency yeast transformation using the LiAc/SS carrier DNA/PEG method. <i>Nature Protocols</i> , 2007, 2, 38-41. | 5.5 | 325 |
| 5085 | Microtiter plate transformation using the LiAc/SS carrier DNA/PEG method. <i>Nature Protocols</i> , 2007, 2, 5-8. | 5.5 | 42 |
| 5086 | Frozen competent yeast cells that can be transformed with high efficiency using the LiAc/SS carrier DNA/PEG method. <i>Nature Protocols</i> , 2007, 2, 1-4. | 5.5 | 285 |
| 5087 | Structure of Sla1p homology domain 1 and interaction with the NPFxD endocytic internalization motif. <i>EMBO Journal</i> , 2007, 26, 1963-1971. | 3.5 | 21 |
| 5088 | ADP/ATP carrier is required for mitochondrial outer membrane permeabilization and cytochrome <i>c</i> release in yeast apoptosis. <i>Molecular Microbiology</i> , 2007, 66, 571-582. | 1.2 | 128 |
| 5089 | The zinc cluster transcription factor Tac1p regulates PDR16 expression in <i>Candida albicans</i> . <i>Molecular Microbiology</i> , 2007, 66, 440-452. | 1.2 | 81 |
| 5090 | The fluorouridine insensitive $\Delta fur1$ mutant is defective in equilibrative nucleoside transporter $\Delta 3$ (ENT3), and thus represents an important pyrimidine nucleoside uptake system in <i>Arabidopsis thaliana</i> . <i>Plant Journal</i> , 2007, 49, 855-864. | 2.8 | 37 |
| 5091 | Two novel AP2/ERF domain proteins interact with cis-element VWRE for wound-induced expression of the Tobacco <i>tpoxN1</i> gene. <i>Plant Journal</i> , 2007, 50, 1079-1092. | 2.8 | 43 |
| 5092 | Snc1p ν SNARE Transport to the Prospore Membrane During Yeast Sporulation is Dependent on Endosomal Retrieval Pathways. <i>Traffic</i> , 2007, 8, 1231-1245. | 1.3 | 19 |
| 5093 | A transcriptome analysis of isoamyl alcohol-induced filamentation in yeast reveals a novel role for Gre2p as isovaleraldehyde reductase. <i>FEMS Yeast Research</i> , 2007, 7, 84-92. | 1.1 | 35 |
| 5094 | Involvement of Rho-type GTPase in control of cell size in <i>Saccharomyces cerevisiae</i> . <i>FEMS Yeast Research</i> , 2007, 7, 569-578. | 1.1 | 12 |
| 5095 | Yeast expression of the <i>Tuber borchii</i> fruiting body specific protein, TBF-1: identification of a noncanonical signal peptide. <i>FEMS Microbiology Letters</i> , 2007, 272, 114-119. | 0.7 | 4 |
| 5096 | Molecular cloning and characterization of methylenedioxy bridge-forming enzymes involved in stylopine biosynthesis in <i>Eschscholzia californica</i> . <i>FEBS Journal</i> , 2007, 274, 1019-1035. | 2.2 | 104 |
| 5097 | Phylogenetic and functional analysis of the Cation Diffusion Facilitator (CDF) family: improved signature and prediction of substrate specificity. <i>BMC Genomics</i> , 2007, 8, 107. | 1.2 | 353 |
| 5098 | Over-expression of GSH1 gene and disruption of PEP4 gene in self-cloning industrial brewer's yeast. <i>International Journal of Food Microbiology</i> , 2007, 119, 192-199. | 2.1 | 30 |

| # | ARTICLE | IF | CITATIONS |
|------|---|-----|-----------|
| 5099 | Rad52 and Rad59 exhibit both overlapping and distinct functions. <i>DNA Repair</i> , 2007, 6, 27-37. | 1.3 | 34 |
| 5100 | Pol32 is required for Pol η -dependent translesion synthesis and prevents double-strand breaks at the replication fork. <i>Mutation Research - Fundamental and Molecular Mechanisms of Mutagenesis</i> , 2007, 625, 164-176. | 0.4 | 26 |
| 5101 | The role of ornithine aminotransferase in fruiting body formation of the mushroom <i>Agaricus bisporus</i> . <i>Mycological Research</i> , 2007, 111, 909-918. | 2.5 | 17 |
| 5102 | Expression of recombinant actin 5C from <i>Drosophila</i> in the methylotrophic yeast <i>Pichia pastoris</i> . <i>Cell and Tissue Biology</i> , 2007, 1, 248-258. | 0.2 | 0 |
| 5103 | 3 Yeast Transformation. <i>Methods in Microbiology</i> , 2007, , 45-54. | 0.4 | 4 |
| 5104 | Surface displayed expression of a neutralizing epitope of spike protein from a Korean strain of porcine epidemic diarrhea virus. <i>Biotechnology and Bioprocess Engineering</i> , 2007, 12, 690-695. | 1.4 | 14 |
| 5105 | Differences in osmotolerant and cell-wall properties of twozygosaccharomyces rouxii strains. <i>Folia Microbiologica</i> , 2007, 52, 241-245. | 1.1 | 11 |
| 5106 | Protective role of mitochondrial superoxide dismutase against high osmolarity, heat and metalloid stress in <i>Saccharomyces cerevisiae</i> . <i>Folia Microbiologica</i> , 2007, 52, 120-126. | 1.1 | 12 |
| 5107 | Construction of proteinase A deficient transformant of industrial brewing yeast. <i>European Food Research and Technology</i> , 2007, 225, 831-835. | 1.6 | 14 |
| 5108 | Detection of estrogenic activity in herbal teas by in vitro reporter assays. <i>European Food Research and Technology</i> , 2007, 225, 913-920. | 1.6 | 4 |
| 5109 | Construction of a novel synergistic system for production and recovery of secreted recombinant proteins by the cell surface engineering. <i>Applied Microbiology and Biotechnology</i> , 2007, 75, 821-828. | 1.7 | 32 |
| 5110 | Development of a yeast biosensorâ€“biocatalyst for the detection and biodegradation of the organophosphate paraoxon. <i>Applied Microbiology and Biotechnology</i> , 2007, 76, 1383-1394. | 1.7 | 21 |
| 5111 | Six new amino acid-auxotrophic markers for targeted gene integration and disruption in fission yeast. <i>Current Genetics</i> , 2007, 52, 97-105. | 0.8 | 28 |
| 5112 | Arc1p is required for cytoplasmic confinement of synthetases and tRNA. <i>Molecular and Cellular Biochemistry</i> , 2007, 300, 47-59. | 1.4 | 16 |
| 5113 | Glucose signaling controls the transcription of retrotransposon Ty2â€“917 in <i>Saccharomyces cerevisiae</i> . <i>Virus Genes</i> , 2007, 33, 279-85. | 0.7 | 12 |
| 5114 | Glucose signaling controls the transcription of retrotransposon Ty2â€“917 in <i>Saccharomyces cerevisiae</i> . <i>Virus Genes</i> , 2007, 35, 713-717. | 0.7 | 2 |
| 5115 | An Na ⁺ /H ⁺ antiporter gene from wheat plays an important role in stress tolerance. <i>Journal of Biosciences</i> , 2007, 32, 1153-1161. | 0.5 | 43 |
| 5116 | Effect of yeast pretreatment on the characteristics of yeast-modified electrodes as mediated amperometric biosensors for lactic acid. <i>Bioelectrochemistry</i> , 2008, 74, 188-194. | 2.4 | 21 |

| # | ARTICLE | IF | CITATIONS |
|------|---|-----|-----------|
| 5117 | Expression and function of two dehydrins under environmental stresses in <i>Brassica juncea</i> L. <i>Molecular Breeding</i> , 2008, 21, 431-438. | 1.0 | 26 |
| 5118 | Robust expression in yeast cells of a reporter gene driven by rumen protozoal promoter sequences. <i>World Journal of Microbiology and Biotechnology</i> , 2008, 24, 1529-1537. | 1.7 | 0 |
| 5119 | The yeast <i>Kluyveromyces marxianus</i> and its biotechnological potential. <i>Applied Microbiology and Biotechnology</i> , 2008, 79, 339-354. | 1.7 | 440 |
| 5120 | Efficient secretion of <i>Bacillus subtilis</i> lipase A in <i>Saccharomyces cerevisiae</i> by translational fusion to the Pir4 cell wall protein. <i>Applied Microbiology and Biotechnology</i> , 2008, 80, 437-445. | 1.7 | 25 |
| 5121 | Development of a yeast protein fragment complementation assay (PCA) system using dihydrofolate reductase (DHFR) with specific additives. <i>Applied Microbiology and Biotechnology</i> , 2008, 80, 735-743. | 1.7 | 16 |
| 5122 | Transformation of Industrialized Strain <i>Candida glycerinogenes</i> with Resistant Gene zeocin via <i>Agrobacterium tumefaciens</i> . <i>Current Microbiology</i> , 2008, 57, 12-17. | 1.0 | 3 |
| 5123 | Transcriptional regulation of <i>Saccharomyces cerevisiae</i> CYS3 encoding cystathionine β -lyase. <i>Current Genetics</i> , 2008, 53, 225-234. | 0.8 | 12 |
| 5124 | A potentially general method for the in vivo selection of inhibitory peptides targeted at a specific protein using yeast. <i>Current Genetics</i> , 2008, 53, 373-379. | 0.8 | 2 |
| 5125 | Heterologous expression of cholera toxin B subunit in <i>Saccharomyces cerevisiae</i> . <i>Biotechnology and Bioprocess Engineering</i> , 2008, 13, 598-605. | 1.4 | 4 |
| 5126 | CDKA and CDKB kinases from <i>Chlamydomonas reinhardtii</i> are able to complement <i>cdc28</i> temperature-sensitive mutants of <i>Saccharomyces cerevisiae</i> . <i>Protoplasma</i> , 2008, 232, 183-191. | 1.0 | 12 |
| 5127 | Heterologous expression of <i>Fusarium oxysporum</i> tomatinase in <i>Saccharomyces cerevisiae</i> increases its resistance to saponins and improves ethanol production during the fermentation of <i>Agave tequilana</i> Weber var. azul and <i>Agave salmiana</i> must. <i>Antonie Van Leeuwenhoek</i> , 2008, 93, 259-266. | 0.7 | 13 |
| 5128 | Cloning and characterization of three genes encoding Qb-SNARE proteins in rice. <i>Molecular Genetics and Genomics</i> , 2008, 279, 291-301. | 1.0 | 20 |
| 5129 | A novel mutant of the Sup35 protein of <i>Saccharomyces cerevisiae</i> defective in translation termination and in GTPase activity still supports cell viability. <i>BMC Molecular Biology</i> , 2008, 9, 22. | 3.0 | 9 |
| 5130 | The FF domains of yeast U1 snRNP protein Prp40 mediate interactions with Luc7 and Snu71. <i>BMC Biochemistry</i> , 2008, 9, 29. | 4.4 | 40 |
| 5131 | Low-copy episomal vector pFY20 and high-saturation coverage genomic libraries for the fission yeast <i>Schizosaccharomyces pombe</i> . <i>Yeast</i> , 2008, 25, 643-650. | 0.8 | 8 |
| 5132 | A small molecule-directed approach to control protein localization and function. <i>Yeast</i> , 2008, 25, 577-594. | 0.8 | 31 |
| 5133 | Enhanced arsenic accumulation by engineered yeast cells expressing <i>Arabidopsis thaliana</i> phytochelatin synthase. <i>Biotechnology and Bioengineering</i> , 2008, 99, 333-340. | 1.7 | 47 |
| 5134 | Human RhoA/RhoGDI complex expressed in yeast: Gtp exchange is sufficient for translocation of RhoA to liposomes. <i>Protein Science</i> , 2000, 9, 376-386. | 3.1 | 14 |

| # | ARTICLE | IF | CITATIONS |
|------|--|------|-----------|
| 5135 | Heterochromatic DNA repeats in <i>Drosophila</i> and unusual gene silencing in yeast cells. <i>Russian Journal of Genetics</i> , 2008, 44, 651-658. | 0.2 | 0 |
| 5136 | Posttranscriptional regulation of <i>FLO11</i> upon amino acid starvation in <i>Saccharomyces cerevisiae</i> . <i>FEMS Yeast Research</i> , 2008, 8, 225-236. | 1.1 | 12 |
| 5137 | Carbonyl cyanide m-chlorophenylhydrazone induced calcium signaling and activation of plasma membrane H ⁺ -ATPase in the yeast <i>Saccharomyces cerevisiae</i> . <i>FEMS Yeast Research</i> , 2008, 8, 622-630. | 1.1 | 28 |
| 5138 | Overexpressed ribosomal proteins suppress defective chaperonins in <i>Saccharomyces cerevisiae</i> . <i>FEMS Yeast Research</i> , 2008, 8, 1236-1244. | 1.1 | 19 |
| 5139 | Human p53 induces cell death and downregulates thioredoxin expression in <i>Saccharomyces cerevisiae</i> . <i>FEMS Yeast Research</i> , 2008, 8, 1254-1262. | 1.1 | 28 |
| 5140 | Mutants of <i>Saccharomyces cerevisiae</i> deficient in acyl-CoA synthetases secrete fatty acids due to interrupted fatty acid recycling. <i>FEBS Journal</i> , 2008, 275, 2765-2778. | 2.2 | 110 |
| 5141 | Comparison of a <i>coq7</i> deletion mutant with other respiration-defective mutants in fission yeast. <i>FEBS Journal</i> , 2008, 275, 5309-5324. | 2.2 | 33 |
| 5142 | The BTG2 protein is a general activator of mRNA deadenylation. <i>EMBO Journal</i> , 2008, 27, 1039-1048. | 3.5 | 100 |
| 5143 | Rga2 is a Rho2 GAP that regulates morphogenesis and cell integrity in <i>S. pombe</i> . <i>Molecular Microbiology</i> , 2008, 70, 867-881. | 1.2 | 55 |
| 5144 | Expression of the <i>AAC2</i> Gene Encoding the Major Mitochondrial ADP/ATP Carrier in <i>Saccharomyces cerevisiae</i> is Controlled at the Transcriptional Level by Oxygen, Heme and HAP2 Factor. <i>FEBS Journal</i> , 1995, 229, 651-657. | 0.2 | 7 |
| 5145 | Protein Phosphatase 2B of <i>Saccharomyces Cerevisiae</i> is Required for Tolerance to Manganese, in Blocking the Entry of ions into the Cells. <i>FEBS Journal</i> , 1995, 232, 712-717. | 0.2 | 27 |
| 5146 | Interaction between fortilin and transforming growth factor- β stimulated clone-22 (TSC-22) prevents apoptosis via the destabilization of TSC-22. <i>FEBS Letters</i> , 2008, 582, 1210-1218. | 1.3 | 27 |
| 5147 | The paclitaxel site in tubulin probed by site-directed mutagenesis of <i>Saccharomyces cerevisiae</i> β -tubulin. <i>FEBS Letters</i> , 2008, 582, 2467-2470. | 1.3 | 12 |
| 5148 | Mutational study of the catalytic tetrad of DNA topoisomerase IB from the hemoflagellate <i>Leishmania donovani</i> : Role of Asp-353 and Asn-221 in camptothecin resistance. <i>Biochemical Pharmacology</i> , 2008, 76, 608-619. | 2.0 | 15 |
| 5149 | An efficient genetic transformation method for glycerol producer <i>Candida glycerinogenes</i> . <i>Microbiological Research</i> , 2008, 163, 531-537. | 2.5 | 18 |
| 5150 | <i>Kluyveromyces lactis</i> β -galactosidase crystallization using full-factorial experimental design. <i>Journal of Molecular Catalysis B: Enzymatic</i> , 2008, 52-53, 178-182. | 1.8 | 8 |
| 5151 | Expression of <i>Brugmansia candida</i> Hyoscyamine 6 β -Hydroxylase gene in <i>Saccharomyces cerevisiae</i> and its potential use as biocatalyst. <i>Microbial Cell Factories</i> , 2008, 7, 17. | 1.9 | 29 |
| 5152 | Membrane transporters and protein traffic networks differentially affecting metal tolerance: a genomic phenotyping study in yeast. <i>Genome Biology</i> , 2008, 9, R67. | 13.9 | 97 |

| # | ARTICLE | IF | CITATIONS |
|------|---|------|-----------|
| 5153 | Enzymatic activities of novel mutant endoglucanases carrying sequential active sites. <i>International Journal of Biological Macromolecules</i> , 2008, 43, 226-231. | 3.6 | 13 |
| 5154 | The Catalytic Subunit of Shiga-like Toxin 1 Interacts with Ribosomal Stalk Proteins and is Inhibited by Their Conserved C-Terminal Domain. <i>Journal of Molecular Biology</i> , 2008, 378, 375-386. | 2.0 | 69 |
| 5155 | The Rat1p 5' to 3' Exonuclease Degrades Telomeric Repeat-Containing RNA and Promotes Telomere Elongation in <i>Saccharomyces cerevisiae</i> . <i>Molecular Cell</i> , 2008, 32, 465-477. | 4.5 | 274 |
| 5156 | Recombinant human fibrinogen expressed in the yeast <i>Pichia pastoris</i> was assembled and biologically active. <i>Protein Expression and Purification</i> , 2008, 59, 289-296. | 0.6 | 18 |
| 5157 | Visualized investigation of yeast transformation induced with Li ⁺ and polyethylene glycol. <i>Talanta</i> , 2008, 77, 262-268. | 2.9 | 24 |
| 5158 | Characterization and regulation of PiDur3, a permease involved in the acquisition of urea by the ectomycorrhizal fungus <i>Paxillus involutus</i> . <i>Fungal Genetics and Biology</i> , 2008, 45, 912-921. | 0.9 | 31 |
| 5159 | UvsF, the large subunit of replication factor C in <i>Aspergillus nidulans</i> , is essential for DNA replication, functions in UV repair and is upregulated in response to MMS-induced DNA damage. <i>Fungal Genetics and Biology</i> , 2008, 45, 1227-1234. | 0.9 | 3 |
| 5160 | Regulatory elements in the KlHEM1 promoter. <i>Biochimica Et Biophysica Acta - Gene Regulatory Mechanisms</i> , 2008, 1779, 128-133. | 0.9 | 9 |
| 5161 | Adaptation to oxidative stress induced by polyunsaturated fatty acids in yeast. <i>Biochimica Et Biophysica Acta - Molecular and Cell Biology of Lipids</i> , 2008, 1781, 283-287. | 1.2 | 48 |
| 5162 | Molecular characterization reveals that YMR278w encoded protein is environmental stress response homologue of <i>Saccharomyces cerevisiae</i> PGM2. <i>Biochemical and Biophysical Research Communications</i> , 2008, 366, 340-345. | 1.0 | 8 |
| 5163 | Incompatibility of Nuclear and Mitochondrial Genomes Causes Hybrid Sterility between Two Yeast Species. <i>Cell</i> , 2008, 135, 1065-1073. | 13.5 | 326 |
| 5164 | Direct ethanol production from barley β -glucan by sake yeast displaying <i>Aspergillus oryzae</i> β -glucosidase and endoglucanase. <i>Journal of Bioscience and Bioengineering</i> , 2008, 105, 622-627. | 1.1 | 88 |
| 5165 | A Recombinant <i>Saccharomyces cerevisiae</i> Strain Overproducing Mannoproteins Stabilizes Wine against Protein Haze. <i>Applied and Environmental Microbiology</i> , 2008, 74, 5533-5540. | 1.4 | 78 |
| 5166 | Microbial production of plant benzyloquinoline alkaloids. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 2008, 105, 7393-7398. | 3.3 | 307 |
| 5167 | Expression of the Novel Wheat Gene TM20 Confers Enhanced Cadmium Tolerance to Bakers' Yeast. <i>Journal of Biological Chemistry</i> , 2008, 283, 15893-15902. | 1.6 | 42 |
| 5168 | Efficient and Direct Fermentation of Starch to Ethanol by Sake Yeast Strains Displaying Fungal Glucoamylases. <i>Bioscience, Biotechnology and Biochemistry</i> , 2008, 72, 1376-1379. | 0.6 | 20 |
| 5169 | Synthesis and Accumulation of Cyanophycin in Transgenic Strains of <i>Saccharomyces cerevisiae</i> . <i>Applied and Environmental Microbiology</i> , 2008, 74, 3410-3418. | 1.4 | 50 |
| 5170 | Impaired ERAD and ER stress are early and specific events in polyglutamine toxicity. <i>Genes and Development</i> , 2008, 22, 3308-3319. | 2.7 | 265 |

| # | ARTICLE | IF | CITATIONS |
|------|--|-----|-----------|
| 5171 | Yeast ribosomal protein L10 helps coordinate tRNA movement through the large subunit. <i>Nucleic Acids Research</i> , 2008, 36, 6187-6198. | 6.5 | 30 |
| 5172 | Distinct acto/myosin-I structures associate with endocytic profiles at the plasma membrane. <i>Journal of Cell Biology</i> , 2008, 180, 1219-1232. | 2.3 | 134 |
| 5173 | Structure/function analysis of yeast ribosomal protein L2. <i>Nucleic Acids Research</i> , 2008, 36, 1826-1835. | 6.5 | 25 |
| 5174 | Regulation of the <i>Saccharomyces cerevisiae</i> CK1-encoded Choline Kinase by Zinc Depletion. <i>Journal of Biological Chemistry</i> , 2008, 283, 10079-10088. | 1.6 | 20 |
| 5175 | Basic-Zipper-Type Transcription Factor FlbB Controls Asexual Development in <i>Aspergillus nidulans</i> . <i>Eukaryotic Cell</i> , 2008, 7, 38-48. | 3.4 | 97 |
| 5176 | Unconventional Genomic Architecture in the Budding Yeast <i>Saccharomyces cerevisiae</i> Masks the Nested Antisense Gene <i>NAG1</i> . <i>Eukaryotic Cell</i> , 2008, 7, 1289-1298. | 3.4 | 17 |
| 5177 | Localization of autophagy-related proteins in yeast using a versatile plasmid-based resource of fluorescent protein fusions. <i>Autophagy</i> , 2008, 4, 792-800. | 4.3 | 11 |
| 5178 | Biochemical and functional characterization of the actin-binding activity of the B subunit of yeast vacuolar H ⁺ -ATPase. <i>Journal of Experimental Biology</i> , 2008, 211, 1102-1108. | 0.8 | 22 |
| 5179 | Oxygen-regulated isoforms of cytochrome <i>c</i> oxidase have differential effects on its nitric oxide production and on hypoxic signaling. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 2008, 105, 8203-8208. | 3.3 | 113 |
| 5180 | A cyclophilin links redox and light signals to cysteine biosynthesis and stress responses in chloroplasts. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 2008, 105, 16386-16391. | 3.3 | 138 |
| 5181 | Yeast Antizyme Mediates Degradation of Yeast Ornithine Decarboxylase by Yeast but Not by Mammalian Proteasome. <i>Journal of Biological Chemistry</i> , 2008, 283, 4528-4534. | 1.6 | 19 |
| 5182 | Characterization of the Yeast DGK1-encoded CTP-dependent Diacylglycerol Kinase. <i>Journal of Biological Chemistry</i> , 2008, 283, 20443-20453. | 1.6 | 82 |
| 5183 | Depurination of Brome Mosaic Virus RNA3 in Vivo Results in Translation-dependent Accelerated Degradation of the Viral RNA. <i>Journal of Biological Chemistry</i> , 2008, 283, 32218-32228. | 1.6 | 42 |
| 5184 | Degradation of a Cytosolic Protein Requires Endoplasmic Reticulum-associated Degradation Machinery. <i>Journal of Biological Chemistry</i> , 2008, 283, 32302-32316. | 1.6 | 125 |
| 5185 | The Gluconeogenic Enzyme Fructose-1,6-Bisphosphatase Is Dispensable for Growth of the Yeast <i>Yarrowia lipolytica</i> in Gluconeogenic Substrates. <i>Eukaryotic Cell</i> , 2008, 7, 1742-1749. | 3.4 | 21 |
| 5186 | Specialization of the paralogue LYS21 determines lysine biosynthesis under respiratory metabolism in <i>Saccharomyces cerevisiae</i> . <i>Microbiology (United Kingdom)</i> , 2008, 154, 1656-1667. | 0.7 | 45 |
| 5187 | Long-Range Communication between the Silencers of <i>HMR</i> . <i>Molecular and Cellular Biology</i> , 2008, 28, 1924-1935. | 1.1 | 58 |
| 5188 | The Pentatricopeptide Repeats Present in Pet309 Are Necessary for Translation but Not for Stability of the Mitochondrial COX1 mRNA in Yeast. <i>Journal of Biological Chemistry</i> , 2008, 283, 1472-1479. | 1.6 | 79 |

| # | ARTICLE | IF | CITATIONS |
|------|--|-----|-----------|
| 5189 | Requirement for the Budding Yeast Polo Kinase Cdc5 in Proper Microtubule Growth and Dynamics. <i>Eukaryotic Cell</i> , 2008, 7, 444-453. | 3.4 | 35 |
| 5190 | Cellular and molecular effects of nonreciprocal chromosome translocations in <i>Saccharomyces cerevisiae</i> . <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 2008, 105, 9703-9708. | 3.3 | 15 |
| 5191 | Immunocytochemical localization of <i>Pisum sativum</i> TRXs f and m in non-photosynthetic tissues. <i>Journal of Experimental Botany</i> , 2008, 59, 1267-1277. | 2.4 | 30 |
| 5192 | Evidence for the Bifunctional Nature of Mitochondrial Phosphatidylserine Decarboxylase: Role in Pdr3-Dependent Retrograde Regulation of <i>PDR5</i> Expression. <i>Molecular and Cellular Biology</i> , 2008, 28, 5851-5864. | 1.1 | 43 |
| 5193 | <i>Trans</i> -Golgi Network and Endosome Dynamics Connect Ceramide Homeostasis with Regulation of the Unfolded Protein Response and TOR Signaling in Yeast. <i>Molecular Biology of the Cell</i> , 2008, 19, 4785-4803. | 0.9 | 47 |
| 5194 | Tissue inhibitor of metalloproteinases-3 interacts with angiotensin II type 2 receptor and additively inhibits angiogenesis. <i>Cardiovascular Research</i> , 2008, 79, 150-160. | 1.8 | 49 |
| 5195 | Sister Chromatid Cohesion Role for <i>CDC28</i> -CDK in <i>Saccharomyces cerevisiae</i> . <i>Genetics</i> , 2008, 180, 7-16. | 1.2 | 10 |
| 5196 | Distinct regions of ATF/CREB proteins Atf1 and Pcr1 control recombination hotspot <i>ade6-M26</i> and the osmotic stress response. <i>Nucleic Acids Research</i> , 2008, 36, 2838-2851. | 6.5 | 40 |
| 5197 | Assembly of a Fab1 Phosphoinositide Kinase Signaling Complex Requires the Fig4 Phosphoinositide Phosphatase. <i>Molecular Biology of the Cell</i> , 2008, 19, 4273-4286. | 0.9 | 120 |
| 5198 | rRNA mutants in the yeast peptidyltransferase center reveal allosteric information networks and mechanisms of drug resistance. <i>Nucleic Acids Research</i> , 2008, 36, 1497-1507. | 6.5 | 38 |
| 5199 | <i>Arabidopsis</i> <i>UEV1D</i> Promotes Lysine-63-Linked Polyubiquitination and Is Involved in DNA Damage Response. <i>Plant Cell</i> , 2008, 20, 213-227. | 3.1 | 79 |
| 5200 | Sorting Signals within the <i>Saccharomyces cerevisiae</i> Sporulation-Specific Dityrosine Transporter, <i>Dtr1p</i> , C Terminus Promote Golgi-to-Prospore Membrane Transport. <i>Eukaryotic Cell</i> , 2008, 7, 1674-1684. | 3.4 | 8 |
| 5201 | G1/S Cyclin-dependent Kinase Regulates Small GTPase Rho1p through Phosphorylation of RhoGEF Tus1p in <i>Saccharomyces cerevisiae</i> . <i>Molecular Biology of the Cell</i> , 2008, 19, 1763-1771. | 0.9 | 47 |
| 5202 | Role of the <i>Saccharomyces cerevisiae</i> <i>Rad51</i> Paralogs in Sister Chromatid Recombination. <i>Genetics</i> , 2008, 178, 113-126. | 1.2 | 63 |
| 5203 | Molecular Cloning and Characterization of CYP80G2, a Cytochrome P450 That Catalyzes an Intramolecular C-C Phenol Coupling of (S)-Reticuline in Magnoflorine Biosynthesis, from Cultured <i>Coptis japonica</i> Cells. <i>Journal of Biological Chemistry</i> , 2008, 283, 8810-8821. | 1.6 | 130 |
| 5204 | Deletion of Many Yeast Introns Reveals a Minority of Genes that Require Splicing for Function. <i>Molecular Biology of the Cell</i> , 2008, 19, 1932-1941. | 0.9 | 99 |
| 5205 | Functional Characterization of the N-terminal Domain of Subunit H (<i>Vma13p</i>) of the Yeast Vacuolar ATPase. <i>Journal of Biological Chemistry</i> , 2008, 283, 29099-29108. | 1.6 | 11 |
| 5206 | Genome-wide screen of <i>Saccharomyces cerevisiae</i> null allele strains identifies genes involved in selenomethionine resistance. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 2008, 105, 17682-17687. | 3.3 | 19 |

| # | ARTICLE | IF | CITATIONS |
|------|---|-----|-----------|
| 5207 | Functional Analysis of Arabidopsis Postprenylation CaaX Processing Enzymes and Their Function in Subcellular Protein Targeting. <i>Plant Physiology</i> , 2008, 148, 119-131. | 2.3 | 50 |
| 5208 | A yeast TDP-43 proteinopathy model: Exploring the molecular determinants of TDP-43 aggregation and cellular toxicity. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 2008, 105, 6439-6444. | 3.3 | 375 |
| 5209 | A Small Stem Loop Element Directs Internal Initiation of the URE2 Internal Ribosome Entry Site in <i>Saccharomyces cerevisiae</i> . <i>Journal of Biological Chemistry</i> , 2008, 283, 19011-19025. | 1.6 | 23 |
| 5210 | RXR heterodimerization allosterically activates LXR binding to the second NR box of activating signal co-integrator-2. <i>Biochemical Journal</i> , 2008, 410, 319-330. | 1.7 | 17 |
| 5211 | Hydrophobins in ectomycorrhizas: heterologous transcription of the <i>Pisolithus HydPt-1</i> gene in yeast and <i>Hebeloma cylindrosporum</i> . <i>European Journal of Histochemistry</i> , 2009, 46, 23. | 0.6 | 8 |
| 5212 | Phosphorylation-Independent Regulation of Atf1-Promoted Meiotic Recombination by Stress-Activated, p38 Kinase Spc1 of Fission Yeast. <i>PLoS ONE</i> , 2009, 4, e5533. | 1.1 | 15 |
| 5213 | Multisite Phosphorylation of the Guanine Nucleotide Exchange Factor Cdc24 during Yeast Cell Polarization. <i>PLoS ONE</i> , 2009, 4, e6563. | 1.1 | 33 |
| 5214 | Transcription Independent Insulation at TFIIIC-Dependent Insulators. <i>Genetics</i> , 2009, 183, 131-148. | 1.2 | 56 |
| 5215 | Evidence that U2/U6 helix I promotes both catalytic steps of pre-mRNA splicing and rearranges in between these steps. <i>Rna</i> , 2009, 15, 1386-1397. | 1.6 | 72 |
| 5216 | Human fungal pathogen <i>Candida albicans</i> in the postgenomic era: an overview. <i>Expert Review of Anti-Infective Therapy</i> , 2009, 7, 121-134. | 2.0 | 11 |
| 5217 | Characterization of the functional role of nucleotides within the URE2 IRES element and the requirements for eIF2A-mediated repression. <i>Rna</i> , 2009, 15, 2264-2277. | 1.6 | 26 |
| 5218 | Phosphatidylinositol-4,5-Bisphosphate and Phospholipase D-Generated Phosphatidic Acid Specify SNARE-Mediated Vesicle Fusion for Prospore Membrane Formation. <i>Eukaryotic Cell</i> , 2009, 8, 1094-1105. | 3.4 | 16 |
| 5219 | Stress-induced Ceramide-activated Protein Phosphatase Can Compensate for Loss of Amphiphysin-like Activity In <i>Saccharomyces cerevisiae</i> and Functions to Reinitiate Endocytosis. <i>Journal of Biological Chemistry</i> , 2009, 284, 11930-11941. | 1.6 | 14 |
| 5220 | Degradation of <i>Saccharomyces cerevisiae</i> Transcription Factor Gcn4 Requires a C-Terminal Nuclear Localization Signal in the Cyclin Pcl5. <i>Eukaryotic Cell</i> , 2009, 8, 496-510. | 3.4 | 9 |
| 5221 | Î²-Glucanase Eng2 Is Required for <i>Ascus</i> Wall Endolysis after Sporulation in the Fission Yeast <i>Schizosaccharomyces pombe</i> . <i>Eukaryotic Cell</i> , 2009, 8, 1278-1286. | 3.4 | 27 |
| 5222 | Uridine-Ribohydrolase Is a Key Regulator in the Uridine Degradation Pathway of <i>Arabidopsis</i> . <i>Plant Cell</i> , 2009, 21, 876-891. | 3.1 | 77 |
| 5223 | Genome-wide Analysis of AP-3-dependent Protein Transport in Yeast. <i>Molecular Biology of the Cell</i> , 2009, 20, 1592-1604. | 0.9 | 43 |
| 5224 | Molecular dissection of translation termination mechanism identifies two new critical regions in eRF1. <i>Nucleic Acids Research</i> , 2009, 37, 1789-1798. | 6.5 | 15 |

| # | ARTICLE | IF | CITATIONS |
|------|--|-----|-----------|
| 5225 | ER membrane-bending proteins are necessary for de novo nuclear pore formation. <i>Journal of Cell Biology</i> , 2009, 184, 659-675. | 2.3 | 137 |
| 5226 | Coordination of Eukaryotic Translation Elongation Factor 1A (eEF1A) Function in Actin Organization and Translation Elongation by the Guanine Nucleotide Exchange Factor eEF1B \pm . <i>Journal of Biological Chemistry</i> , 2009, 284, 4739-4747. | 1.6 | 45 |
| 5227 | Dug1p Is a Cys-Gly Peptidase of the \hat{I}^3 -Glutamyl Cycle of <i>Saccharomyces cerevisiae</i> and Represents a Novel Family of Cys-Gly Peptidases. <i>Journal of Biological Chemistry</i> , 2009, 284, 14493-14502. | 1.6 | 66 |
| 5228 | Accuracy modulating mutations of the ribosomal protein S4-S5 interface do not necessarily destabilize the rps4-rps5 protein-protein interaction. <i>Rna</i> , 2009, 15, 1100-1109. | 1.6 | 18 |
| 5229 | Involvement of SMRT Corepressor in Transcriptional Repression by the Vitamin D Receptor. <i>Molecular Endocrinology</i> , 2009, 23, 251-264. | 3.7 | 36 |
| 5230 | Metabolic Engineering of <i>Saccharomyces cerevisiae</i> for Production of Novel Cyanophycins with an Extended Range of Constituent Amino Acids. <i>Applied and Environmental Microbiology</i> , 2009, 75, 3437-3446. | 1.4 | 49 |
| 5231 | Sgt1 Dimerization Is Negatively Regulated by Protein Kinase CK2-mediated Phosphorylation at Ser361. <i>Journal of Biological Chemistry</i> , 2009, 284, 18692-18698. | 1.6 | 27 |
| 5232 | Cu, Zn Superoxide Dismutase and NADP(H) Homeostasis Are Required for Tolerance of Endoplasmic Reticulum Stress in <i>Saccharomyces cerevisiae</i> . <i>Molecular Biology of the Cell</i> , 2009, 20, 1493-1508. | 0.9 | 57 |
| 5233 | Analysis of Quality Control Substrates in Distinct Cellular Compartments Reveals a Unique Role for Rpn4p in Tolerating Misfolded Membrane Proteins. <i>Molecular Biology of the Cell</i> , 2009, 20, 1006-1019. | 0.9 | 58 |
| 5234 | Rex1p deficiency leads to accumulation of precursor initiator tRNAMet and polyadenylation of substrate RNAs in <i>Saccharomyces cerevisiae</i> . <i>Nucleic Acids Research</i> , 2009, 37, 298-308. | 6.5 | 63 |
| 5235 | Sgt1 Dimerization Is Required for Yeast Kinetochores Assembly. <i>Journal of Biological Chemistry</i> , 2009, 284, 3586-3592. | 1.6 | 26 |
| 5236 | Altering Sphingolipid Metabolism in <i>Saccharomyces cerevisiae</i> Cells Lacking the Amphiphysin Ortholog Rvs161 Reinitiates Sugar Transporter Endocytosis. <i>Eukaryotic Cell</i> , 2009, 8, 779-789. | 3.4 | 6 |
| 5237 | Functional Redundancy of Yeast Proteins Reh1 and Rei1 in Cytoplasmic 60S Subunit Maturation. <i>Molecular and Cellular Biology</i> , 2009, 29, 4014-4023. | 1.1 | 27 |
| 5238 | A protein kinase-phosphatase pair interacts with an ion channel to regulate ABA signaling in plant guard cells. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 2009, 106, 21419-21424. | 3.3 | 565 |
| 5239 | Dynamic Compartmentalization of Base Excision Repair Proteins in Response to Nuclear and Mitochondrial Oxidative Stress. <i>Molecular and Cellular Biology</i> , 2009, 29, 794-807. | 1.1 | 40 |
| 5240 | Multidrug and Toxic Compound Extrusion-Type Transporters Implicated in Vacuolar Sequestration of Nicotine in Tobacco Roots. <i>Plant Physiology</i> , 2009, 149, 708-718. | 2.3 | 184 |
| 5241 | The Yeast HtrA Orthologue Ynm3 Is a Protease with Chaperone Activity that Aids Survival Under Heat Stress. <i>Molecular Biology of the Cell</i> , 2009, 20, 68-77. | 0.9 | 32 |
| 5242 | Aberrant Double-Strand Break Repair Resulting in Half Crossovers in Mutants Defective for Rad51 or the DNA Polymerase \hat{I} Complex. <i>Molecular and Cellular Biology</i> , 2009, 29, 1432-1441. | 1.1 | 82 |

| # | ARTICLE | IF | CITATIONS |
|------|--|-----|-----------|
| 5243 | She3p Possesses a Novel Activity Required for <i>ASH1</i> mRNA Localization in <i>Saccharomyces cerevisiae</i> . <i>Eukaryotic Cell</i> , 2009, 8, 1072-1083. | 3.4 | 18 |
| 5244 | The Mammalian Hsp40 ERdj3 Requires Its Hsp70 Interaction and Substrate-binding Properties to Complement Various Yeast Hsp40-dependent Functions. <i>Journal of Biological Chemistry</i> , 2009, 284, 32462-32471. | 1.6 | 19 |
| 5245 | Pob1 Participates in the Cdc42 Regulation of Fission Yeast Actin Cytoskeleton. <i>Molecular Biology of the Cell</i> , 2009, 20, 4390-4399. | 0.9 | 46 |
| 5246 | Probing the orientation of yeast VDAC1 in vivo. <i>FEBS Letters</i> , 2009, 583, 739-742. | 1.3 | 29 |
| 5247 | Using promoter replacement and selection for loss of heterozygosity to generate an industrially applicable sake yeast strain that homozygously overproduces isoamyl acetate. <i>Journal of Bioscience and Bioengineering</i> , 2009, 108, 359-364. | 1.1 | 14 |
| 5248 | The absence of a mitochondrial genome in rho0 yeast cells extends lifespan independently of retrograde regulation. <i>Experimental Gerontology</i> , 2009, 44, 390-397. | 1.2 | 33 |
| 5249 | <i>Dkk3</i> , downregulated in cervical cancer, functions as a negative regulator of β -catenin. <i>International Journal of Cancer</i> , 2009, 124, 287-297. | 2.3 | 136 |
| 5250 | Transcriptional and metabolic response in yeast <i>Saccharomyces cerevisiae</i> cells during polyethylene glycol-dependent transformation. <i>Journal of Basic Microbiology</i> , 2009, 49, 73-81. | 1.8 | 12 |
| 5251 | Construction of a recombinant autolytic wine yeast strain overexpressing the <i>csc1</i> allele. <i>Biotechnology Progress</i> , 2009, 25, 1598-1604. | 1.3 | 11 |
| 5252 | OsMSRA4.1 and OsMSRB1.1, two rice plastidial methionine sulfoxide reductases, are involved in abiotic stress responses. <i>Planta</i> , 2009, 230, 227-238. | 1.6 | 73 |
| 5253 | A fitness-based interferential genetics approach using hypertoxic/inactive gene alleles as references. <i>Molecular Genetics and Genomics</i> , 2009, 281, 437-445. | 1.0 | 1 |
| 5254 | DNA bridging of yeast chromosomes VIII leads to near-reciprocal translocation and loss of heterozygosity with minor cellular defects. <i>Chromosoma</i> , 2009, 118, 179-191. | 1.0 | 12 |
| 5255 | <i>Penicillium marneffeii</i> SKN7, a Novel Gene, could Complement the Hypersensitivity of <i>S. cerevisiae</i> <i>skn7</i> Disruptant Strain to Oxidative Stress. <i>Mycopathologia</i> , 2009, 168, 23-30. | 1.3 | 22 |
| 5256 | A Bipartite Recombinant Yeast System for the Identification of Subtype-Selective Estrogen Receptor Ligands. <i>Molecular Biotechnology</i> , 2009, 41, 53-62. | 1.3 | 10 |
| 5257 | Gene transfer and protein release of fission yeast by application of a high voltage electric pulse. <i>Analytical and Bioanalytical Chemistry</i> , 2009, 394, 13-16. | 1.9 | 22 |
| 5258 | The cation-efflux transporter BjCET2 mediates zinc and cadmium accumulation in <i>Brassica juncea</i> L. leaves. <i>Plant Cell Reports</i> , 2009, 28, 1235-1242. | 2.8 | 46 |
| 5259 | Cell-surface modification of non-GMO without chemical treatment by novel GMO-coupled and -separated cocultivation method. <i>Applied Microbiology and Biotechnology</i> , 2009, 82, 293-301. | 1.7 | 4 |
| 5260 | Efficient generation of recessive traits in diploid sake yeast by targeted gene disruption and loss of heterozygosity. <i>Applied Microbiology and Biotechnology</i> , 2009, 82, 387-395. | 1.7 | 23 |

| # | ARTICLE | IF | CITATIONS |
|------|--|-----|-----------|
| 5261 | Advances in molecular methods to alter chromosomes and genome in the yeast <i>Saccharomyces cerevisiae</i> . <i>Applied Microbiology and Biotechnology</i> , 2009, 84, 1045-1052. | 1.7 | 13 |
| 5262 | Dse1 may control cross talk between the pheromone and filamentation pathways in yeast. <i>Current Genetics</i> , 2009, 55, 611-621. | 0.8 | 5 |
| 5263 | Production of polyunsaturated fatty acids in yeast <i>Saccharomyces cerevisiae</i> and its relation to alkaline pH tolerance. <i>Yeast</i> , 2009, 26, 167-184. | 0.8 | 39 |
| 5264 | Selection of cell death-deficient p53 mutants in <i>Saccharomyces cerevisiae</i> . <i>Yeast</i> , 2009, 26, 441-450. | 0.8 | 7 |
| 5265 | Efficient production of L-lactic acid by Crabtree-negative yeast <i>Candida boidinii</i> . <i>Yeast</i> , 2009, 26, 485-496. | 0.8 | 40 |
| 5266 | Random and targeted gene integrations through the control of non-homologous end joining in the yeast <i>Kluyveromyces marxianus</i> . <i>Yeast</i> , 2010, 27, 29-39. | 0.8 | 90 |
| 5267 | A novel kanamycin/G418 resistance marker for direct selection of transformants in <i>Escherichia coli</i> and different yeast species. <i>Yeast</i> , 2010, 27, 189-195. | 0.8 | 21 |
| 5268 | cpFBPasell, a novel redox-independent chloroplastic isoform of fructose-1,6-bisphosphatase. <i>Plant, Cell and Environment</i> , 2009, 32, 811-827. | 2.8 | 22 |
| 5269 | AtHMA1 contributes to the detoxification of excess Zn(II) in <i>Arabidopsis</i> . <i>Plant Journal</i> , 2009, 58, 737-753. | 2.8 | 167 |
| 5270 | The yeast Shu complex couples error-free post-replication repair to homologous recombination. <i>Molecular Microbiology</i> , 2009, 73, 89-102. | 1.2 | 88 |
| 5271 | Expression of two human acyl-CoA:diacylglycerol acyltransferase isozymes in yeast and selectivity of microbial inhibitors toward the isozymes. <i>Journal of Antibiotics</i> , 2009, 62, 51-54. | 1.0 | 25 |
| 5272 | A yeast-based genetic screen for identification of pathogenic <i>Salmonella</i> proteins. <i>FEMS Microbiology Letters</i> , 2009, 296, 167-177. | 0.7 | 11 |
| 5273 | Evidence that the assembly of the yeast cytochrome bc ₁ complex involves the formation of a large core structure in the inner mitochondrial membrane. <i>FEBS Journal</i> , 2009, 276, 1900-1914. | 2.2 | 56 |
| 5274 | A large complex mediated by Moc1, Moc2 and Cpc2 regulates sexual differentiation in fission yeast. <i>FEBS Journal</i> , 2009, 276, 5076-5093. | 2.2 | 19 |
| 5275 | CAP2 enhances germination of transgenic tobacco seeds at high temperature and promotes heat stress tolerance in yeast. <i>FEBS Journal</i> , 2009, 276, 5252-5262. | 2.2 | 27 |
| 5276 | Recombinant industrial brewing yeast strains with ADH2 interruption using self-cloning GSH1 + CUP1 cassette. <i>FEMS Yeast Research</i> , 2009, 9, 574-581. | 1.1 | 28 |
| 5277 | Recombinant aryl hydrocarbon receptors for bioassay of aryl hydrocarbon receptor ligands in transgenic tobacco plants. <i>Plant Biotechnology Journal</i> , 2009, 7, 119-128. | 4.1 | 17 |
| 5278 | Establishment of a screening system for essential genes from the pathogenic yeast <i>Candida glabrata</i> : identification of a putative TEM1 homologue. <i>Letters in Applied Microbiology</i> , 2009, 49, 317-323. | 1.0 | 3 |

| # | ARTICLE | IF | CITATIONS |
|------|---|-----|-----------|
| 5279 | Development of a <i>Saccharomyces cerevisiae</i> strain for the production of 1,2-propanediol by gene manipulation. <i>Enzyme and Microbial Technology</i> , 2009, 45, 42-47. | 1.6 | 15 |
| 5280 | Analysis of CMP-sialic acid transporter-like proteins in plants. <i>Phytochemistry</i> , 2009, 70, 1973-1981. | 1.4 | 19 |
| 5281 | Advances in Gene Expression in Non-Conventional Yeasts. , 2009, , 369-403. | | 1 |
| 5282 | Yeast Genetics and Biotechnological Applications. , 2009, , 323-355. | | 3 |
| 5283 | Evidence That β -Synuclein Does Not Inhibit Phospholipase D. <i>Biochemistry</i> , 2009, 48, 1077-1083. | 1.2 | 31 |
| 5284 | The QTK Loop Is Essential for the Communication between the N-Terminal ATPase Domain and the Central Cleavage Ligation Region in Human Topoisomerase II α . <i>Biochemistry</i> , 2009, 48, 6508-6515. | 1.2 | 14 |
| 5285 | Molecular and Biochemical Characterization of a β -Fructofuranosidase from <i>Xanthophyllomyces dendrorhous</i> . <i>Applied and Environmental Microbiology</i> , 2009, 75, 1065-1073. | 1.4 | 87 |
| 5286 | A yeast assay based on the gilthead sea bream (teleost fish) estrogen receptor β for monitoring estrogen mimics. <i>Ecotoxicology and Environmental Safety</i> , 2009, 72, 1529-1537. | 2.9 | 11 |
| 5287 | Two-hybrid analysis identifies PSMD11, a non-ATPase subunit of the proteasome, as a novel interaction partner of AMP-activated protein kinase. <i>International Journal of Biochemistry and Cell Biology</i> , 2009, 41, 2431-2439. | 1.2 | 24 |
| 5288 | Determination of the core promoter regions of the <i>Saccharomyces cerevisiae</i> RPS3 gene. <i>Biochimica Et Biophysica Acta - Gene Regulatory Mechanisms</i> , 2009, 1789, 741-750. | 0.9 | 6 |
| 5289 | Cd ²⁺ , Mn ²⁺ , Ni ²⁺ and Se ²⁺ toxicity to <i>Saccharomyces cerevisiae</i> lacking YPK9p the orthologue of human ATP13A2. <i>Biochemical and Biophysical Research Communications</i> , 2009, 383, 198-202. | 1.0 | 110 |
| 5290 | Expression and characterization of jingzhaotoxin-34, a novel neurotoxin from the venom of the tarantula <i>Chilobrachys jingzhao</i> . <i>Peptides</i> , 2009, 30, 1042-1048. | 1.2 | 18 |
| 5291 | Calcineurin B-like proteins in rice can bind with calcium ion and associate with the Arabidopsis CIPK family members. <i>Plant Science</i> , 2009, 177, 577-583. | 1.7 | 6 |
| 5292 | Structural and Functional Divergence within the Dim1/KsgA Family of rRNA Methyltransferases. <i>Journal of Molecular Biology</i> , 2009, 391, 884-893. | 2.0 | 26 |
| 5293 | Basidiomycetous Yeasts: Current Status. , 2009, , 19-46. | | 6 |
| 5294 | <i>Pichia guilliermondii</i> . , 2009, , 113-134. | | 11 |
| 5296 | Visualization and Quantification of Three-Dimensional Distribution of Yeast in Bread Dough. <i>Bioscience, Biotechnology and Biochemistry</i> , 2009, 73, 1586-1590. | 0.6 | 2 |
| 5297 | Designed Recombinant Transcription Factor with Antibody-Variable Regions. <i>Analytical Chemistry</i> , 2009, 81, 10162-10166. | 3.2 | 3 |

| # | ARTICLE | IF | CITATIONS |
|------|--|-----|-----------|
| 5298 | Glucose Signaling Pathway and Growth Conditions Regulate Gene Expression in Retrotransposon Ty2. <i>Zeitschrift Fur Naturforschung - Section C Journal of Biosciences</i> , 2009, 64, 526-532. | 0.6 | 4 |
| 5299 | Regulatory factors controlling transcription of <i>Saccharomyces cerevisiae</i> IXR1 by oxygen levels: a model of transcriptional adaptation from aerobiosis to hypoxia implicating ROX1 and IXR1 cross-regulation. <i>Biochemical Journal</i> , 2010, 425, 235-243. | 1.7 | 20 |
| 5300 | Molecular design of yeast cell surface for adsorption and recovery of molybdenum, one of rare metals. <i>Applied Microbiology and Biotechnology</i> , 2010, 86, 641-648. | 1.7 | 60 |
| 5301 | Improvement of polyunsaturated fatty acids synthesis by the coexpression of CYB5 with desaturase genes in <i>Saccharomyces cerevisiae</i> . <i>Applied Microbiology and Biotechnology</i> , 2010, 87, 2185-2193. | 1.7 | 16 |
| 5302 | Activation of diacylglycerol acyltransferase expressed in <i>Saccharomyces cerevisiae</i> : overexpression of Dga1p lacking the N-terminal region in the <i>atsnf2</i> disruptant produces a significant increase in its enzyme activity. <i>Applied Microbiology and Biotechnology</i> , 2010, 88, 105-115. | 1.7 | 25 |
| 5303 | Identification and characterization of an acyl-CoA:diacylglycerol acyltransferase 2 (DGAT2) gene from the microalga <i>O. tauri</i> . <i>Plant Physiology and Biochemistry</i> , 2010, 48, 407-416. | 2.8 | 97 |
| 5304 | AtfA bZIP-type transcription factor regulates oxidative and osmotic stress responses in <i>Aspergillus nidulans</i> . <i>Molecular Genetics and Genomics</i> , 2010, 283, 289-303. | 1.0 | 78 |
| 5305 | Enhancement of menadione stress tolerance in yeast by accumulation of hypotaurine and taurine: co-expression of cDNA clones, from <i>Cyprinus carpio</i> , for cysteine dioxygenase and cysteine sulfinate decarboxylase in <i>Saccharomyces cerevisiae</i> . <i>Amino Acids</i> , 2010, 38, 1173-1183. | 1.2 | 11 |
| 5306 | Surface display of active lipase in <i>Saccharomyces cerevisiae</i> using Cwp2 as an anchor protein. <i>Biotechnology Letters</i> , 2010, 32, 255-260. | 1.1 | 30 |
| 5307 | Zebrafish Ubc13 is required for Lys63-linked polyubiquitination and DNA damage tolerance. <i>Molecular and Cellular Biochemistry</i> , 2010, 343, 173-182. | 1.4 | 11 |
| 5308 | Engineering of small sized DNAs by error-prone multiply-primed rolling circle amplification for introduction of random point mutations. <i>Journal of Molecular Catalysis B: Enzymatic</i> , 2010, 67, 92-97. | 1.8 | 3 |
| 5309 | Heterologous expression of glucose oxidase in the yeast <i>Kluyveromyces marxianus</i> . <i>Microbial Cell Factories</i> , 2010, 9, 4. | 1.9 | 40 |
| 5310 | Global expression studies in baker's yeast reveal target genes for the improvement of industrially-relevant traits: the cases of CAF16 and ORC2. <i>Microbial Cell Factories</i> , 2010, 9, 56. | 1.9 | 11 |
| 5311 | His-tagged tRNA ^{His} guanylyltransferase in bacteria. <i>FEBS Letters</i> , 2010, 584, 3567-3572. | 1.3 | 30 |
| 5312 | Enhancement of β -glucosidase activity on the cell-surface of sake yeast by disruption of SED1. <i>Journal of Bioscience and Bioengineering</i> , 2010, 109, 442-446. | 1.1 | 25 |
| 5313 | The construction and application of diploid sake yeast with a homozygous mutation in the FAS2 gene. <i>Journal of Bioscience and Bioengineering</i> , 2010, 110, 675-678. | 1.1 | 16 |
| 5314 | Transcription factor Stb5p is essential for acetaldehyde tolerance in <i>Saccharomyces cerevisiae</i> . <i>Journal of Basic Microbiology</i> , 2010, 50, 494-498. | 1.8 | 14 |
| 5315 | Heterologous expression of an esterase from <i>Thermus thermophilus</i> HB27 in <i>Saccharomyces cerevisiae</i> . <i>Journal of Biotechnology</i> , 2010, 145, 226-232. | 1.9 | 25 |

| # | ARTICLE | IF | CITATIONS |
|------|---|-----|-----------|
| 5316 | Discovery of a steroid 11 β -hydroxylase from <i>Rhizopus oryzae</i> and its biotechnological application. <i>Journal of Biotechnology</i> , 2010, 150, 428-437. | 1.9 | 67 |
| 5317 | Fatty acid production from butter using novel cutinase-displaying yeast. <i>Enzyme and Microbial Technology</i> , 2010, 46, 194-199. | 1.6 | 12 |
| 5318 | Crystallization and preliminary X-ray crystallographic analysis of β -galactosidase from <i>Kluyveromyces lactis</i> . <i>Acta Crystallographica Section F: Structural Biology Communications</i> , 2010, 66, 297-300. | 0.7 | 9 |
| 5319 | Heterologous expression of a tannic acid-inducible laccase3 of <i>Cryphonectria parasitica</i> in <i>Saccharomyces cerevisiae</i> . <i>BMC Biotechnology</i> , 2010, 10, 18. | 1.7 | 22 |
| 5320 | The role of reactive oxygen species in the induction of <i>Ty1</i> retrotransposition in <i>Saccharomyces cerevisiae</i> . <i>Yeast</i> , 2010, 27, 259-267. | 0.8 | 26 |
| 5321 | A functional analysis of <i>Kluyveromyces lactis</i> glutathione reductase. <i>Yeast</i> , 2010, 27, 431-441. | 0.8 | 6 |
| 5322 | Comparative functional analysis of the <i>OCH1</i> mannosyltransferase families in <i>Aspergillus fumigatus</i> and <i>Saccharomyces cerevisiae</i> . <i>Yeast</i> , 2010, 27, 625-636. | 0.8 | 35 |
| 5323 | Isolation and characterization of the carbon catabolite derepressing protein kinase <i>Snf1</i> from the stress tolerant yeast <i>Torulaspora delbrueckii</i> . <i>Yeast</i> , 2010, 27, 1061-1069. | 0.8 | 6 |
| 5324 | Improving the performance of industrial ethanol-producing yeast by expressing the aspartyl protease on the cell surface. <i>Yeast</i> , 2010, 27, 1017-1027. | 0.8 | 12 |
| 5325 | Novel episomal vectors and a highly efficient transformation procedure for the fission yeast <i>Schizosaccharomyces japonicus</i> . <i>Yeast</i> , 2010, 27, 1049-1060. | 0.8 | 40 |
| 5326 | <i>lxr1p</i> regulates oxygen-dependent <i>HEM13</i> transcription. <i>FEMS Yeast Research</i> , 2010, 10, 309-321. | 1.1 | 13 |
| 5327 | The MAP kinase-activated protein kinase <i>Rck2p</i> regulates cellular responses to cell wall stresses, filamentation and virulence in the human fungal pathogen <i>Candida albicans</i> . <i>FEMS Yeast Research</i> , 2010, 10, 441-451. | 1.1 | 21 |
| 5328 | Heterothallism in <i>Saccharomyces cerevisiae</i> isolates from nature: effect of <i>HO</i> locus on the mode of reproduction. <i>Molecular Ecology</i> , 2010, 19, 121-131. | 2.0 | 46 |
| 5329 | Mitochondrial degradation in acetic acid-induced yeast apoptosis: the role of <i>Pep4</i> and the ADP/ATP carrier. <i>Molecular Microbiology</i> , 2010, 76, 1398-1410. | 1.2 | 75 |
| 5330 | <i>FlbC</i> is a putative nuclear C2H2 transcription factor regulating development in <i>Aspergillus nidulans</i> . <i>Molecular Microbiology</i> , 2010, 77, 1203-1219. | 1.2 | 138 |
| 5331 | Calmodulin dissociation regulates <i>Myo5</i> recruitment and function at endocytic sites. <i>EMBO Journal</i> , 2010, 29, 2899-2914. | 3.5 | 25 |
| 5332 | <i>Ku</i> prevents <i>Exo1</i> and <i>Sgs1</i> -dependent resection of DNA ends in the absence of a functional MRX complex or <i>Sae2</i> . <i>EMBO Journal</i> , 2010, 29, 3358-3369. | 3.5 | 262 |
| 5333 | Regulation of clathrin adaptor function in endocytosis: novel role for the SAM domain. <i>EMBO Journal</i> , 2010, 29, 1033-1044. | 3.5 | 38 |

| # | ARTICLE | IF | CITATIONS |
|------|---|-----|-----------|
| 5334 | Conditionally controlling nuclear trafficking in yeast by chemical-induced protein dimerization. <i>Nature Protocols</i> , 2010, 5, 1831-1843. | 5.5 | 25 |
| 5335 | Deletion of <i>GBG1</i> / <i>AYR1</i> Alters Cell Wall Biogenesis in <i>Saccharomyces cerevisiae</i> . <i>Mycobiology</i> , 2010, 38, 102. | 0.6 | 2 |
| 5336 | Multiple Molecular Mechanisms Cause Reproductive Isolation between Three Yeast Species. <i>PLoS Biology</i> , 2010, 8, e1000432. | 2.6 | 131 |
| 5337 | Control of mRNA Export and Translation Termination by Inositol Hexakisphosphate Requires Specific Interaction with Gle1. <i>Journal of Biological Chemistry</i> , 2010, 285, 16683-16692. | 1.6 | 68 |
| 5338 | Transformation of <i>Saccharomyces cerevisiae</i> and other fungi. <i>Bioengineered Bugs</i> , 2010, 1, 395-403. | 2.0 | 174 |
| 5339 | Phosphorylation of Yeast Phosphatidylserine Synthase by Protein Kinase A. <i>Journal of Biological Chemistry</i> , 2010, 285, 11526-11536. | 1.6 | 35 |
| 5340 | A comprehensive analysis of translational missense errors in the yeast <i>Saccharomyces cerevisiae</i> . <i>Rna</i> , 2010, 16, 1797-1808. | 1.6 | 107 |
| 5341 | Regulation of base excision repair: Ntg1 nuclear and mitochondrial dynamic localization in response to genotoxic stress. <i>Nucleic Acids Research</i> , 2010, 38, 3963-3974. | 6.5 | 33 |
| 5342 | Yeast strains with N-terminally truncated ribosomal protein S5: implications for the evolution, structure and function of the Rps5/Rps7 proteins. <i>Nucleic Acids Research</i> , 2010, 38, 1261-1272. | 6.5 | 25 |
| 5343 | Proteinase B Disruption Is Required for High Level Production of Human Mechano-Growth Factor in <i>Saccharomyces cerevisiae</i> . <i>Journal of Molecular Microbiology and Biotechnology</i> , 2010, 18, 188-194. | 1.0 | 3 |
| 5344 | Direct Inhibition of Pumilo Activity by Bam and Bgcn in <i>Drosophila</i> Germ Line Stem Cell Differentiation. <i>Journal of Biological Chemistry</i> , 2010, 285, 4741-4746. | 1.6 | 30 |
| 5345 | Characterization of rice nucleotide sugar transporters capable of transporting UDP-galactose and UDP-glucose. <i>Journal of Biochemistry</i> , 2010, 148, 35-46. | 0.9 | 15 |
| 5346 | A -Mating-Type Gene Expression Can Drive Clamp Formation in the Bipolar Mushroom <i>Pholiota microspora</i> (<i>Pholiota nameko</i>). <i>Eukaryotic Cell</i> , 2010, 9, 1109-1119. | 3.4 | 18 |
| 5347 | Requirements for Transitional Endoplasmic Reticulum Site Structure and Function in <i>Saccharomyces cerevisiae</i> . <i>Molecular Biology of the Cell</i> , 2010, 21, 1530-1545. | 0.9 | 54 |
| 5348 | Constitutive fusion of ubiquitin to PCNA provides DNA damage tolerance independent of translesion polymerase activities. <i>Nucleic Acids Research</i> , 2010, 38, 5047-5058. | 6.5 | 20 |
| 5349 | Analysis of the diffusion of Ras2 in <i>Saccharomyces cerevisiae</i> using fluorescence recovery after photobleaching. <i>Physical Biology</i> , 2010, 7, 026011. | 0.8 | 20 |
| 5350 | Selection systems based on dominant-negative transcription factors for precise genetic engineering. <i>Nucleic Acids Research</i> , 2010, 38, e183-e183. | 6.5 | 6 |
| 5351 | Syt1p promotes activation of Arl1p at the late Golgi to recruit Imh1p. <i>Journal of Cell Science</i> , 2010, 123, 3478-3489. | 1.2 | 24 |

| # | ARTICLE | IF | CITATIONS |
|------|---|-----|-----------|
| 5352 | Identification of novel filament-forming proteins in <i>Saccharomyces cerevisiae</i> and <i>Drosophila melanogaster</i> . <i>Journal of Cell Biology</i> , 2010, 190, 541-551. | 2.3 | 272 |
| 5353 | The Activity of Yeast Hog1 MAPK Is Required during Endoplasmic Reticulum Stress Induced by Tunicamycin Exposure. <i>Journal of Biological Chemistry</i> , 2010, 285, 20088-20096. | 1.6 | 51 |
| 5354 | The Thioredoxin-Thioredoxin Reductase System Can Function in Vivo as an Alternative System to Reduce Oxidized Glutathione in <i>Saccharomyces cerevisiae</i> . <i>Journal of Biological Chemistry</i> , 2010, 285, 6118-6126. | 1.6 | 81 |
| 5355 | Identification of Phosphomethylethanolamine N-Methyltransferase from <i>Arabidopsis</i> and Its Role in Choline and Phospholipid Metabolism. <i>Journal of Biological Chemistry</i> , 2010, 285, 29147-29155. | 1.6 | 33 |
| 5356 | Catabolite Repression of Aox in <i>Pichia pastoris</i> Is Dependent on Hexose Transporter PpHxt1 and Pexophagy. <i>Applied and Environmental Microbiology</i> , 2010, 76, 6108-6118. | 1.4 | 56 |
| 5357 | Regulation of Yeast Nutrient Permease Endocytosis by ATP-binding Cassette Transporters and a Seven-transmembrane Protein, RSB1. <i>Journal of Biological Chemistry</i> , 2010, 285, 35792-35802. | 1.6 | 29 |
| 5358 | Yet1p and Yet3p, the Yeast Homologs of BAP29 and BAP31, Interact with the Endoplasmic Reticulum Translocation Apparatus and Are Required for Inositol Prototrophy. <i>Journal of Biological Chemistry</i> , 2010, 285, 18252-18261. | 1.6 | 28 |
| 5359 | Mutational Analysis of <i>Saccharomyces cerevisiae</i> Erf2 Reveals a Two-step Reaction Mechanism for Protein Palmitoylation by DHHC Enzymes*. <i>Journal of Biological Chemistry</i> , 2010, 285, 38104-38114. | 1.6 | 132 |
| 5360 | Identification, Characterization, and Biosynthesis of a Novel N-Glycan Modification in the Fruiting Body of the Basidiomycete <i>Coprinopsis cinerea</i> . <i>Journal of Biological Chemistry</i> , 2010, 285, 10715-10723. | 1.6 | 24 |
| 5361 | Compartment-specific Synthesis of Phosphatidylethanolamine Is Required for Normal Heavy Metal Resistance. <i>Molecular Biology of the Cell</i> , 2010, 21, 443-455. | 0.9 | 71 |
| 5362 | Differential Roles of Transcriptional Mediator Subunits in Regulation of Multidrug Resistance Gene Expression in <i>Saccharomyces cerevisiae</i> . <i>Molecular Biology of the Cell</i> , 2010, 21, 2469-2482. | 0.9 | 49 |
| 5363 | Extensive DNA End Processing by Exo1 and Sgs1 Inhibits Break-Induced Replication. <i>PLoS Genetics</i> , 2010, 6, e1001007. | 1.5 | 34 |
| 5364 | Multiple Functional Domains of the Yeast 1,3- β -Glucan Synthase Subunit Fks1p Revealed by Quantitative Phenotypic Analysis of Temperature-Sensitive Mutants. <i>Genetics</i> , 2010, 184, 1013-1024. | 1.2 | 56 |
| 5365 | Role of <i>Saccharomyces cerevisiae</i> Oxidoreductases Bdh1p and Ara1p in the Metabolism of Acetoin and 2,3-Butanediol. <i>Applied and Environmental Microbiology</i> , 2010, 76, 670-679. | 1.4 | 38 |
| 5366 | Genetic interaction of RAD53 protein kinase with histones is important for DNA replication. <i>Cell Cycle</i> , 2010, 9, 4735-4747. | 1.3 | 8 |
| 5367 | Components of the multifactor complex needed for internal initiation by the IRES of hepatitis C virus in <i>Saccharomyces cerevisiae</i> . <i>RNA Biology</i> , 2010, 7, 596-605. | 1.5 | 4 |
| 5368 | Improved Transformation of the Halo-Tolerant Yeast <i>Zygosaccharomyces rouxii</i> by Electroporation. <i>Bioscience, Biotechnology and Biochemistry</i> , 2010, 74, 1092-1094. | 0.6 | 20 |
| 5369 | Post-transcriptional regulation in the myo1 ^Δ mutant of <i>Saccharomyces cerevisiae</i> . <i>BMC Genomics</i> , 2010, 11, 690. | 1.2 | 5 |

| # | ARTICLE | IF | CITATIONS |
|------|--|------|-----------|
| 5370 | B-Function Expression in the Flower Center Underlies the Homeotic Phenotype of <i>Lacandonia schismatica</i> (Triuridaceae). <i>Plant Cell</i> , 2010, 22, 3543-3559. | 3.1 | 49 |
| 5371 | Seipin Is a Discrete Homooligomer. <i>Biochemistry</i> , 2010, 49, 10747-10755. | 1.2 | 109 |
| 5372 | Molecular Determinants Required for Selective Interactions between the Thyroid Hormone Receptor Homodimer and the Nuclear Receptor Corepressor N-CoR. <i>Journal of Molecular Biology</i> , 2010, 396, 747-760. | 2.0 | 11 |
| 5373 | Identification and characterisation of <i>eroA</i> and <i>ervA</i> , encoding two putative thiol oxidases from <i>Aspergillus niger</i> . <i>Gene</i> , 2010, 461, 32-41. | 1.0 | 3 |
| 5374 | Lateral reorganization of plasma membrane is involved in the yeast resistance to severe dehydration. <i>Biochimica Et Biophysica Acta - Biomembranes</i> , 2010, 1798, 975-985. | 1.4 | 101 |
| 5375 | Ctf4p facilitates Mcm10p to promote DNA replication in budding yeast. <i>Biochemical and Biophysical Research Communications</i> , 2010, 395, 336-341. | 1.0 | 31 |
| 5376 | Characterization of the developmental regulator FlbE in <i>Aspergillus fumigatus</i> and <i>Aspergillus nidulans</i> . <i>Fungal Genetics and Biology</i> , 2010, 47, 981-993. | 0.9 | 72 |
| 5377 | Photoadaptation in <i>Neurospora</i> by Competitive Interaction of Activating and Inhibitory LOV Domains. <i>Cell</i> , 2010, 142, 762-772. | 13.5 | 149 |
| 5378 | The production, characterisation and enhanced pharmacokinetics of scFv- α 1 albumin fusions expressed in <i>Saccharomyces cerevisiae</i> . <i>Protein Expression and Purification</i> , 2010, 73, 113-124. | 0.6 | 38 |
| 5379 | Site-specific and dose-dependent effects of glucocorticoid receptor phosphorylation in yeast <i>Saccharomyces cerevisiae</i> . <i>Steroids</i> , 2010, 75, 457-465. | 0.8 | 3 |
| 5380 | Establishment of yeast reporter assay systems to detect ligands of thyroid hormone receptors β_1 and β_2 . <i>Toxicology in Vitro</i> , 2010, 24, 638-644. | 1.1 | 33 |
| 5381 | Preparation of a Whole-Cell Biocatalyst of <i>Aspergillus niger</i> Lipase and Its Practical Properties. <i>Journal of Agricultural and Food Chemistry</i> , 2010, 58, 10426-10430. | 2.4 | 17 |
| 5382 | ROS Production and Apoptosis Induction by Formation of Gts1p-Mediated Protein Aggregates. <i>Bioscience, Biotechnology and Biochemistry</i> , 2011, 75, 1546-1553. | 0.6 | 6 |
| 5383 | Analysis of Chloroquine Resistance Transporter (CRT) Isoforms and Orthologues in <i>S. cerevisiae</i> Yeast. <i>Biochemistry</i> , 2011, 50, 6701-6710. | 1.2 | 28 |
| 5384 | Importin Alpha2-Interacting Proteins with Nuclear Roles During Mammalian Spermatogenesis1. <i>Biology of Reproduction</i> , 2011, 85, 1191-1202. | 1.2 | 26 |
| 5385 | Introns within Ribosomal Protein Genes Regulate the Production and Function of Yeast Ribosomes. <i>Cell</i> , 2011, 147, 320-331. | 13.5 | 122 |
| 5386 | Phosphatidylinositol synthase is required for lens structural integrity and photoreceptor cell survival in the zebrafish eye. <i>Experimental Eye Research</i> , 2011, 93, 460-474. | 1.2 | 16 |
| 5387 | Genome-wide inventory of metal homeostasis-related gene products including a functional phytochelatin synthase in the hypogeous mycorrhizal fungus <i>Tuber melanosporum</i> . <i>Fungal Genetics and Biology</i> , 2011, 48, 573-584. | 0.9 | 56 |

| # | ARTICLE | IF | CITATIONS |
|------|--|-----|-----------|
| 5388 | Gln3â€“Gcn4 hybrid transcriptional activator determines catabolic and biosynthetic gene expression in the yeast <i>Saccharomyces cerevisiae</i> . <i>Biochemical and Biophysical Research Communications</i> , 2011, 404, 859-864. | 1.0 | 11 |
| 5389 | Ribosomal protein S3 is stabilized by sumoylation. <i>Biochemical and Biophysical Research Communications</i> , 2011, 414, 523-527. | 1.0 | 21 |
| 5390 | Yeast-based protein delivery to mammalian phagocytic cells is increased by coexpression of bacterial listeriolysin. <i>Microbes and Infection</i> , 2011, 13, 908-913. | 1.0 | 10 |
| 5391 | A Feedback Circuit between Transcriptional Activation and Self-Destruction of Gcn4 Separates Its Metabolic and Morphogenic Response in Diploid Yeasts. <i>Journal of Molecular Biology</i> , 2011, 405, 909-925. | 2.0 | 8 |
| 5392 | Uptake of various yeast genera by antigen-presenting cells and influence of subcellular antigen localization on the activation of ovalbumin-specific CD8 T lymphocytes. <i>Vaccine</i> , 2011, 29, 8165-8173. | 1.7 | 24 |
| 5393 | TDP-43 toxicity in yeast. <i>Methods</i> , 2011, 53, 238-245. | 1.9 | 33 |
| 5394 | Genetic analysis of D-xylose metabolism by endophytic yeast strains of <i>Rhodotorula graminis</i> and <i>Rhodotorula mucilaginosa</i> . <i>Genetics and Molecular Biology</i> , 2011, 34, 471-478. | 0.6 | 12 |
| 5395 | Formaldehyde Oxidizing Enzymes and Genetically Modified Yeast <i>Hansenula polymorpha</i> Cells in Monitoring and Removal of Formaldehyde. , 2011, , . | | 4 |
| 5396 | Extraction of genomic DNA from yeasts for PCR-based applications. <i>BioTechniques</i> , 2011, 50, 325-328. | 0.8 | 413 |
| 5397 | Defects in Very Long Chain Fatty Acid Synthesis Enhance Alpha-Synuclein Toxicity in a Yeast Model of Parkinson's Disease. <i>PLoS ONE</i> , 2011, 6, e15946. | 1.1 | 39 |
| 5398 | A Viable Hypomorphic Allele of the Essential IMP3 Gene Reveals Novel Protein Functions in <i>Saccharomyces cerevisiae</i> . <i>PLoS ONE</i> , 2011, 6, e19500. | 1.1 | 5 |
| 5399 | Heterologous Overexpression and Mutagenesis of the Human Bile Salt Export Pump (ABCB11) Using DREAM (Directed REcombination-Assisted Mutagenesis). <i>PLoS ONE</i> , 2011, 6, e20562. | 1.1 | 11 |
| 5400 | Disruption of <i>Yarrowia lipolytica</i> TPS1 Gene Encoding Trehalose-6-P Synthase Does Not Affect Growth in Glucose but Impairs Growth at High Temperature. <i>PLoS ONE</i> , 2011, 6, e23695. | 1.1 | 23 |
| 5401 | In vivo and in vitro Studies of Adaptor-clathrin Interaction. <i>Journal of Visualized Experiments</i> , 2011, , . | 0.2 | 8 |
| 5402 | Laforin, a dual-specificity phosphatase involved in Lafora disease, is phosphorylated at Ser25 by AMP-activated protein kinase. <i>Biochemical Journal</i> , 2011, 439, 265-275. | 1.7 | 29 |
| 5403 | Red but not dead? Membranes of stressed <i>Saccharomyces cerevisiae</i> are permeable to propidium iodide. <i>Environmental Microbiology</i> , 2011, 13, 163-171. | 1.8 | 175 |
| 5404 | Development of a nondestructive fluorescence-based enzymatic staining of microcolonies for enumerating bacterial contamination in filterable products. <i>Journal of Applied Microbiology</i> , 2011, 110, 69-79. | 1.4 | 10 |
| 5405 | Jasmonic acid perception by CO11 involves inositol polyphosphates in <i>Arabidopsis thaliana</i> . <i>Plant Journal</i> , 2011, 65, 949-957. | 2.8 | 134 |

| # | ARTICLE | IF | CITATIONS |
|------|---|-----|-----------|
| 5406 | Growth defects resulting from inhibiting ERG20 and RAM2 in <i>Candida glabrata</i> . <i>FEMS Microbiology Letters</i> , 2011, 317, 27-33. | 0.7 | 16 |
| 5407 | Vacuolar Import of Phosphatidylcholine Requires the ATP-Binding Cassette Transporter Ybt1. <i>Traffic</i> , 2011, 12, 1257-1268. | 1.3 | 34 |
| 5408 | Cdc42 Regulates Multiple Membrane Traffic Events in Fission Yeast. <i>Traffic</i> , 2011, 12, 1744-1758. | 1.3 | 44 |
| 5409 | Far3p domains involved in the interactions of Far proteins and pheromone-induced cell cycle arrest in budding yeast. <i>FEMS Yeast Research</i> , 2011, 11, 72-79. | 1.1 | 6 |
| 5410 | Calcium/calmodulin kinase1 and its relation to thermotolerance and HSP90 in <i>Sporothrix schenckii</i> : an RNAi and yeast two-hybrid study. <i>BMC Microbiology</i> , 2011, 11, 162. | 1.3 | 38 |
| 5411 | Novel screening system for protein-protein interactions by bimolecular fluorescence complementation in <i>Saccharomyces cerevisiae</i> . <i>Journal of Bioscience and Bioengineering</i> , 2011, 111, 397-401. | 1.1 | 19 |
| 5412 | The retroviral proteinase active site and the N-terminus of Ddi1 are required for repression of protein secretion. <i>FEBS Letters</i> , 2011, 585, 139-142. | 1.3 | 26 |
| 5413 | Bcs1p can rescue a large and productive cytochrome bc1 complex assembly intermediate in the inner membrane of yeast mitochondria. <i>Biochimica Et Biophysica Acta - Molecular Cell Research</i> , 2011, 1813, 91-101. | 1.9 | 15 |
| 5414 | Enhancement of xylitol production by attenuation of intracellular xylitol dehydrogenase activity in <i>Candida tropicalis</i> . <i>Biotechnology Letters</i> , 2011, 33, 1209-1213. | 1.1 | 21 |
| 5415 | Improving the ethanol yield by reducing glycerol formation using cofactor regulation in <i>Saccharomyces cerevisiae</i> . <i>Biotechnology Letters</i> , 2011, 33, 1375-1380. | 1.1 | 32 |
| 5416 | Molecular cloning and characterization of a trehalose-6-phosphate synthase/phosphatase from <i>Dunaliella viridis</i> . <i>Molecular Biology Reports</i> , 2011, 38, 2241-2248. | 1.0 | 7 |
| 5417 | Improving ethanol productivity by modification of glycolytic redox factor generation in glycerol-3-phosphate dehydrogenase mutants of an industrial ethanol yeast. <i>Journal of Industrial Microbiology and Biotechnology</i> , 2011, 38, 935-943. | 1.4 | 24 |
| 5418 | Cellular localization of human p53 expressed in the yeast <i>Saccharomyces cerevisiae</i> : effect of NLSI deletion. <i>Apoptosis: an International Journal on Programmed Cell Death</i> , 2011, 16, 746-756. | 2.2 | 9 |
| 5419 | Phytochelatin synthase of <i>Thlaspi caerulescens</i> enhanced tolerance and accumulation of heavy metals when expressed in yeast and tobacco. <i>Plant Cell Reports</i> , 2011, 30, 1067-1076. | 2.8 | 66 |
| 5420 | The characterization of two peroxiredoxin genes in <i>Dunaliella viridis</i> provides insights into antioxidative response to salt stress. <i>Plant Cell Reports</i> , 2011, 30, 1503-1512. | 2.8 | 9 |
| 5421 | The Role of Cell Wall Revealed by the Visualization of <i>Saccharomyces cerevisiae</i> Transformation. <i>Current Microbiology</i> , 2011, 62, 956-961. | 1.0 | 18 |
| 5422 | Visualization of the synergistic effect of lithium acetate and single-stranded carrier DNA on <i>Saccharomyces cerevisiae</i> transformation. <i>Current Genetics</i> , 2011, 57, 233-239. | 0.8 | 26 |
| 5423 | Heterologous expression of a thermophilic esterase in <i>Kluyveromyces</i> yeasts. <i>Applied Microbiology and Biotechnology</i> , 2011, 89, 375-385. | 1.7 | 34 |

| # | ARTICLE | IF | CITATIONS |
|------|--|-----|-----------|
| 5424 | Efficient accumulation of oleic acid in <i>Saccharomyces cerevisiae</i> caused by expression of rat elongase 2 gene (rELO2) and its contribution to tolerance to alcohols. <i>Applied Microbiology and Biotechnology</i> , 2011, 91, 1593-1600. | 1.7 | 31 |
| 5425 | Intermolecular Interactions of Chrysanthemum virus B Coat Protein: Implications for Capsid Assembly. <i>Indian Journal of Virology: an Official Organ of Indian Virological Society</i> , 2011, 22, 111-116. | 0.7 | 6 |
| 5426 | Elastic properties of chemically modified baker's yeast cells studied by AFM. <i>Surface and Interface Analysis</i> , 2011, 43, 1636-1640. | 0.8 | 11 |
| 5427 | TCA cycle-independent acetate metabolism via the glyoxylate cycle in <i>Saccharomyces cerevisiae</i> . <i>Yeast</i> , 2011, 28, 153-166. | 0.8 | 63 |
| 5428 | Expression of aspartic protease from <i>Neurospora crassa</i> in industrial ethanol-producing yeast and its application in ethanol production. <i>Enzyme and Microbial Technology</i> , 2011, 48, 148-154. | 1.6 | 15 |
| 5429 | Expression of bacterial levanase in yeast enables simultaneous saccharification and fermentation of grass juice to bioethanol. <i>Bioresource Technology</i> , 2011, 102, 1503-1508. | 4.8 | 13 |
| 5430 | Development of an industrial ethanol-producing yeast strain for efficient utilization of cellobiose. <i>Enzyme and Microbial Technology</i> , 2011, 49, 105-112. | 1.6 | 26 |
| 5431 | Minimization of glycerol synthesis in industrial ethanol yeast without influencing its fermentation performance. <i>Metabolic Engineering</i> , 2011, 13, 49-59. | 3.6 | 62 |
| 5432 | Inhibition of Heat Tolerance and Nuclear Import of Gts1p by Ssa1p and Ssa2p. <i>Bioscience, Biotechnology and Biochemistry</i> , 2011, 75, 323-330. | 0.6 | 5 |
| 5433 | GTS1 Induction Causes Derepression of Tup1-Cyc8-Repressing Genes and Chromatin Remodeling through the Interaction of Gts1p with Cyc8p. <i>Bioscience, Biotechnology and Biochemistry</i> , 2011, 75, 740-747. | 0.6 | 4 |
| 5434 | Identification of Yeast Genes Involved in K ⁺ Homeostasis: Loss of Membrane Traffic Genes Affects K ⁺ Uptake. <i>G3: Genes, Genomes, Genetics</i> , 2011, 1, 43-56. | 0.8 | 13 |
| 5435 | Meiotic recombination protein Rec12: functional conservation, crossover homeostasis and early crossover/non-crossover decision. <i>Nucleic Acids Research</i> , 2011, 39, 1460-1472. | 6.5 | 23 |
| 5436 | The N-Terminal Domain of the Flo1 Flocculation Protein from <i>Saccharomyces cerevisiae</i> Binds Specifically to Mannose Carbohydrates. <i>Eukaryotic Cell</i> , 2011, 10, 110-117. | 3.4 | 35 |
| 5437 | Functional Pentameric Formation via Coexpression of the <i>Escherichia coli</i> Heat-Labile Enterotoxin B Subunit and Its Fusion Protein Subunit with a Neutralizing Epitope of ApxIIA Exotoxin Improves the Mucosal Immunogenicity and Protection against Challenge by <i>Actinobacillus pleuropneumoniae</i> . <i>Vaccine Journal</i> , 2011, 18, 2168-2177. | 3.2 | 15 |
| 5438 | Putative Role of Cellulosomal Protease Inhibitors in <i>Clostridium cellulovorans</i> Based on Gene Expression and Measurement of Activities. <i>Journal of Bacteriology</i> , 2011, 193, 5527-5530. | 1.0 | 14 |
| 5439 | Hap2-3-5-Gln3 determine transcriptional activation of GDH1 and ASN1 under repressive nitrogen conditions in the yeast <i>Saccharomyces cerevisiae</i> . <i>Microbiology (United Kingdom)</i> , 2011, 157, 879-889. | 0.7 | 14 |
| 5440 | Structural and functional analysis of Nro1/Ett1: a protein involved in translation termination in <i>S. cerevisiae</i> and in O ₂ -mediated gene control in <i>S. pombe</i> . <i>Rna</i> , 2011, 17, 1213-1224. | 1.6 | 11 |
| 5441 | Mutations in Genes <i>patA</i> and <i>patL</i> of <i>Anabaena</i> sp. Strain PCC 7120 Result in Similar Phenotypes, and the Proteins Encoded by Those Genes May Interact. <i>Journal of Bacteriology</i> , 2011, 193, 6070-6074. | 1.0 | 13 |

| # | ARTICLE | IF | CITATIONS |
|------|--|-----|-----------|
| 5442 | Processing of HEBP1 by Cathepsin D Gives Rise to F2L, the Agonist of Formyl Peptide Receptor 3. <i>Journal of Immunology</i> , 2011, 187, 1475-1485. | 0.4 | 20 |
| 5443 | Mutations of highly conserved bases in the peptidyltransferase center induce compensatory rearrangements in yeast ribosomes. <i>Rna</i> , 2011, 17, 855-864. | 1.6 | 23 |
| 5444 | The Putative Lipid Transporter, Arv1, Is Required for Activating Pheromone-Induced MAP Kinase Signaling in <i>Saccharomyces cerevisiae</i> . <i>Genetics</i> , 2011, 187, 455-465. | 1.2 | 15 |
| 5445 | Yet1p and Yet3p interacts with Scs2p and Opi1p to regulate ER localization of the Opi1p repressor. <i>Molecular Biology of the Cell</i> , 2011, 22, 1430-1439. | 0.9 | 15 |
| 5446 | Multicopy Suppression Screening of <i>Saccharomyces cerevisiae</i> Identifies the Ubiquitination Machinery as a Main Target for Improving Growth at Low Temperatures. <i>Applied and Environmental Microbiology</i> , 2011, 77, 7517-7525. | 1.4 | 14 |
| 5447 | Two Proteins with Different Functions Are Derived from the <i>KIHEM13</i> Gene. <i>Eukaryotic Cell</i> , 2011, 10, 1331-1339. | 3.4 | 1 |
| 5448 | HIV proteinase inhibitors target the Ddi1-like protein of <i>Leishmania</i> parasites. <i>FASEB Journal</i> , 2011, 25, 1729-1736. | 0.2 | 43 |
| 5449 | DGK1-encoded Diacylglycerol Kinase Activity Is Required for Phospholipid Synthesis during Growth Resumption from Stationary Phase in <i>Saccharomyces cerevisiae</i> . <i>Journal of Biological Chemistry</i> , 2011, 286, 1464-1474. | 1.6 | 63 |
| 5450 | Loss of Subcellular Lipid Transport Due to ARV1 Deficiency Disrupts Organelle Homeostasis and Activates the Unfolded Protein Response. <i>Journal of Biological Chemistry</i> , 2011, 286, 11951-11959. | 1.6 | 34 |
| 5451 | Differential Oxidant Tolerance Determined by the Key Transcription Factor Yap1 Is Controlled by Levels of the Yap1-binding Protein, Ybp1. <i>Journal of Biological Chemistry</i> , 2011, 286, 34071-34081. | 1.6 | 35 |
| 5452 | Phosphorylation of Phosphatidate Phosphatase Regulates Its Membrane Association and Physiological Functions in <i>Saccharomyces cerevisiae</i> . <i>Journal of Biological Chemistry</i> , 2011, 286, 1486-1498. | 1.6 | 106 |
| 5453 | Endogenous ribosomal frameshift signals operate as mRNA destabilizing elements through at least two molecular pathways in yeast. <i>Nucleic Acids Research</i> , 2011, 39, 2799-2808. | 6.5 | 62 |
| 5454 | 5'UTR: Identification and Analysis of Translationally Regulative 5' Untranslated Regions in Amino Acid Starved Yeast Cells. <i>Molecular and Cellular Proteomics</i> , 2011, 10, M110.003350. | 2.5 | 5 |
| 5455 | The yeast lipin orthologue Pah1p is important for biogenesis of lipid droplets. <i>Journal of Cell Biology</i> , 2011, 192, 1043-1055. | 2.3 | 264 |
| 5456 | Functional expression of l-fucokinase/guanosine 5'-diphosphate-l-fucose pyrophosphorylase from <i>Bacteroides fragilis</i> in <i>Saccharomyces cerevisiae</i> for the production of nucleotide sugars from exogenous monosaccharides. <i>Glycobiology</i> , 2011, 21, 1228-1236. | 1.3 | 25 |
| 5457 | Mechanistic Analysis of AKT1 Regulation by the CBL-CIPK-PP2CA Interactions. <i>Molecular Plant</i> , 2011, 4, 527-536. | 3.9 | 136 |
| 5458 | Nuclear localization of the meiosis-specific transcription factor Ndt80 is regulated by the pachytene checkpoint. <i>Molecular Biology of the Cell</i> , 2011, 22, 1878-1886. | 0.9 | 23 |
| 5459 | Regulation of the CgPdr1 Transcription Factor from the Pathogen <i>Candida glabrata</i> . <i>Eukaryotic Cell</i> , 2011, 10, 187-197. | 3.4 | 65 |

| # | ARTICLE | IF | CITATIONS |
|------|---|-----|-----------|
| 5460 | Molecular Determinants and Genetic Modifiers of Aggregation and Toxicity for the ALS Disease Protein FUS/TLS. <i>PLoS Biology</i> , 2011, 9, e1000614. | 2.6 | 396 |
| 5461 | Impaired dimer assembly and decreased stability of naturally recurring R260C mutant A subunit for coagulation factor XIII. <i>Journal of Biochemistry</i> , 2012, 152, 471-478. | 0.9 | 7 |
| 5462 | TERRA Promotes Telomere Shortening through Exonuclease 1-Mediated Resection of Chromosome Ends. <i>PLoS Genetics</i> , 2012, 8, e1002747. | 1.5 | 132 |
| 5463 | Heme Uptake by <i>Leishmania amazonensis</i> Is Mediated by the Transmembrane Protein LHR1. <i>PLoS Pathogens</i> , 2012, 8, e1002795. | 2.1 | 88 |
| 5464 | Protein Kinase A-mediated Phosphorylation of Pah1p Phosphatidate Phosphatase Functions in Conjunction with the Pho85p-Pho80p and Cdc28p-Cyclin B Kinases to Regulate Lipid Synthesis in Yeast. <i>Journal of Biological Chemistry</i> , 2012, 287, 33364-33376. | 1.6 | 70 |
| 5465 | Protein phosphatase CaPpz1 is involved in cation homeostasis, cell wall integrity and virulence of <i>Candida albicans</i> . <i>Microbiology (United Kingdom)</i> , 2012, 158, 1258-1267. | 0.7 | 34 |
| 5466 | Rbg1-Tma46 dimer structure reveals new functional domains and their role in polysome recruitment. <i>Nucleic Acids Research</i> , 2012, 40, 11100-11114. | 6.5 | 23 |
| 5467 | Structural Studies and Protein Engineering of Inositol Phosphate Multikinase. <i>Journal of Biological Chemistry</i> , 2012, 287, 35360-35369. | 1.6 | 28 |
| 5468 | The <i>Salmonella</i> Typhimurium effector SteC inhibits Cdc42-mediated signaling through binding to the exchange factor Cdc24 in <i>Saccharomyces cerevisiae</i> . <i>Molecular Biology of the Cell</i> , 2012, 23, 4430-4443. | 0.9 | 14 |
| 5469 | The Putative Guanine Nucleotide Exchange Factor RicA Mediates Upstream Signaling for Growth and Development in <i>Aspergillus</i> . <i>Eukaryotic Cell</i> , 2012, 11, 1399-1412. | 3.4 | 25 |
| 5470 | Application of Novel Nanotechnology Strategies in Plant Biotransformation: A Contemporary Overview. <i>Recent Patents on Biotechnology</i> , 2012, 6, 69-79. | 0.4 | 23 |
| 5471 | Ca ²⁺ Induces Spontaneous Dephosphorylation of a Novel P5A-type ATPase. <i>Journal of Biological Chemistry</i> , 2012, 287, 28336-28348. | 1.6 | 17 |
| 5472 | Nuclear Envelope Phosphatase 1-Regulatory Subunit 1 (Formerly TMEM188) Is the Metazoan Spo7p Ortholog and Functions in the Lipin Activation Pathway. <i>Journal of Biological Chemistry</i> , 2012, 287, 3123-3137. | 1.6 | 86 |
| 5473 | Phosphatidate Phosphatase Plays Role in Zinc-mediated Regulation of Phospholipid Synthesis in Yeast. <i>Journal of Biological Chemistry</i> , 2012, 287, 968-977. | 1.6 | 42 |
| 5474 | Proteolytic Degradation of the Yap1 Transcription Factor Is Regulated by Subcellular Localization and the E3 Ubiquitin Ligase Not4. <i>Journal of Biological Chemistry</i> , 2012, 287, 26796-26805. | 1.6 | 41 |
| 5475 | The <i>Saccharomyces cerevisiae</i> Actin Patch Protein App1p Is a Phosphatidate Phosphatase Enzyme. <i>Journal of Biological Chemistry</i> , 2012, 287, 40186-40196. | 1.6 | 48 |
| 5476 | Pho85p-Pho80p Phosphorylation of Yeast Pah1p Phosphatidate Phosphatase Regulates Its Activity, Location, Abundance, and Function in Lipid Metabolism. <i>Journal of Biological Chemistry</i> , 2012, 287, 11290-11301. | 1.6 | 89 |
| 5477 | Competition between the golgin Imh1p and the Gcs1p GAP stabilizes activated Arl1p at the late-Golgi. <i>Journal of Cell Science</i> , 2012, 125, 4586-96. | 1.2 | 9 |

| # | ARTICLE | IF | CITATIONS |
|------|---|-----|-----------|
| 5478 | Integrity and Function of the <i>Saccharomyces cerevisiae</i> Spindle Pole Body Depends on Connections Between the Membrane Proteins Ndc1, Rtn1, and Yop1. <i>Genetics</i> , 2012, 192, 441-455. | 1.2 | 27 |
| 5479 | Topologically Conserved Residues Direct Heme Transport in HRC-1-related Proteins. <i>Journal of Biological Chemistry</i> , 2012, 287, 4914-4924. | 1.6 | 55 |
| 5480 | SLAC, a complex between Sla1 and Las17, regulates actin polymerization during clathrin-mediated endocytosis. <i>Molecular Biology of the Cell</i> , 2012, 23, 4256-4272. | 0.9 | 50 |
| 5481 | Evaluating the role of the FUS/TLS-related gene EWSR1 in amyotrophic lateral sclerosis. <i>Human Molecular Genetics</i> , 2012, 21, 2899-2911. | 1.4 | 246 |
| 5482 | Interaction of two photoreceptors in the regulation of bacterial photosynthesis genes. <i>Nucleic Acids Research</i> , 2012, 40, 5901-5909. | 6.5 | 20 |
| 5483 | Impaired coenzyme A synthesis in fission yeast causes defective mitosis, quiescence-exit failure, histone hypoacetylation and fragile DNA. <i>Open Biology</i> , 2012, 2, 120117. | 1.5 | 32 |
| 5484 | Mitochondrial type 2C protein phosphatases CaPtc5p, CaPtc6p, and CaPtc7p play vital roles in cellular responses to antifungal drugs and cadmium in <i>Candida albicans</i> . <i>FEMS Yeast Research</i> , 2012, 12, 897-906. | 1.1 | 16 |
| 5485 | Development and characterization of a vector set with regulated promoters for systematic metabolic engineering in <i>Saccharomyces cerevisiae</i> . <i>Yeast</i> , 2012, 29, 495-503. | 0.8 | 31 |
| 5486 | The Erf4 Subunit of the Yeast Ras Palmitoyl Acyltransferase Is Required for Stability of the Acyl-Erf2 Intermediate and Palmitoyl Transfer to a Ras2 Substrate. <i>Journal of Biological Chemistry</i> , 2012, 287, 34337-34348. | 1.6 | 41 |
| 5487 | Involvement of GDH3-encoded NADP ⁺ -dependent Glutamate Dehydrogenase in Yeast Cell Resistance to Stress-induced Apoptosis in Stationary Phase Cells. <i>Journal of Biological Chemistry</i> , 2012, 287, 44221-44233. | 1.6 | 23 |
| 5488 | Arabidopsis ABCB21 is a Facultative Auxin Importer/Exporter Regulated by Cytoplasmic Auxin Concentration. <i>Plant and Cell Physiology</i> , 2012, 53, 2090-2100. | 1.5 | 132 |
| 5489 | Delivery of functional DNA and messenger RNA to mammalian phagocytic cells by recombinant yeast. <i>Gene Therapy</i> , 2012, 19, 237-245. | 2.3 | 35 |
| 5490 | Functional roles of the protein phosphatase 2C, AtAIP1, in abscisic acid signaling and sugar tolerance in Arabidopsis. <i>Plant Science</i> , 2012, 187, 83-88. | 1.7 | 35 |
| 5491 | Analysis of the subcellular targeting of the smaller replicase protein of Pelargonium flower break virus. <i>Virus Research</i> , 2012, 163, 580-591. | 1.1 | 5 |
| 5492 | Molar concentrations of sorbitol and polyethylene glycol inhibit the Plasmodium aquaglyceroporin but not that of E. coli: Involvement of the channel vestibules. <i>Biochimica Et Biophysica Acta - Biomembranes</i> , 2012, 1818, 1218-1224. | 1.4 | 25 |
| 5493 | Arv1 lipid transporter function is conserved between pathogenic and nonpathogenic fungi. <i>Fungal Genetics and Biology</i> , 2012, 49, 101-113. | 0.9 | 9 |
| 5494 | Conserved regulators of the cell separation process in Schizosaccharomyces. <i>Fungal Genetics and Biology</i> , 2012, 49, 235-249. | 0.9 | 9 |
| 5495 | KlRox1p contributes to yeast resistance to metals and is necessary for KLYCF1 expression in the presence of cadmium. <i>Gene</i> , 2012, 497, 27-37. | 1.0 | 14 |

| # | ARTICLE | IF | CITATIONS |
|------|--|-----|-----------|
| 5496 | Specific adsorption of tungstate by cell surface display of the newly designed ModE mutant. <i>Applied Microbiology and Biotechnology</i> , 2012, 96, 153-159. | 1.7 | 24 |
| 5497 | Extracellular glutathione fermentation using engineered <i>Saccharomyces cerevisiae</i> expressing a novel glutathione exporter. <i>Applied Microbiology and Biotechnology</i> , 2012, 96, 1021-1027. | 1.7 | 24 |
| 5498 | Functional and genetic interactions of TOR in the budding yeast <i>Saccharomyces cerevisiae</i> with myosin type II-deficiency (<i>myo1^Δ</i>). <i>BMC Cell Biology</i> , 2012, 13, 13. | 3.0 | 4 |
| 5499 | A stress inducible SUMO conjugating enzyme gene (<i>SaSce9</i>) from a grass halophyte <i>Spartina alterniflora</i> enhances salinity and drought stress tolerance in <i>Arabidopsis</i> . <i>BMC Plant Biology</i> , 2012, 12, 187. | 1.6 | 61 |
| 5500 | Rice UBC13, a candidate housekeeping gene, is required for K63-linked polyubiquitination and tolerance to DNA damage. <i>Rice</i> , 2012, 5, 24. | 1.7 | 22 |
| 5501 | Construction of a novel selection system for endoglucanases exhibiting carbohydrate-binding modules optimized for biomass using yeast cell-surface engineering. <i>AMB Express</i> , 2012, 2, 56. | 1.4 | 10 |
| 5502 | Characterization of <i>GCY1</i> in <i>Saccharomyces cerevisiae</i> by metabolic profiling. <i>Journal of Applied Microbiology</i> , 2012, 113, 1468-1478. | 1.4 | 18 |
| 5503 | A pentapeptide signature motif plays a pivotal role in <i>Leishmania</i> DNA topoisomerase IB activity and camptothecin sensitivity. <i>Biochimica Et Biophysica Acta - General Subjects</i> , 2012, 1820, 2062-2071. | 1.1 | 14 |
| 5504 | Interaction network of antimicrobial peptides of <i>Arabidopsis thaliana</i> , based on high-throughput yeast two-hybrid screening. <i>Plant Physiology and Biochemistry</i> , 2012, 58, 245-252. | 2.8 | 10 |
| 5505 | Low temperature highlights the functional role of the cell wall integrity pathway in the regulation of growth in <i>Saccharomyces cerevisiae</i> . <i>Biochemical Journal</i> , 2012, 446, 477-488. | 1.7 | 19 |
| 5506 | Yeasts as a Tool for Heterologous Gene Expression. <i>Methods in Molecular Biology</i> , 2012, 824, 359-370. | 0.4 | 22 |
| 5507 | Cloning the <i>Acholeplasma laidlawii</i> PG-8A Genome in <i>Saccharomyces cerevisiae</i> as a Yeast Centromeric Plasmid. <i>ACS Synthetic Biology</i> , 2012, 1, 22-28. | 1.9 | 43 |
| 5508 | Air-drying kinetics affect yeast membrane organization and survival. <i>Applied Microbiology and Biotechnology</i> , 2012, 96, 471-480. | 1.7 | 39 |
| 5509 | Mitochondrial-Nuclear DNA Interactions Contribute to the Regulation of Nuclear Transcript Levels as Part of the Inter-Organelle Communication System. <i>PLoS ONE</i> , 2012, 7, e30943. | 1.1 | 21 |
| 5510 | Cationic Amphiphilic Drugs Are Potent Inhibitors of Yeast Sporulation. <i>PLoS ONE</i> , 2012, 7, e42853. | 1.1 | 8 |
| 5511 | MRN1 Implicates Chromatin Remodeling Complexes and Architectural Factors in mRNA Maturation. <i>PLoS ONE</i> , 2012, 7, e44373. | 1.1 | 6 |
| 5512 | Chloroplast NADPH-Dependent Thioredoxin Reductase from <i>Chlorella vulgaris</i> Alleviates Environmental Stresses in Yeast Together with 2-Cys Peroxiredoxin. <i>PLoS ONE</i> , 2012, 7, e45988. | 1.1 | 8 |
| 5513 | The Co-Chaperone Hch1 Regulates Hsp90 Function Differently than Its Homologue Aha1 and Confers Sensitivity to Yeast to the Hsp90 Inhibitor NVP-AUY922. <i>PLoS ONE</i> , 2012, 7, e49322. | 1.1 | 28 |

| # | ARTICLE | IF | CITATIONS |
|------|--|-----|-----------|
| 5514 | Effect of pretreatment of hydrothermally processed rice straw with laccase-displaying yeast on ethanol fermentation. <i>Applied Microbiology and Biotechnology</i> , 2012, 94, 939-948. | 1.7 | 39 |
| 5515 | Enhanced arsenate uptake in <i>Saccharomyces cerevisiae</i> overexpressing the Pho84 phosphate transporter. <i>Biotechnology Progress</i> , 2012, 28, 654-661. | 1.3 | 24 |
| 5516 | Expressing Î ² -glucosidase from <i>Saccharomycopsis fibuligera</i> in industrial ethanol producing yeast and evaluation of the expressing sufficiency. <i>Annals of Microbiology</i> , 2012, 62, 539-544. | 1.1 | 4 |
| 5517 | Protective effects of flavonoids and extract from <i>Vellozia kolbekii</i> Alves against oxidative stress induced by hydrogen peroxide in yeast. <i>Journal of Natural Medicines</i> , 2012, 66, 367-372. | 1.1 | 10 |
| 5518 | The N-terminal domain of the Flo11 protein from <i>Saccharomyces cerevisiae</i> is an adhesin without mannose-binding activity. <i>FEMS Yeast Research</i> , 2012, 12, 78-87. | 1.1 | 28 |
| 5519 | Zebrafish Mms2 promotes K63-linked polyubiquitination and is involved in p53-mediated DNA-damage response. <i>DNA Repair</i> , 2012, 11, 157-166. | 1.3 | 14 |
| 5520 | Î-Aminolevulinate synthase is required for apical transcellular barrier formation in the skin of the <i>Drosophila</i> larva. <i>European Journal of Cell Biology</i> , 2012, 91, 204-215. | 1.6 | 23 |
| 5521 | Novel genetic tools for <i>Hansenula polymorpha</i> . <i>FEMS Yeast Research</i> , 2012, 12, 271-278. | 1.1 | 71 |
| 5522 | Mon2 is a negative regulator of the monomeric G protein, Arl1. <i>FEMS Yeast Research</i> , 2012, 12, 637-650. | 1.1 | 5 |
| 5523 | Direct fermentation of newspaper after laccase-treatment using yeast codisplaying endoglucanase, cellobiohydrolase, and Î ² -glucosidase. <i>Renewable Energy</i> , 2012, 44, 199-205. | 4.3 | 15 |
| 5524 | RWDD1 interacts with the ligand binding domain of the androgen receptor and acts as a coactivator of androgen-dependent transactivation. <i>Molecular and Cellular Endocrinology</i> , 2012, 358, 53-62. | 1.6 | 9 |
| 5525 | Construction of the industrial ethanol-producing strain of <i>Saccharomyces cerevisiae</i> able to ferment cellobiose and melibiose. <i>Applied Biochemistry and Microbiology</i> , 2012, 48, 216-221. | 0.3 | 6 |
| 5526 | Co-expression of <i>Arabidopsis thaliana</i> phytochelatin synthase and <i>Treponema denticola</i> cysteine desulfhydrase for enhanced arsenic accumulation. <i>Biotechnology and Bioengineering</i> , 2012, 109, 605-608. | 1.7 | 18 |
| 5527 | Plant ornithine decarboxylase is not post-transcriptionally feedback regulated by polyamines but can interact with a cytosolic ribosomal protein S15 polypeptide. <i>Amino Acids</i> , 2012, 42, 519-527. | 1.2 | 10 |
| 5528 | Overexpression of the phosphatidylinositol synthase gene from <i>Zea mays</i> in tobacco plants alters the membrane lipids composition and improves drought stress tolerance. <i>Planta</i> , 2012, 235, 69-84. | 1.6 | 28 |
| 5529 | Efficient expression of a <i>Paenibacillus barcinonensis</i> endoglucanase in <i>Saccharomyces cerevisiae</i> . <i>Journal of Industrial Microbiology and Biotechnology</i> , 2012, 39, 115-123. | 1.4 | 13 |
| 5530 | An eYFP Reporter Gene for the Yeast Two-hybrid System. <i>Protein Journal</i> , 2013, 32, 126-130. | 0.7 | 0 |
| 5531 | Functional analyses of <i>TaHMA2</i> , a <i>P1B</i> -type <i>ATPase</i> in wheat. <i>Plant Biotechnology Journal</i> , 2013, 11, 420-431. | 4.1 | 82 |

| # | ARTICLE | IF | CITATIONS |
|------|---|------|-----------|
| 5532 | Profilin is required for Ca ²⁺ homeostasis and Ca ²⁺ -modulated bud formation in yeast. <i>Molecular Genetics and Genomics</i> , 2013, 288, 317-328. | 1.0 | 3 |
| 5533 | Construction of a Novel D ^o -Agglutinin Expression System on the Surface of Wild-Type <i>Saccharomyces cerevisiae</i> Y5 and Genetic Immobilization of I ² -Glucosidase1. <i>Bioenergy Research</i> , 2013, 6, 1205-1211. | 2.2 | 3 |
| 5534 | Engineering of the glycerol decomposition pathway and cofactor regulation in an industrial yeast improves ethanol production. <i>Journal of Industrial Microbiology and Biotechnology</i> , 2013, 40, 1153-1160. | 1.4 | 22 |
| 5535 | ERG9 and COQ1 disruption reveals isoprenoids biosynthesis is closely related to mitochondrial function in <i>Saccharomyces cerevisiae</i> . <i>Integrative Biology (United Kingdom)</i> , 2013, 5, 1282. | 0.6 | 7 |
| 5536 | Comparison of pyruvate decarboxylases from <i>Saccharomyces cerevisiae</i> and <i>Komagataella pastoris</i> (<i>Pichia pastoris</i>). <i>Applied Microbiology and Biotechnology</i> , 2013, 97, 9439-9449. | 1.7 | 17 |
| 5537 | Overexpression of the active diacylglycerol acyltransferase variant transforms <i>Saccharomyces cerevisiae</i> into an oleaginous yeast. <i>Applied Microbiology and Biotechnology</i> , 2013, 97, 7345-7355. | 1.7 | 56 |
| 5538 | Oxidized glutathione fermentation using <i>Saccharomyces cerevisiae</i> engineered for glutathione metabolism. <i>Applied Microbiology and Biotechnology</i> , 2013, 97, 7399-7404. | 1.7 | 18 |
| 5539 | Implementation of a transhydrogenase-like shunt to counter redox imbalance during xylose fermentation in <i>Saccharomyces cerevisiae</i> . <i>Applied Microbiology and Biotechnology</i> , 2013, 97, 1669-1678. | 1.7 | 26 |
| 5540 | Overcoming recalcitrant transformation and gene manipulation in Pucciniomycotina yeasts. <i>Applied Microbiology and Biotechnology</i> , 2013, 97, 283-295. | 1.7 | 39 |
| 5541 | Overexpression of a <i>Zea mays</i> phospholipase C1 gene enhances drought tolerance in tobacco in part by maintaining stability in the membrane lipid composition. <i>Plant Cell, Tissue and Organ Culture</i> , 2013, 115, 253-262. | 1.2 | 17 |
| 5542 | Cytochrome c peroxidase is a mitochondrial heme-based H ₂ O ₂ sensor that modulates antioxidant defense. <i>Free Radical Biology and Medicine</i> , 2013, 65, 541-551. | 1.3 | 36 |
| 5543 | Cocktail $\hat{\Gamma}$ -integration of xylose assimilation genes for efficient ethanol production from xylose in <i>Saccharomyces cerevisiae</i> . <i>Journal of Bioscience and Bioengineering</i> , 2013, 116, 333-336. | 1.1 | 32 |
| 5544 | An oral vaccine against candidiasis generated by a yeast molecular display system. <i>Pathogens and Disease</i> , 2013, 69, 262-268. | 0.8 | 44 |
| 5545 | Negative Functional Interaction Between Cell Integrity MAPK Pathway and Rho1 GTPase in Fission Yeast. <i>Genetics</i> , 2013, 195, 421-432. | 1.2 | 27 |
| 5546 | The THO complex component Thp2 counteracts telomeric R-loops and telomere shortening. <i>EMBO Journal</i> , 2013, 32, 2861-2871. | 3.5 | 125 |
| 5547 | Crystal structure of the 14-subunit RNA polymerase $\hat{\text{A}}$. <i>Nature</i> , 2013, 502, 644-649. | 13.7 | 179 |
| 5548 | Relationship between ethanol and oxidative stress in laboratory and brewing yeast strains. <i>Journal of Bioscience and Bioengineering</i> , 2013, 116, 697-705. | 1.1 | 42 |
| 5549 | Biocatalytic production of alpha-hydroxy ketones and vicinal diols by yeast and human aldo $\hat{\text{e}}$ “keto reductases. <i>Chemico-Biological Interactions</i> , 2013, 202, 195-203. | 1.7 | 16 |

| # | ARTICLE | IF | CITATIONS |
|------|--|------|-----------|
| 5550 | Increased isobutanol production in <i>Saccharomyces cerevisiae</i> by eliminating competing pathways and resolving cofactor imbalance. <i>Microbial Cell Factories</i> , 2013, 12, 119. | 1.9 | 126 |
| 5551 | A New Simple Method for Isolating Multistress-Tolerant Semidominant Mutants of <i>Saccharomyces cerevisiae</i> by One-Step Selection under Lethal Hydrogen Peroxide Stress Condition. <i>Bioscience, Biotechnology and Biochemistry</i> , 2013, 77, 224-228. | 0.6 | 5 |
| 5552 | Yeast eIF4B binds to the head of the 40S ribosomal subunit and promotes mRNA recruitment through its N-terminal and internal repeat domains. <i>Rna</i> , 2013, 19, 191-207. | 1.6 | 66 |
| 5553 | Characterization of a Methionine Sulfoxide Reductase B from Tomato (<i>Solanum lycopersicum</i>), and Its Protecting Role in <i>Saccharomyces cerevisiae</i> . <i>Protein Journal</i> , 2013, 32, 39-47. | 0.7 | 6 |
| 5554 | Nucleotide-Specific Recognition of Iron-Responsive Elements by Iron Regulatory Protein 1. <i>Journal of Molecular Biology</i> , 2013, 425, 3301-3310. | 2.0 | 14 |
| 5555 | Functional analysis of an ATP-binding cassette transporter protein from <i>Aspergillus fumigatus</i> by heterologous expression in <i>Saccharomyces cerevisiae</i> . <i>Fungal Genetics and Biology</i> , 2013, 57, 85-91. | 0.9 | 14 |
| 5556 | Rice DEP1, encoding a highly cysteine-rich G protein β^3 subunit, confers cadmium tolerance on yeast cells and plants. <i>Journal of Experimental Botany</i> , 2013, 64, 4517-4527. | 2.4 | 64 |
| 5557 | Characterization of a trifunctional fatty acid desaturase from oleaginous filamentous fungus <i>Mortierella alpina</i> 1S-4 using a yeast expression system. <i>Journal of Bioscience and Bioengineering</i> , 2013, 116, 672-676. | 1.1 | 30 |
| 5558 | ABC transporters and cell wall proteins involved in organic solvent tolerance in <i>Saccharomyces cerevisiae</i> . <i>Journal of Biotechnology</i> , 2013, 165, 145-152. | 1.9 | 34 |
| 5559 | Enhancement of plasmid DNA transformation efficiencies in early stationary phase yeast cell cultures. <i>Yeast</i> , 2013, 30, 191-200. | 0.8 | 24 |
| 5560 | Arp2/3 complex ATP hydrolysis promotes lamellipodial actin network disassembly but is dispensable for assembly. <i>Journal of Cell Biology</i> , 2013, 200, 619-633. | 2.3 | 33 |
| 5561 | Mutations in prion-like domains in hnRNP2B1 and hnRNP1 cause multisystem proteinopathy and ALS. <i>Nature</i> , 2013, 495, 467-473. | 13.7 | 1,249 |
| 5562 | Redox-dependent stability of the β^3 -glutamylcysteine synthetase enzyme of <i>Escherichia coli</i> : a novel means of redox regulation. <i>Biochemical Journal</i> , 2013, 449, 783-794. | 1.7 | 8 |
| 5563 | Indoleamine 2,3-dioxygenases with very low catalytic activity are well conserved across kingdoms: IDOs of Basidiomycota. <i>Fungal Genetics and Biology</i> , 2013, 56, 98-106. | 0.9 | 26 |
| 5564 | Direct transfer of whole genomes from bacteria to yeast. <i>Nature Methods</i> , 2013, 10, 410-412. | 9.0 | 64 |
| 5565 | Molecular cloning and characterization of a cytochrome P450 in sanguinarine biosynthesis from <i>Eschscholzia californica</i> cells. <i>Phytochemistry</i> , 2013, 91, 100-108. | 1.4 | 64 |
| 5566 | Display of <i>Clostridium cellulovorans</i> xylose isomerase on the cell surface of <i>Saccharomyces cerevisiae</i> and its direct application to xylose fermentation. <i>Biotechnology Progress</i> , 2013, 29, 346-351. | 1.3 | 31 |
| 5567 | PP2A ^{Cdc55} regulates G ₁ cyclin stability. <i>Cell Cycle</i> , 2013, 12, 1201-1210. | 1.3 | 12 |

| # | ARTICLE | IF | CITATIONS |
|------|--|-----|-----------|
| 5568 | Yeast competence for exogenous DNA uptake: towards understanding its genetic component. Antonie Van Leeuwenhoek, 2013, 103, 1181-1207. | 0.7 | 15 |
| 5569 | Gene expression and function involved in polyol biosynthesis of <i>Trichosporonoides megachiliensis</i> under hyper-osmotic stress. Journal of Bioscience and Bioengineering, 2013, 115, 645-650. | 1.1 | 26 |
| 5570 | Soy peptides enhance yeast cell growth at low temperatures. Biotechnology Letters, 2013, 35, 375-382. | 1.1 | 16 |
| 5571 | Interaction with Cfd1 Increases the Kinetic Lability of FeS on the Nbp35 Scaffold. Journal of Biological Chemistry, 2013, 288, 23358-23367. | 1.6 | 35 |
| 5572 | Dynamic Large-Scale Chromosomal Rearrangements Fuel Rapid Adaptation in Yeast Populations. PLoS Genetics, 2013, 9, e1003232. | 1.5 | 106 |
| 5573 | Rad52 Sumoylation Prevents the Toxicity of Unproductive Rad51 Filaments Independently of the Anti-Recombinase Srs2. PLoS Genetics, 2013, 9, e1003833. | 1.5 | 20 |
| 5574 | Mono and Dual Cofactor Dependence of Human Cystathionine β -Synthase Enzyme Variants <i>In Vivo</i> and <i>In Vitro</i> . G3: Genes, Genomes, Genetics, 2013, 3, 1619-1628. | 0.8 | 7 |
| 5575 | Din7 and Mhr1 expression levels regulate double-strand-break-induced replication and recombination of mtDNA at ori5 in yeast. Nucleic Acids Research, 2013, 41, 5799-5816. | 6.5 | 28 |
| 5576 | Chemical Transformation of Yeast. Methods in Enzymology, 2013, 529, 311-320. | 0.4 | 11 |
| 5577 | Yeast telomere maintenance is globally controlled by programmed ribosomal frameshifting and the nonsense-mediated mRNA decay pathway. Translation, 2013, 1, e24418. | 2.9 | 27 |
| 5578 | RACK1/Asc1p, a Ribosomal Node in Cellular Signaling. Molecular and Cellular Proteomics, 2013, 12, 87-105. | 2.5 | 37 |
| 5579 | Mutual Cross Talk between the Regulators Hac1 of the Unfolded Protein Response and Gcn4 of the General Amino Acid Control of <i>Saccharomyces cerevisiae</i> . Eukaryotic Cell, 2013, 12, 1142-1154. | 3.4 | 17 |
| 5580 | TEAK: Topology Enrichment Analysis framework for detecting activated biological subpathways. Nucleic Acids Research, 2013, 41, 1425-1437. | 6.5 | 59 |
| 5581 | ABA Signaling in Guard Cells Entails a Dynamic Protein-Protein Interaction Relay from the PYL-RCAR Family Receptors to Ion Channels. Molecular Plant, 2013, 6, 528-538. | 3.9 | 139 |
| 5582 | Independence of symmetry breaking on Bem1-mediated autocatalytic activation of Cdc42. Journal of Cell Biology, 2013, 202, 1091-1106. | 2.3 | 61 |
| 5583 | Abscisic Acid-Dependent Regulation of Small Rubber Particle Protein Gene Expression in <i>Taraxacum brevicorniculatum</i> is Mediated by TbbZIP1. Plant and Cell Physiology, 2013, 54, 448-464. | 1.5 | 29 |
| 5584 | Transkingdom Genetic Transfer from <i>Escherichia coli</i> to <i>Saccharomyces cerevisiae</i> as a Simple Gene Introduction Tool. Applied and Environmental Microbiology, 2013, 79, 4393-4400. | 1.4 | 11 |
| 5585 | The Yeast Anaerobic Response Element AR1b Regulates Aerobic Antifungal Drug-dependent Sterol Gene Expression. Journal of Biological Chemistry, 2013, 288, 35466-35477. | 1.6 | 15 |

| # | ARTICLE | IF | CITATIONS |
|------|---|-----|-----------|
| 5586 | The 2-oxoglutarate supply exerts significant control on the lysine synthesis flux in <i>Saccharomyces cerevisiae</i> . <i>FEBS Journal</i> , 2013, 280, 5737-5749. | 2.2 | 5 |
| 5587 | Aspirin-induced apoptosis of yeast cells is associated with mitochondrial superoxide radical accumulation and NAD(P)H oxidation. <i>FEMS Yeast Research</i> , 2013, 13, 755-768. | 1.1 | 11 |
| 5588 | Transcription Factor Reb1p Regulates DGK1-encoded Diacylglycerol Kinase and Lipid Metabolism in <i>Saccharomyces cerevisiae</i> . <i>Journal of Biological Chemistry</i> , 2013, 288, 29124-29133. | 1.6 | 10 |
| 5589 | PAH1-encoded Phosphatidate Phosphatase Plays a Role in the Growth Phase- and Inositol-mediated Regulation of Lipid Synthesis in <i>Saccharomyces cerevisiae</i> . <i>Journal of Biological Chemistry</i> , 2013, 288, 35781-35792. | 1.6 | 57 |
| 5590 | Pepper Arginine Decarboxylase Is Required for Polyamine and β -Aminobutyric Acid Signaling in Cell Death and Defense Response. <i>Plant Physiology</i> , 2013, 162, 2067-2083. | 2.3 | 79 |
| 5591 | A Powerful Yeast Model to Investigate the Synergistic Interaction of β -Synuclein and Tau in Neurodegeneration. <i>PLoS ONE</i> , 2013, 8, e55848. | 1.1 | 27 |
| 5592 | Histatin 5 Resistance of <i>Candida glabrata</i> Can Be Reversed by Insertion of <i>Candida albicans</i> Polyamine Transporter-Encoding Genes DUR3 and DUR31. <i>PLoS ONE</i> , 2013, 8, e61480. | 1.1 | 27 |
| 5593 | Vanillin Inhibits Translation and Induces Messenger Ribonucleoprotein (mRNP) Granule Formation in <i>Saccharomyces cerevisiae</i> : Application and Validation of High-Content, Image-Based Profiling. <i>PLoS ONE</i> , 2013, 8, e61748. | 1.1 | 71 |
| 5594 | Identification and Functional Analysis of Delta-9 Desaturase, a Key Enzyme in PUFA Synthesis, Isolated from the Oleaginous Diatom <i>Fistulifera</i> . <i>PLoS ONE</i> , 2013, 8, e73507. | 1.1 | 20 |
| 5595 | The Yeast Shu Complex Utilizes Homologous Recombination Machinery for Error-free Lesion Bypass via Physical Interaction with a Rad51 Paralogue. <i>PLoS ONE</i> , 2013, 8, e81371. | 1.1 | 28 |
| 5596 | A novel method for genetic transformation of yeast cells using oligoelectrolyte polymeric nanoscale carriers. <i>BioTechniques</i> , 2013, 54, 35-43. | 0.8 | 15 |
| 5597 | Cloning and Characterization of a <i>Phragmites australis</i> Phytochelatin Synthase (PaPCS) and Achieving Cd Tolerance in Tall Fescue. <i>PLoS ONE</i> , 2014, 9, e103771. | 1.1 | 28 |
| 5598 | Topoisomerase 2 Is Dispensable for the Replication and Segregation of Small Yeast Artificial Chromosomes (YACs). <i>PLoS ONE</i> , 2014, 9, e104995. | 1.1 | 3 |
| 5599 | The Mre11-Rad50-Xrs2 Complex Is Required for Yeast DNA Postreplication Repair. <i>PLoS ONE</i> , 2014, 9, e109292. | 1.1 | 9 |
| 5600 | Investigating Acid Stress Response in Different <i>Saccharomyces</i> Strains. <i>Journal of Mycology</i> , 2014, 2014, 1-9. | 0.5 | 16 |
| 5601 | The Lap3p aminopeptidase is not solely responsible for bleomycin resistance in <i>Candida albicans</i> . <i>Journal of Yeast and Fungal Research</i> , 2014, 5, 39-49. | 0.6 | 0 |
| 5602 | Requirement of the spindle pole body for targeting and/or tethering proteins to the inner nuclear membrane. <i>Nucleus</i> , 2014, 5, 352-366. | 0.6 | 6 |
| 5603 | Activities of elements of the yeast β -factor precursor leader at different stages of somatropin secretion by <i>Saccharomyces cerevisiae</i> . <i>Applied Biochemistry and Microbiology</i> , 2014, 50, 829-834. | 0.3 | 3 |

| # | ARTICLE | IF | CITATIONS |
|------|---|-----|-----------|
| 5604 | Pepper mildew resistance locus O interacts with pepper calmodulin and suppresses Xanthomonas AvrBsT-triggered cell death and defense responses. <i>Planta</i> , 2014, 240, 827-839. | 1.6 | 22 |
| 5605 | <i>Saccharomyces cerevisiae</i> Gpi2, an accessory subunit of the enzyme catalyzing the first step of glycosylphosphatidylinositol (GPI) anchor biosynthesis, selectively complements some of the functions of its homolog in <i>Candida albicans</i> . <i>Glycoconjugate Journal</i> , 2014, 31, 497-507. | 1.4 | 8 |
| 5606 | Mutations in the X-linked Intellectual Disability Gene, zDHHC9, Alter Autopalmitoylation Activity by Distinct Mechanisms. <i>Journal of Biological Chemistry</i> , 2014, 289, 18582-18592. | 1.6 | 46 |
| 5607 | Ribosomes in the balance: structural equilibrium ensures translational fidelity and proper gene expression. <i>Nucleic Acids Research</i> , 2014, 42, 13384-13392. | 6.5 | 5 |
| 5608 | A Heavy Metal-Associated Protein (AchMA1) from the Halophyte, <i>Atriplex canescens</i> (Pursh) Nutt., Confers Tolerance to Iron and Other Abiotic Stresses When Expressed in <i>Saccharomyces cerevisiae</i> . <i>International Journal of Molecular Sciences</i> , 2014, 15, 14891-14906. | 1.8 | 31 |
| 5609 | Reversible disassembly of the yeast V-ATPase revisited under <i>in vivo</i> conditions. <i>Biochemical Journal</i> , 2014, 462, 185-197. | 1.7 | 47 |
| 5610 | Expression of an endoglucanase from <i>Tribolium castaneum</i> (TcEG1) in <i>Saccharomyces cerevisiae</i> . <i>Insect Science</i> , 2014, 21, 609-618. | 1.5 | 7 |
| 5611 | Amino acid residues required for Ctr1p-Gtr2p complex formation and its interactions with the Ego1p-Ego3p complex and TORC1 components in yeast. <i>Genes To Cells</i> , 2014, 19, 449-463. | 0.5 | 20 |
| 5612 | Serum albumin promotes ATP-binding cassette transporter-dependent sterol uptake in yeast. <i>FEMS Yeast Research</i> , 2014, 14, 1223-1233. | 1.1 | 16 |
| 5613 | Regulation of PKA activity by an autophosphorylation mechanism in <i>Saccharomyces cerevisiae</i> . <i>Biochemical Journal</i> , 2014, 462, 567-579. | 1.7 | 9 |
| 5614 | Ethylene production in relation to nitrogen metabolism in <i>Saccharomyces cerevisiae</i> . <i>FEMS Yeast Research</i> , 2014, 14, n/a-n/a. | 1.1 | 4 |
| 5615 | Negative control glucose dependent mediated by the PreS2 region on the translation efficiency of the reporter <i>Sh-bleomycin</i> gene in <i>Saccharomyces cerevisiae</i> . <i>FEMS Yeast Research</i> , 2014, 14, 357-363. | 1.1 | 0 |
| 5616 | The DExD/H-box ATPase Prp2p destabilizes and proofreads the catalytic RNA core of the spliceosome. <i>Rna</i> , 2014, 20, 282-294. | 1.6 | 37 |
| 5617 | C-Terminal Truncations of the <i>Saccharomyces cerevisiae</i> PMA1 H ⁺ -ATPase Have Major Impacts on Protein Conformation, Trafficking, Quality Control, and Function. <i>Eukaryotic Cell</i> , 2014, 13, 43-52. | 3.4 | 27 |
| 5618 | Impact of Photocatalysis on Fungal Cells: Depiction of Cellular and Molecular Effects on <i>Saccharomyces cerevisiae</i> . <i>Applied and Environmental Microbiology</i> , 2014, 80, 7527-7535. | 1.4 | 44 |
| 5619 | The analysis of <i>S. cerevisiae</i> cells deleted for mitotic cyclin Clb2 reveals a novel requirement of Sgs1 DNA helicase and Exonuclease 1 when replication forks break in the presence of alkylation damage. <i>Mutation Research - Fundamental and Molecular Mechanisms of Mutagenesis</i> , 2014, 769, 80-92. | 0.4 | 5 |
| 5620 | Template Switching During Break-Induced Replication Is Promoted by the Mph1 Helicase in <i>Saccharomyces cerevisiae</i> . <i>Genetics</i> , 2014, 196, 1017-1028. | 1.2 | 56 |
| 5621 | F-BAR domain protein Rga7 collaborates with Cdc15 and Imp2 to ensure proper cytokinesis in fission yeast. <i>Journal of Cell Science</i> , 2014, 127, 4146-58. | 1.2 | 26 |

| # | ARTICLE | IF | CITATIONS |
|------|--|-----|-----------|
| 5622 | Enhanced Adsorption and Recovery of Uranyl Ions by NikR Mutant-Displaying Yeast. <i>Biomolecules</i> , 2014, 4, 390-401. | 1.8 | 14 |
| 5623 | Bypass of the pre-60S ribosomal quality control as a pathway to oncogenesis. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 2014, 111, 5640-5645. | 3.3 | 71 |
| 5624 | Rps5-Rps16 communication is essential for efficient translation initiation in yeast <i>S. cerevisiae</i> . <i>Nucleic Acids Research</i> , 2014, 42, 8537-8555. | 6.5 | 14 |
| 5625 | DIPA-family coiled-coils bind conserved isoform-specific head domain of p120-catenin family: potential roles in hydrocephalus and heterotopia. <i>Molecular Biology of the Cell</i> , 2014, 25, 2592-2603. | 0.9 | 29 |
| 5626 | Osmoporation: a simple way to internalize hydrophilic molecules into yeast. <i>Applied Microbiology and Biotechnology</i> , 2014, 98, 1271-1280. | 1.7 | 32 |
| 5627 | Phylogenetic and comparative functional analysis of the cell-separation β -glucanase Agn1p in <i>Schizosaccharomyces</i> . <i>Microbiology (United Kingdom)</i> , 2014, 160, 1063-1074. | 0.7 | 5 |
| 5628 | Physical methods for genetic transformation of fungi and yeast. <i>Physics of Life Reviews</i> , 2014, 11, 184-203. | 1.5 | 50 |
| 5629 | Mutations in the Chromodomain-like Insertion of Translation Elongation Factor 3 Compromise Protein Synthesis through Reduced ATPase Activity. <i>Journal of Biological Chemistry</i> , 2014, 289, 4853-4860. | 1.6 | 12 |
| 5630 | Molecular and biochemical identification of inositol 1,3,4,5,6-pentakisphosphate 2-kinase encoding mRNA variants in castor bean (<i>Ricinus communis</i> L.) seeds. <i>Planta</i> , 2014, 239, 965-977. | 1.6 | 3 |
| 5631 | Rapid, efficient and precise allele replacement in the fission yeast <i>Schizosaccharomyces pombe</i> . <i>Current Genetics</i> , 2014, 60, 109-119. | 0.8 | 24 |
| 5632 | Pathogenesis-related protein 4b interacts with leucine-rich repeat protein 1 to suppress <i>PR4</i> -triggered cell death and defense response in pepper. <i>Plant Journal</i> , 2014, 77, 521-533. | 2.8 | 44 |
| 5633 | Heat treatment improves antigen-specific T cell activation after protein delivery by several but not all yeast genera. <i>Vaccine</i> , 2014, 32, 2591-2598. | 1.7 | 20 |
| 5634 | Chemical evaluation of white wines elaborated with a recombinant <i>Saccharomyces cerevisiae</i> strain overproducing mannoproteins. <i>Food Chemistry</i> , 2014, 147, 84-91. | 4.2 | 4 |
| 5635 | Pepper Suppressor of the G2 Allele of <i>skp1</i> Interacts with the Receptor-Like Cytoplasmic Kinase1 and Type III Effector AvrBsT and Promotes the Hypersensitive Cell Death Response in a Phosphorylation-Dependent Manner. <i>Plant Physiology</i> , 2014, 165, 76-91. | 2.3 | 32 |
| 5636 | Role of Arabidopsis UV RESISTANCE LOCUS 8 in Plant Growth Reduction under Osmotic Stress and Low Levels of UV-B. <i>Molecular Plant</i> , 2014, 7, 773-791. | 3.9 | 57 |
| 5637 | Novel Antifungal Drug Discovery Based on Targeting Pathways Regulating the Fungus-Conserved Upc2 Transcription Factor. <i>Antimicrobial Agents and Chemotherapy</i> , 2014, 58, 258-266. | 1.4 | 29 |
| 5638 | Improvement of the transformation efficiency of <i>Saccharomyces cerevisiae</i> by altering carbon sources in pre-culture. <i>Bioscience, Biotechnology and Biochemistry</i> , 2014, 78, 1090-1093. | 0.6 | 2 |
| 5639 | Translation Elongation Factor 1A Mutants with Altered Actin Bundling Activity Show Reduced Aminoacyl-tRNA Binding and Alter Initiation via eIF2 β Phosphorylation. <i>Journal of Biological Chemistry</i> , 2014, 289, 20928-20938. | 1.6 | 20 |

| # | ARTICLE | IF | CITATIONS |
|------|--|-----|-----------|
| 5640 | Cross-talk Phosphorylations by Protein Kinase C and Pho85p-Pho80p Protein Kinase Regulate Pah1p Phosphatidate Phosphatase Abundance in <i>Saccharomyces cerevisiae</i> . <i>Journal of Biological Chemistry</i> , 2014, 289, 18818-18830. | 1.6 | 44 |
| 5641 | The covalent bioconjugate of multiwalled carbon nanotube and amino- ϵ -modified linearized plasmid DNA for gene delivery. <i>Biotechnology Progress</i> , 2014, 30, 224-232. | 1.3 | 28 |
| 5642 | Improving industrial yeast strains: exploiting natural and artificial diversity. <i>FEMS Microbiology Reviews</i> , 2014, 38, 947-995. | 3.9 | 403 |
| 5643 | A Prominent Role for RCAR3-Mediated ABA Signaling in Response to <i>Pseudomonas syringae</i> pv. tomato DC3000 Infection in <i>Arabidopsis</i> . <i>Plant and Cell Physiology</i> , 2014, 55, 1691-1703. | 1.5 | 83 |
| 5644 | Identification and characterization of CYP79D16 and CYP71AN24 catalyzing the first and second steps in l-phenylalanine-derived cyanogenic glycoside biosynthesis in the Japanese apricot, <i>Prunus mume</i> Sieb. et Zucc.. <i>Plant Molecular Biology</i> , 2014, 86, 215-223. | 2.0 | 63 |
| 5645 | Secretion expression of SOD1 and its overlapping function with GSH in brewing yeast strain for better flavor and anti-aging ability. <i>Journal of Industrial Microbiology and Biotechnology</i> , 2014, 41, 1415-1424. | 1.4 | 10 |
| 5646 | Mechanism of action of flippase Drs2p in modulating GTP hydrolysis of Arl1p. <i>Journal of Cell Science</i> , 2014, 127, 2615-20. | 1.2 | 6 |
| 5647 | Over-expression of heat shock factor gene (<i>AtHsfA1d</i>) from <i>Arabidopsis thaliana</i> confers formaldehyde tolerance in tobacco. <i>Acta Physiologiae Plantarum</i> , 2014, 36, 1455-1462. | 1.0 | 6 |
| 5648 | Yeast Transformation by the LiAc/SS Carrier DNA/PEG Method. <i>Methods in Molecular Biology</i> , 2014, 1163, 33-44. | 0.4 | 215 |
| 5649 | A Mutation in the Catalytic Loop of Hsp90 Specifically Impairs ATPase Stimulation by Aha1p, But Not Hch1p. <i>Journal of Molecular Biology</i> , 2014, 426, 2379-2392. | 2.0 | 16 |
| 5650 | The Rad5 helicase activity is dispensable for error-free DNA post-replication repair. <i>DNA Repair</i> , 2014, 16, 74-83. | 1.3 | 26 |
| 5651 | Development of yeast reporter assay for screening specific ligands of retinoic acid and retinoid X receptor subtypes. <i>Journal of Pharmacological and Toxicological Methods</i> , 2014, 69, 245-252. | 0.3 | 18 |
| 5652 | Chromatin regulates DNA torsional energy via topoisomerase II ϵ -mediated relaxation of positive supercoils. <i>EMBO Journal</i> , 2014, 33, 1492-1501. | 3.5 | 35 |
| 5653 | Rice <i>MtN3/saliva/SWEET</i> gene family: Evolution, expression profiling, and sugar transport. <i>Journal of Integrative Plant Biology</i> , 2014, 56, 559-570. | 4.1 | 71 |
| 5654 | A fluorescence-based assay to monitor autopalmitylation of zDHHC proteins applicable to high-throughput screening. <i>Analytical Biochemistry</i> , 2014, 460, 1-8. | 1.1 | 30 |
| 5655 | Genetic Engineering Tools for <i>Saccharomyces cerevisiae</i> . , 2014, , 287-301. | | 2 |
| 5656 | Requirement of peroxiredoxin on the stationary phase of yeast cell growth. <i>Journal of Toxicological Sciences</i> , 2014, 39, 51-58. | 0.7 | 5 |
| 5657 | Human hepatic metabolism of the anti-osteoporosis drug eldecacitol involves sterol C4 α -methyl oxidase. <i>Pharmacology Research and Perspectives</i> , 2015, 3, e00120. | 1.1 | 6 |

| # | ARTICLE | IF | CITATIONS |
|------|---|-----|-----------|
| 5658 | Redox engineering by ectopic expression of glutamate dehydrogenase genes links NADPH availability and NADH oxidation with cold growth in <i>Saccharomyces cerevisiae</i> . <i>Microbial Cell Factories</i> , 2015, 14, 100. | 1.9 | 20 |
| 5659 | The laforin/malin E3-ubiquitin ligase complex ubiquitinates pyruvate kinase M1/M2. <i>BMC Biochemistry</i> , 2015, 16, 24. | 4.4 | 24 |
| 5660 | The Gene YALIOE20207g from <i>Yarrowia lipolytica</i> Encodes an N-Acetylglucosamine Kinase Implicated in the Regulated Expression of the Genes from the N-Acetylglucosamine Assimilatory Pathway. <i>PLoS ONE</i> , 2015, 10, e0122135. | 1.1 | 7 |
| 5661 | Metagenomics of an Alkaline Hot Spring in Galicia (Spain): Microbial Diversity Analysis and Screening for Novel Lipolytic Enzymes. <i>Frontiers in Microbiology</i> , 2015, 6, 1291. | 1.5 | 54 |
| 5662 | Fission Yeast Scp3 Potentially Maintains Microtubule Orientation through Bundling. <i>PLoS ONE</i> , 2015, 10, e0120109. | 1.1 | 1 |
| 5663 | Eukaryotic rRNA Modification by Yeast 5-Methylcytosine-Methyltransferases and Human Proliferation-Associated Antigen p120. <i>PLoS ONE</i> , 2015, 10, e0133321. | 1.1 | 73 |
| 5664 | Transcription Factor SomA Is Required for Adhesion, Development and Virulence of the Human Pathogen <i>Aspergillus fumigatus</i> . <i>PLoS Pathogens</i> , 2015, 11, e1005205. | 2.1 | 57 |
| 5665 | Xylose fermentation by <i>Saccharomyces cerevisiae</i> using endogenous xylose-assimilating genes. <i>Biotechnology Letters</i> , 2015, 37, 1623-1630. | 1.1 | 18 |
| 5666 | Transformation of <i>Saccharomyces cerevisiae</i> : Spheroplast Method. <i>Fungal Biology</i> , 2015, , 61-63. | 0.3 | 0 |
| 5667 | Application of Novel Polymeric Carrier of Plasmid DNA for Transformation of Yeast Cells. <i>Fungal Biology</i> , 2015, , 201-207. | 0.3 | 2 |
| 5668 | Protoplast Transformation for Genome Manipulation in Fungi. <i>Fungal Biology</i> , 2015, , 21-40. | 0.3 | 8 |
| 5669 | Nanoparticles-Based Delivery Systems in Plant Genetic Transformation. , 2015, , 209-239. | | 7 |
| 5670 | Two Duplicated Genes DDI2 and DDI3 in Budding Yeast Encode a Cyanamide Hydratase and Are Induced by Cyanamide. <i>Journal of Biological Chemistry</i> , 2015, 290, 12664-12675. | 1.6 | 15 |
| 5671 | Yeast red pigment modifies Amyloid beta growth in Alzheimer disease models in both <i>Saccharomyces cerevisiae</i> and <i>Drosophila melanogaster</i> . <i>Amyloid: the International Journal of Experimental and Clinical Investigation: the Official Journal of the International Society of Amyloidosis</i> , 2015, 22, 100-111. | 1.4 | 8 |
| 5672 | Pepper aldehyde dehydrogenase CaALDH1 interacts with <i>Xanthomonas</i> effector AvrBsT and promotes effector-triggered cell death and defence responses. <i>Journal of Experimental Botany</i> , 2015, 66, 3367-3380. | 2.4 | 26 |
| 5673 | Multiple abiotic stress tolerance of the transformants yeast cells and the transgenic <i>Arabidopsis</i> plants expressing a novel durum wheat catalase. <i>Plant Physiology and Biochemistry</i> , 2015, 97, 420-431. | 2.8 | 26 |
| 5674 | Overexpression of yeast thioredoxin TRX2 reduces p53-mediated cell death in yeast. <i>Applied Microbiology and Biotechnology</i> , 2015, 99, 8619-8628. | 1.7 | 5 |
| 5675 | Nitrate-dependent control of shoot K homeostasis by NPF7.3/NRT1.5 and SKOR in <i>Arabidopsis</i> . <i>Plant Physiology</i> , 2015, 169, pp.01152.2015. | 2.3 | 83 |

| # | ARTICLE | IF | CITATIONS |
|------|--|-----|-----------|
| 5676 | Addition of methionine and low cultivation temperatures increase palmitoleic acid production by engineered <i>Saccharomyces cerevisiae</i> . <i>Applied Microbiology and Biotechnology</i> , 2015, 99, 201-210. | 1.7 | 11 |
| 5677 | Overexpression of ubiquitinâ€like <i>HUB1</i> gene confers drought tolerance in perennial ryegrass. <i>Plant Biotechnology Journal</i> , 2015, 13, 689-699. | 4.1 | 25 |
| 5678 | Yeast-based methods to assess PTEN phosphoinositide phosphatase activity in vivo. <i>Methods</i> , 2015, 77-78, 172-179. | 1.9 | 13 |
| 5679 | ¹³ C-based metabolic flux analysis of <i>Saccharomyces cerevisiae</i> with a reduced Crabtree effect. <i>Journal of Bioscience and Bioengineering</i> , 2015, 120, 140-144. | 1.1 | 18 |
| 5680 | Effect of an Ala81His Mutation on the Met80 Loop Dynamics of Iso-1-cytochrome <i>c</i> . <i>Biochemistry</i> , 2015, 54, 1729-1742. | 1.2 | 13 |
| 5681 | High-throughput plasmid construction using homologous recombination in yeast: its mechanisms and application to protein production for X-ray crystallography. <i>Bioscience, Biotechnology and Biochemistry</i> , 2015, 79, 1-10. | 0.6 | 28 |
| 5683 | GLYCINEâ€RICH RNAâ€BINDING PROTEIN1 interacts with RECEPTORâ€LIKE CYTOPLASMIC PROTEIN KINASE1 and suppresses cell death and defense responses in pepper (<i>Capsicum annuum</i>). <i>New Phytologist</i> , 2015, 205, 786-800. | 3.5 | 28 |
| 5684 | Nonsense codon suppression in fission yeast due to mutations of tRNA ^{Ser.11} and translation release factor Sup35 (eRF3). <i>Current Genetics</i> , 2015, 61, 165-173. | 0.8 | 4 |
| 5685 | Efficient tryptophanâ€catabolizing activity is consistently conserved through evolution of TDO enzymes, but not IDO enzymes. <i>Journal of Experimental Zoology Part B: Molecular and Developmental Evolution</i> , 2015, 324, 128-140. | 0.6 | 26 |
| 5686 | The Heme Transport Capacity of LHR1 Determines the Extent of Virulence in <i>Leishmania amazonensis</i> . <i>PLoS Neglected Tropical Diseases</i> , 2015, 9, e0003804. | 1.3 | 27 |
| 5687 | Engineering of yeast pyruvate decarboxylase for enhanced selectivity towards carboligation. <i>Bioresource Technology</i> , 2015, 192, 90-96. | 4.8 | 9 |
| 5688 | Improvement of oxidized glutathione fermentation by thiol redox metabolism engineering in <i>Saccharomyces cerevisiae</i> . <i>Applied Microbiology and Biotechnology</i> , 2015, 99, 9771-9778. | 1.7 | 13 |
| 5689 | UV-B induction of the E3 ligase ARIADNE12 depends on CONSTITUTIVELY PHOTOMORPHOGENIC 1. <i>Plant Physiology and Biochemistry</i> , 2015, 93, 18-28. | 2.8 | 12 |
| 5690 | High Efficiency DNA Transformation of <i>Saccharomyces cerevisiae</i> with the LiAc/SS-DNA/PEG Method. <i>Fungal Biology</i> , 2015, , 177-186. | 0.3 | 8 |
| 5691 | Involvement of glycolysis/gluconeogenesis and signaling regulatory pathways in <i>Saccharomyces cerevisiae</i> biofilms during fermentation. <i>Frontiers in Microbiology</i> , 2015, 6, 139. | 1.5 | 36 |
| 5692 | Defining the cytosolic pathway of glutathione degradation in <i>Arabidopsis thaliana</i> : role of the ChaC/GCG family of Î³-glutamyl cyclotransferases as glutathione-degrading enzymes and AtLAP1 as the Cys-Gly peptidase. <i>Biochemical Journal</i> , 2015, 468, 73-85. | 1.7 | 54 |
| 5693 | Enhanced butanol production by eukaryotic <i>Saccharomyces cerevisiae</i> engineered to contain an improved pathway. <i>Bioscience, Biotechnology and Biochemistry</i> , 2015, 79, 314-320. | 0.6 | 24 |
| 5694 | <i>Arabidopsis NIP3;1</i> Plays an Important Role in Arsenic Uptake and Root-to-Shoot Translocation under Arsenite Stress Conditions. <i>Molecular Plant</i> , 2015, 8, 722-733. | 3.9 | 166 |

| # | ARTICLE | IF | CITATIONS |
|------|--|-----|-----------|
| 5695 | Rho4 interaction with exocyst and septins regulates cell separation in fission yeast. <i>Microbiology (United Kingdom)</i> , 2015, 161, 948-959. | 0.7 | 17 |
| 5696 | Diversification of Paralogous Î±-Isopropylmalate Synthases by Modulation of Feedback Control and Hetero-Oligomerization in <i>Saccharomyces cerevisiae</i> . <i>Eukaryotic Cell</i> , 2015, 14, 564-577. | 3.4 | 29 |
| 5697 | Analysis of the essentiality of ROM2 genes in the pathogenic yeasts <i>Candida glabrata</i> and <i>Candida albicans</i> using temperature-sensitive mutants. <i>Journal of Applied Microbiology</i> , 2015, 118, 851-863. | 1.4 | 4 |
| 5698 | <i>Arabidopsis</i> abscisic acid receptors play an important role in disease resistance. <i>Plant Molecular Biology</i> , 2015, 88, 313-324. | 2.0 | 40 |
| 5699 | Integrated Automation for Continuous High-Throughput Synthetic Chromosome Assembly and Transformation to Identify Improved Yeast Strains for Industrial Production of Biofuels and Bio-based Chemicals. <i>Fungal Biology</i> , 2015, , 183-199. | 0.3 | 0 |
| 5700 | Cooperation between COA6 and SCO2 in COX2 Maturation during Cytochrome c Oxidase Assembly Links Two Mitochondrial Cardiomyopathies. <i>Cell Metabolism</i> , 2015, 21, 823-833. | 7.2 | 68 |
| 5701 | A ginseng PgTIP1 gene whose protein biological activity related to Ser128 residue confers faster growth and enhanced salt stress tolerance in <i>Arabidopsis</i> . <i>Plant Science</i> , 2015, 234, 74-85. | 1.7 | 12 |
| 5702 | Limited Portability of G-Patch Domains in Regulators of the Prp43 RNA Helicase Required for Pre-mRNA Splicing and Ribosomal RNA Maturation in <i>Saccharomyces cerevisiae</i> . <i>Genetics</i> , 2015, 200, 135-147. | 1.2 | 14 |
| 5703 | A Second Las17 Monomeric Actinâ€œBinding Motif Functions in Arp2/3â€œDependent Actin Polymerization During Endocytosis. <i>Traffic</i> , 2015, 16, 379-397. | 1.3 | 14 |
| 5704 | Inhibition of peroxisome fission, but not mitochondrial fission, increases yeast chronological lifespan. <i>Cell Cycle</i> , 2015, 14, 1698-1703. | 1.3 | 24 |
| 5705 | Pepper Heat Shock Protein 70a Interacts with the Type III Effector AvrBsT and Triggers Plant Cell Death and Immunity Å. <i>Plant Physiology</i> , 2015, 167, 307-322. | 2.3 | 61 |
| 5706 | Protein Kinase Ymr291w/Tda1 Is Essential for Glucose Signaling in <i>Saccharomyces cerevisiae</i> on the Level of Hexokinase Isoenzyme Schxk2 Phosphorylation*. <i>Journal of Biological Chemistry</i> , 2015, 290, 6243-6255. | 1.6 | 18 |
| 5707 | Altered Lipid Synthesis by Lack of Yeast Pah1 Phosphatidate Phosphatase Reduces Chronological Life Span. <i>Journal of Biological Chemistry</i> , 2015, 290, 25382-25394. | 1.6 | 47 |
| 5708 | Assembly and dynamics of the U4/U6 di-snRNP by single-molecule FRET. <i>Nucleic Acids Research</i> , 2015, 43, 10963-10974. | 6.5 | 35 |
| 5709 | Unlocking the steric gate of DNA polymerase Î· leads to increased genomic instability in <i>Saccharomyces cerevisiae</i> . <i>DNA Repair</i> , 2015, 35, 1-12. | 1.3 | 22 |
| 5710 | PKA-chromatin association at stress responsive target genes from <i>Saccharomyces cerevisiae</i> . <i>Biochimica Et Biophysica Acta - Gene Regulatory Mechanisms</i> , 2015, 1849, 1329-1339. | 0.9 | 11 |
| 5711 | The Intracellular Domain of the Frazzled/DCC Receptor Is a Transcription Factor Required for Commissural Axon Guidance. <i>Neuron</i> , 2015, 87, 751-763. | 3.8 | 44 |
| 5712 | Ribosomal protein uS19 mutants reveal its role in coordinating ribosome structure and function. <i>Translation</i> , 2015, 3, e1117703. | 2.9 | 13 |

| # | ARTICLE | IF | CITATIONS |
|------|--|-----|-----------|
| 5713 | Isolation of the GFA1 gene encoding glucosamine-6-phosphate synthase of <i>Sporothrix schenckii</i> and its expression in <i>Saccharomyces cerevisiae</i> . <i>Protein Expression and Purification</i> , 2015, 110, 57-64. | 0.6 | 5 |
| 5714 | Important Role of Catalase in the Cellular Response of the Budding Yeast <i>Saccharomyces cerevisiae</i> Exposed to Ionizing Radiation. <i>Current Microbiology</i> , 2015, 70, 404-407. | 1.0 | 17 |
| 5715 | Transcriptional and antioxidative responses to endogenous polyunsaturated fatty acid accumulation in yeast. <i>Molecular and Cellular Biochemistry</i> , 2015, 399, 27-37. | 1.4 | 12 |
| 5716 | African swine fever virus ORF P1192R codes for a functional type II DNA topoisomerase. <i>Virology</i> , 2015, 474, 82-93. | 1.1 | 27 |
| 5717 | <i>Cryptococcus neoformans</i> Yap1 is required for normal fluconazole and oxidative stress resistance. <i>Fungal Genetics and Biology</i> , 2015, 74, 1-9. | 0.9 | 32 |
| 5718 | Application of Carbon Nanotubes for Plant Genetic Transformation. <i>Springer Proceedings in Physics</i> , 2015, , 233-255. | 0.1 | 5 |
| 5719 | Improved ethanol production from xylose in the presence of acetic acid by the overexpression of the HAA1 gene in <i>Saccharomyces cerevisiae</i> . <i>Journal of Bioscience and Bioengineering</i> , 2015, 119, 297-302. | 1.1 | 47 |
| 5720 | Identification of Protein Palmitoylation Inhibitors from a Scaffold Ranking Library. <i>Combinatorial Chemistry and High Throughput Screening</i> , 2016, 19, 262-274. | 0.6 | 19 |
| 5721 | Presence of an isoform of H ⁺ -pyrophosphatase located in the alveolar sacs of a scuticociliate parasite of turbot: physiological consequences. <i>Parasitology</i> , 2016, 143, 576-587. | 0.7 | 3 |
| 5722 | Overexpression of a Water-Forming NADH Oxidase Improves the Metabolism and Stress Tolerance of <i>Saccharomyces cerevisiae</i> in Aerobic Fermentation. <i>Frontiers in Microbiology</i> , 2016, 7, 1427. | 1.5 | 8 |
| 5723 | Characteristics of Three Thioredoxin Genes and Their Role in Chilling Tolerance of Harvested Banana Fruit. <i>International Journal of Molecular Sciences</i> , 2016, 17, 1526. | 1.8 | 17 |
| 5724 | Autophagy-Associated Protein SmATG12 Is Required for Fruiting-Body Formation in the Filamentous Ascomycete <i>Sordaria macrospora</i> . <i>PLoS ONE</i> , 2016, 11, e0157960. | 1.1 | 10 |
| 5725 | Ubiquitin-Dependent Modification of Skeletal Muscle by the Parasitic Nematode, <i>Trichinella spiralis</i> . <i>PLoS Pathogens</i> , 2016, 12, e1005977. | 2.1 | 24 |
| 5726 | Molecular Cloning and Functional Characterization of Two <i>Brachypodium distachyon</i> UBC13 Genes Whose Products Promote K63-Linked Polyubiquitination. <i>Frontiers in Plant Science</i> , 2015, 6, 1222. | 1.7 | 10 |
| 5727 | Three <i>Brachypodium distachyon</i> Uev1s Promote Ubc13-Mediated Lys63-Linked Polyubiquitination and Confer Different Functions. <i>Frontiers in Plant Science</i> , 2016, 7, 1551. | 1.7 | 16 |
| 5728 | Detection of Acyl-CoA Derivatized with Butylamide for <i>in vitro</i> Fatty Acid Desaturase Assay. <i>Journal of Oleo Science</i> , 2016, 65, 161-167. | 0.6 | 1 |
| 5729 | Trypanosomatid parasites rescue heme from endocytosed hemoglobin through lysosomal HRG transporters. <i>Molecular Microbiology</i> , 2016, 101, 895-908. | 1.2 | 29 |
| 5730 | The catalytic subunit of the first mannosyltransferase in the GPI biosynthetic pathway affects growth, cell wall integrity and hyphal morphogenesis in <i>Candida albicans</i> . <i>Yeast</i> , 2016, 33, 365-383. | 0.8 | 6 |

| # | ARTICLE | IF | CITATIONS |
|------|---|-----|-----------|
| 5731 | Identification of Genes in <i>Candida glabrata</i> Conferring Altered Responses to Caspofungin, a Cell Wall Synthesis Inhibitor. <i>G3: Genes, Genomes, Genetics</i> , 2016, 6, 2893-2907. | 0.8 | 19 |
| 5732 | Gene Function Analysis in the Ubiquitous Human Commensal and Pathogen <i>Malassezia</i> Genus. <i>MBio</i> , 2016, 7, . | 1.8 | 44 |
| 5733 | Phosphorylation of Dgk1 Diacylglycerol Kinase by Casein Kinase II Regulates Phosphatidic Acid Production in <i>Saccharomyces cerevisiae</i> . <i>Journal of Biological Chemistry</i> , 2016, 291, 26455-26467. | 1.6 | 20 |
| 5734 | Breeding of high malate-producing diploid sake yeast with a homozygous mutation in the <i>VID24</i> gene. <i>Journal of the Institute of Brewing</i> , 2016, 122, 605-611. | 0.8 | 4 |
| 5735 | <i>Rga6</i> is a fission yeast Rho GAP involved in Cdc42 regulation of polarized growth. <i>Molecular Biology of the Cell</i> , 2016, 27, 1524-1535. | 0.9 | 38 |
| 5736 | A water-forming NADH oxidase regulates metabolism in anaerobic fermentation. <i>Biotechnology for Biofuels</i> , 2016, 9, 103. | 6.2 | 15 |
| 5737 | Effect of aluminum stress on the expression of calmodulin and the role of \hat{A} calmodulin in aluminum tolerance. <i>Journal of Bioscience and Bioengineering</i> , 2016, 122, 558-562. | 1.1 | 9 |
| 5738 | Functional characterization and inhibition of the type II DNA topoisomerase coded by African swine fever virus. <i>Virology</i> , 2016, 493, 209-216. | 1.1 | 22 |
| 5739 | A Novel Inositol Pyrophosphate Phosphatase in <i>Saccharomyces cerevisiae</i> . <i>Journal of Biological Chemistry</i> , 2016, 291, 6772-6783. | 1.6 | 55 |
| 5740 | Structural elements that govern Sec14-like PTP sensitivities to potent small molecule inhibitors. <i>Journal of Lipid Research</i> , 2016, 57, 650-662. | 2.0 | 15 |
| 5741 | Overexpression of Sly41 suppresses COPII vesicle tethering deficiencies by elevating intracellular calcium levels. <i>Molecular Biology of the Cell</i> , 2016, 27, 1635-1649. | 0.9 | 4 |
| 5742 | Oral Vaccine Development by Molecular Display Methods Using Microbial Cells. <i>Methods in Molecular Biology</i> , 2016, 1404, 497-509. | 0.4 | 11 |
| 5743 | Responses of transgenic <i>Arabidopsis</i> plants and recombinant yeast cells expressing a novel durum wheat manganese superoxide dismutase TdMnSOD to various abiotic stresses. <i>Journal of Plant Physiology</i> , 2016, 198, 56-68. | 1.6 | 42 |
| 5744 | Modeling Metazoan Apoptotic Pathways in Yeast. <i>Methods in Molecular Biology</i> , 2016, 1419, 161-183. | 0.4 | 5 |
| 5745 | Phosphorylation of Yeast Pah1 Phosphatidate Phosphatase by Casein Kinase II Regulates Its Function in Lipid Metabolism. <i>Journal of Biological Chemistry</i> , 2016, 291, 9974-9990. | 1.6 | 41 |
| 5746 | Role of H ⁺ -pyrophosphatase activity in the regulation of intracellular pH in a scuticociliate parasite of turbot: Physiological effects. <i>Experimental Parasitology</i> , 2016, 169, 59-68. | 0.5 | 6 |
| 5747 | Heme acquisition in the parasitic filarial nematode <i>Brugia malayi</i> . <i>FASEB Journal</i> , 2016, 30, 3501-3514. | 0.2 | 20 |
| 5748 | <i>Botrytis cinerea</i> can import and utilize nucleosides in salvage and catabolism and BcENT functions as high affinity nucleoside transporter. <i>Fungal Biology</i> , 2016, 120, 904-916. | 1.1 | 5 |

| # | ARTICLE | IF | CITATIONS |
|------|--|-----|-----------|
| 5749 | Identification of SUMO modification sites in the base excision repair protein, Ntg1. <i>DNA Repair</i> , 2016, 48, 51-62. | 1.3 | 8 |
| 5750 | Fluorescence Anisotropy as a Tool to Study Protein-protein Interactions. <i>Journal of Visualized Experiments</i> , 2016, , . | 0.2 | 23 |
| 5751 | Transformation of Probiotic Yeast and Their Recovery from Gastrointestinal Immune Tissues Following Oral Gavage in Mice. <i>Journal of Visualized Experiments</i> , 2016, , e53453. | 0.2 | 2 |
| 5752 | Modulation of gluconeogenesis and lipid production in an engineered oleaginous <i>Saccharomyces cerevisiae</i> transformant. <i>Applied Microbiology and Biotechnology</i> , 2016, 100, 8147-8157. | 1.7 | 4 |
| 5753 | Involvement of budding yeast Rad5 in translesion DNA synthesis through physical interaction with Rev1. <i>Nucleic Acids Research</i> , 2016, 44, 5231-5245. | 6.5 | 71 |
| 5754 | LmABC3, an atypical mitochondrial ABC transporter essential for <i>Leishmania major</i> virulence, acts in heme and cytosolic iron/sulfur clusters biogenesis. <i>Parasites and Vectors</i> , 2016, 9, 7. | 1.0 | 22 |
| 5755 | Direct ethanol fermentation of the algal storage polysaccharide laminarin with an optimized combination of engineered yeasts. <i>Journal of Biotechnology</i> , 2016, 231, 129-135. | 1.9 | 30 |
| 5756 | Pepper protein phosphatase type 2C, CaADIP1 and its interacting partner CaRLP1 antagonistically regulate ABA signalling and drought response. <i>Plant, Cell and Environment</i> , 2016, 39, 1559-1575. | 2.8 | 70 |
| 5757 | The complete chemical structure of <i>Saccharomyces cerevisiae</i> rRNA: partial pseudouridylation of U2345 in 25S rRNA by snoRNA snR9. <i>Nucleic Acids Research</i> , 2016, 44, 8951-8961. | 6.5 | 87 |
| 5758 | Cytokine-dependent activation of JAK-STAT pathway in <i>Saccharomyces cerevisiae</i> . <i>Biotechnology and Bioengineering</i> , 2016, 113, 1796-1804. | 1.7 | 3 |
| 5759 | Surface-modified yeast cells: A novel eukaryotic carrier for oral application. <i>Journal of Controlled Release</i> , 2016, 224, 1-7. | 4.8 | 18 |
| 5760 | Gbp2 interacts with THO/TREX through a novel type of RRM domain. <i>Nucleic Acids Research</i> , 2016, 44, 437-448. | 6.5 | 26 |
| 5761 | Unfolded protein response regulates yeast small GTPase Arl1p activation at late Golgi via phosphorylation of Arf GEF Syt1p. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 2016, 113, E1683-E1690. | 3.3 | 13 |
| 5762 | Enhancement of malate-production and increase in sensitivity to dimethyl succinate by mutation of the VID24 gene in <i>Saccharomyces cerevisiae</i> . <i>Journal of Bioscience and Bioengineering</i> , 2016, 121, 665-671. | 1.1 | 19 |
| 5763 | RipAY, a Plant Pathogen Effector Protein, Exhibits Robust ³ -Glutamyl Cyclotransferase Activity When Stimulated by Eukaryotic Thioredoxins. <i>Journal of Biological Chemistry</i> , 2016, 291, 6813-6830. | 1.6 | 50 |
| 5764 | Near-freezing effects on the proteome of industrial yeast strains of <i>Saccharomyces cerevisiae</i> . <i>Journal of Biotechnology</i> , 2016, 221, 70-77. | 1.9 | 9 |
| 5765 | A Droplet Microfluidic Platform for Automating Genetic Engineering. <i>ACS Synthetic Biology</i> , 2016, 5, 426-433. | 1.9 | 63 |
| 5766 | Identification of amino acid residues that determine the substrate specificity of mammalian membrane-bound front-end fatty acid desaturases. <i>Journal of Lipid Research</i> , 2016, 57, 89-99. | 2.0 | 22 |

| # | ARTICLE | IF | CITATIONS |
|------|--|-----|-----------|
| 5767 | Yeast <i>MRX</i> deletions have short chronological life span and more triacylglycerols. <i>FEMS Yeast Research</i> , 2016, 16, fov109. | 1.1 | 4 |
| 5768 | Enantioselective Synthesis of Vicinal (<i>R</i> , <i>R</i>)-Diols by <i>Saccharomyces cerevisiae</i> Butanediol Dehydrogenase. <i>Applied and Environmental Microbiology</i> , 2016, 82, 1706-1721. | 1.4 | 14 |
| 5769 | Engineered yeast whole-cell biocatalyst for direct degradation of alginate from macroalgae and production of non-commercialized useful monosaccharide from alginate. <i>Applied Microbiology and Biotechnology</i> , 2016, 100, 1723-1732. | 1.7 | 37 |
| 5770 | Ceramide signals for initiation of yeast mating-specific cell cycle arrest. <i>Cell Cycle</i> , 2016, 15, 441-454. | 1.3 | 12 |
| 5771 | Display of phytase on the cell surface of <i>Saccharomyces cerevisiae</i> to degrade phytate phosphorus and improve bioethanol production. <i>Applied Microbiology and Biotechnology</i> , 2016, 100, 2449-2458. | 1.7 | 18 |
| 5772 | Development of a system for multicopy gene integration in <i>Saccharomyces cerevisiae</i> . <i>Journal of Microbiological Methods</i> , 2016, 120, 44-49. | 0.7 | 8 |
| 5773 | Coexpression of human perforin improves yeast-mediated delivery of DNA and mRNA to mammalian antigen-presenting cells. <i>Gene Therapy</i> , 2016, 23, 103-107. | 2.3 | 6 |
| 5774 | Immature Core protein of hepatitis C virus induces an unfolded protein response through inhibition of <i>ERAD</i> in a yeast model system. <i>Genes To Cells</i> , 2017, 22, 160-173. | 0.5 | 5 |
| 5775 | A <i>Cdc48</i> Retrochaperone Function Is Required for the Solubility of Retrotranslocated, Integral Membrane Endoplasmic Reticulum-associated Degradation (ERAD-M) Substrates. <i>Journal of Biological Chemistry</i> , 2017, 292, 3112-3128. | 1.6 | 38 |
| 5776 | Involvement of <i>Chlamydomonas</i> DNA damage tolerance gene <i>UBC2</i> in lipid accumulation. <i>Algal Research</i> , 2017, 22, 148-159. | 2.4 | 13 |
| 5777 | Acquisition of accurate data from intramolecular quenched fluorescence protease assays. <i>Analytical Biochemistry</i> , 2017, 522, 30-36. | 1.1 | 7 |
| 5778 | <i>ChaC2</i> , an Enzyme for Slow Turnover of Cytosolic Glutathione. <i>Journal of Biological Chemistry</i> , 2017, 292, 638-651. | 1.6 | 56 |
| 5779 | A Ribosomopathy Reveals Decoding Defective Ribosomes Driving Human Dysmorphism. <i>American Journal of Human Genetics</i> , 2017, 100, 506-522. | 2.6 | 69 |
| 5780 | The Pepper RING-Type E3 Ligase <i>CaAIRF1</i> Regulates ABA and Drought Signaling via <i>CaADIP1</i> Protein Phosphatase Degradation. <i>Plant Physiology</i> , 2017, 173, 2323-2339. | 2.3 | 56 |
| 5781 | Enhanced direct ethanol production by cofactor optimization of cell surface-displayed xylose isomerase in yeast. <i>Biotechnology Progress</i> , 2017, 33, 1068-1076. | 1.3 | 12 |
| 5782 | <i>CTT1</i> overexpression increases the replicative lifespan of MMS-sensitive <i>Saccharomyces cerevisiae</i> deficient in <i>KSP1</i> . <i>Mechanisms of Ageing and Development</i> , 2017, 164, 27-36. | 2.2 | 11 |
| 5783 | Differential Control of Asexual Development and Sterigmatocystin Biosynthesis by a Novel Regulator in <i>Aspergillus nidulans</i> . <i>Scientific Reports</i> , 2017, 7, 46340. | 1.6 | 10 |
| 5784 | <i>Schizosaccharomyces pombe</i> <i>MutS</i> and <i>MutL</i> Maintain Stability of Tetra-Nucleotide Repeats and <i>Msh3</i> of Hepta-Nucleotide Repeats. <i>G3: Genes, Genomes, Genetics</i> , 2017, 7, 1463-1473. | 0.8 | 3 |

| # | ARTICLE | IF | CITATIONS |
|------|--|-----|-----------|
| 5785 | Diversification of the kinetic properties of yeast NADP ⁺ glutamate ⁺ dehydrogenase isozymes proceeds independently of their evolutionary origin. <i>MicrobiologyOpen</i> , 2017, 6, e00419. | 1.2 | 9 |
| 5786 | New data on flatfish scuticociliatosis reveal that <i>Miamiensis avidus</i> and <i>Philasterides dicentrarchi</i> are different species. <i>Parasitology</i> , 2017, 144, 1394-1411. | 0.7 | 19 |
| 5787 | The Arl3 and Arl1 GTPases cooperate with Cog8 to regulate selective autophagy via Atg9 trafficking. <i>Traffic</i> , 2017, 18, 580-589. | 1.3 | 15 |
| 5788 | Heavy metal ATPase 3 (HMA3) confers cadmium hypertolerance on the cadmium/zinc hyperaccumulator <i>Sedum plumbizincicola</i> . <i>New Phytologist</i> , 2017, 215, 687-698. | 3.5 | 191 |
| 5790 | Glutathione production from mannan-based bioresource by mannanase/mannosidase expressing <i>Saccharomyces cerevisiae</i> . <i>Bioresource Technology</i> , 2017, 245, 1400-1406. | 4.8 | 15 |
| 5791 | A Ras GTPase associated protein is involved in the phototropic and circadian photobiology responses in fungi. <i>Scientific Reports</i> , 2017, 7, 44790. | 1.6 | 22 |
| 5792 | A novel cytochrome P450, CYP3201B1, is involved in mandelonitrile biosynthesis in a cyanogenic millipede. <i>FEBS Open Bio</i> , 2017, 7, 335-347. | 1.0 | 26 |
| 5793 | Yeast-mediated mRNA delivery polarizes immuno-suppressive macrophages towards an immuno-stimulatory phenotype. <i>European Journal of Pharmaceutics and Biopharmaceutics</i> , 2017, 117, 1-13. | 2.0 | 18 |
| 5794 | Mitochondrial nuclear evolution leads to hybrid incompatibility through pentatricopeptide repeat proteins. <i>EMBO Reports</i> , 2017, 18, 87-101. | 2.0 | 41 |
| 5795 | High-Throughput Analysis of Mammalian Receptor Tyrosine Kinase Activation in Yeast Cells. <i>Methods in Molecular Biology</i> , 2017, 1487, 35-52. | 0.4 | 2 |
| 5796 | General Amino Acid Control and 14-3-3 Proteins Bmh1/2 Are Required for Nitrogen Catabolite Repression-Sensitive Regulation of Gln3 and Gat1 Localization. <i>Genetics</i> , 2017, 205, 633-655. | 1.2 | 27 |
| 5797 | Inappropriate translation inhibition and P-body formation cause cold-sensitivity in tryptophan-auxotroph yeast mutants. <i>Biochimica Et Biophysica Acta - Molecular Cell Research</i> , 2017, 1864, 314-323. | 1.9 | 3 |
| 5798 | Lanosterol synthase-like is involved with differential accumulation of steroidal glycoalkaloids in potato. <i>Planta</i> , 2017, 246, 1189-1202. | 1.6 | 19 |
| 5799 | Rice inositol polyphosphate kinase gene (OsIPK2), a putative new player of gibberellic acid signaling, involves in modulation of shoot elongation and fertility. <i>Plant Cell, Tissue and Organ Culture</i> , 2017, 131, 471-482. | 1.2 | 4 |
| 5800 | Diversification of Transcriptional Regulation Determines Subfunctionalization of Paralogous Branched Chain Aminotransferases in the Yeast <i>Saccharomyces cerevisiae</i> . <i>Genetics</i> , 2017, 207, 975-991. | 1.2 | 27 |
| 5801 | Targeted gene replacement at the <i>URA3</i> locus of the basidiomycetous yeast <i>Pseudozyma antarctica</i> and its transformation using lithium acetate treatment. <i>Yeast</i> , 2017, 34, 483-494. | 0.8 | 10 |
| 5802 | Insights into the molecular basis for substrate binding and specificity of the fungal cystine transporter CgCYN1. <i>Biochimica Et Biophysica Acta - Biomembranes</i> , 2017, 1859, 2259-2268. | 1.4 | 8 |
| 5803 | Mating response and construction of heterothallic strains of the fission yeast <i>Schizosaccharomyces octosporus</i> . <i>FEMS Yeast Research</i> , 2017, 17, . | 1.1 | 14 |

| # | ARTICLE | IF | CITATIONS |
|------|--|-----|-----------|
| 5804 | Regulation of Small Mitochondrial DNA Replicative Advantage by Ribonucleotide Reductase in <i>Saccharomyces cerevisiae</i> . <i>G3: Genes, Genomes, Genetics</i> , 2017, 7, 3083-3090. | 0.8 | 10 |
| 5805 | Novel function of a dynein light chain in actin assembly during clathrin-mediated endocytosis. <i>Journal of Cell Biology</i> , 2017, 216, 2565-2580. | 2.3 | 14 |
| 5806 | Unravelling the Function of a Bacterial Effector from a Non-cultivable Plant Pathogen Using a Yeast Two-hybrid Screen. <i>Journal of Visualized Experiments</i> , 2017, , . | 0.2 | 6 |
| 5807 | A TUBB6 mutation is associated with autosomal dominant non-progressive congenital facial palsy, bilateral ptosis and velopharyngeal dysfunction. <i>Human Molecular Genetics</i> , 2017, 26, 4055-4066. | 1.4 | 13 |
| 5808 | A link between very long chain fatty acid elongation and mating-specific yeast cell cycle arrest. <i>Cell Cycle</i> , 2017, 16, 2192-2203. | 1.3 | 5 |
| 5809 | Elp3 and Dph3 of <i>Schizosaccharomyces pombe</i> mediate cellular stress responses through tRNA ^{Lys} UUU modifications. <i>Scientific Reports</i> , 2017, 7, 7225. | 1.6 | 8 |
| 5810 | The golgin protein Coy1 functions in intra-Golgi retrograde transport and interacts with the COG complex and Golgi SNAREs. <i>Molecular Biology of the Cell</i> , 2017, 28, 2686-2700. | 0.9 | 12 |
| 5811 | A conserved tryptophan within the WRDPLVDID domain of yeast Pah1 phosphatidate phosphatase is required for its <i>in vivo</i> function in lipid metabolism. <i>Journal of Biological Chemistry</i> , 2017, 292, 19580-19589. | 1.6 | 17 |
| 5812 | ORP-Mediated ER Contact with Endocytic Sites Facilitates Actin Polymerization. <i>Developmental Cell</i> , 2017, 43, 588-602.e6. | 3.1 | 41 |
| 5813 | Early emergence of negative regulation of the tyrosine kinase Src by the C-terminal Src kinase. <i>Journal of Biological Chemistry</i> , 2017, 292, 18518-18529. | 1.6 | 7 |
| 5814 | Wheat Vacuolar Iron Transporter TaVIT2 Transports Fe and Mn and Is Effective for Biofortification. <i>Plant Physiology</i> , 2017, 174, 2434-2444. | 2.3 | 206 |
| 5815 | Nucleolin directly mediates Epstein-Barr virus immune evasion through binding to G-quadruplexes of EBNA1 mRNA. <i>Nature Communications</i> , 2017, 8, 16043. | 5.8 | 94 |
| 5816 | Enzymatic improvement of mitochondrial thiol oxidase Erv1 for oxidized glutathione fermentation by <i>Saccharomyces cerevisiae</i> . <i>Microbial Cell Factories</i> , 2017, 16, 44. | 1.9 | 3 |
| 5817 | Protein-Protein Interactions: Yeast Two-Hybrid System. <i>Methods in Molecular Biology</i> , 2017, 1615, 177-187. | 0.4 | 30 |
| 5818 | Isolation of conditional mutations in genes essential for viability of <i>Cryptococcus neoformans</i> . <i>Current Genetics</i> , 2017, 63, 519-530. | 0.8 | 23 |
| 5819 | Development of a sufficient and effective procedure for transformation of an oleaginous yeast, <i>Rhodospiridium toruloides</i> DMKU3-TK16. <i>Current Genetics</i> , 2017, 63, 359-371. | 0.8 | 21 |
| 5820 | Heterologous expression of carnation Italian ringspot virus p36 protein enhances necrotic cell death in response to acetic acid in <i>Saccharomyces cerevisiae</i> . <i>Mechanisms of Ageing and Development</i> , 2017, 161, 255-261. | 2.2 | 2 |
| 5821 | A Simple Fluorescence-based Reporter Assay to Identify Cellular Components Required for Ricin Toxin A Chain (RTA) Trafficking in Yeast. <i>Journal of Visualized Experiments</i> , 2017, , . | 0.2 | 1 |

| # | ARTICLE | IF | CITATIONS |
|------|--|-----|-----------|
| 5822 | Homozygous EEF1A2 mutation causes dilated cardiomyopathy, failure to thrive, global developmental delay, epilepsy and early death. <i>Human Molecular Genetics</i> , 2017, 26, 3545-3552. | 1.4 | 27 |
| 5823 | Expression of K1 Toxin Derivatives in <i>Saccharomyces cerevisiae</i> Mimics Treatment with Exogenous Toxin and Provides a Useful Tool for Elucidating K1 Mechanisms of Action and Immunity. <i>Toxins</i> , 2017, 9, 345. | 1.5 | 15 |
| 5824 | Principles of Genetic Engineering. , 2017, , 81-127. | | 3 |
| 5825 | Purification and characterisation of the yeast plasma membrane ATP binding cassette transporter Pdr11p. <i>PLoS ONE</i> , 2017, 12, e0184236. | 1.1 | 9 |
| 5826 | Characterization of four rice UEV1 genes required for Lys63-linked polyubiquitination and distinct functions. <i>BMC Plant Biology</i> , 2017, 17, 126. | 1.6 | 12 |
| 5827 | A silver bullet in a golden age of functional genomics: the impact of <i>Agrobacterium</i> -mediated transformation of fungi. <i>Fungal Biology and Biotechnology</i> , 2017, 4, 6. | 2.5 | 63 |
| 5828 | The abundance of Fob1 modulates the efficiency of rRFBs to stall replication forks. <i>Nucleic Acids Research</i> , 2017, 45, 10089-10102. | 6.5 | 10 |
| 5829 | A systematic assessment of chemical, genetic, and epigenetic factors influencing the activity of anticancer drug KP1019 (FFC14A). <i>Oncotarget</i> , 2017, 8, 98426-98454. | 0.8 | 21 |
| 5830 | AMPK Protein Interaction Analyses by Yeast Two-Hybrid. <i>Methods in Molecular Biology</i> , 2018, 1732, 143-157. | 0.4 | 3 |
| 5831 | Cloning and expression of an endoglucanase gene from the thermotolerant fungus <i>Aspergillus fumigatus</i> DBiNU-1 in <i>Kluyveromyces lactis</i> . <i>Brazilian Journal of Microbiology</i> , 2018, 49, 647-655. | 0.8 | 14 |
| 5832 | Extramitochondrial Assembly of Mitochondrial Targeting Signal Disrupted Mitochondrial Enzyme Aldehyde Dehydrogenase. <i>Scientific Reports</i> , 2018, 8, 6186. | 1.6 | 5 |
| 5833 | A DEAD-box RNA helicase, RH8, is critical for regulation of ABA signalling and the drought stress response via inhibition of PP2CA activity. <i>Plant, Cell and Environment</i> , 2018, 41, 1593-1604. | 2.8 | 45 |
| 5834 | Advances, Problems, and Prospects of Genetic Transformation of Fungi. <i>Cytology and Genetics</i> , 2018, 52, 139-154. | 0.2 | 11 |
| 5835 | Expression of <i>Saccharomyces cerevisiae</i> cDNAs to enhance the growth of non-ethanol-producing <i>S. cerevisiae</i> strains lacking pyruvate decarboxylases. <i>Journal of Bioscience and Bioengineering</i> , 2018, 126, 317-321. | 1.1 | 1 |
| 5836 | The Dfm1 Derlin Is Required for ERAD Retrotranslocation of Integral Membrane Proteins. <i>Molecular Cell</i> , 2018, 69, 306-320.e4. | 4.5 | 76 |
| 5837 | Genetic Evidence for Roles of Yeast Mitotic Cyclins at Single-Stranded Gaps Created by DNA Replication. <i>C3: Genes, Genomes, Genetics</i> , 2018, 8, 737-752. | 0.8 | 1 |
| 5838 | More than One Way in: Three Gln3 Sequences Required To Relieve Negative Ure2 Regulation and Support Nuclear Gln3 Import in <i>Saccharomyces cerevisiae</i> . <i>Genetics</i> , 2018, 208, 207-227. | 1.2 | 12 |
| 5839 | Metabolic engineering of the 2-ketobutyrate biosynthetic pathway for 1-propanol production in <i>Saccharomyces cerevisiae</i> . <i>Microbial Cell Factories</i> , 2018, 17, 38. | 1.9 | 22 |

| # | ARTICLE | IF | CITATIONS |
|------|--|-----|-----------|
| 5840 | Transfer of plasmid into the pentose-fermenting yeast <i>Pachysolen tannophilus</i> . <i>Journal of Microbiological Methods</i> , 2018, 148, 97-103. | 0.7 | 6 |
| 5841 | <i>Drosophila</i> bendless catalyzes K63-linked polyubiquitination and is involved in the response to DNA damage. <i>Mutation Research - Fundamental and Molecular Mechanisms of Mutagenesis</i> , 2018, 808, 39-47. | 0.4 | 8 |
| 5842 | Sgs1 helicase is required for efficient PCNA monoubiquitination and translesion DNA synthesis in <i>Saccharomyces cerevisiae</i> . <i>Current Genetics</i> , 2018, 64, 459-468. | 0.8 | 10 |
| 5843 | Mutation in the peroxin-coding gene PEX22 contributing to high malate production in <i>Saccharomyces cerevisiae</i> . <i>Journal of Bioscience and Bioengineering</i> , 2018, 125, 211-217. | 1.1 | 10 |
| 5844 | The pepper dehydration-responsive homeobox 1, CaDRHB1, plays a positive role in the dehydration response. <i>Environmental and Experimental Botany</i> , 2018, 147, 104-115. | 2.0 | 10 |
| 5845 | The Sla1 adaptor- ϵ -clathrin interaction regulates coat formation and progression of endocytosis. <i>Traffic</i> , 2018, 19, 446-462. | 1.3 | 9 |
| 5846 | Optimization of chemico-physical transformation methods for various bacterial species using diverse chemical compounds and nanomaterials. <i>Journal of Biotechnology</i> , 2018, 288, 55-60. | 1.9 | 12 |
| 5847 | Construction and characterization of a <i>Saccharomyces cerevisiae</i> strain able to grow on glucosamine as sole carbon and nitrogen source. <i>Scientific Reports</i> , 2018, 8, 16949. | 1.6 | 7 |
| 5848 | Fission Yeast CENP-C (Cnp3) Plays a Role in Restricting the Site of CENP-A Accumulation. <i>G3: Genes, Genomes, Genetics</i> , 2018, 8, 2723-2733. | 0.8 | 4 |
| 5849 | Utilization of a Strongly Inducible DDI2 Promoter to Control Gene Expression in <i>Saccharomyces cerevisiae</i> . <i>Frontiers in Microbiology</i> , 2018, 9, 2736. | 1.5 | 9 |
| 5850 | Phosphoramidon inhibits the integral membrane protein zinc metalloprotease ZMPSTE24. <i>Acta Crystallographica Section D: Structural Biology</i> , 2018, 74, 739-747. | 1.1 | 4 |
| 5851 | Role of phosphate limitation and pyruvate decarboxylase in rewiring of the metabolic network for increasing flux towards isoprenoid pathway in a TATA binding protein mutant of <i>Saccharomyces cerevisiae</i> . <i>Microbial Cell Factories</i> , 2018, 17, 152. | 1.9 | 7 |
| 5852 | Highly cited papers in Microbiology: identification and conceptual analysis. <i>FEMS Microbiology Letters</i> , 2018, 365, . | 0.7 | 16 |
| 5853 | Hrg1 promotes heme-iron recycling during hemolysis in the zebrafish kidney. <i>PLoS Genetics</i> , 2018, 14, e1007665. | 1.5 | 21 |
| 5854 | Functional Characterization of Cryptococcal Genes: Then and Now. <i>Frontiers in Microbiology</i> , 2018, 9, 2263. | 1.5 | 1 |
| 5855 | Paxillin-Mediated Recruitment of Calcineurin to the Contractile Ring Is Required for the Correct Progression of Cytokinesis in Fission Yeast. <i>Cell Reports</i> , 2018, 25, 772-783.e4. | 2.9 | 23 |
| 5856 | <i>Saccharomyces cerevisiae</i> mitochondrial Por1/yVDAC1 (voltage-dependent anion channel 1) interacts physically with the MBOAT O-acyltransferase Gup1/HHATL in the control of cell wall integrity and programmed cell death. <i>FEMS Yeast Research</i> , 2018, 18, . | 1.1 | 3 |
| 5857 | Protein kinase A phosphorylates the Nem1-Spo7 protein phosphatase complex that regulates the phosphorylation state of the phosphatidate phosphatase Pah1 in yeast. <i>Journal of Biological Chemistry</i> , 2018, 293, 15801-15814. | 1.6 | 16 |

| # | ARTICLE | IF | CITATIONS |
|------|--|-----|-----------|
| 5858 | Identification and characterization of Δ^{12} and $\Delta^{12/15}$ bifunctional fatty acid desaturases in the oleaginous yeast <i>Lipomyces starkeyi</i> . <i>Applied Microbiology and Biotechnology</i> , 2018, 102, 8817-8826. | 1.7 | 10 |
| 5859 | Valuation of agro-industrial wastes as substrates for heterologous production of β -galactosidase. <i>Microbial Cell Factories</i> , 2018, 17, 137. | 1.9 | 16 |
| 5860 | Construction and Use of a Recyclable Marker To Examine the Role of Major Facilitator Superfamily Protein Members in <i>Candida glabrata</i> Drug Resistance Phenotypes. <i>MSphere</i> , 2018, 3, . | 1.3 | 13 |
| 5861 | Wheat F-Box Protein Gene TaFBA1 Is Involved in Plant Tolerance to Heat Stress. <i>Frontiers in Plant Science</i> , 2018, 9, 521. | 1.7 | 123 |
| 5862 | Cloning and characterization of SucNHX1, a novel vacuolar Na ⁺ /H ⁺ antiporter from the halophyte <i>Suaeda corniculata</i> that enhances the saline-alkali tolerance in <i>Arabidopsis</i> by its overexpression. <i>Plant Cell, Tissue and Organ Culture</i> , 2018, 134, 395-407. | 1.2 | 7 |
| 5863 | A novel EPM2A mutation yields a slow progression form of Lafora disease. <i>Epilepsy Research</i> , 2018, 145, 169-177. | 0.8 | 10 |
| 5864 | Overexpression of THI4 and HAP4 Improves Glucose Metabolism and Ethanol Production in <i>Saccharomyces cerevisiae</i> . <i>Frontiers in Microbiology</i> , 2018, 9, 1444. | 1.5 | 19 |
| 5865 | Tribute to Marcelle Grenson (1925–1996), A Pioneer in the Study of Amino Acid Transport in Yeast. <i>International Journal of Molecular Sciences</i> , 2018, 19, 1207. | 1.8 | 3 |
| 5866 | Intermolecular RNA Recombination Occurs at Different Frequencies in Alternate Forms of Brome Mosaic Virus RNA Replication Compartments. <i>Viruses</i> , 2018, 10, 131. | 1.5 | 15 |
| 5867 | Analysis of β -tubulin-carbendazim interaction reveals that binding site for MBC fungicides does not include residues involved in fungicide resistance. <i>Scientific Reports</i> , 2018, 8, 7161. | 1.6 | 51 |
| 5868 | Negative regulation of <i>Candida glabrata</i> Pdr1 by the deubiquitinase subunit Bre5 occurs in a ubiquitin independent manner. <i>Molecular Microbiology</i> , 2018, 110, 309-323. | 1.2 | 9 |
| 5869 | Low Ctr1p, due to lack of Sco1p results in lowered cisplatin uptake and mediates insensitivity of rho ⁰ yeast to cisplatin. <i>Journal of Inorganic Biochemistry</i> , 2018, 187, 14-24. | 1.5 | 7 |
| 5870 | Limited expression of functional cytochrome p450 2c subtypes in the liver and small intestine of domestic cats. <i>Xenobiotica</i> , 2019, 49, 627-635. | 0.5 | 13 |
| 5871 | The NADP ⁺ -dependent glutamate dehydrogenase Gdh1 is subjected to glucose starvation-induced reversible aggregation that affects stress resistance in yeast. <i>Journal of Microbiology</i> , 2019, 57, 884-892. | 1.3 | 2 |
| 5872 | Quantitative proteomics reveals a G β /MAPK signaling hub that controls pheromone-induced cellular polarization in yeast. <i>Journal of Proteomics</i> , 2019, 207, 103467. | 1.2 | 4 |
| 5873 | Carboxypeptidase Y activity and maintenance is modulated by a large helical structure. <i>FEBS Open Bio</i> , 2019, 9, 1337-1343. | 1.0 | 24 |
| 5874 | Modulation of ruthenium anticancer drugs analogs with tolfenamic acid: Reactivity, biological interactions and growth inhibition of yeast cell. <i>Journal of Inorganic Biochemistry</i> , 2019, 199, 110769. | 1.5 | 13 |
| 5875 | COP9 Signalosome Interaction with UspA/Usp15 Deubiquitinase Controls VeA-Mediated Fungal Multicellular Development. <i>Biomolecules</i> , 2019, 9, 238. | 1.8 | 15 |

| # | ARTICLE | IF | CITATIONS |
|------|---|------|-----------|
| 5876 | Roles of pepper <i>bZIP</i> transcription factor <i>CaATBZ1</i> and its interacting partner <i>RING</i> -type E3 ligase <i>CaASRF1</i> in modulation of <i>ABA</i> signalling and drought tolerance. <i>Plant Journal</i> , 2019, 100, 399-410. | 2.8 | 52 |
| 5877 | Experimental evolution reveals a general role for the methyltransferase <i>Hmt1</i> in noise buffering. <i>PLoS Biology</i> , 2019, 17, e3000433. | 2.6 | 7 |
| 5878 | Critical Roles of the Pentose Phosphate Pathway and <i>GLN3</i> in Isobutanol-Specific Tolerance in Yeast. <i>Cell Systems</i> , 2019, 9, 534-547.e5. | 2.9 | 28 |
| 5879 | The RING finger E3 ligases <i>PIR1</i> and <i>PIR2</i> mediate <i>PP2CA</i> degradation to enhance abscisic acid response in <i>Arabidopsis</i> . <i>Plant Journal</i> , 2019, 100, 473-486. | 2.8 | 29 |
| 5880 | A quantitative screen for metabolic enzyme structures reveals patterns of assembly across the yeast metabolic network. <i>Molecular Biology of the Cell</i> , 2019, 30, 2721-2736. | 0.9 | 49 |
| 5881 | Fine-tuning the expression of target genes using a <i>DDI2</i> promoter gene switch in budding yeast. <i>Scientific Reports</i> , 2019, 9, 12538. | 1.6 | 9 |
| 5882 | Protein kinase C mediates the phosphorylation of the <i>Nem1</i> - <i>Spo7</i> protein phosphatase complex in yeast. <i>Journal of Biological Chemistry</i> , 2019, 294, 15997-16009. | 1.6 | 8 |
| 5883 | Characterization of two <i>Vitis vinifera</i> carotenoid cleavage dioxygenases by heterologous expression in <i>Saccharomyces cerevisiae</i> . <i>Molecular Biology Reports</i> , 2019, 46, 6311-6323. | 1.0 | 14 |
| 5884 | Substitution of cysteines in the yeast viral killer toxin <i>K1</i> precursor reveals novel insights in heterodimer formation and immunity. <i>Scientific Reports</i> , 2019, 9, 13127. | 1.6 | 6 |
| 5885 | <i>eIF5B</i> gates the transition from translation initiation to elongation. <i>Nature</i> , 2019, 573, 605-608. | 13.7 | 60 |
| 5886 | Methods for studying the regulation of membrane traffic by ubiquitin and the ESCRT pathway. <i>Methods in Enzymology</i> , 2019, 619, 269-291. | 0.4 | 1 |
| 5887 | <i>amdS</i> as a dominant recyclable marker in <i>Cryptococcus neoformans</i> . <i>Fungal Genetics and Biology</i> , 2019, 131, 103241. | 0.9 | 10 |
| 5888 | <i>Sit4</i> and <i>PP2A</i> Dephosphorylate Nitrogen Catabolite Repression-Sensitive <i>Gln3</i> When <i>TorC1</i> Is Up- as Well as Downregulated. <i>Genetics</i> , 2019, 212, 1205-1225. | 1.2 | 9 |
| 5889 | Conserved juxtamembrane domains in the yeast golgin <i>Coy1</i> drive assembly of a megadalton-sized complex and mediate binding to tethering and SNARE proteins. <i>Journal of Biological Chemistry</i> , 2019, 294, 9690-9705. | 1.6 | 2 |
| 5890 | Evidence that Ergosterol Biosynthesis Modulates Activity of the <i>Pdr1</i> Transcription Factor in <i>Candida glabrata</i> . <i>MBio</i> , 2019, 10, . | 1.8 | 42 |
| 5891 | <i>Cus2</i> enforces the first ATP-dependent step of splicing by binding to yeast <i>SF3b1</i> through a UHM-UJM interaction. <i>Rna</i> , 2019, 25, 1020-1037. | 1.6 | 19 |
| 5892 | <i>Ser625</i> of <i>msn2</i> transcription factor is indispensable for ethanol tolerance and alcoholic fermentation process. <i>Biotechnology Progress</i> , 2019, 35, e2837. | 1.3 | 5 |
| 5893 | Integrating after CEN Excision (ICE) Plasmids: Combining the ease of yeast recombination cloning with the stability of genomic integration. <i>Yeast</i> , 2019, 36, 593-605. | 0.8 | 11 |

| # | ARTICLE | IF | CITATIONS |
|------|--|------|-----------|
| 5894 | CoQ10 production in <i>Schizosaccharomyces pombe</i> is increased by reduction of glucose levels or deletion of <i>pka1</i> . <i>Applied Microbiology and Biotechnology</i> , 2019, 103, 4899-4915. | 1.7 | 9 |
| 5895 | Communication between RACK1/Asc1 and uS3 (Rps3) is essential for RACK1/Asc1 function in yeast <i>Saccharomyces cerevisiae</i> . <i>Gene</i> , 2019, 706, 69-76. | 1.0 | 3 |
| 5896 | Bioconversion of Beet Molasses to Alpha-Galactosidase and Ethanol. <i>Frontiers in Microbiology</i> , 2019, 10, 405. | 1.5 | 22 |
| 5897 | Novel Antifungal Compound Z-705 Specifically Inhibits Protein Kinase C of Filamentous Fungi. <i>Applied and Environmental Microbiology</i> , 2019, 85, . | 1.4 | 11 |
| 5898 | Prevention of mitochondrial genomic instability in yeast by the mitochondrial recombinase Mhr1. <i>Scientific Reports</i> , 2019, 9, 5433. | 1.6 | 16 |
| 5899 | Simplified protocol for faster transformation of (a large number of) <i>Pichia pastoris</i> strains. <i>Yeast</i> , 2019, 36, 399-410. | 0.8 | 8 |
| 5900 | The Glycosylphosphatidylinositol-Anchored DFG Family Is Essential for the Insertion of Galactomannan into the Î²-(1,3)-Glucan Chitin Core of the Cell Wall of <i>Aspergillus fumigatus</i> . <i>MSphere</i> , 2019, 4, . | 1.3 | 28 |
| 5901 | Functional Analysis of Pepper F-box Protein CaDIF1 and Its Interacting Partner CaDIS1: Modulation of ABA Signaling and Drought Stress Response. <i>Frontiers in Plant Science</i> , 2019, 10, 1365. | 1.7 | 14 |
| 5902 | Meiotic chromosomal recombination defect in sake yeasts. <i>Journal of Bioscience and Bioengineering</i> , 2019, 127, 190-196. | 1.1 | 11 |
| 5903 | Yeast-inspired drug delivery: biotechnology meets bioengineering and synthetic biology. <i>Expert Opinion on Drug Delivery</i> , 2019, 16, 27-41. | 2.4 | 34 |
| 5904 | Role of the uS9/yS16 C-terminal tail in translation initiation and elongation in <i>Saccharomyces cerevisiae</i> . <i>Nucleic Acids Research</i> , 2019, 47, 806-823. | 6.5 | 20 |
| 5905 | Sustainable production of glutathione from lignocellulose-derived sugars using engineered <i>Saccharomyces cerevisiae</i> . <i>Applied Microbiology and Biotechnology</i> , 2019, 103, 1243-1254. | 1.7 | 8 |
| 5906 | Cbs overdosage is necessary and sufficient to induce cognitive phenotypes in mouse models of Down syndrome and interacts genetically with <i>Dyrk1a</i> . <i>Human Molecular Genetics</i> , 2019, 28, 1561-1577. | 1.4 | 41 |
| 5907 | Introns are mediators of cell response to starvation. <i>Nature</i> , 2019, 565, 612-617. | 13.7 | 161 |
| 5908 | CYP81A P45Os are involved in concomitant cross-resistance to acetolactate synthase and acetyl-CoA carboxylase herbicides in <i>Echinochloa phyllopogon</i> . <i>New Phytologist</i> , 2019, 221, 2112-2122. | 3.5 | 112 |
| 5909 | Delta-9 fatty acid desaturase overexpression enhanced lipid production and oleic acid content in <i>Rhodospiridium toruloides</i> for preferable yeast lipid production. <i>Journal of Bioscience and Bioengineering</i> , 2019, 127, 430-440. | 1.1 | 36 |
| 5910 | A pepper RING-type E3 ligase, CaASRF1, plays a positive role in drought tolerance via modulation of CaAIBZ1 stability. <i>Plant Journal</i> , 2019, 98, 5-18. | 2.8 | 55 |
| 5911 | Specific serine residues of Msn2/4 are responsible for regulation of alcohol fermentation rates and ethanol resistance. <i>Biotechnology Progress</i> , 2019, 35, e2759. | 1.3 | 7 |

| # | ARTICLE | IF | CITATIONS |
|------|---|-----|-----------|
| 5912 | Leishmania heme uptake involves LmFLVCRb, a novel porphyrin transporter essential for the parasite. Cellular and Molecular Life Sciences, 2020, 77, 1827-1845. | 2.4 | 22 |
| 5913 | Arabidopsis <i>OTU1</i> , a linkage-specific deubiquitinase, is required for endoplasmic reticulum-associated protein degradation. Plant Journal, 2020, 101, 141-155. | 2.8 | 16 |
| 5914 | The K ⁺ and NO ₃ ⁻ Interaction Mediated by NITRATE TRANSPORTER1.1 Ensures Better Plant Growth under K ⁺ -Limiting Conditions. Plant Physiology, 2020, 184, 1900-1916. | 2.3 | 22 |
| 5915 | An Erg11 lanosterol 14- β -demethylase-Arv1 complex is required for <i>Candida albicans</i> virulence. PLoS ONE, 2020, 15, e0235746. | 1.1 | 6 |
| 5916 | Functional Studies of Plasmodium falciparum's Prohibitin1 and Prohibitin 2 in Yeast. Indian Journal of Medical Microbiology, 2020, 38, 213-215. | 0.3 | 6 |
| 5917 | In <i>Kluyveromyces lactis</i> a Pair of Paralogous Isozymes Catalyze the First Committed Step of Leucine Biosynthesis in Either the Mitochondria or the Cytosol. Frontiers in Microbiology, 2020, 11, 1843. | 1.5 | 4 |
| 5918 | Functional characterization of <i>Candida glabrata</i> ORF, CAGL0M02233g for its role in stress tolerance and virulence. Microbial Pathogenesis, 2020, 149, 104469. | 1.3 | 9 |
| 5919 | Cargo-Mediated Recruitment of the Endocytic Adaptor Protein Sla1. Journal of Cell Science, 2020, 133, . | 1.2 | 0 |
| 5920 | A novel reverse two-hybrid method for the identification of missense mutations that disrupt protein-protein binding. Scientific Reports, 2020, 10, 21043. | 1.6 | 3 |
| 5921 | Yeast Viral Killer Toxin K1 Induces Specific Host Cell Adaptions via Intrinsic Selection Pressure. Applied and Environmental Microbiology, 2020, 86, . | 1.4 | 8 |
| 5922 | The pepper <i>RING</i> type <i>E3</i> ligase, <i>CaATIR1</i> , positively regulates abscisic acid signalling and drought response by modulating the stability of <i>CaATBZ1</i> . Plant, Cell and Environment, 2020, 43, 1911-1924. | 2.8 | 23 |
| 5923 | Identification and functional characterization of two acyl CoA:diacylglycerol acyltransferase 1 (DGAT1) genes from forage sorghum (<i>Sorghum bicolor</i>) embryo. Phytochemistry, 2020, 176, 112405. | 1.4 | 7 |
| 5924 | Modified chemical method for efficient transformation and diagnosis in <i>Pichia pastoris</i> . Protein Expression and Purification, 2020, 174, 105685. | 0.6 | 8 |
| 5925 | The Spo7 sequence LLI is required for Nem1-Spo7/Pah1 phosphatase cascade function in yeast lipid metabolism. Journal of Biological Chemistry, 2020, 295, 11473-11485. | 1.6 | 13 |
| 5926 | Damaging de novo missense variants in <i>EEF1A2</i> lead to a developmental and degenerative epileptic dyskinetic encephalopathy. Human Mutation, 2020, 41, 1263-1279. | 1.1 | 24 |
| 5927 | A Transcriptional Regulatory System of the <i>S. cerevisiae</i> OLE1 Gene Responds to Fatty Acid Species and Intracellular Amount, and not Simply Membrane Status. Journal of Lipids, 2020, 2020, 1-7. | 1.9 | 2 |
| 5928 | Cloning of the SPO11 gene that complements a meiotic recombination defect in sake yeast. Journal of Bioscience and Bioengineering, 2020, 130, 367-373. | 1.1 | 4 |
| 5929 | Evolutionary and Functional Analysis of a Chara Plasma Membrane H ⁺ -ATPase. Frontiers in Plant Science, 2019, 10, 1707. | 1.7 | 10 |

| # | ARTICLE | IF | CITATIONS |
|------|---|-----|-----------|
| 5930 | Characterization of the mechanism of thioredoxin-dependent activation of Γ^3 -glutamylcyclotransferase, RipAY, from <i>Ralstonia solanacearum</i> . <i>Biochemical and Biophysical Research Communications</i> , 2020, 523, 759-765. | 1.0 | 9 |
| 5931 | Exploiting the NADPH pool for xylitol production using recombinant <i>Saccharomyces cerevisiae</i> . <i>Biotechnology Progress</i> , 2020, 36, e2972. | 1.3 | 13 |
| 5932 | Effects of mutations of GID protein-coding genes on malate production and enzyme expression profiles in <i>Saccharomyces cerevisiae</i> . <i>Applied Microbiology and Biotechnology</i> , 2020, 104, 4971-4983. | 1.7 | 7 |
| 5933 | Analysis of Yeast Killer Toxin K1 Precursor Processing via Site-Directed Mutagenesis: Implications for Toxicity and Immunity. <i>MSphere</i> , 2020, 5, . | 1.3 | 4 |
| 5934 | Functional characterization of two type-1 diacylglycerol acyltransferase (DGAT1) genes from rice (<i>Oryza sativa</i>) embryo restoring the triacylglycerol accumulation in yeast. <i>Plant Molecular Biology</i> , 2021, 105, 247-262. | 2.0 | 9 |
| 5935 | Hypoosmolality impedes cytoophidium integrity during nitrogen starvation. <i>Yeast</i> , 2021, 38, 276-289. | 0.8 | 10 |
| 5936 | Cac1 WHD and PIP domains have distinct roles in replisome progression and genomic stability. <i>Current Genetics</i> , 2021, 67, 129-139. | 0.8 | 7 |
| 5937 | Lipid production by oleaginous yeasts. <i>Advances in Applied Microbiology</i> , 2021, 116, 1-98. | 1.3 | 14 |
| 5938 | Metabolic and Transcriptional Analysis of Recombinant <i>Saccharomyces Cerevisiae</i> for Xylose Fermentation: A Feasible and Efficient Approach. <i>IEEE Journal of Biomedical and Health Informatics</i> , 2022, 26, 2425-2434. | 3.9 | 1 |
| 5939 | <i>Saccharomyces cerevisiae</i> as a microbial cell factory. , 2021, , 319-333. | | 5 |
| 5940 | Variation among <i>Metschnikowia pulcherrima</i> Isolates for Genetic Modification and Homologous Recombination. <i>Microorganisms</i> , 2021, 9, 290. | 1.6 | 7 |
| 5941 | Enhancing the chemical transformation of <i>Candida parapsilosis</i> . <i>Virulence</i> , 2021, 12, 937-950. | 1.8 | 7 |
| 5942 | N- and C-terminal Gln3-Tor1 interaction sites: one acting negatively and the other positively to regulate nuclear Gln3 localization. <i>Genetics</i> , 2021, 217, . | 1.2 | 6 |
| 5944 | A conserved, buried cysteine near the P-site is accessible to cysteine modifications and increases ROS stability in the P-type plasma membrane H ⁺ -ATPase. <i>Biochemical Journal</i> , 2021, 478, 619-632. | 1.7 | 9 |
| 5945 | Organic nitrogen nutrition: LHT1.2 protein from hybrid aspen (<i>Populus tremula</i> L. x <i>tremuloides</i>) Tj ETQq0 0 0 rgBT /Overlock 10 2021, 41, 1479-1496. | 1.4 | 9 |
| 5946 | Mutation in gene coding for glucose-induced degradation-deficient protein contributes to high malate production in yeast strain No. 28 and No. 77 used for industrial brewing of sake. <i>Bioscience, Biotechnology and Biochemistry</i> , 2021, 85, 1283-1289. | 0.6 | 2 |
| 5947 | Improved yeast-screening assay for phytoestrogens. <i>Bioscience, Biotechnology and Biochemistry</i> , 2021, 85, 1422-1432. | 0.6 | 1 |
| 5948 | Combining Nanopore and Illumina Sequencing Permits Detailed Analysis of Insertion Mutations and Structural Variations Produced by PEG-Mediated Transformation in <i>Ostreococcus tauri</i> . <i>Cells</i> , 2021, 10, 664. | 1.8 | 3 |

| # | ARTICLE | IF | CITATIONS |
|------|--|-----|-----------|
| 5949 | Inner-nuclear-membrane-associated degradation employs Dfm1-independent retrotranslocation and alleviates misfolded transmembrane-protein toxicity. <i>Molecular Biology of the Cell</i> , 2021, 32, 521-537. | 0.9 | 6 |
| 5950 | Negative regulation of diminutive cancer regulator through differentiation and microRNA pathway components in <i>Drosophila</i> cells. <i>Turkish Journal of Biology</i> , 2021, 45, 180-186. | 2.1 | 0 |
| 5951 | HBD1 protein with a tandem repeat of two HMG-box domains is a DNA clip to organize chloroplast nucleoids in <i>Chlamydomonas reinhardtii</i> . <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 2021, 118, . | 3.3 | 7 |
| 5953 | The mitochondrial aspartate/glutamate carrier (AGC or Aralar1) isoforms in <i>D. melanogaster</i> : biochemical characterization, gene structure, and evolutionary analysis. <i>Biochimica Et Biophysica Acta - General Subjects</i> , 2021, 1865, 129854. | 1.1 | 9 |
| 5954 | Cantharidin downregulates PSD1 expression and inhibits autophagic flux in yeast cells. <i>FEBS Open Bio</i> , 2021, , . | 1.0 | 3 |
| 5955 | ABC transporter Pdr5 is required for cantharidin resistance in <i>Saccharomyces cerevisiae</i> . <i>Biochemical and Biophysical Research Communications</i> , 2021, 553, 141-147. | 1.0 | 3 |
| 5957 | The dynein light chain protein Tda2 functions as a dimerization engine to regulate actin capping protein during endocytosis. <i>Molecular Biology of the Cell</i> , 2021, 32, mbc.E21-01-0032. | 0.9 | 8 |
| 5958 | Characterization of HNRNPA1 mutations defines diversity in pathogenic mechanisms and clinical presentation. <i>JCI Insight</i> , 2021, 6, . | 2.3 | 38 |
| 5960 | Coupled sterol synthesis and transport machineries at ER-endocytic contact sites. <i>Journal of Cell Biology</i> , 2021, 220, . | 2.3 | 12 |
| 5961 | CaADIP1-dependent CaADIK1-kinase activation is required for abscisic acid signalling and drought stress response in <i>Capsicum annuum</i> . <i>New Phytologist</i> , 2021, 231, 2247-2261. | 3.5 | 7 |
| 5962 | Homology length dictates the requirement for Rad51 and Rad52 in gene targeting in the Basidiomycota yeast <i>Naganishia liquefaciens</i> . <i>Current Genetics</i> , 2021, 67, 919-936. | 0.8 | 3 |
| 5963 | Experimental evolution improves mitochondrial genome quality control in <i>Saccharomyces cerevisiae</i> and extends its replicative lifespan. <i>Current Biology</i> , 2021, 31, 3663-3670.e4. | 1.8 | 5 |
| 5965 | The <i>Candida glabrata</i> Upc2A transcription factor is a global regulator of antifungal drug resistance pathways. <i>PLoS Genetics</i> , 2021, 17, e1009582. | 1.5 | 22 |
| 5966 | Haem-responsive gene transporter enables mobilization of host haem in ticks. <i>Open Biology</i> , 2021, 11, 210048. | 1.5 | 6 |
| 5967 | The HMGB Protein Kllxr1, a DNA Binding Regulator of <i>Kluyveromyces lactis</i> Gene Expression Involved in Oxidative Metabolism, Growth, and dNTP Synthesis. <i>Biomolecules</i> , 2021, 11, 1392. | 1.8 | 2 |
| 5968 | Regulation of sphingolipid biosynthesis in the endoplasmic reticulum via signals from the plasma membrane in budding yeast. <i>FEBS Journal</i> , 2022, 289, 457-472. | 2.2 | 8 |
| 5969 | Production of transglutaminase in glutathione-producing recombinant <i>Saccharomyces cerevisiae</i> . <i>AMB Express</i> , 2021, 11, 13. | 1.4 | 4 |
| 5970 | Derlin Rhomboid Pseudoproteases Employ Substrate Engagement and Lipid Distortion Function for Retrotranslocation of ER Multi-Spanning Membrane Substrates. <i>SSRN Electronic Journal</i> , 0, , . | 0.4 | 0 |

| # | ARTICLE | IF | CITATIONS |
|------|--|-----|-----------|
| 5972 | Opi1p translocation to the nucleus is regulated by hydrogen peroxide in <i>Saccharomyces cerevisiae</i> . <i>Yeast</i> , 2017, 34, 383-395. | 0.8 | 1 |
| 5973 | CRISPR Nickase-Mediated Base Editing in Yeast. <i>Methods in Molecular Biology</i> , 2021, 2196, 27-37. | 0.4 | 3 |
| 5974 | Recombinant Formaldehyde Dehydrogenase and Gene-Engineered Methylotrophic Yeasts as Bioanalytical Instruments for Assay of Toxic Formaldehyde. <i>NATO Science for Peace and Security Series A: Chemistry and Biology</i> , 2008, , 311-333. | 0.5 | 3 |
| 5975 | Attempts to Transform for Kanamycin-Resistance in Mature Pollen of Tobacco. , 1986, , 65-70. | | 22 |
| 5976 | A Single Gene Change Can Extend Yeast Life Span: The Role of RAS in Cellular Senescence. <i>Advances in Experimental Medicine and Biology</i> , 1993, 330, 45-53. | 0.8 | 14 |
| 5977 | Baker's yeast. , 1998, , 158-173. | | 2 |
| 5978 | Novel Techniques for Breeding of Microorganisms and Application of the Microorganisms to Production of Glutathione. , 1985, , 35-47. | | 2 |
| 5979 | Yeast genetics. , 1996, , 43-82. | | 5 |
| 5980 | The Evolutionary Origin and Maintenance of Sexual Recombination: A Review of Contemporary Models. , 2003, , 27-138. | | 29 |
| 5981 | Generation of Arming Yeasts with Active Proteins and Peptides via Cell Surface Display System: Cell Surface Engineering, Bio-arming Technology. <i>Methods in Molecular Biology</i> , 2014, 1152, 137-155. | 0.4 | 14 |
| 5982 | Evaluation of Unconventional Protein Secretion in <i>Saccharomyces cerevisiae</i> . <i>Methods in Molecular Biology</i> , 2015, 1270, 51-70. | 0.4 | 2 |
| 5983 | Enzyme Evolution by Yeast Cell Surface Engineering. <i>Methods in Molecular Biology</i> , 2015, 1319, 217-232. | 0.4 | 2 |
| 5984 | Preparation of an Oral Vaccine by Proteome Analysis and Molecular Display Technology. <i>Methods in Molecular Biology</i> , 2017, 1625, 237-245. | 0.4 | 6 |
| 5985 | Identification of Determinants of Sensitivity to Antitumor Drugs. , 2003, , 319-330. | | 3 |
| 5986 | Identification of DNA-Binding Proteins and Protein-Protein Interactions by Yeast One-Hybrid and Yeast Two-Hybrid Screen. <i>Methods in Molecular Biology</i> , 2010, 639, 171-192. | 0.4 | 6 |
| 5987 | Assays for Effector-Mediated Suppression of Programmed Cell Death in Yeast. <i>Methods in Molecular Biology</i> , 2011, 712, 173-180. | 0.4 | 4 |
| 5988 | Identification and Assays of Polyamine Transport Systems in <i>Escherichia coli</i> and <i>Saccharomyces cerevisiae</i> . <i>Methods in Molecular Biology</i> , 2011, 720, 295-308. | 0.4 | 21 |
| 5989 | Finding Modulators of Stochasticity Levels by Quantitative Genetics. <i>Methods in Molecular Biology</i> , 2011, 734, 223-240. | 0.4 | 2 |

| # | ARTICLE | IF | CITATIONS |
|------|--|-----|-----------|
| 5990 | Genetic Transformation of Filamentous Fungi: Achievements and Challenges. <i>Grand Challenges in Biology and Biotechnology</i> , 2020, , 123-164. | 2.4 | 8 |
| 5991 | Pathways and Mechanisms of Yeast Competence: A New Frontier of Yeast Genetics. <i>Fungal Biology</i> , 2015, , 223-237. | 0.3 | 3 |
| 5992 | Fungal Cell Walls – A Review. , 1990, , 5-30. | | 86 |
| 5993 | Formation of Jumbo Yeast Cells by Introduction of Methylglyoxal Resistant Genes, Development of New Vector for Wild or Industrially Used Cells. , 1989, , 173-180. | | 3 |
| 5994 | A Role for Sorbitol in Desiccation Tolerance of Developing Maize Kernels: Inference from the Properties of Maize Sorbitol Dehydrogenase. , 1994, , 345-368. | | 2 |
| 5995 | Yeast Halotolerance Genes: Crucial Ion Transport and Metabolic Reactions in Salt Tolerance. , 1994, , 371-380. | | 3 |
| 5996 | <i>Arxula adenivorans</i> . , 1996, , 389-409. | | 17 |
| 5997 | Genetic Analysis and Manipulation of <i>Mucor</i> Species by DNA-Mediated Transformation. <i>Modern Methods of Plant Analysis</i> , 1988, , 207-222. | 0.1 | 3 |
| 5998 | Genetic Manipulation of Fungi by DNA-Mediated Transformation. , 1995, , 109-139. | | 5 |
| 5999 | Yeast plasmids and transformation. , 1987, , 80-103. | | 2 |
| 6000 | Fungal enzymes used in oriental food and beverage industries. , 1992, , 54-99. | | 35 |
| 6001 | Yeast Genetics. , 1991, , 37-87. | | 1 |
| 6002 | Enhancement of Aroma in Grapes and Wines: Biotechnological Approaches. , 2001, , 225-240. | | 3 |
| 6003 | Metabolic engineering of <i>Saccharomyces cerevisiae</i> for efficient production of pure α -lactate. <i>Applied Biochemistry and Biotechnology</i> , 1996, 131, 795-807. | 1.4 | 1 |
| 6004 | The upstream region of the isocitrate lyase gene (UPR-ICL) of <i>Candida tropicalis</i> induces gene expression in both <i>Saccharomyces cerevisiae</i> and <i>Escherichia coli</i> by acetate via two distinct promoters. <i>Archives of Microbiology</i> , 1995, 163, 322-328. | 1.0 | 3 |
| 6005 | Construction of bioengineered yeast platform for direct bioethanol production from alginate and mannitol. <i>Applied Microbiology and Biotechnology</i> , 2017, 101, 6627-6636. | 1.7 | 29 |
| 6006 | Characterization of the catalase of the n-alkane-utilizing yeast <i>Candida tropicalis</i> functionally expressed in <i>Saccharomyces cerevisiae</i> . <i>Applied Microbiology and Biotechnology</i> , 1994, 40, 682-686. | 1.7 | 2 |
| 6007 | One-step enzymatic hydrolysis of starch using a recombinant strain of <i>Saccharomyces cerevisiae</i> producing α -amylase, glucoamylase and pullulanase. <i>Applied Microbiology and Biotechnology</i> , 1995, 42, 878-883. | 1.7 | 4 |

| # | ARTICLE | IF | CITATIONS |
|------|--|-----|-----------|
| 6008 | A novel promoter, derived from the isocitrate lyase gene of <i>Candida tropicalis</i> , inducible with acetate in <i>Saccharomyces cerevisiae</i> . <i>Applied Microbiology and Biotechnology</i> , 1995, 43, 489-492. | 1.7 | 4 |
| 6009 | yAP-1- and yAP-2-mediated, heat shock-induced transcriptional activation of the multidrug resistance ABC transporter genes in <i>Saccharomyces cerevisiae</i> . <i>Current Genetics</i> , 1996, 29, 103-105. | 0.8 | 6 |
| 6010 | Molecular and functional characterization of a mutant allele of the mitogen-activated protein-kinase gene SLT2(MPK1) rescued from yeast autolytic mutants. <i>Current Genetics</i> , 1996, 29, 516-522. | 0.8 | 4 |
| 6011 | Plasmid transformation in <i>Bacillus subtilis</i> NB22, an antifungal-antibiotic iturin producer. <i>FEMS Microbiology Letters</i> , 1990, 67, 227-229. | 0.7 | 2 |
| 6012 | Yeast Transformation. , 1985, , 161-195. | | 2 |
| 6013 | Molecular Biology of the qa Gene Cluster of <i>Neurospora</i> . , 1985, , 229-244. | | 6 |
| 6014 | Cloning and Transformation in <i>Aspergillus</i> . , 1985, , 259-278. | | 29 |
| 6015 | Molecular Genetics of Human Fungal Pathogens. , 1991, , 342-363. | | 5 |
| 6016 | Gene Cloning and Expression in Fission Yeast. , 1989, , 243-271. | | 69 |
| 6017 | Transformation. , 1992, 21, 113-156. | | 21 |
| 6018 | Casein kinase II phosphorylates the C-terminal region of Lif1 to promote the Lif1-Xrs2 interaction needed for non-homologous end joining. <i>Biochemical and Biophysical Research Communications</i> , 2018, 501, 1080-1084. | 1.0 | 2 |
| 6019 | <i>Saccharomyces cerevisiae</i> STE14 gene is required for COOH-terminal methylation of a-factor mating pheromone. <i>Journal of Biological Chemistry</i> , 1990, 265, 20057-20060. | 1.6 | 60 |
| 6020 | Sorting of peroxisomal membrane protein PMP47 from <i>Candida boidinii</i> into peroxisomal membranes of <i>Saccharomyces cerevisiae</i> . <i>Journal of Biological Chemistry</i> , 1990, 265, 20098-20105. | 1.6 | 54 |
| 6021 | The OLE1 gene of <i>Saccharomyces cerevisiae</i> encodes the delta 9 fatty acid desaturase and can be functionally replaced by the rat stearoyl-CoA desaturase gene. <i>Journal of Biological Chemistry</i> , 1990, 265, 20144-20149. | 1.6 | 375 |
| 6022 | A mutation in the largest subunit of yeast TFIIC affects tRNA and 5 S RNA synthesis. Identification of two classes of suppressors.. <i>Journal of Biological Chemistry</i> , 1994, 269, 23374-23381. | 1.6 | 51 |
| 6023 | Suppressors of the Ca(2+)-sensitive yeast mutant (<i>csg2</i>) identify genes involved in sphingolipid biosynthesis. Cloning and characterization of SCS1, a gene required for serine palmitoyltransferase activity.. <i>Journal of Biological Chemistry</i> , 1994, 269, 21480-21488. | 1.6 | 102 |
| 6024 | A polybasic domain allows nonprenylated Ras proteins to function in <i>Saccharomyces cerevisiae</i> .. <i>Journal of Biological Chemistry</i> , 1994, 269, 21540-21546. | 1.6 | 44 |
| 6025 | Human apolipoprotein B signal sequence variants confer a secretion-defective phenotype when expressed in yeast.. <i>Journal of Biological Chemistry</i> , 1994, 269, 21670-21675. | 1.6 | 41 |

| # | ARTICLE | IF | CITATIONS |
|------|---|-----|-----------|
| 6026 | Quantitative analysis of mutually competitive binding of human Raf-1 and yeast adenylyl cyclase to Ras proteins.. Journal of Biological Chemistry, 1994, 269, 20845-20851. | 1.6 | 41 |
| 6027 | Incorrect targeting of plant vacuolar lectins in yeast.. Journal of Biological Chemistry, 1994, 269, 20866-20871. | 1.6 | 8 |
| 6028 | Characterization of a glucose-repressible ADP-ribosylation factor 3 (ARF3) from <i>Saccharomyces cerevisiae</i> .. Journal of Biological Chemistry, 1994, 269, 20931-20937. | 1.6 | 55 |
| 6029 | Dominance of metallothionein in metal ion buffering in yeast capable of synthesis of (gamma EC)nG isopeptides.. Journal of Biological Chemistry, 1994, 269, 21010-21015. | 1.6 | 19 |
| 6030 | Extent of in vivo binding by an upstream activation factor and the role of multiple binding sites in synergistic transcriptional activation.. Journal of Biological Chemistry, 1994, 269, 20771-20779. | 1.6 | 3 |
| 6031 | The <i>Saccharomyces cerevisiae</i> PLB1 gene encodes a protein required for lysophospholipase and phospholipase B activity.. Journal of Biological Chemistry, 1994, 269, 19725-19730. | 1.6 | 115 |
| 6032 | Divergent roles of RAS1 and RAS2 in yeast longevity. Journal of Biological Chemistry, 1994, 269, 18638-18645. | 1.6 | 171 |
| 6033 | Transcriptional control of yeast plasma membrane H(+)-ATPase by glucose. Cloning and characterization of a new gene involved in this regulation.. Journal of Biological Chemistry, 1994, 269, 18076-18082. | 1.6 | 57 |
| 6034 | High yield expression of the <i>Neurospora crassa</i> plasma membrane H(+)-ATPase in <i>Saccharomyces cerevisiae</i> . Journal of Biological Chemistry, 1994, 269, 17705-17712. | 1.6 | 20 |
| 6035 | Functional expression of P-glycoproteins in secretory vesicles.. Journal of Biological Chemistry, 1994, 269, 12277-12284. | 1.6 | 121 |
| 6036 | Molecular mechanism of regulation of yeast plasma membrane H(+)-ATPase by glucose. Interaction between domains and identification of new regulatory sites.. Journal of Biological Chemistry, 1994, 269, 10393-10399. | 1.6 | 100 |
| 6037 | The OPI1 gene of <i>Saccharomyces cerevisiae</i> , a negative regulator of phospholipid biosynthesis, encodes a protein containing polyglutamine tracts and a leucine zipper.. Journal of Biological Chemistry, 1991, 266, 863-872. | 1.6 | 162 |
| 6038 | Isolation and characterization of the structural gene for secreted acid phosphatase from <i>Schizosaccharomyces pombe</i> .. Journal of Biological Chemistry, 1986, 261, 2936-2941. | 1.6 | 82 |
| 6039 | <i>Saccharomyces cerevisiae</i> tRNA ligase. Purification of the protein and isolation of the structural gene.. Journal of Biological Chemistry, 1986, 261, 2978-2986. | 1.6 | 169 |
| 6040 | Nuclear genes coding the yeast mitochondrial adenosine triphosphatase complex. Primary sequence analysis of ATP2 encoding the F1-ATPase beta-subunit precursor.. Journal of Biological Chemistry, 1985, 260, 15458-15465. | 1.6 | 105 |
| 6041 | INO2 and INO4 gene products, positive regulators of phospholipid biosynthesis in <i>Saccharomyces cerevisiae</i> , form a complex that binds to the INO1 promoter.. Journal of Biological Chemistry, 1994, 269, 15344-15349. | 1.6 | 179 |
| 6042 | Transcriptional activation mediated by the yeast AP-1 protein is required for normal cadmium tolerance.. Journal of Biological Chemistry, 1994, 269, 14690-14697. | 1.6 | 81 |
| 6043 | Cloning and sequencing of flavin-containing monooxygenases FMO3 and FMO4 from rabbit and characterization of FMO3.. Journal of Biological Chemistry, 1994, 269, 14314-14322. | 1.6 | 48 |

| # | ARTICLE | IF | CITATIONS |
|------|---|-----|-----------|
| 6044 | NAT2, an essential gene encoding methionine N alpha-acetyltransferase in the yeast <i>Saccharomyces cerevisiae</i> .. <i>Journal of Biological Chemistry</i> , 1994, 269, 13141-13147. | 1.6 | 21 |
| 6045 | Isolation of a cDNA encoding chloroplast ferrochelatase from <i>Arabidopsis thaliana</i> by functional complementation of a yeast mutant.. <i>Journal of Biological Chemistry</i> , 1994, 269, 13405-13413. | 1.6 | 64 |
| 6046 | The prosequence of <i>Rhizopus niveus</i> aspartic proteinase-I supports correct folding and secretion of its mature part in <i>Saccharomyces cerevisiae</i> . <i>Journal of Biological Chemistry</i> , 1994, 269, 9556-9561. | 1.6 | 38 |
| 6047 | Cloning of the <i>pka1</i> gene encoding the catalytic subunit of the cAMP-dependent protein kinase in <i>Schizosaccharomyces pombe</i> . <i>Journal of Biological Chemistry</i> , 1994, 269, 9632-9637. | 1.6 | 139 |
| 6048 | Systematic mutagenesis of the yeast mating pheromone receptor third intracellular loop.. <i>Journal of Biological Chemistry</i> , 1994, 269, 8831-8841. | 1.6 | 45 |
| 6049 | Heterologous expression of human prostatic acid phosphatase and site-directed mutagenesis of the enzyme active site.. <i>Journal of Biological Chemistry</i> , 1994, 269, 8971-8978. | 1.6 | 54 |
| 6050 | A novel protein, CSG2p, is required for Ca ²⁺ regulation in <i>Saccharomyces cerevisiae</i> .. <i>Journal of Biological Chemistry</i> , 1994, 269, 7279-7284. | 1.6 | 100 |
| 6051 | Functional characterization of the <i>INO2</i> gene of <i>Saccharomyces cerevisiae</i> . A positive regulator of phospholipid biosynthesis.. <i>Journal of Biological Chemistry</i> , 1994, 269, 7402-7411. | 1.6 | 58 |
| 6052 | Leptomycin B targets a regulatory cascade of <i>crm1</i> , a fission yeast nuclear protein, involved in control of higher order chromosome structure and gene expression.. <i>Journal of Biological Chemistry</i> , 1994, 269, 6320-6324. | 1.6 | 406 |
| 6053 | The influence of beta subunit structure on the stability of Na ⁺ /K ⁽⁺⁾ -ATPase complexes and interaction with K ⁺ .. <i>Journal of Biological Chemistry</i> , 1994, 269, 6550-6557. | 1.6 | 103 |
| 6054 | Requirement of the propeptide for in vivo formation of active yeast carboxypeptidase Y.. <i>Journal of Biological Chemistry</i> , 1994, 269, 7006-7012. | 1.6 | 52 |
| 6055 | Enhanced bactericidal action of lysozyme to <i>Escherichia coli</i> by inserting a hydrophobic pentapeptide into its C terminus.. <i>Journal of Biological Chemistry</i> , 1994, 269, 5059-5063. | 1.6 | 111 |
| 6056 | Yeast mRNA splicing in vitro.. <i>Journal of Biological Chemistry</i> , 1985, 260, 14780-14792. | 1.6 | 291 |
| 6057 | Cloning and characterization of the yeast nuclear gene for subunit 5 of cytochrome oxidase.. <i>Journal of Biological Chemistry</i> , 1985, 260, 9513-9515. | 1.6 | 51 |
| 6058 | Cloning and characterization of <i>LAG1</i> , a longevity-assurance gene in yeast.. <i>Journal of Biological Chemistry</i> , 1994, 269, 15451-15459. | 1.6 | 213 |
| 6059 | Ribosomal protein P0, contrary to phosphoproteins P1 and P2, is required for ribosome activity and <i>Saccharomyces cerevisiae</i> viability.. <i>Journal of Biological Chemistry</i> , 1994, 269, 15689-15696. | 1.6 | 108 |
| 6060 | Mutagenic alteration of the distal switch II region of RAS blocks CDC25-dependent signaling functions.. <i>Journal of Biological Chemistry</i> , 1994, 269, 15740-15748. | 1.6 | 13 |
| 6061 | Effect of acidic ribosomal phosphoprotein mRNA 5'-untranslated region on gene expression and protein accumulation.. <i>Journal of Biological Chemistry</i> , 1994, 269, 3968-3975. | 1.6 | 12 |

| # | ARTICLE | IF | CITATIONS |
|------|--|-----|-----------|
| 6062 | Functional comparisons between plant plasma membrane H(+)-ATPase isoforms expressed in yeast.. Journal of Biological Chemistry, 1994, 269, 3027-3033. | 1.6 | 119 |
| 6063 | A novel gene, STT4, encodes a phosphatidylinositol 4-kinase in the PKC1 protein kinase pathway of <i>Saccharomyces cerevisiae</i> .. Journal of Biological Chemistry, 1994, 269, 1166-1172. | 1.6 | 176 |
| 6064 | PIM1 encodes a mitochondrial ATP-dependent protease that is required for mitochondrial function in the yeast <i>Saccharomyces cerevisiae</i> .. Journal of Biological Chemistry, 1994, 269, 238-242. | 1.6 | 204 |
| 6065 | Nuclear genes for mitochondrial proteins. Identification and isolation of a structural gene for subunit V of yeast cytochrome c oxidase.. Journal of Biological Chemistry, 1983, 258, 13418-13421. | 1.6 | 47 |
| 6066 | Isolation and characterization of QCR9, a nuclear gene encoding the 7.3-kDa subunit 9 of the <i>Saccharomyces cerevisiae</i> ubiquinol-cytochrome c oxidoreductase complex. An intron-containing gene with a conserved sequence occurring in the intron of COX4.. Journal of Biological Chemistry, 1990, 265, 20813-20821. | 1.6 | 46 |
| 6067 | Isolation of the GSY1 gene encoding yeast glycogen synthase and evidence for the existence of a second gene.. Journal of Biological Chemistry, 1990, 265, 20879-20886. | 1.6 | 74 |
| 6068 | The specificities of yeast methionine aminopeptidase and acetylation of amino-terminal methionine in vivo. Processing of altered iso-1-cytochromes c created by oligonucleotide transformation.. Journal of Biological Chemistry, 1990, 265, 19638-19643. | 1.6 | 184 |
| 6069 | Isolation of a temperature-sensitive mutant with an altered tRNA nucleotidyltransferase and cloning of the gene encoding tRNA nucleotidyltransferase in the yeast <i>Saccharomyces cerevisiae</i> .. Journal of Biological Chemistry, 1990, 265, 16216-16220. | 1.6 | 106 |
| 6070 | Yeast cytochrome c oxidase subunit VII is essential for assembly of an active enzyme. Cloning, sequencing, and characterization of the nuclear-encoded gene.. Journal of Biological Chemistry, 1990, 265, 16389-16393. | 1.6 | 43 |
| 6071 | Proteasomes are essential for yeast proliferation. cDNA cloning and gene disruption of two major subunits.. Journal of Biological Chemistry, 1990, 265, 16604-16613. | 1.6 | 120 |
| 6072 | The top2-5 mutant of yeast topoisomerase II encodes an enzyme resistant to etoposide and amsacrine.. Journal of Biological Chemistry, 1993, 268, 18586-18592. | 1.6 | 54 |
| 6073 | Subunit VIa of yeast cytochrome c oxidase is not necessary for assembly of the enzyme complex but modulates the enzyme activity. Isolation and characterization of the nuclear-coded gene.. Journal of Biological Chemistry, 1993, 268, 18754-18761. | 1.6 | 61 |
| 6074 | Molecular cloning of the gene for bilirubin oxidase from <i>Myrothecium verrucaria</i> and its expression in yeast.. Journal of Biological Chemistry, 1993, 268, 18801-18809. | 1.6 | 54 |
| 6075 | The propeptide is required for in vivo formation of stable active yeast proteinase A and can function even when not covalently linked to the mature region.. Journal of Biological Chemistry, 1993, 268, 18002-18007. | 1.6 | 82 |
| 6076 | VMA13 encodes a 54-kDa vacuolar H(+)-ATPase subunit required for activity but not assembly of the enzyme complex in <i>Saccharomyces cerevisiae</i> .. Journal of Biological Chemistry, 1993, 268, 18286-18292. | 1.6 | 148 |
| 6077 | Cadmium tolerance mediated by the yeast AP-1 protein requires the presence of an ATP-binding cassette transporter-encoding gene, YCF1.. Journal of Biological Chemistry, 1994, 269, 32592-32597. | 1.6 | 187 |
| 6078 | ERG10 from <i>Saccharomyces cerevisiae</i> encodes acetoacetyl-CoA thiolase.. Journal of Biological Chemistry, 1994, 269, 31383-31389. | 1.6 | 52 |
| 6079 | Vps34p required for yeast vacuolar protein sorting is a multiple specificity kinase that exhibits both protein kinase and phosphatidylinositol-specific PI 3-kinase activities.. Journal of Biological Chemistry, 1994, 269, 31552-31562. | 1.6 | 214 |

| # | ARTICLE | IF | CITATIONS |
|------|--|-----|-----------|
| 6080 | Human and Giardia ADP-ribosylation factors (ARFs) complement ARF function in <i>Saccharomyces cerevisiae</i> .. <i>Journal of Biological Chemistry</i> , 1992, 267, 24441-24445. | 1.6 | 37 |
| 6081 | Isolation and characterization of the yeast gene encoding the MDH3 isozyme of malate dehydrogenase.. <i>Journal of Biological Chemistry</i> , 1992, 267, 24708-24715. | 1.6 | 87 |
| 6082 | In vivo phosphorylation of the yeast uracil permease.. <i>Journal of Biological Chemistry</i> , 1992, 267, 23767-23771. | 1.6 | 55 |
| 6083 | SEC7 encodes an unusual, high molecular weight protein required for membrane traffic from the yeast Golgi apparatus.. <i>Journal of Biological Chemistry</i> , 1988, 263, 11711-11717. | 1.6 | 134 |
| 6084 | The MAS-encoded processing protease of yeast mitochondria. Interaction of the purified enzyme with signal peptides and a purified precursor protein.. <i>Journal of Biological Chemistry</i> , 1991, 266, 6416-6423. | 1.6 | 73 |
| 6085 | Mechanisms of camptothecin resistance in yeast DNA topoisomerase I mutants.. <i>Journal of Biological Chemistry</i> , 1993, 268, 22322-22330. | 1.6 | 82 |
| 6086 | Isolation and characterization of COX12, the nuclear gene for a previously unrecognized subunit of <i>Saccharomyces cerevisiae</i> cytochrome c oxidase.. <i>Journal of Biological Chemistry</i> , 1992, 267, 22473-22480. | 1.6 | 69 |
| 6087 | Glucose-induced degradation of the MDH2 isozyme of malate dehydrogenase in yeast.. <i>Journal of Biological Chemistry</i> , 1992, 267, 17458-17464. | 1.6 | 42 |
| 6088 | Yeast NSR1 protein that has structural similarity to mammalian nucleolin is involved in pre-rRNA processing.. <i>Journal of Biological Chemistry</i> , 1992, 267, 16252-16258. | 1.6 | 102 |
| 6089 | Cloning and characterization of the gene encoding the IDH1 subunit of NAD(+)-dependent isocitrate dehydrogenase from <i>Saccharomyces cerevisiae</i> .. <i>Journal of Biological Chemistry</i> , 1992, 267, 16417-16423. | 1.6 | 93 |
| 6090 | The mitochondrial F1ATPase alpha-subunit is necessary for efficient import of mitochondrial precursors.. <i>Journal of Biological Chemistry</i> , 1992, 267, 14697-14702. | 1.6 | 20 |
| 6091 | Cytoplasmic and mitochondrial tRNA nucleotidyltransferase activities are derived from the same gene in the yeast <i>Saccharomyces cerevisiae</i> .. <i>Journal of Biological Chemistry</i> , 1992, 267, 14879-14883. | 1.6 | 48 |
| 6092 | Coordinate regulation of glycogen metabolism in the yeast <i>Saccharomyces cerevisiae</i> . Induction of glycogen branching enzyme.. <i>Journal of Biological Chemistry</i> , 1992, 267, 15224-15228. | 1.6 | 58 |
| 6093 | Expression of cytochrome b5 in yeast and characterization of mutants of the membrane-anchoring domain.. <i>Journal of Biological Chemistry</i> , 1992, 267, 12583-12591. | 1.6 | 45 |
| 6094 | Biogenesis of the yeast vacuole (lysosome). The precursor forms of the soluble hydrolase carboxypeptidase yscS are associated with the vacuolar membrane.. <i>Journal of Biological Chemistry</i> , 1992, 267, 8021-8029. | 1.6 | 52 |
| 6095 | Structure and function of MRP20 and MRP49, the nuclear genes for two proteins of the 54 S subunit of the yeast mitochondrial ribosome.. <i>Journal of Biological Chemistry</i> , 1992, 267, 5162-5170. | 1.6 | 55 |
| 6096 | Yeast tRNA ligase mutants are nonviable and accumulate tRNA splicing intermediates.. <i>Journal of Biological Chemistry</i> , 1992, 267, 4577-4582. | 1.6 | 71 |
| 6097 | Systematic mutational analysis of cAMP-dependent protein kinase identifies unregulated catalytic subunits and defines regions important for the recognition of the regulatory subunit.. <i>Journal of Biological Chemistry</i> , 1992, 267, 4806-4814. | 1.6 | 110 |

| # | ARTICLE | IF | CITATIONS |
|------|--|-----|-----------|
| 6098 | Structural and functional studies of the amino terminus of yeast metallothionein.. Journal of Biological Chemistry, 1987, 262, 12912-12919. | 1.6 | 32 |
| 6099 | Transcriptional elements of the yeast ribosomal protein gene CYH2.. Journal of Biological Chemistry, 1987, 262, 5690-5695. | 1.6 | 46 |
| 6100 | Spermidine biosynthesis in <i>Saccharomyces cerevisiae</i> . Biosynthesis and processing of a proenzyme form of S-adenosylmethionine decarboxylase.. Journal of Biological Chemistry, 1990, 265, 22321-22328. | 1.6 | 60 |
| 6101 | Coexpression of human cAMP-specific phosphodiesterase activity and high affinity rolipram binding in yeast.. Journal of Biological Chemistry, 1992, 267, 1798-1804. | 1.6 | 131 |
| 6102 | A single amino acid mutation (Ser180→Cys) determines the polymorphism in cytochrome P450g (P4502C13) by altering protein stability.. Journal of Biological Chemistry, 1992, 267, 2032-2037. | 1.6 | 22 |
| 6103 | Uncoupling of mRNA 3' cleavage and polyadenylation by expression of a hammerhead ribozyme in yeast.. Journal of Biological Chemistry, 1994, 269, 27378-27383. | 1.6 | 16 |
| 6104 | VMA7 encodes a novel 14-kDa subunit of the <i>Saccharomyces cerevisiae</i> vacuolar H(+)-ATPase complex. Journal of Biological Chemistry, 1994, 269, 25974-25977. | 1.6 | 68 |
| 6105 | Molecular analysis of the interaction of calcineurin with drug-immunophilin complexes. Journal of Biological Chemistry, 1994, 269, 26431-26437. | 1.6 | 103 |
| 6106 | Features of vacuolar H(+)-ATPase revealed by yeast suppressor mutants. Journal of Biological Chemistry, 1994, 269, 26479-26485. | 1.6 | 27 |
| 6107 | Activators of protein kinase C stimulate association of Shc and the PEST tyrosine phosphatase.. Journal of Biological Chemistry, 1994, 269, 25243-25246. | 1.6 | 58 |
| 6108 | Isolation and characterization of the nuclear gene encoding the Rieske iron-sulfur protein (RIP1) from <i>Saccharomyces cerevisiae</i> .. Journal of Biological Chemistry, 1987, 262, 8901-8909. | 1.6 | 115 |
| 6109 | Isolation of the yeast gene encoding elongation factor 3 for protein synthesis.. Journal of Biological Chemistry, 1987, 262, 7802-7807. | 1.6 | 30 |
| 6110 | The role of protein structure in the mitochondrial import pathway. Unfolding of mitochondrially bound precursors is required for membrane translocation.. Journal of Biological Chemistry, 1987, 262, 15605-15609. | 1.6 | 114 |
| 6111 | Cloning and mutational analysis of the gene encoding subunit C of yeast vacuolar H(+)-ATPase.. Journal of Biological Chemistry, 1992, 267, 774-779. | 1.6 | 72 |
| 6112 | Assembly and targeting of peripheral and integral membrane subunits of the yeast vacuolar H(+)-ATPase.. Journal of Biological Chemistry, 1992, 267, 447-454. | 1.6 | 172 |
| 6113 | Interaction between the <i>Saccharomyces cerevisiae</i> CDC25 gene product and mammalian ras. Journal of Biological Chemistry, 1992, 267, 22747-22751. | 1.6 | 10 |
| 6114 | Biochemical and genetic dissection of the <i>Saccharomyces cerevisiae</i> RNA polymerase C53 subunit through the analysis of a mitochondrially mis-sorted mutant construct. Journal of Biological Chemistry, 1992, 267, 23099-23107. | 1.6 | 15 |
| 6115 | Ubiquitin-specific proteases of <i>Saccharomyces cerevisiae</i> . Cloning of UBP2 and UBP3, and functional analysis of the UBP gene family. Journal of Biological Chemistry, 1992, 267, 23364-23375. | 1.6 | 227 |

| # | ARTICLE | IF | CITATIONS |
|------|--|-----|-----------|
| 6116 | Isolation, nucleotide sequence, and disruption of the <i>Saccharomyces cerevisiae</i> gene encoding mitochondrial NADP(H)-specific isocitrate dehydrogenase. <i>Journal of Biological Chemistry</i> , 1991, 266, 2339-2345. | 1.6 | 83 |
| 6117 | Human ADP-ribosylation factors. A functionally conserved family of GTP-binding proteins. <i>Journal of Biological Chemistry</i> , 1991, 266, 2606-2614. | 1.6 | 179 |
| 6118 | Isolation of human glycogen branching enzyme cDNAs by screening complementation in yeast.. <i>Journal of Biological Chemistry</i> , 1993, 268, 7509-7513. | 1.6 | 87 |
| 6119 | Old Yellow Enzyme. The discovery of multiple isozymes and a family of related proteins.. <i>Journal of Biological Chemistry</i> , 1993, 268, 6097-6106. | 1.6 | 140 |
| 6120 | A low-K _m , rolipram-sensitive, cAMP-specific phosphodiesterase from human brain. Cloning and expression of cDNA, biochemical characterization of recombinant protein, and tissue distribution of mRNA.. <i>Journal of Biological Chemistry</i> , 1993, 268, 6470-6476. | 1.6 | 120 |
| 6121 | Mutagenesis of conserved residues in the phosphorylation domain of the yeast plasma membrane H(+)-ATPase. Effects on structure and function.. <i>Journal of Biological Chemistry</i> , 1993, 268, 6708-6713. | 1.6 | 38 |
| 6122 | Brefeldin A reversibly inhibits secretion in <i>Saccharomyces cerevisiae</i> .. <i>Journal of Biological Chemistry</i> , 1993, 268, 5345-5348. | 1.6 | 47 |
| 6123 | Functional characterization of flavin-containing monooxygenase 1B1 expressed in <i>Saccharomyces cerevisiae</i> and <i>Escherichia coli</i> and analysis of proposed FAD- and membrane-binding domains.. <i>Journal of Biological Chemistry</i> , 1993, 268, 5728-5734. | 1.6 | 53 |
| 6124 | Functional expression of human cathepsin S in <i>Saccharomyces cerevisiae</i> . Purification and characterization of the recombinant enzyme.. <i>Journal of Biological Chemistry</i> , 1993, 268, 4832-4838. | 1.6 | 151 |
| 6125 | The VPS16 gene product associates with a sedimentable protein complex and is essential for vacuolar protein sorting in yeast.. <i>Journal of Biological Chemistry</i> , 1993, 268, 4953-4962. | 1.6 | 101 |
| 6126 | Stimulation of calcium uptake in <i>Saccharomyces cerevisiae</i> by bovine protein kinase C alpha.. <i>Journal of Biological Chemistry</i> , 1993, 268, 3456-3462. | 1.6 | 16 |
| 6127 | Enhanced stimulation of myosin subfragment 1 ATPase activity by addition of negatively charged residues to the yeast actin NH ₂ terminus.. <i>Journal of Biological Chemistry</i> , 1993, 268, 2410-2415. | 1.6 | 89 |
| 6128 | Purification and structural characterization of transcriptional regulator Leu3 of yeast.. <i>Journal of Biological Chemistry</i> , 1993, 268, 2505-2512. | 1.6 | 25 |
| 6129 | Cloning and sequencing of a cDNA encoding <i>Saccharomyces cerevisiae</i> carnitine acetyltransferase. Use of the cDNA in gene disruption studies.. <i>Journal of Biological Chemistry</i> , 1993, 268, 1824-1829. | 1.6 | 39 |
| 6130 | VMA12 is essential for assembly of the vacuolar H(+)-ATPase subunits onto the vacuolar membrane in <i>Saccharomyces cerevisiae</i> .. <i>Journal of Biological Chemistry</i> , 1993, 268, 961-967. | 1.6 | 67 |
| 6131 | The gene PPG encodes a novel yeast protein phosphatase involved in glycogen accumulation.. <i>Journal of Biological Chemistry</i> , 1993, 268, 1349-1354. | 1.6 | 53 |
| 6132 | Cloning and characterization of the <i>Saccharomyces cerevisiae</i> gene encoding NAD-dependent 5,10-methylenetetrahydrofolate dehydrogenase.. <i>Journal of Biological Chemistry</i> , 1993, 268, 153-160. | 1.6 | 24 |
| 6133 | ATP synthase of yeast mitochondria. Isolation and disruption of the ATP epsilon gene.. <i>Journal of Biological Chemistry</i> , 1993, 268, 161-167. | 1.6 | 54 |

| # | ARTICLE | IF | CITATIONS |
|------|---|-----|-----------|
| 6134 | Identity of <i>Saccharomyces cerevisiae</i> tRNA(Trp) is not changed by an anticodon mutation that creates an amber suppressor.. <i>Journal of Biological Chemistry</i> , 1993, 268, 217-220. | 1.6 | 8 |
| 6135 | Genetic and biochemical studies of a mutant <i>Saccharomyces cerevisiae</i> myristoyl-CoA:protein N-myristoyltransferase, nmt72pLeu99â€”>Pro, that produces temperature-sensitive myristic acid auxotrophy.. <i>Journal of Biological Chemistry</i> , 1993, 268, 483-494. | 1.6 | 34 |
| 6136 | The yeast VPS17 gene encodes a membrane-associated protein required for the sorting of soluble vacuolar hydrolases.. <i>Journal of Biological Chemistry</i> , 1993, 268, 559-569. | 1.6 | 44 |
| 6137 | Mutation of invariant cysteines of mammalian metallothionein alters metal binding capacity, cadmium resistance, and ¹¹³ Cd NMR spectrum.. <i>Journal of Biological Chemistry</i> , 1991, 266, 24390-24397. | 1.6 | 28 |
| 6138 | Evidence for coupling between membrane and cytoplasmic domains of the yeast plasma membrane H(+)-ATPase. An analysis of intragenic revertants of pma1-105.. <i>Journal of Biological Chemistry</i> , 1991, 266, 24439-24445. | 1.6 | 61 |
| 6139 | The yeast GLC7 gene required for glycogen accumulation encodes a type 1 protein phosphatase.. <i>Journal of Biological Chemistry</i> , 1991, 266, 23796-23801. | 1.6 | 135 |
| 6140 | Activation of the proteinase B precursor of the yeast <i>Saccharomyces cerevisiae</i> by autocatalysis and by an internal sequence.. <i>Journal of Biological Chemistry</i> , 1991, 266, 22851-22857. | 1.6 | 54 |
| 6141 | NAD(+)-dependent isocitrate dehydrogenase. Cloning, nucleotide sequence, and disruption of the IDH2 gene from <i>Saccharomyces cerevisiae</i> .. <i>Journal of Biological Chemistry</i> , 1991, 266, 22199-22205. | 1.6 | 123 |
| 6142 | Expression of duck lens delta-crystallin cDNAs in yeast and bacterial hosts. Delta 2-crystallin is an active argininosuccinate lyase.. <i>Journal of Biological Chemistry</i> , 1991, 266, 22319-22322. | 1.6 | 55 |
| 6143 | Mutational analysis of the mitochondrial Rieske iron-sulfur protein of <i>Saccharomyces cerevisiae</i> . III. Import, protease processing, and assembly into the cytochrome bc ₁ complex of iron-sulfur protein lacking the iron-sulfur cluster.. <i>Journal of Biological Chemistry</i> , 1991, 266, 22485-22492. | 1.6 | 58 |
| 6144 | The yeast mitochondrial RNA polymerase specificity factor, MTF1, is similar to bacterial sigma factors.. <i>Journal of Biological Chemistry</i> , 1991, 266, 22671-22677. | 1.6 | 141 |
| 6145 | The SV40 core sequence functions as a repressor element in yeast.. <i>Journal of Biological Chemistry</i> , 1991, 266, 21362-21367. | 1.6 | 4 |
| 6146 | Zinc-binding subunits of yeast RNA polymerases.. <i>Journal of Biological Chemistry</i> , 1991, 266, 21971-21976. | 1.6 | 67 |
| 6147 | Mitochondrial phosphate transport. The <i>Saccharomyces cerevisiae</i> (threonine 43 to cysteine) mutant protein explicitly identifies transport with genomic sequence.. <i>Journal of Biological Chemistry</i> , 1991, 266, 19882-19885. | 1.6 | 27 |
| 6148 | Transcriptional activation of yeast nucleotide biosynthetic gene ADE4 by GCN4.. <i>Journal of Biological Chemistry</i> , 1991, 266, 20453-20456. | 1.6 | 47 |
| 6149 | Chitinase is required for cell separation during growth of <i>Saccharomyces cerevisiae</i> .. <i>Journal of Biological Chemistry</i> , 1991, 266, 19758-19767. | 1.6 | 524 |
| 6150 | Transactivation functions facilitate the disruption of chromatin structure by estrogen receptor derivatives in vivo.. <i>Journal of Biological Chemistry</i> , 1991, 266, 18179-18187. | 1.6 | 48 |
| 6151 | The pma1 and pma2 H(+)-ATPases from <i>Schizosaccharomyces pombe</i> are functionally interchangeable.. <i>Journal of Biological Chemistry</i> , 1991, 266, 18276-18279. | 1.6 | 39 |

| # | ARTICLE | IF | CITATIONS |
|------|---|-----|-----------|
| 6152 | Secretion and lipid association of human apolipoprotein E in <i>Saccharomyces cerevisiae</i> requires a host mutation in sterol esterification and uptake.. <i>Journal of Biological Chemistry</i> , 1991, 266, 16273-16276. | 1.6 | 8 |
| 6153 | Expression of a hybrid Cu/Zn-type superoxide dismutase which has high affinity for heparin-like proteoglycans on vascular endothelial cells.. <i>Journal of Biological Chemistry</i> , 1991, 266, 16409-16414. | 1.6 | 34 |
| 6154 | ATP synthase of yeast mitochondria. Characterization of subunit d and sequence analysis of the structural gene ATP7.. <i>Journal of Biological Chemistry</i> , 1991, 266, 16541-16549. | 1.6 | 66 |
| 6155 | Ubiquinone biosynthesis in <i>Saccharomyces cerevisiae</i> . Isolation and sequence of COQ3, the 3,4-dihydroxy-5-hexaprenylbenzoate methyltransferase gene.. <i>Journal of Biological Chemistry</i> , 1991, 266, 16636-16644. | 1.6 | 73 |
| 6156 | Unusual metabolism of the yeast actin amino terminus.. <i>Journal of Biological Chemistry</i> , 1991, 266, 16825-16833. | 1.6 | 61 |
| 6157 | Gene-enzyme relationship in the sulfate assimilation pathway of <i>Saccharomyces cerevisiae</i> . Study of the 3'-phosphoadenylylsulfate reductase structural gene.. <i>Journal of Biological Chemistry</i> , 1990, 265, 15518-15524. | 1.6 | 67 |
| 6158 | Protein synthesis in yeast. Structural and functional analysis of the gene encoding elongation factor 3.. <i>Journal of Biological Chemistry</i> , 1990, 265, 15838-15844. | 1.6 | 63 |
| 6159 | Primary structure of the yeast choline transport gene and regulation of its expression.. <i>Journal of Biological Chemistry</i> , 1990, 265, 15996-16003. | 1.6 | 79 |
| 6160 | Metabolic Studies on Citrate Synthase Mutants of Yeast. <i>Journal of Biological Chemistry</i> , 1989, 264, 11204-11210. | 1.6 | 53 |
| 6161 | Carboxyl groups near the active site zinc contribute to catalysis in yeast alcohol dehydrogenase.. <i>Journal of Biological Chemistry</i> , 1988, 263, 5446-5454. | 1.6 | 75 |
| 6162 | Construction and analysis of deletions in the amino-terminal extension of glutamine tRNA synthetase of <i>Saccharomyces cerevisiae</i> .. <i>Journal of Biological Chemistry</i> , 1987, 262, 10807-10813. | 1.6 | 31 |
| 6163 | Molecular characterization of the cell cycle-regulated thymidylate synthase gene of <i>Saccharomyces cerevisiae</i> .. <i>Journal of Biological Chemistry</i> , 1987, 262, 5298-5307. | 1.6 | 51 |
| 6164 | Primary structure and disruption of the phosphatidylinositol synthase gene of <i>Saccharomyces cerevisiae</i> .. <i>Journal of Biological Chemistry</i> , 1987, 262, 4876-4881. | 1.6 | 139 |
| 6165 | Isolation and characterization of the yeast gene coding for the alpha subunit of mitochondrial phenylalanyl-tRNA synthetase.. <i>Journal of Biological Chemistry</i> , 1987, 262, 3690-3696. | 1.6 | 46 |
| 6166 | Rat testosterone 7 alpha-hydroxylase. Isolation, sequence, and expression of cDNA and its developmental regulation and induction by 3-methylcholanthrene.. <i>Journal of Biological Chemistry</i> , 1987, 262, 2787-2793. | 1.6 | 94 |
| 6167 | Molecular characterization of the yeast PRT1 gene in which mutations affect translation initiation and regulation of cell proliferation.. <i>Journal of Biological Chemistry</i> , 1987, 262, 2845-2851. | 1.6 | 63 |
| 6168 | Expression of the Four Subunits of the Torpedo Californica Nicotinic Acetylcholine Receptor in <i>Saccharomyces cerevisiae</i> . <i>Journal of Biological Chemistry</i> , 1989, 264, 15022-15027. | 1.6 | 19 |
| 6169 | Nuclear genes encoding the yeast mitochondrial ATPase complex. Analysis of ATP1 coding the F1-ATPase alpha-subunit and its assembly.. <i>Journal of Biological Chemistry</i> , 1986, 261, 15126-15133. | 1.6 | 93 |

| # | ARTICLE | IF | CITATIONS |
|------|---|-----|-----------|
| 6170 | Isolation and characterization of the TRM1 locus, a gene essential for the N2,N2-dimethylguanosine modification of both mitochondrial and cytoplasmic tRNA in <i>Saccharomyces cerevisiae</i> .. Journal of Biological Chemistry, 1986, 261, 9703-9709. | 1.6 | 130 |
| 6171 | Separate genes encode functionally equivalent ADP/ATP carrier proteins in <i>Saccharomyces cerevisiae</i> . Isolation and analysis of AAC2.. Journal of Biological Chemistry, 1988, 263, 14812-14818. | 1.6 | 160 |
| 6172 | Mutations of the yeast plasma membrane H ⁺ -ATPase which cause thermosensitivity and altered regulation of the enzyme.. Journal of Biological Chemistry, 1988, 263, 14134-14139. | 1.6 | 48 |
| 6173 | Intron sequence and structure requirements for tRNA splicing in <i>Saccharomyces cerevisiae</i> .. Journal of Biological Chemistry, 1988, 263, 13839-13847. | 1.6 | 35 |
| 6174 | The presequence of yeast 5-aminolevulinic synthase is not required for targeting to mitochondria.. Journal of Biological Chemistry, 1988, 263, 8294-8299. | 1.6 | 25 |
| 6175 | Construction, expression, and function of a new yeast amber suppressor, tRNATrpA.. Journal of Biological Chemistry, 1988, 263, 7316-7321. | 1.6 | 25 |
| 6176 | Characterization of the gene for fructose-1,6-bisphosphatase from <i>Saccharomyces cerevisiae</i> and <i>Schizosaccharomyces pombe</i> . Sequence, protein homology, and expression during growth on glucose.. Journal of Biological Chemistry, 1988, 263, 6051-6057. | 1.6 | 67 |
| 6177 | Structure and function of the yeast tRNA ligase gene.. Journal of Biological Chemistry, 1988, 263, 3171-3176. | 1.6 | 41 |
| 6178 | RPA190, the gene coding for the largest subunit of yeast RNA polymerase A.. Journal of Biological Chemistry, 1988, 263, 2830-2839. | 1.6 | 107 |
| 6179 | A subunit of yeast site-specific endonuclease Scel is a mitochondrial version of the 70-kDa heat shock protein.. Journal of Biological Chemistry, 1990, 265, 15189-15197. | 1.6 | 58 |
| 6180 | Cloning, sequence analysis, and expression of ligninolytic phenoloxidase genes of the white-rot basidiomycete <i>Coriolus hirsutus</i> .. Journal of Biological Chemistry, 1990, 265, 15224-15230. | 1.6 | 182 |
| 6181 | Dispensable presequence for cellular localization and function of mitochondrial malate dehydrogenase from <i>Saccharomyces cerevisiae</i> . Journal of Biological Chemistry, 1989, 264, 12091-12096. | 1.6 | 32 |
| 6182 | The Two <i>Xenopus laevis</i> SRC Genes Are Co-expressed and Each Produces Functional pp60. Journal of Biological Chemistry, 1989, 264, 10649-10653. | 1.6 | 46 |
| 6183 | Purification of the Yeast Centromere Binding Protein CP1 and a Mutational Analysis of Its Binding Site. Journal of Biological Chemistry, 1989, 264, 10843-10850. | 1.6 | 105 |
| 6184 | Molecular Cloning of <i>Saccharomyces cerevisiae</i> CDC6 Gene. Journal of Biological Chemistry, 1989, 264, 9022-9029. | 1.6 | 56 |
| 6185 | An RNA-binding protein gene (RBP1) of <i>Saccharomyces cerevisiae</i> encodes a putative glucose-repressible protein containing two RNA recognition motifs. Journal of Biological Chemistry, 1993, 268, 15080-15087. | 1.6 | 30 |
| 6186 | Transcription of Yeast COX6, the Gene for Cytochrome c Oxidase Subunit VI, Is Dependent On Heme and on the HAP2 Gene. Journal of Biological Chemistry, 1989, 264, 7005-7008. | 1.6 | 44 |
| 6187 | Cloning and characterization of the iron-sulfur subunit gene of succinate dehydrogenase from <i>Saccharomyces cerevisiae</i> .. Journal of Biological Chemistry, 1990, 265, 10419-10423. | 1.6 | 75 |

| # | ARTICLE | IF | CITATIONS |
|------|--|-----|-----------|
| 6188 | Cloning and Characterization of the Yeast CKI Gene Encoding Choline Kinase and Its Expression in Escherichia coli. Journal of Biological Chemistry, 1989, 264, 2053-2059. | 1.6 | 89 |
| 6189 | Characterization of a yeast nuclear gene (MST1) coding for the mitochondrial threonyl-tRNA1 synthetase.. Journal of Biological Chemistry, 1985, 260, 15362-15370. | 1.6 | 79 |
| 6190 | Two glycogen synthase isoforms in Saccharomyces cerevisiae are coded by distinct genes that are differentially controlled. Journal of Biological Chemistry, 1991, 266, 15602-15607. | 1.6 | 140 |
| 6191 | Missense mutation serine106→proline causes 17 alpha-hydroxylase deficiency. Journal of Biological Chemistry, 1991, 266, 15992-15998. | 1.6 | 125 |
| 6192 | RPC19, the gene for a subunit common to yeast RNA polymerases A (I) and C (III). Journal of Biological Chemistry, 1991, 266, 15300-15307. | 1.6 | 107 |
| 6193 | Cloning and characterization of a Saccharomyces cerevisiae gene encoding a new member of the ubiquitin-conjugating protein family. Journal of Biological Chemistry, 1991, 266, 15549-15554. | 1.6 | 42 |
| 6194 | Cloning and expression of a yeast protein tyrosine phosphatase. Journal of Biological Chemistry, 1991, 266, 12964-12970. | 1.6 | 59 |
| 6195 | Heterologous expression of peptide hormone precursors in the yeast Saccharomyces cerevisiae. Evidence for a novel prohormone endoprotease with specificity for monobasic amino acids. Journal of Biological Chemistry, 1991, 266, 13203-13209. | 1.6 | 18 |
| 6196 | Isolation and deduced amino acid sequence of the gene encoding gp115, a yeast glycopospholipid-anchored protein containing a serine-rich region. Journal of Biological Chemistry, 1991, 266, 12242-12248. | 1.6 | 80 |
| 6197 | Heat shock factor-independent heat control of transcription of the CTT1 gene encoding the cytosolic catalase T of Saccharomyces cerevisiae. Journal of Biological Chemistry, 1991, 266, 12406-12411. | 1.6 | 133 |
| 6198 | Two yeast genes encoding calmodulin-dependent protein kinases. Isolation, sequencing and bacterial expressions of CMK1 and CMK2. Journal of Biological Chemistry, 1991, 266, 12784-12794. | 1.6 | 73 |
| 6199 | Cloning and functional analysis of the ubiquitin-specific protease gene UBP1 of Saccharomyces cerevisiae. Journal of Biological Chemistry, 1991, 266, 12021-12028. | 1.6 | 146 |
| 6200 | Isolation and characterization of two distinct myo-inositol transporter genes of Saccharomyces cerevisiae. Journal of Biological Chemistry, 1991, 266, 11184-11191. | 1.6 | 127 |
| 6201 | The Saccharomyces cerevisiae LOS1 gene involved in pre-tRNA splicing encodes a nuclear protein that behaves as a component of the nuclear matrix.. Journal of Biological Chemistry, 1993, 268, 19436-19444. | 1.6 | 37 |
| 6202 | Transcriptional activation in yeast by the proline-rich activation domain of human CTF1.. Journal of Biological Chemistry, 1993, 268, 20866-20869. | 1.6 | 41 |
| 6203 | Yeast MCK1 protein kinase autophosphorylates at tyrosine and serine but phosphorylates exogenous substrates at serine and threonine.. Journal of Biological Chemistry, 1993, 268, 21155-21164. | 1.6 | 36 |
| 6204 | Function of the hydrophilic carboxyl terminus of type II DNA topoisomerase from Drosophila melanogaster. I. In vitro studies.. Journal of Biological Chemistry, 1993, 268, 21328-21334. | 1.6 | 48 |
| 6205 | Function of the hydrophilic carboxyl terminus of type II DNA topoisomerase from Drosophila melanogaster. II. In vivo studies.. Journal of Biological Chemistry, 1993, 268, 21335-21343. | 1.6 | 36 |

| # | ARTICLE | IF | CITATIONS |
|------|---|-----|-----------|
| 6206 | A third ADP/ATP translocator gene in yeast.. Journal of Biological Chemistry, 1990, 265, 12711-12716. | 1.6 | 190 |
| 6207 | Regulation of yeast LEU2. Total deletion of regulatory gene LEU3 unmasks GCN4-dependent basal level expression of LEU2.. Journal of Biological Chemistry, 1990, 265, 11667-11675. | 1.6 | 44 |
| 6208 | Yeast cAMP-dependent protein kinase regulatory subunit mutations display a variety of phenotypes.. Journal of Biological Chemistry, 1990, 265, 11897-11904. | 1.6 | 41 |
| 6209 | The CDC42 homologue from <i>Caenorhabditis elegans</i> . Complementation of yeast mutation. Journal of Biological Chemistry, 1993, 268, 13280-13285. | 1.6 | 58 |
| 6210 | Characterization of human tissue factor pathway inhibitor variants expressed in <i>Saccharomyces cerevisiae</i> . Journal of Biological Chemistry, 1993, 268, 13344-13351. | 1.6 | 32 |
| 6211 | Binding and hydrolysis of guanine nucleotides by Sec4p, a yeast protein involved in the regulation of vesicular traffic.. Journal of Biological Chemistry, 1990, 265, 9366-9372. | 1.6 | 95 |
| 6212 | Rat C1-tetrahydrofolate synthase. cDNA isolation, tissue-specific levels of the mRNA, and expression of the protein in yeast.. Journal of Biological Chemistry, 1990, 265, 7907-7913. | 1.6 | 39 |
| 6213 | The ferrochelatase from <i>Saccharomyces cerevisiae</i> . Sequence, disruption, and expression of its structural gene HEM15.. Journal of Biological Chemistry, 1990, 265, 7278-7283. | 1.6 | 90 |
| 6214 | Molecular structure of a gene, VMA1, encoding the catalytic subunit of H(+)-translocating adenosine triphosphatase from vacuolar membranes of <i>Saccharomyces cerevisiae</i> .. Journal of Biological Chemistry, 1990, 265, 6726-6733. | 1.6 | 441 |
| 6215 | Identification and purification of a yeast transcriptional trans-activator. The yeast homolog of the Rous sarcoma virus internal enhancer binding factor.. Journal of Biological Chemistry, 1990, 265, 6131-6138. | 1.6 | 5 |
| 6216 | Activation of ATP hydrolysis by an uncoupler in mutant mitochondria lacking an intrinsic ATPase inhibitor in yeast.. Journal of Biological Chemistry, 1990, 265, 6274-6278. | 1.6 | 46 |
| 6217 | Synthesis and assembly of functional mammalian Na,K-ATPase in yeast.. Journal of Biological Chemistry, 1990, 265, 4189-4192. | 1.6 | 148 |
| 6218 | The gene encoding the phosphatidylinositol transfer protein is essential for cell growth.. Journal of Biological Chemistry, 1990, 265, 4711-4717. | 1.6 | 110 |
| 6219 | The <i>Saccharomyces cerevisiae</i> INO4 gene encodes a small, highly basic protein required for derepression of phospholipid biosynthetic enzymes.. Journal of Biological Chemistry, 1990, 265, 4736-4745. | 1.6 | 86 |
| 6220 | Characterization of a nonglycosylated single chain urinary plasminogen activator secreted from yeast.. Journal of Biological Chemistry, 1990, 265, 801-807. | 1.6 | 38 |
| 6221 | Farnesyl diphosphate synthetase. Journal of Biological Chemistry, 1989, 264, 19176-19184. | 1.6 | 146 |
| 6222 | TIP 1, a cold shock-inducible gene of <i>Saccharomyces cerevisiae</i> .. Journal of Biological Chemistry, 1991, 266, 17537-17544. | 1.6 | 107 |
| 6223 | Rat procathepsin B. Proteolytic processing to the mature form in vitro.. Journal of Biological Chemistry, 1992, 267, 15993-15999. | 1.6 | 162 |

| # | ARTICLE | IF | CITATIONS |
|------|---|-----|-----------|
| 6224 | Molecular cloning and analysis of a yeast protein phosphatase with an unusual amino-terminal region.. Journal of Biological Chemistry, 1992, 267, 11734-11740. | 1.6 | 64 |
| 6225 | Functional expression of plant plasma membrane H(+)-ATPase in yeast endoplasmic reticulum.. Journal of Biological Chemistry, 1992, 267, 12341-12349. | 1.6 | 166 |
| 6226 | Removal of the amino-terminal acidic residues of yeast actin. Studies in vitro and in vivo.. Journal of Biological Chemistry, 1992, 267, 9430-9436. | 1.6 | 111 |
| 6227 | The <i>Saccharomyces cerevisiae</i> VMA7 gene encodes a 14-kDa subunit of the vacuolar H(+)-ATPase catalytic sector.. Journal of Biological Chemistry, 1994, 269, 24150-24155. | 1.6 | 61 |
| 6228 | Yeast beta- and beta'-coat proteins (COP). Two coatomer subunits essential for endoplasmic reticulum-to-Golgi protein traffic.. Journal of Biological Chemistry, 1994, 269, 24486-24495. | 1.6 | 124 |
| 6229 | The MAK11 protein is essential for cell growth and replication of M double-stranded RNA and is apparently a membrane-associated protein.. Journal of Biological Chemistry, 1988, 263, 1467-1475. | 1.6 | 45 |
| 6230 | Autoregulation of the yeast copper metallothionein gene depends on metal binding.. Journal of Biological Chemistry, 1988, 263, 1570-1574. | 1.6 | 35 |
| 6231 | Yeast LEU4 encodes mitochondrial and nonmitochondrial forms of alpha-isopropylmalate synthase.. Journal of Biological Chemistry, 1988, 263, 368-374. | 1.6 | 82 |
| 6232 | Fibrillar Genes Encode Both a Conserved Nucleolar Protein and a Novel Small Nucleolar RNA Involved in Ribosomal RNA Methylation in <i>Arabidopsis thaliana</i> . Journal of Biological Chemistry, 2000, 275, 27212-27220. | 1.6 | 101 |
| 6233 | Structure of <i>Saccharomyces cerevisiae</i> alg3, sec18 mutant oligosaccharides.. Journal of Biological Chemistry, 1991, 266, 5547-5551. | 1.6 | 33 |
| 6234 | Analysis of naturally occurring and site-directed mutations in the argininosuccinate lyase gene.. Journal of Biological Chemistry, 1991, 266, 5286-5290. | 1.6 | 37 |
| 6235 | Evidence that an ATPase functions in the maintenance of the acidic pH of the hamster sperm acrosome.. Journal of Biological Chemistry, 1981, 256, 4708-4711. | 1.6 | 26 |
| 6236 | Kinetic alpha-deuterium isotope effects for <i>Escherichia coli</i> purine nucleoside phosphorylase-catalyzed phosphorolysis of adenosine and inosine.. Journal of Biological Chemistry, 1981, 256, 767-772. | 1.6 | 21 |
| 6237 | Purification and characterization of a new DNA polymerase from budding yeast <i>Saccharomyces cerevisiae</i> . A probable homolog of mammalian DNA polymerase beta.. Journal of Biological Chemistry, 1993, 268, 27148-27153. | 1.6 | 40 |
| 6238 | Cloning and expression of <i>Geotrichum candidum</i> lipase II gene in yeast. Probing of the enzyme active site by site-directed mutagenesis.. Journal of Biological Chemistry, 1993, 268, 26212-26219. | 1.6 | 29 |
| 6239 | Absence of electron transport (Rho 0 state) restores growth of a manganese-superoxide dismutase-deficient <i>Saccharomyces cerevisiae</i> in hyperoxia. Evidence for electron transport as a major source of superoxide generation in vivo.. Journal of Biological Chemistry, 1993, 268, 26699-26703. | 1.6 | 139 |
| 6240 | How do zinc finger proteins distinguish between similar upstream activation sites? Comparison of the DNA-binding specificity of the GAL4 protein in vitro and in vivo.. Journal of Biological Chemistry, 1993, 268, 24699-24706. | 1.6 | 73 |
| 6241 | Amino-terminal deletions in the presequence of an imported mitochondrial protein block the targeting function and proteolytic cleavage of the presequence at the carboxy terminus.. Journal of Biological Chemistry, 1987, 262, 1420-1424. | 1.6 | 58 |

| # | ARTICLE | IF | CITATIONS |
|------|---|-----|-----------|
| 6242 | Use of directed mutagenesis to probe the role of tyrosine 198 in the catalytic mechanism of carboxypeptidase A.. Journal of Biological Chemistry, 1987, 262, 576-582. | 1.6 | 33 |
| 6243 | Isolation, complementary DNA sequence, and regulation of rat hepatic lauric acid omega-hydroxylase (cytochrome P-450LA omega). Identification of a new cytochrome P-450 gene family.. Journal of Biological Chemistry, 1987, 262, 801-810. | 1.6 | 284 |
| 6244 | Sequences distal to the mitochondrial targeting sequences are necessary for the maturation of the F1-ATPase beta-subunit precursor in mitochondria.. Journal of Biological Chemistry, 1987, 262, 411-418. | 1.6 | 64 |
| 6245 | Characterization of COX9, the nuclear gene encoding the yeast mitochondrial protein cytochrome c oxidase subunit VIIa. Subunit VIIa lacks a leader peptide and is an essential component of the holoenzyme.. Journal of Biological Chemistry, 1986, 261, 17183-17191. | 1.6 | 74 |
| 6246 | Analysis and in vivo disruption of the gene coding for adenylate kinase (ADK1) in the yeast <i>Saccharomyces cerevisiae</i> .. Journal of Biological Chemistry, 1988, 263, 19468-19474. | 1.6 | 35 |
| 6247 | The sn-1,2-diacylglycerol ethanolaminophosphotransferase activity of <i>Saccharomyces cerevisiae</i> . Isolation of mutants and cloning of the EPT1 gene.. Journal of Biological Chemistry, 1988, 263, 19748-19757. | 1.6 | 70 |
| 6248 | Isolation, sequence, and regulation by oxygen of the yeast HEM13 gene coding for coproporphyrinogen oxidase.. Journal of Biological Chemistry, 1988, 263, 9718-9724. | 1.6 | 109 |
| 6249 | Regulation of <i>Saccharomyces cerevisiae</i> ornithine decarboxylase expression in response to polyamine. Journal of Biological Chemistry, 1989, 264, 18110-18118. | 1.6 | 35 |
| 6250 | Isolation and characterization of OLE1, a gene affecting fatty acid desaturation from <i>Saccharomyces cerevisiae</i> . Journal of Biological Chemistry, 1989, 264, 16537-16544. | 1.6 | 271 |
| 6251 | Secretion by yeast of the zymogen form of Mucor rennin, an aspartic proteinase of <i>Mucor pusillus</i> , and its conversion to the mature form. Journal of Biological Chemistry, 1989, 264, 16862-16866. | 1.6 | 41 |
| 6252 | The <i>Saccharomyces cerevisiae</i> SOC8-1 gene and its relationship to a nucleotide kinase. Journal of Biological Chemistry, 1989, 264, 15593-15599. | 1.6 | 7 |
| 6253 | Mutational Analysis of the Mitochondrial Rieske Iron-Sulfur Protein of <i>Saccharomyces cerevisiae</i> . Journal of Biological Chemistry, 1989, 264, 3713-3722. | 1.6 | 56 |
| 6254 | Localized mutagenesis and evidence for post-transcriptional regulation of MAK3. A putative N-acetyltransferase required for double-stranded RNA virus propagation in <i>Saccharomyces cerevisiae</i> .. Journal of Biological Chemistry, 1992, 267, 20270-20276. | 1.6 | 74 |
| 6255 | <i>Saccharomyces cerevisiae</i> PKC1 encodes a protein kinase C (PKC) homolog with a substrate specificity similar to that of mammalian PKC. Journal of Biological Chemistry, 1994, 269, 16829-16836. | 1.6 | 130 |
| 6256 | Genetic and biochemical characterization of Cu,Zn superoxide dismutase mutants in <i>Saccharomyces cerevisiae</i> . Journal of Biological Chemistry, 1991, 266, 4417-4424. | 1.6 | 67 |
| 6257 | Control of yeast glycogen synthase-2 by COOH-terminal phosphorylation.. Journal of Biological Chemistry, 1993, 268, 23799-23805. | 1.6 | 90 |
| 6258 | Purification and characterization of VDE, a site-specific endonuclease from the yeast <i>Saccharomyces cerevisiae</i> .. Journal of Biological Chemistry, 1993, 268, 21844-21853. | 1.6 | 63 |
| 6259 | The nuclear-coded subunits of yeast cytochrome c oxidase. II. The amino acid sequence of subunit VIII and a model for its disposition in the inner mitochondrial membrane.. Journal of Biological Chemistry, 1984, 259, 6571-6574. | 1.6 | 31 |

| # | ARTICLE | IF | CITATIONS |
|------|--|-----|-----------|
| 6260 | Subunit function in eukaryote cytochrome c oxidase. A mutation in the nuclear-coded subunit IV allows assembly but alters the function and stability of yeast cytochrome c oxidase.. Journal of Biological Chemistry, 1991, 266, 7688-7693. | 1.6 | 18 |
| 6261 | Expression of the yeast plasma membrane [H ⁺]ATPase in secretory vesicles. A new strategy for directed mutagenesis.. Journal of Biological Chemistry, 1991, 266, 7940-7949. | 1.6 | 146 |
| 6262 | Half-calmodulin is sufficient for cell proliferation. Expressions of N- and C-terminal halves of calmodulin in the yeast <i>Saccharomyces cerevisiae</i> .. Journal of Biological Chemistry, 1991, 266, 7008-7015. | 1.6 | 28 |
| 6263 | Identification of potential substrate-binding sites in yeast and human acyl-CoA sterol acyltransferases by mutagenesis of conserved sequences. Journal of Lipid Research, 2001, 42, 1282-1291. | 2.0 | 49 |
| 6264 | Comprehensive evaluation of isoprenoid biosynthesis regulation in <i>Saccharomyces cerevisiae</i> utilizing the Genome Reporter Matrix [®] . Journal of Lipid Research, 1999, 40, 850-860. | 2.0 | 71 |
| 6265 | Structural and functional similarities between the central eukaryotic initiation factor (eIF)4A-binding domain of mammalian eIF4G and the eIF4A-binding domain of yeast eIF4G. Biochemical Journal, 2001, 355, 223. | 1.7 | 19 |
| 6266 | A carbon-source-responsive element is required for regulation of the hypoxic ADP/ATP carrier (AAC3) isoform in <i>Saccharomyces cerevisiae</i> . Biochemical Journal, 2000, 352, 893-898. | 1.7 | 7 |
| 6267 | Cloning, sequencing and functional expression of a cDNA encoding porcine pancreatic proprocarboxypeptidase A1. FEBS Journal, 1999, 259, 719-726. | 0.2 | 8 |
| 6268 | Translation and a 42-nucleotide segment within the coding region of the mRNA encoded by the MAT alpha 1 gene are involved in promoting rapid mRNA decay in yeast.. Proceedings of the National Academy of Sciences of the United States of America, 1990, 87, 2780-2784. | 3.3 | 131 |
| 6269 | TEMPERATURE-SENSITIVE LETHAL MUTATIONS ON YEAST CHROMOSOME <i>1</i> APPEAR TO DEFINE ONLY A SMALL NUMBER OF GENES. Genetics, 1984, 108, 67-90. | 1.2 | 71 |
| 6270 | ISOLATION AND CHARACTERIZATION OF MUTATIONS IN THE β -TUBULIN GENE OF <i>SACCHAROMYCES CEREVISIAE</i> . Genetics, 1985, 111, 715-734. | 1.2 | 252 |
| 6271 | GENETIC AND MOLECULAR ANALYSIS OF THE <i>GAL3</i> GENE IN THE EXPRESSION OF THE GALACTOSE/MELIBIOSE REGULON OF <i>SACCHAROMYCES CEREVISIAE</i> . Genetics, 1986, 113, 229-246. | 1.2 | 74 |
| 6272 | ANALYSIS OF MEIOSIS-DEFECTIVE MUTATIONS IN YEAST BY PHYSICAL MONITORING OF RECOMBINATION. Genetics, 1986, 113, 551-567. | 1.2 | 129 |
| 6273 | INDUCTION OF INTRACHROMOSOMAL RECOMBINATION IN YEAST BY INHIBITION OF THYMIDYLATE BIOSYNTHESIS. Genetics, 1986, 114, 375-392. | 1.2 | 29 |
| 6274 | THYMIDINE UTILIZATION BY <i>tut</i> MUTANTS AND FACILE CLONING OF MUTANT ALLELES BY PLASMID CONVERSION IN <i>S. CEREVISIAE</i> . Genetics, 1986, 114, 753-767. | 1.2 | 41 |
| 6275 | Cloning, Disruption and Chromosomal Mapping of Yeast LEU3, a Putative Regulatory Gene. Genetics, 1987, 115, 91-99. | 1.2 | 28 |
| 6276 | Meiotic Gene Conversion and Crossing Over Between Dispersed Homologous Sequences Occurs Frequently in <i>Saccharomyces cerevisiae</i> . Genetics, 1987, 115, 233-246. | 1.2 | 215 |
| 6277 | Protease B of <i>Saccharomyces cerevisiae</i> : Isolation and Regulation of the <i>PRB1</i> Structural Gene. Genetics, 1987, 115, 255-263. | 1.2 | 87 |

| # | ARTICLE | IF | CITATIONS |
|------|--|-----|-----------|
| 6278 | The Constitutive, Glucose-Repression-Insensitive Mutation of the Yeast MAL4 Locus Is an Alteration of the MAL43 Gene. <i>Genetics</i> , 1987, 116, 23-31. | 1.2 | 40 |
| 6279 | The Cloning and Mapping of <i>ADR6</i> , a Gene Required for Sporulation and for Expression of the Alcohol Dehydrogenase II Isozyme From <i>Saccharomyces cerevisiae</i> . <i>Genetics</i> , 1987, 116, 531-540. | 1.2 | 20 |
| 6280 | Characterization of Null Mutants of the <i>RAD55</i> Gene of <i>Saccharomyces cerevisiae</i> : Effects of Temperature, Osmotic Strength and Mating Type. <i>Genetics</i> , 1987, 116, 547-553. | 1.2 | 113 |
| 6281 | A New Type of Fusion Analysis Applicable to Many Organisms: Protein Fusions to the <i>URA3</i> Gene of Yeast. <i>Genetics</i> , 1987, 117, 5-12. | 1.2 | 46 |
| 6282 | Mutational Analysis of Meiotic and Mitotic Centromere Function in <i>Saccharomyces cerevisiae</i> . <i>Genetics</i> , 1987, 117, 203-212. | 1.2 | 119 |
| 6283 | Interactions Between Positive and Negative Regulators of <i>GCN4</i> Controlling Gene Expression and Entry Into the Yeast Cell Cycle. <i>Genetics</i> , 1987, 117, 409-419. | 1.2 | 51 |
| 6284 | Evidence for Related Functions of the <i>RNA</i> Genes of <i>Saccharomyces cerevisiae</i> . <i>Genetics</i> , 1987, 117, 619-631. | 1.2 | 59 |
| 6285 | Isolation and analysis of a novel class of suppressor of Ty insertion mutations in <i>Saccharomyces cerevisiae</i> . <i>Genetics</i> , 1988, 118, 203-212. | 1.2 | 134 |
| 6286 | Mutations affecting the tRNA-splicing endonuclease activity of <i>Saccharomyces cerevisiae</i> . <i>Genetics</i> , 1988, 118, 609-617. | 1.2 | 90 |
| 6287 | Analysis of the mechanism for reversion of a disrupted gene. <i>Genetics</i> , 1988, 119, 237-247. | 1.2 | 119 |
| 6288 | Unlinked noncomplementation: isolation of new conditional-lethal mutations in each of the tubulin genes of <i>Saccharomyces cerevisiae</i> . <i>Genetics</i> , 1988, 119, 249-260. | 1.2 | 210 |
| 6289 | Isolation and characterization of mutants which show an oversecretion phenotype in <i>Saccharomyces cerevisiae</i> . <i>Genetics</i> , 1988, 119, 499-506. | 1.2 | 62 |
| 6290 | The Ade6-M26 Mutation of <i>Schizosaccharomyces Pombe</i> Increases the Frequency of Crossing over. <i>Genetics</i> , 1988, 119, 507-515. | 1.2 | 72 |
| 6291 | Molecular genetics of serine and threonine catabolism in <i>Saccharomyces cerevisiae</i> . <i>Genetics</i> , 1988, 119, 527-534. | 1.2 | 51 |
| 6292 | Suppressors of a <i>gpa1</i> mutation cause sterility in <i>Saccharomyces cerevisiae</i> . <i>Genetics</i> , 1988, 119, 797-804. | 1.2 | 14 |
| 6293 | Mutational analysis of the GAL4-encoded transcriptional activator protein of <i>Saccharomyces cerevisiae</i> . <i>Genetics</i> , 1988, 120, 63-74. | 1.2 | 63 |
| 6294 | The naturally occurring alleles of MAL1 in <i>Saccharomyces</i> species evolved by various mutagenic processes including chromosomal rearrangement. <i>Genetics</i> , 1988, 120, 83-93. | 1.2 | 59 |
| 6295 | Transposon tagging using Ty elements in yeast. <i>Genetics</i> , 1988, 120, 95-108. | 1.2 | 77 |

| # | ARTICLE | IF | CITATIONS |
|------|--|-----|-----------|
| 6296 | Transformation of yeast by agitation with glass beads.. Genetics, 1988, 120, 667-670. | 1.2 | 77 |
| 6297 | Identification of REO1, a gene involved in negative regulation of COX5b and ANB1 in aerobically grown <i>Saccharomyces cerevisiae</i> .. Genetics, 1988, 120, 671-680. | 1.2 | 27 |
| 6298 | <i>Saccharomyces cerevisiae</i> cho2 mutants are deficient in phospholipid methylation and cross-pathway regulation of inositol synthesis.. Genetics, 1988, 120, 909-922. | 1.2 | 160 |
| 6299 | Mutations in elongation factor EF-1 alpha affect the frequency of frameshifting and amino acid misincorporation in <i>Saccharomyces cerevisiae</i> .. Genetics, 1988, 120, 923-934. | 1.2 | 135 |
| 6300 | HOP1: a yeast meiotic pairing gene.. Genetics, 1989, 121, 445-462. | 1.2 | 288 |
| 6301 | Suppressors of yeast actin mutations.. Genetics, 1989, 121, 659-674. | 1.2 | 290 |
| 6302 | Dominant suppressors of yeast actin mutations that are reciprocally suppressed.. Genetics, 1989, 121, 675-683. | 1.2 | 98 |
| 6303 | A system of shuttle vectors and yeast host strains designed for efficient manipulation of DNA in <i>Saccharomyces cerevisiae</i> .. Genetics, 1989, 122, 19-27. | 1.2 | 8,595 |
| 6304 | Molecular evolution of the telomere-associated MAL loci of <i>Saccharomyces</i> .. Genetics, 1989, 122, 307-316. | 1.2 | 167 |
| 6305 | Mutations in the <i>Saccharomyces cerevisiae</i> opi3 gene: effects on phospholipid methylation, growth and cross-pathway regulation of inositol synthesis.. Genetics, 1989, 122, 317-330. | 1.2 | 149 |
| 6306 | Genetic and physical analysis of double-strand break repair and recombination in <i>Saccharomyces cerevisiae</i> .. Genetics, 1989, 122, 519-534. | 1.2 | 224 |
| 6307 | Disruption of the RAD52 gene alters the spectrum of spontaneous SUP4-o mutations in <i>Saccharomyces cerevisiae</i> .. Genetics, 1989, 122, 535-542. | 1.2 | 43 |
| 6308 | gcd12 mutations are gcn3-dependent alleles of GCD2, a negative regulator of GCN4 in the general amino acid control of <i>Saccharomyces cerevisiae</i> .. Genetics, 1989, 122, 543-550. | 1.2 | 31 |
| 6309 | Extrachromosomal elements cause a reduced division potential in nib 1 strains of <i>Saccharomyces cerevisiae</i> .. Genetics, 1989, 122, 749-757. | 1.2 | 9 |
| 6310 | A general screen for mutant of <i>Saccharomyces cerevisiae</i> deficient in tRNA biosynthesis.. Genetics, 1989, 123, 55-68. | 1.2 | 51 |
| 6311 | Analysis of yeast retrotransposon Ty insertions at the CAN1 locus.. Genetics, 1989, 123, 655-665. | 1.2 | 60 |
| 6312 | MEI4, a yeast gene required for meiotic recombination.. Genetics, 1989, 123, 675-682. | 1.2 | 88 |
| 6313 | Intragenic and extragenic suppressors of mutations in the heptapeptide repeat domain of <i>Saccharomyces cerevisiae</i> RNA polymerase II.. Genetics, 1989, 123, 715-724. | 1.2 | 253 |

| # | ARTICLE | IF | CITATIONS |
|------|---|-----|-----------|
| 6314 | The genetic control of direct-repeat recombination in <i>Saccharomyces</i> : the effect of rad52 and rad1 on mitotic recombination at GAL10, a transcriptionally regulated gene.. <i>Genetics</i> , 1989, 123, 725-738. | 1.2 | 256 |
| 6315 | Isolation and characterization of temperature-sensitive mutations in the RAS2 and CYR1 genes of <i>Saccharomyces cerevisiae</i> .. <i>Genetics</i> , 1989, 123, 739-748. | 1.2 | 30 |
| 6316 | Genetic instability of clathrin-deficient strains of <i>Saccharomyces cerevisiae</i> .. <i>Genetics</i> , 1990, 124, 27-38. | 1.2 | 26 |
| 6317 | Consequences of growth media, gene copy number, and regulatory mutations on the expression of the PRB1 gene of <i>Saccharomyces cerevisiae</i> .. <i>Genetics</i> , 1990, 124, 39-55. | 1.2 | 48 |
| 6318 | Mitotic chromosome transmission fidelity mutants in <i>Saccharomyces cerevisiae</i> .. <i>Genetics</i> , 1990, 124, 237-249. | 1.2 | 342 |
| 6319 | Yeast mutants sensitive to antimicrotubule drugs define three genes that affect microtubule function.. <i>Genetics</i> , 1990, 124, 251-262. | 1.2 | 168 |
| 6320 | Distance-independence of mitotic intrachromosomal recombination in <i>Saccharomyces cerevisiae</i> .. <i>Genetics</i> , 1990, 124, 263-273. | 1.2 | 51 |
| 6321 | Identification and characterization of a mutation affecting the division arrest signaling of the pheromone response pathway in <i>Saccharomyces cerevisiae</i> .. <i>Genetics</i> , 1990, 124, 275-282. | 1.2 | 23 |
| 6322 | The CCR4 gene from <i>Saccharomyces cerevisiae</i> is required for both nonfermentative and spt-mediated gene expression.. <i>Genetics</i> , 1990, 124, 283-291. | 1.2 | 118 |
| 6323 | Isolation of omnipotent suppressors in an [eta+] yeast strain.. <i>Genetics</i> , 1990, 124, 505-514. | 1.2 | 27 |
| 6324 | The subtelomeric Y' repeat family in <i>Saccharomyces cerevisiae</i> : an experimental system for repeated sequence evolution.. <i>Genetics</i> , 1990, 124, 533-545. | 1.2 | 116 |
| 6325 | Mitotic recombination among subtelomeric Y' repeats in <i>Saccharomyces cerevisiae</i> .. <i>Genetics</i> , 1990, 124, 547-559. | 1.2 | 122 |
| 6326 | A chromosome containing HOT1 preferentially receives information during mitotic interchromosomal gene conversion.. <i>Genetics</i> , 1990, 124, 561-572. | 1.2 | 60 |
| 6327 | Comparison of thermosensitive alleles of the CDC25 gene involved in the cAMP metabolism of <i>Saccharomyces cerevisiae</i> .. <i>Genetics</i> , 1990, 124, 797-806. | 1.2 | 62 |
| 6328 | Isolation and characterization of mutants constitutive for expression of the fbp1 gene of <i>Schizosaccharomyces pombe</i> .. <i>Genetics</i> , 1990, 124, 807-816. | 1.2 | 112 |
| 6329 | The SRS2 suppressor of rad6 mutations of <i>Saccharomyces cerevisiae</i> acts by channeling DNA lesions into the RAD52 DNA repair pathway.. <i>Genetics</i> , 1990, 124, 817-831. | 1.2 | 170 |
| 6330 | GAL4 mutations that separate the transcriptional activation and GAL80-interactive functions of the yeast GAL4 protein.. <i>Genetics</i> , 1990, 125, 21-27. | 1.2 | 56 |
| 6331 | Analysis of the GAL3 signal transduction pathway activating GAL4 protein-dependent transcription in <i>Saccharomyces cerevisiae</i> .. <i>Genetics</i> , 1990, 125, 281-291. | 1.2 | 100 |

| # | ARTICLE | IF | CITATIONS |
|------|---|-----|-----------|
| 6332 | TRK2 is required for low affinity K ⁺ transport in <i>Saccharomyces cerevisiae</i> .. Genetics, 1990, 125, 305-312. | 1.2 | 99 |
| 6333 | Centromeric regions control autonomous segregation tendencies in single-division meiosis of <i>Saccharomyces cerevisiae</i> .. Genetics, 1990, 125, 487-494. | 1.2 | 25 |
| 6334 | Isolation and characterization of PEP5, a gene essential for vacuolar biogenesis in <i>Saccharomyces cerevisiae</i> .. Genetics, 1990, 125, 739-752. | 1.2 | 54 |
| 6335 | Mitotic transmission of artificial chromosomes in cdc mutants of the yeast, <i>Saccharomyces cerevisiae</i> .. Genetics, 1990, 125, 763-774. | 1.2 | 68 |
| 6336 | <i>Drosophila</i> nonsense suppressors: functional analysis in <i>Saccharomyces cerevisiae</i> , <i>Drosophila</i> tissue culture cells and <i>Drosophila melanogaster</i> .. Genetics, 1990, 126, 625-637. | 1.2 | 10 |
| 6337 | kem mutations affect nuclear fusion in <i>Saccharomyces cerevisiae</i> .. Genetics, 1990, 126, 799-812. | 1.2 | 120 |
| 6338 | Decreasing gradients of gene conversion on both sides of the initiation site for meiotic recombination at the ARG4 locus in yeast.. Genetics, 1990, 126, 813-822. | 1.2 | 112 |
| 6339 | An RME1-independent pathway for sporulation control in <i>Saccharomyces cerevisiae</i> acts through IME1 transcript accumulation.. Genetics, 1990, 126, 823-835. | 1.2 | 15 |
| 6340 | Transfer RNA genes are genomic targets for de Novo transposition of the yeast retrotransposon Ty3.. Genetics, 1990, 126, 837-850. | 1.2 | 114 |
| 6341 | Gene conversion tracts stimulated by HOT1-promoted transcription are long and continuous.. Genetics, 1990, 126, 851-867. | 1.2 | 80 |
| 6342 | Allelic and ectopic interactions in recombination-defective yeast strains.. Genetics, 1991, 127, 53-60. | 1.2 | 43 |
| 6343 | The hyper-gene conversion hpr5-1 mutation of <i>Saccharomyces cerevisiae</i> is an allele of the SRS2/RADH gene.. Genetics, 1991, 127, 75-85. | 1.2 | 157 |
| 6344 | Distributive disjunction of authentic chromosomes in <i>Saccharomyces cerevisiae</i> .. Genetics, 1991, 127, 475-488. | 1.2 | 74 |
| 6345 | The SNF2, SNF5 and SNF6 genes are required for Ty transcription in <i>Saccharomyces cerevisiae</i> .. Genetics, 1991, 128, 69-77. | 1.2 | 101 |
| 6346 | The mechanism of inducer formation in gal3 mutants of the yeast galactose system is independent of normal galactose metabolism and mitochondrial respiratory function.. Genetics, 1991, 128, 233-239. | 1.2 | 32 |
| 6347 | Interaction of the yeast omnipotent suppressors SUP1(SUP45) and SUP2(SUP35) with non-mendelian factors.. Genetics, 1991, 128, 513-520. | 1.2 | 51 |
| 6348 | Genetic analysis of the gyrase A-like domain of DNA topoisomerase II of <i>Saccharomyces cerevisiae</i> .. Genetics, 1991, 128, 703-716. | 1.2 | 25 |
| 6349 | Analysis of a circular derivative of <i>Saccharomyces cerevisiae</i> chromosome III: a physical map and identification and location of ARS elements.. Genetics, 1991, 129, 343-357. | 1.2 | 136 |

| # | ARTICLE | IF | CITATIONS |
|------|---|-----|-----------|
| 6350 | New SNF genes, GAL11 and GRR1 affect SUC2 expression in <i>Saccharomyces cerevisiae</i> .. Genetics, 1991, 129, 675-684. | 1.2 | 79 |
| 6351 | SUM1-1: a suppressor of silencing defects in <i>Saccharomyces cerevisiae</i> .. Genetics, 1991, 129, 685-696. | 1.2 | 46 |
| 6352 | Extrachromosomal recombination is deranged in the rec2 mutant of <i>Ustilago maydis</i> .. Genetics, 1991, 129, 1053-1060. | 1.2 | 18 |
| 6353 | The rec102 mutant of yeast is defective in meiotic recombination and chromosome synapsis.. Genetics, 1992, 130, 59-69. | 1.2 | 59 |
| 6354 | Relationship of the cAMP-dependent protein kinase pathway to the SNF1 protein kinase and invertase expression in <i>Saccharomyces cerevisiae</i> .. Genetics, 1992, 130, 71-80. | 1.2 | 74 |
| 6355 | Seventeen complementation groups of mutations decreasing meiotic recombination in <i>Schizosaccharomyces pombe</i> .. Genetics, 1992, 130, 251-262. | 1.2 | 115 |
| 6356 | Yeast KRE2 defines a new gene family encoding probable secretory proteins, and is required for the correct N-glycosylation of proteins.. Genetics, 1992, 130, 273-283. | 1.2 | 105 |
| 6357 | Elimination of the yeast RAD6 ubiquitin conjugase enhances base-pair transitions and G.C→T.A transversions as well as transposition of the Ty element: implications for the control of spontaneous mutation.. Genetics, 1992, 130, 285-294. | 1.2 | 64 |
| 6358 | Analysis of URSG-mediated glucose repression of the GAL1 promoter of <i>Saccharomyces cerevisiae</i> .. Genetics, 1992, 130, 295-304. | 1.2 | 59 |
| 6359 | Site-specific recombination determined by I-SceI, a mitochondrial group I intron-encoded endonuclease expressed in the yeast nucleus.. Genetics, 1992, 130, 451-460. | 1.2 | 213 |
| 6360 | Temperature-sensitive cdc7 mutations of <i>Saccharomyces cerevisiae</i> are suppressed by the DBF4 gene, which is required for the G1/S cell cycle transition.. Genetics, 1992, 131, 21-29. | 1.2 | 101 |
| 6361 | Fitness effects of Ty transposition in <i>Saccharomyces cerevisiae</i> .. Genetics, 1992, 131, 31-42. | 1.2 | 98 |
| 6362 | Meiosis in <i>Saccharomyces cerevisiae</i> mutants lacking the centromere-binding protein CP1.. Genetics, 1992, 131, 43-53. | 1.2 | 22 |
| 6363 | Isolation and characterization of extragenic suppressors of mutations in the SSA hsp70 genes of <i>Saccharomyces cerevisiae</i> .. Genetics, 1992, 131, 277-285. | 1.2 | 38 |
| 6364 | Isolation and characterization of SUA5, a novel gene required for normal growth in <i>Saccharomyces cerevisiae</i> .. Genetics, 1992, 131, 791-801. | 1.2 | 45 |
| 6365 | Rearrangements occurring adjacent to a single Ty1 yeast retrotransposon in the presence and absence of full-length Ty1 transcription.. Genetics, 1992, 131, 833-850. | 1.2 | 12 |
| 6366 | Analysis of mitotic and meiotic defects in <i>Saccharomyces cerevisiae</i> SRS2 DNA helicase mutants.. Genetics, 1992, 132, 23-37. | 1.2 | 99 |
| 6367 | Possible cross-regulation of phosphate and sulfate metabolism in <i>Saccharomyces cerevisiae</i> .. Genetics, 1992, 132, 63-73. | 1.2 | 50 |

| # | ARTICLE | IF | CITATIONS |
|------|---|-----|-----------|
| 6368 | Meiotically induced <i>rec7</i> and <i>rec8</i> genes of <i>Schizosaccharomyces pombe</i> .. <i>Genetics</i> , 1992, 132, 75-85. | 1.2 | 60 |
| 6369 | <i>MAK10</i> , a glucose-repressible gene necessary for replication of a dsRNA virus of <i>Saccharomyces cerevisiae</i> , has T cell receptor alpha-subunit motifs.. <i>Genetics</i> , 1992, 132, 87-96. | 1.2 | 34 |
| 6370 | <i>cis-</i> and <i>trans-</i> acting suppressors of a translation initiation defect at the <i>cyc1</i> locus of <i>Saccharomyces cerevisiae</i> .. <i>Genetics</i> , 1992, 132, 97-112. | 1.2 | 38 |
| 6371 | <i>SPT4</i> , <i>SPT5</i> and <i>SPT6</i> interactions: effects on transcription and viability in <i>Saccharomyces cerevisiae</i> .. <i>Genetics</i> , 1992, 132, 325-336. | 1.2 | 174 |
| 6372 | Systematic mutational analysis of the yeast <i>ACT1</i> gene.. <i>Genetics</i> , 1992, 132, 337-350. | 1.2 | 330 |
| 6373 | Identification and characterization of three genes that affect expression of <i>ADH2</i> in <i>Saccharomyces cerevisiae</i> .. <i>Genetics</i> , 1992, 132, 351-359. | 1.2 | 20 |
| 6374 | A mutant tRNA affects delta-mediated transcription in <i>Saccharomyces cerevisiae</i> .. <i>Genetics</i> , 1992, 132, 361-374. | 1.2 | 15 |
| 6375 | Sister chromatids are preferred over homologs as substrates for recombinational repair in <i>Saccharomyces cerevisiae</i> .. <i>Genetics</i> , 1992, 132, 387-402. | 1.2 | 474 |
| 6376 | N-terminal mutations modulate yeast <i>SNF1</i> protein kinase function.. <i>Genetics</i> , 1992, 132, 639-650. | 1.2 | 114 |
| 6377 | <i>XRS2</i> , a DNA repair gene of <i>Saccharomyces cerevisiae</i> , is needed for meiotic recombination.. <i>Genetics</i> , 1992, 132, 651-664. | 1.2 | 204 |
| 6378 | Donor locus selection during <i>Saccharomyces cerevisiae</i> mating type interconversion responds to distant regulatory signals.. <i>Genetics</i> , 1992, 132, 929-942. | 1.2 | 40 |
| 6379 | The <i>CCR4</i> protein from <i>Saccharomyces cerevisiae</i> contains a leucine-rich repeat region which is required for its control of <i>ADH2</i> gene expression.. <i>Genetics</i> , 1992, 132, 951-962. | 1.2 | 51 |
| 6380 | Characterization of insertion mutations in the <i>Saccharomyces cerevisiae</i> <i>MSH1</i> and <i>MSH2</i> genes: evidence for separate mitochondrial and nuclear functions.. <i>Genetics</i> , 1992, 132, 975-985. | 1.2 | 357 |
| 6381 | <i>HTS1</i> encodes both the cytoplasmic and mitochondrial histidyl-tRNA synthetase of <i>Saccharomyces cerevisiae</i> : mutations alter the specificity of compartmentation.. <i>Genetics</i> , 1992, 132, 987-1001. | 1.2 | 48 |
| 6382 | Identification of <i>Aspergillus</i> <i>brlA</i> response elements (BREs) by genetic selection in yeast.. <i>Genetics</i> , 1993, 133, 29-38. | 1.2 | 138 |
| 6383 | Telomere-mediated plasmid segregation in <i>Saccharomyces cerevisiae</i> involves gene products required for transcriptional repression at silencers and telomeres.. <i>Genetics</i> , 1993, 133, 171-182. | 1.2 | 30 |
| 6384 | α -control of DNA repair in the yeast <i>Saccharomyces cerevisiae</i> : genetic and physiological aspects.. <i>Genetics</i> , 1993, 133, 489-498. | 1.2 | 84 |
| 6385 | A ubiquitin-conjugating enzyme, <i>RAD6</i> , affects the distribution of <i>Ty1</i> retrotransposon integration positions.. <i>Genetics</i> , 1993, 133, 499-508. | 1.2 | 58 |

| # | ARTICLE | IF | CITATIONS |
|------|---|-----|-----------|
| 6386 | A mutational analysis of killer toxin resistance in <i>Saccharomyces cerevisiae</i> identifies new genes involved in cell wall (1->6)-beta-glucan synthesis.. <i>Genetics</i> , 1993, 133, 837-849. | 1.2 | 110 |
| 6387 | Stimulation of meiotic recombination in yeast by an ARS element.. <i>Genetics</i> , 1993, 134, 175-188. | 1.2 | 24 |
| 6388 | Mutations in POL1 increase the mitotic instability of tandem inverted repeats in <i>Saccharomyces cerevisiae</i> .. <i>Genetics</i> , 1993, 134, 43-56. | 1.2 | 54 |
| 6389 | Isolation and characterization of SGE1: a yeast gene that partially suppresses the gal11 mutation in multiple copies.. <i>Genetics</i> , 1993, 134, 675-683. | 1.2 | 19 |
| 6390 | Positive and negative regulatory elements control expression of the yeast retrotransposon Ty3.. <i>Genetics</i> , 1993, 134, 685-700. | 1.2 | 56 |
| 6391 | Directionality of fission yeast mating-type interconversion is controlled by the location of the donor loci.. <i>Genetics</i> , 1993, 134, 1045-1054. | 1.2 | 84 |
| 6392 | Molecular and genetic analysis of the SNF7 gene in <i>Saccharomyces cerevisiae</i> .. <i>Genetics</i> , 1993, 135, 17-23. | 1.2 | 39 |
| 6393 | Screens for extragenic mutations that fail to complement act1 alleles identify genes that are important for actin function in <i>Saccharomyces cerevisiae</i> .. <i>Genetics</i> , 1993, 135, 265-274. | 1.2 | 52 |
| 6394 | Genetic evidence for functional interactions between actin noncomplementing (Anc) gene products and actin cytoskeletal proteins in <i>Saccharomyces cerevisiae</i> .. <i>Genetics</i> , 1993, 135, 275-286. | 1.2 | 33 |
| 6395 | Point mutations that separate the role of <i>Saccharomyces cerevisiae</i> centromere binding factor 1 in chromosome segregation from its role in transcriptional activation.. <i>Genetics</i> , 1993, 135, 287-296. | 1.2 | 27 |
| 6396 | Mixed segregation and recombination of chromosomes and YACs during single-division meiosis in spo13 strains of <i>Saccharomyces cerevisiae</i> .. <i>Genetics</i> , 1993, 135, 297-308. | 1.2 | 65 |
| 6397 | A rare tRNA-Arg(CCU) that regulates Ty1 element ribosomal frameshifting is essential for Ty1 retrotransposition in <i>Saccharomyces cerevisiae</i> .. <i>Genetics</i> , 1993, 135, 309-320. | 1.2 | 127 |
| 6398 | Genetic and molecular characterization of GAL83: its interaction and similarities with other genes involved in glucose repression in <i>Saccharomyces cerevisiae</i> .. <i>Genetics</i> , 1993, 135, 655-664. | 1.2 | 67 |
| 6399 | Mutations that suppress the deletion of an upstream activating sequence in yeast: involvement of a protein kinase and histone H3 in repressing transcription in vivo.. <i>Genetics</i> , 1993, 135, 665-676. | 1.2 | 148 |
| 6400 | Isolation and characterization of chromosome-gain and increase-in-ploidy mutants in yeast.. <i>Genetics</i> , 1993, 135, 677-691. | 1.2 | 281 |
| 6401 | Molecular genetics of cryptopleurine resistance in <i>Saccharomyces cerevisiae</i> : expression of a ribosomal protein gene family.. <i>Genetics</i> , 1993, 135, 719-730. | 1.2 | 23 |
| 6402 | The pheromone receptors inhibit the pheromone response pathway in <i>Saccharomyces cerevisiae</i> by a process that is independent of their associated G alpha protein.. <i>Genetics</i> , 1993, 135, 943-953. | 1.2 | 22 |
| 6403 | Spo12 is a limiting factor that interacts with the cell cycle protein kinases Dbf2 and Dbf20, which are involved in mitotic chromatid disjunction.. <i>Genetics</i> , 1993, 135, 963-971. | 1.2 | 34 |

| # | ARTICLE | IF | CITATIONS |
|------|--|-----|-----------|
| 6404 | Translational maintenance of frame: mutants of <i>Saccharomyces cerevisiae</i> with altered -1 ribosomal frameshifting efficiencies.. <i>Genetics</i> , 1994, 136, 75-86. | 1.2 | 86 |
| 6405 | The SPT10 and SPT21 genes of <i>Saccharomyces cerevisiae</i> .. <i>Genetics</i> , 1994, 136, 93-105. | 1.2 | 45 |
| 6406 | The Yak1 protein kinase of <i>Saccharomyces cerevisiae</i> moderates thermotolerance and inhibits growth by an Sch9 protein kinase-independent mechanism.. <i>Genetics</i> , 1994, 136, 465-474. | 1.2 | 51 |
| 6407 | Characterization of glycogen-deficient <i>glc</i> mutants of <i>Saccharomyces cerevisiae</i> .. <i>Genetics</i> , 1994, 136, 485-503. | 1.2 | 157 |
| 6408 | Transient, meiosis-induced expression of the <i>rec6</i> and <i>rec12</i> genes of <i>Schizosaccharomyces pombe</i> .. <i>Genetics</i> , 1994, 136, 769-779. | 1.2 | 101 |
| 6409 | Extragenic suppressors of <i>Saccharomyces cerevisiae</i> <i>prp4</i> mutations identify a negative regulator of PRP genes.. <i>Genetics</i> , 1994, 136, 833-847. | 1.2 | 29 |
| 6410 | Suppressors reveal two classes of glucose repression genes in the yeast <i>Saccharomyces cerevisiae</i> .. <i>Genetics</i> , 1994, 136, 1271-1278. | 1.2 | 44 |
| 6411 | Isolation and characterization of a mutant of <i>Saccharomyces cerevisiae</i> with pleiotropic deficiencies in transcriptional activation and repression.. <i>Genetics</i> , 1994, 137, 55-65. | 1.2 | 52 |
| 6412 | Overexpression of yeast homologs of the mammalian checkpoint gene <i>RCC1</i> suppresses the class of alpha-tubulin mutations that arrest with excess microtubules.. <i>Genetics</i> , 1994, 137, 381-392. | 1.2 | 50 |
| 6413 | Unrepaired heteroduplex DNA in <i>Saccharomyces cerevisiae</i> is decreased in <i>RAD1 RAD52</i> -independent recombination.. <i>Genetics</i> , 1994, 137, 393-405. | 1.2 | 94 |
| 6414 | <i>BFR1</i> , a multicopy suppressor of brefeldin A-induced lethality, is implicated in secretion and nuclear segregation in <i>Saccharomyces cerevisiae</i> .. <i>Genetics</i> , 1994, 137, 423-437. | 1.2 | 43 |
| 6415 | Specificity of the yeast <i>rev3 delta</i> antimutator and <i>REV3</i> dependency of the mutator resulting from a defect (<i>rad1 delta</i>) in nucleotide excision repair.. <i>Genetics</i> , 1994, 137, 637-646. | 1.2 | 96 |
| 6416 | The <i>Saccharomyces cerevisiae</i> <i>SPT8</i> gene encodes a very acidic protein that is functionally related to <i>SPT3</i> and TATA-binding protein.. <i>Genetics</i> , 1994, 137, 647-657. | 1.2 | 86 |
| 6417 | The dominant <i>PNM2-</i> mutation which eliminates the <i>psi</i> factor of <i>Saccharomyces cerevisiae</i> is the result of a missense mutation in the <i>SUP35</i> gene.. <i>Genetics</i> , 1994, 137, 659-670. | 1.2 | 194 |
| 6418 | The <i>SUP35</i> omnipotent suppressor gene is involved in the maintenance of the non-Mendelian determinant [<i>psi+</i>] in the yeast <i>Saccharomyces cerevisiae</i> .. <i>Genetics</i> , 1994, 137, 671-676. | 1.2 | 373 |
| 6419 | <i>SIP1</i> is a catabolite repression-specific negative regulator of <i>GAL</i> gene expression.. <i>Genetics</i> , 1994, 137, 689-700. | 1.2 | 11 |
| 6420 | Characterization of mutations that suppress the temperature-sensitive growth of the <i>hpr1 delta</i> mutant of <i>Saccharomyces cerevisiae</i> .. <i>Genetics</i> , 1994, 137, 945-956. | 1.2 | 51 |
| 6421 | High-copy suppression of glucose transport defects by <i>HXT4</i> and regulatory elements in the promoters of the <i>HXT</i> genes in <i>Saccharomyces cerevisiae</i> .. <i>Genetics</i> , 1994, 137, 957-966. | 1.2 | 42 |

| # | ARTICLE | IF | CITATIONS |
|------|--|-----|-----------|
| 6422 | Three additional linkage groups that repress transcription and meiotic recombination in the mating-type region of <i>Schizosaccharomyces pombe</i> .. <i>Genetics</i> , 1994, 138, 29-38. | 1.2 | 117 |
| 6423 | Glucose repression of <i>fbp1</i> transcription of <i>Schizosaccharomyces pombe</i> is partially regulated by adenylate cyclase activation by a G protein alpha subunit encoded by <i>gpa2</i> (<i>git8</i>).. <i>Genetics</i> , 1994, 138, 39-45. | 1.2 | 65 |
| 6424 | Use of a chromosomal inverted repeat to demonstrate that the <i>RAD51</i> and <i>RAD52</i> genes of <i>Saccharomyces cerevisiae</i> have different roles in mitotic recombination.. <i>Genetics</i> , 1994, 138, 587-595. | 1.2 | 132 |
| 6425 | Structure-based systematic isolation of conditional-lethal mutations in the single yeast calmodulin gene.. <i>Genetics</i> , 1994, 138, 1041-1054. | 1.2 | 47 |
| 6426 | <i>CHL12</i> , a gene essential for the fidelity of chromosome transmission in the yeast <i>Saccharomyces cerevisiae</i> .. <i>Genetics</i> , 1994, 138, 1067-1079. | 1.2 | 44 |
| 6427 | Identification of target genes regulated by homeotic proteins in <i>Drosophila melanogaster</i> through genetic selection of Ultrabithorax protein-binding sites in yeast.. <i>Genetics</i> , 1995, 139, 349-363. | 1.2 | 81 |
| 6428 | Isolation and genetic analysis of extragenic suppressors of the hyper-deletion phenotype of the <i>Saccharomyces cerevisiae</i> <i>hpr1</i> delta mutation.. <i>Genetics</i> , 1995, 139, 57-66. | 1.2 | 28 |
| 6429 | The <i>sep1</i> mutant of <i>Saccharomyces cerevisiae</i> arrests in pachytene and is deficient in meiotic recombination.. <i>Genetics</i> , 1995, 139, 495-509. | 1.2 | 49 |
| 6430 | Mutations in <i>GCR1</i> , a transcriptional activator of <i>Saccharomyces cerevisiae</i> glycolytic genes, function as suppressors of <i>gcr2</i> mutations.. <i>Genetics</i> , 1995, 139, 511-521. | 1.2 | 33 |
| 6431 | The <i>Saccharomyces cerevisiae</i> <i>SPT7</i> gene encodes a very acidic protein important for transcription in vivo.. <i>Genetics</i> , 1995, 139, 523-536. | 1.2 | 91 |
| 6432 | Regulation of <i>HIS4</i> expression by the <i>Saccharomyces cerevisiae</i> <i>SIN4</i> transcriptional regulator.. <i>Genetics</i> , 1995, 140, 103-114. | 1.2 | 74 |
| 6433 | Alterations in ribosomal protein <i>RPS28</i> can diversely affect translational accuracy in <i>Saccharomyces cerevisiae</i> .. <i>Genetics</i> , 1995, 140, 1247-1258. | 1.2 | 27 |
| 6434 | The yeast <i>MER2</i> gene is required for chromosome synapsis and the initiation of meiotic recombination.. <i>Genetics</i> , 1995, 141, 49-59. | 1.2 | 73 |
| 6435 | 5 S rRNA is involved in fidelity of translational reading frame.. <i>Genetics</i> , 1995, 141, 95-105. | 1.2 | 45 |
| 6436 | Isolation of mutants of <i>Saccharomyces cerevisiae</i> requiring DNA topoisomerase I.. <i>Genetics</i> , 1995, 141, 465-479. | 1.2 | 59 |
| 6437 | Increased expression of <i>Saccharomyces cerevisiae</i> translation elongation factor 1 alpha bypasses the lethality of a <i>TEF5</i> null allele encoding elongation factor 1 beta.. <i>Genetics</i> , 1995, 141, 481-489. | 1.2 | 36 |
| 6438 | Unigenic evolution: a novel genetic method localizes a putative leucine zipper that mediates dimerization of the <i>Saccharomyces cerevisiae</i> regulator <i>Gcr1p</i> .. <i>Genetics</i> , 1995, 141, 1263-1274. | 1.2 | 34 |
| 6439 | Suppressors of a <i>Saccharomyces cerevisiae</i> <i>pkc1</i> mutation identify alleles of the phosphatase gene <i>PTC1</i> and of a novel gene encoding a putative basic leucine zipper protein.. <i>Genetics</i> , 1995, 141, 1275-1285. | 1.2 | 71 |

| # | ARTICLE | IF | CITATIONS |
|------|---|-----|-----------|
| 6440 | Potential RNA Binding Proteins in <i>Saccharomyces cerevisiae</i> Identified as Suppressors of Temperature-Sensitive Mutations in <i>NPL3</i> . <i>Genetics</i> , 1996, 142, 103-115. | 1.2 | 90 |
| 6441 | Genetic Analysis of the <i>Saccharomyces cerevisiae</i> <i>RHO3</i> Gene, Encoding a Rho-Type Small GTPase, Provides Evidence for a Role in Bud Formation. <i>Genetics</i> , 1996, 142, 359-369. | 1.2 | 77 |
| 6442 | Genetic Requirements for the Single-Strand Annealing Pathway of Double-Strand Break Repair in <i>Saccharomyces cerevisiae</i> . <i>Genetics</i> , 1996, 142, 693-704. | 1.2 | 368 |
| 6443 | Mutations in the RNA Polymerase II Transcription Machinery Suppress the Hyperrecombination Mutant <i>hpr1</i> of <i>Saccharomyces cerevisiae</i> . <i>Genetics</i> , 1996, 142, 749-759. | 1.2 | 145 |
| 6444 | Genetic Relationships Between the G Protein $\beta\gamma$ Complex, Ste5p, Ste20p and Cdc42p: Investigation of Effector Roles in the Yeast Pheromone Response Pathway. <i>Genetics</i> , 1996, 143, 103-117. | 1.2 | 62 |
| 6445 | Genetic Interactions Between <i>REG1/HEX2</i> and <i>GLC7</i> , the Gene Encoding the Protein Phosphatase Type 1 Catalytic Subunit in <i>Saccharomyces cerevisiae</i> . <i>Genetics</i> , 1996, 143, 119-127. | 1.2 | 46 |
| 6446 | Genetic Studies of the <i>PRP17</i> Gene of <i>Saccharomyces cerevisiae</i> : A Domain Essential for Function Maps to a Nonconserved Region of the Protein. <i>Genetics</i> , 1996, 143, 45-55. | 1.2 | 24 |
| 6447 | Deletion of the Gene Encoding the Cyclin-Dependent Protein Kinase <i>Pho85</i> Alters Glycogen Metabolism in <i>Saccharomyces cerevisiae</i> . <i>Genetics</i> , 1996, 143, 57-66. | 1.2 | 76 |
| 6448 | The Yeast <i>BSD2-1</i> Mutation Influences Both the Requirement for Phosphatidylinositol Transfer Protein Function and Derepression of Phospholipid Biosynthetic Gene Expression in Yeast. <i>Genetics</i> , 1996, 143, 685-697. | 1.2 | 19 |
| 6449 | <i>Loslp</i> , Involved in Yeast Pre-tRNA Splicing, Positively Regulates Members of the <i>SOL</i> Gene Family. <i>Genetics</i> , 1996, 143, 699-712. | 1.2 | 18 |
| 6450 | Isolation and Identification of Genes Activating UAS2-Dependent <i>ADH2</i> Expression in <i>Saccharomyces cerevisiae</i> . <i>Genetics</i> , 1996, 143, 1137-1148. | 1.2 | 13 |
| 6451 | Mutations in the Yeast <i>SRB2</i> General Transcription Factor Suppress <i>hpr1</i> -Induced Recombination and Show Defects in DNA Repair. <i>Genetics</i> , 1996, 143, 1533-1542. | 1.2 | 29 |
| 6452 | A Test of the Double-Strand Break Repair Model for Meiotic Recombination in <i>Saccharomyces cerevisiae</i> . <i>Genetics</i> , 1996, 144, 27-41. | 1.2 | 131 |
| 6453 | Analysis of Meiotic Recombination Pathways in the Yeast <i>Saccharomyces cerevisiae</i> . <i>Genetics</i> , 1996, 144, 71-86. | 1.2 | 130 |
| 6454 | Yeast Mutants That Produce a Novel Type of Ascus Containing Asci Instead of Spores. <i>Genetics</i> , 1996, 144, 979-989. | 1.2 | 14 |
| 6455 | Yeast Mutants Deficient in ER-Associated Degradation of the Z Variant of Alpha-1-Protease Inhibitor. <i>Genetics</i> , 1996, 144, 1355-1362. | 1.2 | 33 |
| 6456 | <i>CDP1</i> , a Novel <i>Saccharomyces cerevisiae</i> Gene Required for Proper Nuclear Division and Chromosome Segregation. <i>Genetics</i> , 1996, 144, 1387-1397. | 1.2 | 26 |
| 6457 | <i>Saccharomyces cerevisiae</i> <i>HOC1</i> , a Suppressor of <i>pkc1</i> , Encodes a Putative Glycosyltransferase. <i>Genetics</i> , 1997, 145, 637-645. | 1.2 | 56 |

| # | ARTICLE | IF | CITATIONS |
|------|--|-----|-----------|
| 6458 | Dissection of Filamentous Growth by Transposon Mutagenesis in <i>Saccharomyces cerevisiae</i> . <i>Genetics</i> , 1997, 145, 671-684. | 1.2 | 233 |
| 6459 | A WD Repeat Protein, Rec14, Essential for Meiotic Recombination in <i>Schizosaccharomyces pombe</i> . <i>Genetics</i> , 1997, 146, 1253-1264. | 1.2 | 43 |
| 6460 | The Role of Gcr1p in the Transcriptional Activation of Glycolytic Genes in Yeast <i>Saccharomyces cerevisiae</i> . <i>Genetics</i> , 1997, 147, 521-532. | 1.2 | 48 |
| 6461 | Mutations in <i>GSF1</i> and <i>GSF2</i> Alter Glucose Signaling in <i>Saccharomyces cerevisiae</i> . <i>Genetics</i> , 1997, 147, 557-566. | 1.2 | 22 |
| 6462 | <i>SRO9</i> , a Multicopy Suppressor of the Bud Growth Defect in the <i>Saccharomyces cerevisiae rho3</i> -Deficient Cells, Shows Strong Genetic Interactions With Tropomyosin Genes, Suggesting Its Role in Organization of the Actin Cytoskeleton. <i>Genetics</i> , 1997, 147, 1003-1016. | 1.2 | 33 |
| 6463 | The <i>Saccharomyces cerevisiae RAD30</i> Gene, a Homologue of <i>Escherichia coli dinB</i> and <i>umuC</i> , Is DNA Damage Inducible and Functions in a Novel Error-Free Postreplication Repair Mechanism. <i>Genetics</i> , 1997, 147, 1557-1568. | 1.2 | 305 |
| 6464 | Control of Amino Acid Permease Sorting in the Late Secretory Pathway of <i>Saccharomyces cerevisiae</i> by <i>SEC13</i> , <i>LST4</i> , <i>LST7</i> and <i>LST78</i> . <i>Genetics</i> , 1997, 147, 1569-1584. | 1.2 | 132 |
| 6465 | Mutations Synthetically Lethal With <i>tpm1</i> Lie in Genes Involved in Morphogenesis. <i>Genetics</i> , 1997, 147, 1595-1607. | 1.2 | 13 |
| 6466 | Dhh1p, a Putative RNA Helicase, Associates with the General Transcription Factors Pop2p and Ccr4p from <i>Saccharomyces cerevisiae</i> . <i>Genetics</i> , 1998, 148, 571-579. | 1.2 | 127 |
| 6467 | Identification of <i>GCD14</i> and <i>GCD15</i> , Novel Genes Required for Translational Repression of <i>GCN4</i> mRNA in <i>Saccharomyces cerevisiae</i> . <i>Genetics</i> , 1998, 148, 1007-1020. | 1.2 | 22 |
| 6468 | Substitutions in the Pheromone-Responsive <i>Gα2</i> Protein of <i>Saccharomyces cerevisiae</i> Confer a Defect in Recovery from Pheromone Treatment. <i>Genetics</i> , 1998, 148, 947-961. | 1.2 | 10 |
| 6469 | The Yeast <i>HSM3</i> Gene Acts in One of the Mismatch Repair Pathways. <i>Genetics</i> , 1998, 148, 963-973. | 1.2 | 28 |
| 6470 | Yeast <i>dom34</i> Mutants Are Defective in Multiple Developmental Pathways and Exhibit Decreased Levels of Polyribosomes. <i>Genetics</i> , 1998, 149, 45-56. | 1.2 | 59 |
| 6471 | Regulation of Yeast Glycogen Metabolism and Sporulation by <i>Glc7p</i> Protein Phosphatase. <i>Genetics</i> , 1998, 149, 57-72. | 1.2 | 61 |
| 6472 | A Nucleolar Protein That Affects Mating Efficiency in <i>Saccharomyces cerevisiae</i> by Altering the Morphological Response to Pheromone. <i>Genetics</i> , 1998, 149, 795-805. | 1.2 | 40 |
| 6473 | Meiotic Chromosome Morphology and Behavior in <i>zip1</i> Mutants of <i>Saccharomyces cerevisiae</i> . <i>Genetics</i> , 1998, 149, 817-832. | 1.2 | 114 |
| 6474 | Sir- and Silencer-Independent Disruption of Silencing in <i>Saccharomyces</i> by <i>Sas10p</i> . <i>Genetics</i> , 1998, 149, 903-914. | 1.2 | 35 |
| 6475 | <i>Sro7p</i> , a <i>Saccharomyces cerevisiae</i> Counterpart of the Tumor Suppressor <i>h(2)gl</i> Protein, Is Related to Myosins in Function. <i>Genetics</i> , 1998, 149, 1717-1727. | 1.2 | 60 |

| # | ARTICLE | IF | CITATIONS |
|------|--|-----|-----------|
| 6476 | Characterization of an Allele-Nonspecific Intragenic Suppressor in the Yeast Plasma Membrane H ⁺ -ATPase Gene (PMA1). <i>Genetics</i> , 1998, 150, 11-19. | 1.2 | 11 |
| 6477 | Identification of Functional Connections Between Calmodulin and the Yeast Actin Cytoskeleton. <i>Genetics</i> , 1998, 150, 43-58. | 1.2 | 21 |
| 6478 | Novel Mutations in the RAD3 and SSL1 Genes Perturb Genome Stability by Stimulating Recombination Between Short Repeats in <i>Saccharomyces cerevisiae</i> . <i>Genetics</i> , 1998, 150, 963-976. | 1.2 | 29 |
| 6479 | Sth1p, a <i>Saccharomyces cerevisiae</i> Snf2p/Swi2p Homolog, Is an Essential ATPase in RSC and Differs From Snf/Swi in Its Interactions With Histones and Chromatin-Associated Proteins. <i>Genetics</i> , 1998, 150, 987-1005. | 1.2 | 104 |
| 6480 | A Genetic Study of Signaling Processes for Repression of PHO5 Transcription in <i>Saccharomyces cerevisiae</i> . <i>Genetics</i> , 1998, 150, 1349-1359. | 1.2 | 43 |
| 6481 | RAD53 Regulates DBF4 Independently of Checkpoint Function in <i>Saccharomyces cerevisiae</i> . <i>Genetics</i> , 1999, 151, 965-977. | 1.2 | 49 |
| 6482 | Suppressor Analysis of the <i>Saccharomyces cerevisiae</i> Gene REC104 Reveals a Genetic Interaction With REC102. <i>Genetics</i> , 1999, 151, 1261-1272. | 1.2 | 24 |
| 6483 | Host Genes That Affect the Target-Site Distribution of the Yeast Retrotransposon Ty1. <i>Genetics</i> , 1999, 151, 1393-1407. | 1.2 | 22 |
| 6484 | Relative Dependence of Different Outputs of the <i>Saccharomyces cerevisiae</i> Pheromone Response Pathway on the MAP Kinase Fus3p. <i>Genetics</i> , 1999, 151, 1425-1444. | 1.2 | 56 |
| 6485 | Evidence for the Involvement of the Glc7-Reg1 Phosphatase and the Snf1-Snf4 Kinase in the Regulation of INO1 Transcription in <i>Saccharomyces cerevisiae</i> . <i>Genetics</i> , 1999, 152, 73-87. | 1.2 | 46 |
| 6486 | Identification and Characterization of <i>Schizosaccharomyces pombe</i> asp1+, a Gene That Interacts with Mutations in the Arp2/3 Complex and Actin. <i>Genetics</i> , 1999, 152, 895-908. | 1.2 | 34 |
| 6487 | Chromosome Size-Dependent Control of Meiotic Reciprocal Recombination in <i>Saccharomyces cerevisiae</i> : The Role of Crossover Interference. <i>Genetics</i> , 1999, 152, 1475-1486. | 1.2 | 95 |
| 6488 | Characterization of the Wsc1 Protein, a Putative Receptor in the Stress Response of <i>Saccharomyces cerevisiae</i> . <i>Genetics</i> , 1999, 152, 1487-1499. | 1.2 | 112 |
| 6489 | The TamA Protein Fused to a DNA-Binding Domain Can Recruit AreA, the Major Nitrogen Regulatory Protein, to Activate Gene Expression in <i>Aspergillus nidulans</i> . <i>Genetics</i> , 1999, 153, 95-105. | 1.2 | 38 |
| 6490 | Genetic Control of Recombination Partner Preference in Yeast Meiosis: Isolation and Characterization of Mutants Elevated for Meiotic Unequal Sister-Chromatid Recombination. <i>Genetics</i> , 1999, 153, 621-641. | 1.2 | 119 |
| 6491 | Regulation of Mitotic Homeologous Recombination in Yeast: Functions of Mismatch Repair and Nucleotide Excision Repair Genes. <i>Genetics</i> , 2000, 154, 133-146. | 1.2 | 107 |
| 6492 | Expression of the <i>Saccharomyces cerevisiae</i> Gene YME1 in the Petite-Negative Yeast <i>Schizosaccharomyces pombe</i> Converts It to Petite-Positive. <i>Genetics</i> , 2000, 154, 147-154. | 1.2 | 25 |
| 6493 | The <i>Saccharomyces cerevisiae</i> DNA Recombination and Repair Functions of the RAD52 Epistasis Group Inhibit Ty1 Transposition. <i>Genetics</i> , 2000, 154, 543-556. | 1.2 | 43 |

| # | ARTICLE | IF | CITATIONS |
|------|--|-----|-----------|
| 6494 | Highly Diverged Homologs of <i>Saccharomyces cerevisiae</i> Mitochondrial mRNA-Specific Translational Activators Have Orthologous Functions in Other Budding Yeasts. <i>Genetics</i> , 2000, 154, 999-1012. | 1.2 | 33 |
| 6495 | Involvement of the PP2C-Like Phosphatase Ptc2p in the DNA Checkpoint Pathways of <i>Saccharomyces cerevisiae</i> . <i>Genetics</i> , 2000, 154, 1523-1532. | 1.2 | 36 |
| 6496 | The <i>Drosophila</i> <i>embargoed</i> Gene Is Required for Larval Progression and Encodes the Functional Homolog of <i>Schizosaccharomyces</i> Crm1. <i>Genetics</i> , 2000, 155, 1799-1807. | 1.2 | 34 |
| 6497 | A Family of cAMP-Response-Element-Related DNA Sequences With Meiotic Recombination Hotspot Activity in <i>Schizosaccharomyces pombe</i> . <i>Genetics</i> , 2000, 156, 59-68. | 1.2 | 45 |
| 6498 | Accumulation of Phosphorylated Sphingoid Long Chain Bases Results in Cell Growth Inhibition in <i>Saccharomyces cerevisiae</i> . <i>Genetics</i> , 2000, 156, 1519-1529. | 1.2 | 95 |
| 6499 | Characterization of <i>rec7</i> , an Early Meiotic Recombination Gene in <i>Schizosaccharomyces pombe</i> . <i>Genetics</i> , 2001, 157, 519-532. | 1.2 | 66 |
| 6500 | The <i>Saccharomyces cerevisiae</i> MUM2 Gene Interacts With the DNA Replication Machinery and Is Required for Meiotic Levels of Double Strand Breaks. <i>Genetics</i> , 2001, 157, 1179-1189. | 1.2 | 36 |
| 6501 | Overexpression of Translation Elongation Factor 1A Affects the Organization and Function of the Actin Cytoskeleton in Yeast. <i>Genetics</i> , 2001, 157, 1425-1436. | 1.2 | 78 |
| 6502 | A Novel Functional Domain of Cdc15 Kinase Is Required for Its Interaction With Tem1 GTPase in <i>Saccharomyces cerevisiae</i> . <i>Genetics</i> , 2001, 157, 1437-1450. | 1.2 | 57 |
| 6503 | Fidelity of Mitotic Double-Strand-Break Repair in <i>Saccharomyces cerevisiae</i> : A Role for <i>SAE2</i> / <i>COM1</i> . <i>Genetics</i> , 2001, 158, 109-122. | 1.2 | 135 |
| 6504 | Requirement for Msh6, but Not for Swi4 (Msh3), in Msh2-Dependent Repair of Base-Base Mismatches and Mononucleotide Loops in <i>Schizosaccharomyces pombe</i> . <i>Genetics</i> , 2001, 158, 65-75. | 1.2 | 20 |
| 6505 | <i>SPO14</i> Separation-of-Function Mutations Define Unique Roles for Phospholipase D in Secretion and Cellular Differentiation in <i>Saccharomyces cerevisiae</i> . <i>Genetics</i> , 2001, 158, 1431-1444. | 1.2 | 35 |
| 6506 | The Yeast Recombinational Repair Protein Rad59 Interacts With Rad52 and Stimulates Single-Strand Annealing. <i>Genetics</i> , 2001, 159, 515-525. | 1.2 | 126 |
| 6507 | Overlapping Functions of the <i>Saccharomyces cerevisiae</i> Mre11, Exo1 and Rad27 Nucleases in DNA Metabolism. <i>Genetics</i> , 2001, 159, 1423-1433. | 1.2 | 151 |
| 6508 | The Ess1 Prolyl Isomerase Is Required for Growth and Morphogenetic Switching in <i>Candida albicans</i> . <i>Genetics</i> , 2002, 160, 37-48. | 1.2 | 43 |
| 6509 | Differential Regulation of <i>Saccharomyces cerevisiae</i> Phospholipase D in Sporulation and Sec14-Independent Secretion. <i>Genetics</i> , 2002, 160, 1353-1361. | 1.2 | 29 |
| 6510 | The <i>mcm5-bob1</i> Bypass of Cdc7p/Dbf4p in DNA Replication Depends on Both Cdk1-Independent and Cdk1-Dependent Steps in <i>Saccharomyces cerevisiae</i> . <i>Genetics</i> , 2002, 161, 47-57. | 1.2 | 36 |
| 6511 | Yeast RSC Function Is Required for Organization of the Cellular Cytoskeleton via an Alternative <i>PKC1</i> Pathway. <i>Genetics</i> , 2002, 161, 575-584. | 1.2 | 29 |

| # | ARTICLE | IF | CITATIONS |
|------|--|-----|-----------|
| 6512 | Gene Overexpression as a Tool for Identifying New <i>trans</i> -Acting Factors Involved in Translation Termination in <i>Saccharomyces cerevisiae</i> . <i>Genetics</i> , 2002, 161, 585-594. | 1.2 | 19 |
| 6513 | A Novel Yeast Silencer: The 2 ^{1/4} Origin of <i>Saccharomyces cerevisiae</i> Has HST3-, MIG1- and SIR-Dependent Silencing Activity. <i>Genetics</i> , 2002, 162, 59-71. | 1.2 | 27 |
| 6514 | Dissection of Upstream Regulatory Components of the Rho1p Effector, 1,3- β -Glucan Synthase, in <i>Saccharomyces cerevisiae</i> . <i>Genetics</i> , 2002, 162, 663-676. | 1.2 | 112 |
| 6515 | The Roles of <i>REV3</i> and <i>RAD57</i> in Double-Strand-Break-Repair-Induced Mutagenesis of <i>Saccharomyces cerevisiae</i> . <i>Genetics</i> , 2002, 162, 1063-1077. | 1.2 | 98 |
| 6516 | A Defect of Kap104 Alleviates the Requirement of Mitotic Exit Network Gene Functions in <i>Saccharomyces cerevisiae</i> . <i>Genetics</i> , 2002, 162, 1545-1556. | 1.2 | 19 |
| 6517 | Differential Processing of Leading- and Lagging-Strand Ends at <i>Saccharomyces cerevisiae</i> Telomeres Revealed by the Absence of Rad27p Nuclease. <i>Genetics</i> , 2002, 162, 1583-1594. | 1.2 | 41 |
| 6518 | Loss of CDC5 Function in <i>Saccharomyces cerevisiae</i> Leads to Defects in Swe1p Regulation and Bfa1p/Bub2p-Independent Cytokinesis. <i>Genetics</i> , 2003, 163, 21-33. | 1.2 | 38 |
| 6519 | The External Amino Acid Signaling Pathway Promotes Activation of Stp1 and Uga35/Dal81 Transcription Factors for Induction of the <i>AGP1</i> Gene in <i>Saccharomyces cerevisiae</i> . <i>Genetics</i> , 2004, 166, 1727-1739. | 1.2 | 20 |
| 6520 | A cellular protein which binds hepatitis B virus but not hepatitis B surface antigen.. <i>Journal of General Virology</i> , 1999, 80, 607-615. | 1.3 | 4 |
| 6521 | Involvement of the CDC25 gene product in the signal transmission pathway of the glucose-induced RAS-mediated cAMP signal in the yeast <i>Saccharomyces cerevisiae</i> . <i>Microbiology (United Kingdom)</i> , 1991, 137, 341-349. | 0.7 | 60 |
| 6522 | A <i>Candida albicans</i> 37 kDa polypeptide with homology to the laminin receptor is a component of the translational machinery. <i>Microbiology (United Kingdom)</i> , 1998, 144, 839-847. | 0.7 | 12 |
| 6523 | The <i>Candida albicans</i> gene for mRNA 5' cap methyltransferase: identification of additional residues essential for catalysis The GenBank accession numbers for the nucleotide sequence of CaABD1 and hMet are AB020965 and AB020966, respectively.. <i>Microbiology (United Kingdom)</i> , 1999, 145, 3023-3033. | 0.7 | 11 |
| 6524 | The yeast Chs4 protein stimulates the trypsin-sensitive activity of chitin synthase 3 through an apparent protein-protein interaction. <i>Microbiology (United Kingdom)</i> , 2000, 146, 385-391. | 0.7 | 53 |
| 6525 | Evaluation of in vivo activation of protein kinase A under non-dissociable conditions through the overexpression of wild-type and mutant regulatory subunits in <i>Saccharomyces cerevisiae</i> . <i>Microbiology (United Kingdom)</i> , 2001, 147, 1149-1159. | 0.7 | 11 |
| 6526 | Cell cycle control of septin ring dynamics in the budding yeast. <i>Microbiology (United Kingdom)</i> , 2001, 147, 1437-1450. | 0.7 | 89 |
| 6527 | The <i>Candida albicans</i> Sup35p protein (CaSup35p): function, prion-like behaviour and an associated polyglutamine length polymorphism The GenBank accession numbers for the sequences determined in this work are AF020554 (CaSUP35 gene) and AY028660 AY028673 (CaSUP35 N domains).. <i>Microbiology (United Kingdom)</i> , 2002, 148, 1049-1060. | 0.7 | 30 |
| 6528 | Cdc42 activation state affects its localization and protein levels in fission yeast. <i>Microbiology (United Kingdom)</i> , 2017, 163, 1156-1166. | 0.7 | 9 |
| 6531 | Nuclear export of Far1p in response to pheromones requires the export receptor Msn5p/Ste21p. <i>Genes and Development</i> , 1999, 13, 2284-2300. | 2.7 | 90 |

| # | ARTICLE | IF | CITATIONS |
|------|--|-----|-----------|
| 6532 | Luc7p, a novel yeast U1 snRNP protein with a role in 5' splice site recognition. <i>Genes and Development</i> , 1999, 13, 2425-2438. | 2.7 | 98 |
| 6533 | Ctf7p is essential for sister chromatid cohesion and links mitotic chromosome structure to the DNA replication machinery. <i>Genes and Development</i> , 1999, 13, 307-319. | 2.7 | 420 |
| 6534 | Regulation of <i>Saccharomyces cerevisiae</i> kinetochores by the type 1 phosphatase Glc7p. <i>Genes and Development</i> , 1999, 13, 545-555. | 2.7 | 135 |
| 6535 | A stem/loop in U6 RNA defines a conformational switch required for pre-mRNA splicing. <i>Genes and Development</i> , 1994, 8, 221-233. | 2.7 | 95 |
| 6536 | Expression of the AAC2 Gene Encoding the Major Mitochondrial ADP/ATP Carrier in <i>Saccharomyces cerevisiae</i> is Controlled at the Transcriptional Level by Oxygen, Heme and HAP2 Factor. <i>FEBS Journal</i> , 1995, 229, 651-657. | 0.2 | 28 |
| 6537 | Identification of Residues Potentially Involved in the Interactions Between Subunits in Yeast Alcohol Dehydrogenases. <i>FEBS Journal</i> , 1995, 231, 214-219. | 0.2 | 12 |
| 6538 | Protein Phosphatase 2B of <i>Saccharomyces Cerevisiae</i> is Required for Tolerance to Manganese, in Blocking the Entry of ions into the Cells. <i>FEBS Journal</i> , 1995, 232, 712-717. | 0.2 | 55 |
| 6539 | Production and Regeneration of <i>Lactobacillus casei</i> Protoplasts. <i>Applied and Environmental Microbiology</i> , 1984, 48, 994-1000. | 1.4 | 66 |
| 6540 | Optimum conditions for ursodeoxycholic acid production from lithocholic acid by <i>Fusarium equiseti</i> M41. <i>Applied and Environmental Microbiology</i> , 1985, 49, 338-344. | 1.4 | 22 |
| 6541 | Phenotype character of the methylglyoxal resistance gene in <i>Saccharomyces cerevisiae</i> : expression in <i>Escherichia coli</i> and application to breeding wild-type yeast strains. <i>Applied and Environmental Microbiology</i> , 1985, 50, 1200-1207. | 1.4 | 61 |
| 6542 | Transformation of <i>Saccharomyces cerevisiae</i> by electroporation. <i>Applied and Environmental Microbiology</i> , 1989, 55, 2242-2246. | 1.4 | 96 |
| 6543 | The prepro-peptide of <i>Mucor rennin</i> directs the secretion of human growth hormone by <i>Saccharomyces cerevisiae</i> . <i>Applied and Environmental Microbiology</i> , 1990, 56, 2125-2132. | 1.4 | 21 |
| 6544 | Inducible overexpression of the FUM1 gene in <i>Saccharomyces cerevisiae</i> : localization of fumarase and efficient fumaric acid bioconversion to L-malic acid. <i>Applied and Environmental Microbiology</i> , 1990, 56, 2777-2783. | 1.4 | 39 |
| 6545 | Genetic engineering of a sake yeast producing no urea by successive disruption of arginase gene. <i>Applied and Environmental Microbiology</i> , 1991, 57, 301-306. | 1.4 | 102 |
| 6546 | The secretion leader of <i>Mucor pusillus rennin</i> which possesses an artificial Lys-Arg sequence directs the secretion of mature human growth hormone by <i>Saccharomyces cerevisiae</i> . <i>Applied and Environmental Microbiology</i> , 1991, 57, 2052-2056. | 1.4 | 6 |
| 6547 | Influence of Polyethylene Glycol on the Size of <i>Schizosaccharomyces pombe</i> Electropores. <i>Applied and Environmental Microbiology</i> , 1992, 58, 1201-1206. | 1.4 | 14 |
| 6548 | Expression and secretion of <i>Bacillus amyloliquefaciens</i> alpha-amylase by using the yeast pheromone alpha-factor promoter and leader sequence in <i>Saccharomyces cerevisiae</i> . <i>Applied and Environmental Microbiology</i> , 1993, 59, 1253-1258. | 1.4 | 10 |
| 6549 | Transformation of <i>Kluyveromyces lactis</i> by Electroporation. <i>Applied and Environmental Microbiology</i> , 1993, 59, 2087-2092. | 1.4 | 55 |

| # | ARTICLE | IF | CITATIONS |
|------|--|-----|-----------|
| 6550 | Cloning and expression in <i>Saccharomyces cerevisiae</i> of a <i>Trichoderma reesei</i> beta-mannanase gene containing a cellulose binding domain. <i>Applied and Environmental Microbiology</i> , 1995, 61, 1090-1097. | 1.4 | 152 |
| 6551 | Combined Expression of <i>Aspergillus nidulans</i> Endoxylanase X24 and <i>Aspergillus oryzae</i> (alpha)-Amylase in Industrial Baker's Yeasts and Their Use in Bread Making. <i>Applied and Environmental Microbiology</i> , 1996, 62, 3712-3715. | 1.4 | 27 |
| 6552 | Recombinant thermostable cyclinulo-oligosaccharide fructanotransferase produced by <i>Saccharomyces cerevisiae</i> . <i>Applied and Environmental Microbiology</i> , 1997, 63, 4956-4960. | 1.4 | 16 |
| 6553 | Expression and secretion of a thermostable bacterial xylanase in <i>Kluyveromyces lactis</i> . <i>Applied and Environmental Microbiology</i> , 1997, 63, 3297-3300. | 1.4 | 31 |
| 6554 | Balance of Activities of Alcohol Acetyltransferase and Esterase in <i>Saccharomyces cerevisiae</i> Is Important for Production of Isoamyl Acetate. <i>Applied and Environmental Microbiology</i> , 1998, 64, 4076-4078. | 1.4 | 111 |
| 6555 | Expression of an <i>Aspergillus niger</i> Phytase Gene (<i>phyA</i>) in <i>Saccharomyces cerevisiae</i> . <i>Applied and Environmental Microbiology</i> , 1999, 65, 1915-1918. | 1.4 | 118 |
| 6556 | Transcriptional Control of <i>ADH</i> Genes in the Xylose-Fermenting Yeast <i>Pichia stipitis</i> . <i>Applied and Environmental Microbiology</i> , 1999, 65, 2363-2368. | 1.4 | 37 |
| 6557 | The Carboxy-Terminal Portion of the Aflatoxin Pathway Regulatory Protein AFLR of <i>Aspergillus parasiticus</i> Activates <i>GAL1</i> :: <i>lacZ</i> Gene Expression in <i>Saccharomyces cerevisiae</i> . <i>Applied and Environmental Microbiology</i> , 1999, 65, 2508-2512. | 1.4 | 45 |
| 6558 | Expression of the Extracellular Domain of the Human Immunodeficiency Virus Type 1 Envelope Protein and Its Fusion with β -Galactosidase in <i>Saccharomyces cerevisiae</i> . <i>Vaccine Journal</i> , 1998, 5, 592-594. | 2.6 | 1 |
| 6559 | Gene isolation by complementation in <i>Candida albicans</i> and applications to physical and genetic mapping. <i>Infection and Immunity</i> , 1992, 60, 876-884. | 1.0 | 62 |
| 6560 | Disruption of the SNF1 gene abolishes trehalose utilization in the pathogenic yeast <i>Candida glabrata</i> . <i>Infection and Immunity</i> , 1996, 64, 5269-5273. | 1.0 | 27 |
| 6561 | A gene homologous to <i>Saccharomyces cerevisiae</i> SNF1 appears to be essential for the viability of <i>Candida albicans</i> . <i>Infection and Immunity</i> , 1997, 65, 4909-4917. | 1.0 | 70 |
| 6562 | Prohibitin, a putative negative control element present in <i>Pneumocystis carinii</i> . <i>Infection and Immunity</i> , 1997, 65, 5125-5130. | 1.0 | 13 |
| 6563 | Expression, cloning, and characterization of a <i>Candida albicans</i> gene, ALA1, that confers adherence properties upon <i>Saccharomyces cerevisiae</i> for extracellular matrix proteins. <i>Infection and Immunity</i> , 1997, 65, 5289-5294. | 1.0 | 178 |
| 6564 | Transformation of <i>Kluyveromyces fragilis</i> . <i>Journal of Bacteriology</i> , 1984, 158, 1165-1167. | 1.0 | 47 |
| 6565 | Molecular cloning of chromosome I DNA from <i>Saccharomyces cerevisiae</i> : isolation of the ADE1 gene. <i>Journal of Bacteriology</i> , 1984, 159, 413-417. | 1.0 | 38 |
| 6566 | Deletion of plasmid sequences during <i>Saccharomyces cerevisiae</i> transformation. <i>Journal of Bacteriology</i> , 1984, 159, 1065-1067. | 1.0 | 33 |
| 6567 | Characterization of <i>Saccharomyces cerevisiae</i> mutants supersensitive to aminoglycoside antibiotics. <i>Journal of Bacteriology</i> , 1985, 163, 8-14. | 1.0 | 26 |

| # | ARTICLE | IF | CITATIONS |
|------|---|-----|-----------|
| 6568 | Cloning of GLN4: an essential gene that encodes glutaminyl-tRNA synthetase in <i>Saccharomyces cerevisiae</i> . <i>Journal of Bacteriology</i> , 1985, 163, 763-768. | 1.0 | 13 |
| 6569 | Stimulation of Transcription by Mutations Affecting Conserved Regions of RNA Polymerase II. <i>Journal of Bacteriology</i> , 1998, 180, 2590-2598. | 1.0 | 20 |
| 6570 | Altered Na ⁺ and Li ⁺ Homeostasis in <i>Saccharomyces cerevisiae</i> Cells Expressing the Bacterial Cation Antiporter NhaA. <i>Journal of Bacteriology</i> , 1998, 180, 3131-3136. | 1.0 | 23 |
| 6571 | Regulation of Expression of <i>GLT1</i> , the Gene Encoding Glutamate Synthase in <i>Saccharomyces cerevisiae</i> . <i>Journal of Bacteriology</i> , 1998, 180, 3533-3540. | 1.0 | 49 |
| 6572 | Disruption of the <i>Candida albicans</i> TPS1 Gene Encoding Trehalose-6-Phosphate Synthase Impairs Formation of Hyphae and Decreases Infectivity. <i>Journal of Bacteriology</i> , 1998, 180, 3809-3815. | 1.0 | 121 |
| 6573 | Novel <i>Escherichia coli</i> umuD ^Δ Mutants: Structure-Function Insights into SOS Mutagenesis. <i>Journal of Bacteriology</i> , 1998, 180, 4658-4666. | 1.0 | 18 |
| 6574 | Isolation of <i>Candida glabrata</i> Homologs of the <i>Saccharomyces cerevisiae</i> KRE9 and <i>KNH1</i> Genes and Their Involvement in Cell Wall β -1,6-Glucan Synthesis. <i>Journal of Bacteriology</i> , 1998, 180, 5020-5029. | 1.0 | 26 |
| 6575 | Disruption of PMR1, Encoding a Ca ²⁺ -ATPase Homolog in <i>Yarrowia lipolytica</i> , Affects Secretion and Processing of Homologous and Heterologous Proteins. <i>Journal of Bacteriology</i> , 1998, 180, 6736-6742. | 1.0 | 19 |
| 6576 | Loss of the Plasma Membrane-Bound Protein Gas1p in <i>Saccharomyces cerevisiae</i> Results in the Release of β -1,3-Glucan into the Medium and Induces a Compensation Mechanism To Ensure Cell Wall Integrity. <i>Journal of Bacteriology</i> , 1998, 180, 1418-1424. | 1.0 | 184 |
| 6577 | The TATA-Binding Protein (TBP) from the Human Fungal Pathogen <i>Candida albicans</i> Can Complement Defects in Human and Yeast TBPs. <i>Journal of Bacteriology</i> , 1998, 180, 1771-1776. | 1.0 | 4 |
| 6578 | A Small Protein (Ags1p) and the Pho80p-Pho85p Kinase Complex Contribute to Aminoglycoside Antibiotic Resistance of the Yeast <i>Saccharomyces cerevisiae</i> . <i>Journal of Bacteriology</i> , 1998, 180, 1887-1894. | 1.0 | 12 |
| 6579 | Control of 5 ^{â€²} ,5 ^{â€²} -Dinucleoside Triphosphate Catabolism by APH1, a <i>Saccharomyces cerevisiae</i> Analog of Human FHIT. <i>Journal of Bacteriology</i> , 1998, 180, 2345-2349. | 1.0 | 23 |
| 6580 | Identification of Two Mannoproteins Released from Cell Walls of a <i>Saccharomyces cerevisiae</i> mnn1 mnn9 Double Mutant by Reducing Agents. <i>Journal of Bacteriology</i> , 1999, 181, 4741-4745. | 1.0 | 67 |
| 6581 | Synergistic Operation of the <i>CAR2</i> (Ornithine Transaminase) Promoter Elements in <i>Saccharomyces cerevisiae</i> . <i>Journal of Bacteriology</i> , 1999, 181, 7052-7064. | 1.0 | 14 |
| 6582 | Molecular Analysis of the <i>Candida albicans</i> Homolog of <i>Saccharomyces cerevisiae</i> MNN9, Required for Glycosylation of Cell Wall Mannoproteins. <i>Journal of Bacteriology</i> , 1999, 181, 7439-7448. | 1.0 | 45 |
| 6583 | Cyclic AMP Can Decrease Expression of Genes Subject to Catabolite Repression in <i>Saccharomyces cerevisiae</i> . <i>Journal of Bacteriology</i> , 1999, 181, 2640-2642. | 1.0 | 23 |
| 6584 | Expression of hepatitis B virus core antigen gene in <i>Saccharomyces cerevisiae</i> : synthesis of two polypeptides translated from different initiation codons. <i>Journal of Virology</i> , 1986, 59, 176-180. | 1.5 | 127 |
| 6585 | Trans activation by the bovine papillomavirus E2 protein in <i>Saccharomyces cerevisiae</i> . <i>Journal of Virology</i> , 1989, 63, 4422-4425. | 1.5 | 50 |

| # | ARTICLE | IF | CITATIONS |
|------|--|-----|-----------|
| 6586 | Structure-function studies of the herpes simplex virus type 1 DNA polymerase. <i>Journal of Virology</i> , 1990, 64, 5008-5018. | 1.5 | 40 |
| 6587 | Rous sarcoma virus expression in <i>Saccharomyces cerevisiae</i> : processing and membrane targeting of the gag gene product. <i>Journal of Virology</i> , 1990, 64, 5628-5632. | 1.5 | 8 |
| 6588 | Infectious foot-and-mouth disease virus derived from a cloned full-length cDNA. <i>Journal of Virology</i> , 1990, 64, 2467-2473. | 1.5 | 101 |
| 6589 | Characterization of a transpositionally active Ty3 element and identification of the Ty3 integrase protein. <i>Journal of Virology</i> , 1990, 64, 2599-2607. | 1.5 | 93 |
| 6590 | Proteolytic processing of pol-TYB proteins from the yeast retrotransposon Ty1. <i>Journal of Virology</i> , 1991, 65, 4573-4581. | 1.5 | 77 |
| 6591 | Expression and characterization of the trans-activating protein Tax of human T-cell leukemia virus type I in <i>Saccharomyces cerevisiae</i> . <i>Journal of Virology</i> , 1992, 66, 7253-7261. | 1.5 | 11 |
| 6592 | Ty3 GAG3 and POL3 genes encode the components of intracellular particles. <i>Journal of Virology</i> , 1992, 66, 1414-1424. | 1.5 | 72 |
| 6593 | Ribosomal frameshifting efficiency and gag/gag-pol ratio are critical for yeast M1 double-stranded RNA virus propagation. <i>Journal of Virology</i> , 1992, 66, 3669-3676. | 1.5 | 219 |
| 6594 | Comparing regions of the Epstein-Barr virus ZEBRA protein which function as transcriptional activating sequences in <i>Saccharomyces cerevisiae</i> and in B cells. <i>Journal of Virology</i> , 1993, 67, 7472-7481. | 1.5 | 17 |
| 6595 | Activation of the Epstein-Barr virus BMRF1 and BZLF1 promoters by ZEBRA in <i>Saccharomyces cerevisiae</i> . <i>Journal of Virology</i> , 1994, 68, 7628-7633. | 1.5 | 16 |
| 6596 | Synthesis of Semliki Forest virus RNA polymerase components nsP1 through nsP4 in <i>Saccharomyces cerevisiae</i> by expression of cDNA encoding the nonstructural polyprotein. <i>Journal of Virology</i> , 1996, 70, 4086-4089. | 1.5 | 2 |
| 6597 | A conserved domain of the Epstein-Barr virus nuclear antigens 3A and 3C binds to a discrete domain of Jkappa. <i>Journal of Virology</i> , 1996, 70, 4228-4236. | 1.5 | 106 |
| 6598 | Interaction cloning of NS1-I, a human protein that binds to the nonstructural NS1 proteins of influenza A and B viruses. <i>Journal of Virology</i> , 1996, 70, 5363-5372. | 1.5 | 52 |
| 6599 | In vivo DNA expression of functional brome mosaic virus RNA replicons in <i>Saccharomyces cerevisiae</i> . <i>Journal of Virology</i> , 1997, 71, 7781-7790. | 1.5 | 83 |
| 6600 | The NPI-1/NPI-3 (karyopherin alpha) binding site on the influenza A virus nucleoprotein NP is a nonconventional nuclear localization signal. <i>Journal of Virology</i> , 1997, 71, 1850-1856. | 1.5 | 227 |
| 6601 | Specific repression of Tax trans-activation by TAR RNA-binding protein TRBP. <i>Journal of Virology</i> , 1997, 71, 2628-2635. | 1.5 | 16 |
| 6602 | Self-association of herpes simplex virus type 1 ICP35 is via coiled-coil interactions and promotes stable interaction with the major capsid protein. <i>Journal of Virology</i> , 1997, 71, 5197-5208. | 1.5 | 44 |
| 6603 | CREB-2, a Cellular CRE-Dependent Transcription Repressor, Functions in Association with Tax as an Activator of the Human T-Cell Leukemia Virus Type 1 Promoter. <i>Journal of Virology</i> , 1998, 72, 8332-8337. | 1.5 | 88 |

| # | ARTICLE | IF | CITATIONS |
|------|---|-----|-----------|
| 6604 | Putative RNA Capping Activities Encoded by Brome Mosaic Virus: Methylation and Covalent Binding of Guanylate by Replicase Protein 1a. <i>Journal of Virology</i> , 1999, 73, 10061-10069. | 1.5 | 94 |
| 6605 | Brome Mosaic Virus RNA Replication Proteins 1a and 2a Colocalize and 1a Independently Localizes on the Yeast Endoplasmic Reticulum. <i>Journal of Virology</i> , 1999, 73, 10303-10309. | 1.5 | 119 |
| 6606 | A Brome Mosaic Virus Intergenic RNA3 Replication Signal Functions with Viral Replication Protein 1a To Dramatically Stabilize RNA In Vivo. <i>Journal of Virology</i> , 1999, 73, 2622-2632. | 1.5 | 132 |
| 6607 | The Carboxy-Terminal Acidic Domain of Rift Valley Fever Virus NSs Protein Is Essential for the Formation of Filamentous Structures but Not for the Nuclear Localization of the Protein. <i>Journal of Virology</i> , 1999, 73, 5018-5025. | 1.5 | 68 |
| 6608 | Overexpression of the STE4 gene leads to mating response in haploid <i>Saccharomyces cerevisiae</i> . <i>Molecular and Cellular Biology</i> , 1990, 10, 217-222. | 1.1 | 99 |
| 6609 | PRP18, a protein required for the second reaction in pre-mRNA splicing. <i>Molecular and Cellular Biology</i> , 1990, 10, 324-332. | 1.1 | 48 |
| 6610 | Farnesyl Cysteine C-Terminal Methyltransferase Activity Is Dependent upon the <i>STE14</i> Gene Product in <i>Saccharomyces cerevisiae</i> . <i>Molecular and Cellular Biology</i> , 1990, 10, 5071-5076. | 1.1 | 48 |
| 6611 | Ribosomal Protein L30 Is Dispensable in the Yeast <i>Saccharomyces cerevisiae</i> . <i>Molecular and Cellular Biology</i> , 1990, 10, 5235-5243. | 1.1 | 35 |
| 6612 | <i>Drosophila</i> Scaffold-Attached Regions Bind Nuclear Scaffolds and Can Function as ARS Elements in Both Budding and Fission Yeasts. <i>Molecular and Cellular Biology</i> , 1990, 10, 5442-5454. | 1.1 | 39 |
| 6613 | The SNF5 Protein of <i>Saccharomyces cerevisiae</i> Is a Glutamine- and Proline-Rich Transcriptional Activator That Affects Expression of a Broad Spectrum of Genes. <i>Molecular and Cellular Biology</i> , 1990, 10, 5616-5625. | 1.1 | 94 |
| 6614 | The <i>Saccharomyces cerevisiae SIN3</i> Gene, a Negative Regulator of <i>HO</i> , Contains Four Paired Amphipathic Helix Motifs. <i>Molecular and Cellular Biology</i> , 1990, 10, 5927-5936. | 1.1 | 104 |
| 6615 | Role of <i>IME1</i> Expression in Regulation of Meiosis in <i>Saccharomyces cerevisiae</i> . <i>Molecular and Cellular Biology</i> , 1990, 10, 6103-6113. | 1.1 | 87 |
| 6616 | A Suppressor of an RNA Polymerase II Mutation of <i>Saccharomyces cerevisiae</i> Encodes a Subunit Common to RNA Polymerases I, II, and III. <i>Molecular and Cellular Biology</i> , 1990, 10, 6123-6131. | 1.1 | 32 |
| 6617 | Role of Arginine 180 and Glutamic Acid 177 of Ricin Toxin A Chain in Enzymatic Inactivation of Ribosomes. <i>Molecular and Cellular Biology</i> , 1990, 10, 6257-6263. | 1.1 | 33 |
| 6618 | Transcriptional Derepression of the <i>Saccharomyces cerevisiae HSP26</i> Gene during Heat Shock. <i>Molecular and Cellular Biology</i> , 1990, 10, 6362-6373. | 1.1 | 22 |
| 6619 | <i>gcr2</i> , a new mutation affecting glycolytic gene expression in <i>Saccharomyces cerevisiae</i> . <i>Molecular and Cellular Biology</i> , 1990, 10, 6389-6396. | 1.1 | 51 |
| 6620 | The Yeast <i>PRP6</i> Gene Encodes a U4/U6 Small Nuclear Ribonucleoprotein Particle (snRNP) Protein, and the <i>PRP9</i> Gene Encodes a Protein Required for U2 snRNP Binding. <i>Molecular and Cellular Biology</i> , 1990, 10, 6417-6425. | 1.1 | 72 |
| 6621 | Characterization of <i>TUP1</i> , a Mediator of Glucose Repression in <i>Saccharomyces cerevisiae</i> . <i>Molecular and Cellular Biology</i> , 1990, 10, 6500-6511. | 1.1 | 134 |

| # | ARTICLE | IF | CITATIONS |
|------|---|-----|-----------|
| 6622 | Characterization of <i>RAD9</i> of <i>Saccharomyces cerevisiae</i> and Evidence that Its Function Acts Posttranslationally in Cell Cycle Arrest after DNA Damage. <i>Molecular and Cellular Biology</i> , 1990, 10, 6554-6564. | 1.1 | 111 |
| 6623 | ADP Ribosylation Factor Is an Essential Protein in <i>Saccharomyces cerevisiae</i> and Is Encoded by Two Genes. <i>Molecular and Cellular Biology</i> , 1990, 10, 6690-6699. | 1.1 | 94 |
| 6624 | Characterization of <i>VPS34</i> , a Gene Required for Vacuolar Protein Sorting and Vacuole Segregation in <i>Saccharomyces cerevisiae</i> . <i>Molecular and Cellular Biology</i> , 1990, 10, 6742-6754. | 1.1 | 216 |
| 6625 | Stoichiometry of G Protein Subunits Affects the <i>Saccharomyces cerevisiae</i> Mating Pheromone Signal Transduction Pathway. <i>Molecular and Cellular Biology</i> , 1990, 10, 510-517. | 1.1 | 92 |
| 6626 | <i>HOL1</i> Mutations Confer Novel Ion Transport in <i>Saccharomyces cerevisiae</i> . <i>Molecular and Cellular Biology</i> , 1990, 10, 643-652. | 1.1 | 12 |
| 6627 | Mutations in <i>RAD6</i> , a Yeast Gene Encoding a Ubiquitin-Conjugating Enzyme, Stimulate Retrotransposition. <i>Molecular and Cellular Biology</i> , 1990, 10, 1017-1022. | 1.1 | 46 |
| 6628 | Mutational Analysis of the SNF3 Glucose Transporter of <i>Saccharomyces cerevisiae</i> . <i>Molecular and Cellular Biology</i> , 1990, 10, 1105-1115. | 1.1 | 35 |
| 6629 | Depletion of U14 Small Nuclear RNA (snR128) Disrupts Production of 18S rRNA in <i>Saccharomyces cerevisiae</i> . <i>Molecular and Cellular Biology</i> , 1990, 10, 1145-1152. | 1.1 | 170 |
| 6630 | In vivo analysis of the <i>Saccharomyces cerevisiae</i> HO nuclease recognition site by site-directed mutagenesis. <i>Molecular and Cellular Biology</i> , 1990, 10, 1174-1179. | 1.1 | 53 |
| 6631 | A Yeast <i>ARS</i> -Binding Protein Activates Transcription Synergistically in Combination with Other Weak Activating Factors. <i>Molecular and Cellular Biology</i> , 1990, 10, 887-897. | 1.1 | 94 |
| 6632 | Vanadate-Resistant Mutants of <i>Saccharomyces cerevisiae</i> Show Alterations in Protein Phosphorylation and Growth Control. <i>Molecular and Cellular Biology</i> , 1990, 10, 898-909. | 1.1 | 20 |
| 6633 | The Product of the <i>Saccharomyces cerevisiae</i> Cell Cycle Gene <i>DBF2</i> Has Homology with Protein Kinases and Is Periodically Expressed in the Cell Cycle. <i>Molecular and Cellular Biology</i> , 1990, 10, 1358-1366. | 1.1 | 72 |
| 6634 | Transcription of a Yeast Phosphoglucosyltransferase Isozyme Gene Is Galactose Inducible and Glucose Repressible. <i>Molecular and Cellular Biology</i> , 1990, 10, 1415-1422. | 1.1 | 28 |
| 6635 | <i>HPR1</i> , a Novel Yeast Gene That Prevents Intrachromosomal Excision Recombination, Shows Carboxy-Terminal Homology to the <i>Saccharomyces cerevisiae</i> <i>TOP1</i> Gene. <i>Molecular and Cellular Biology</i> , 1990, 10, 1439-1451. | 1.1 | 92 |
| 6636 | Self-Regulation of 70-Kilodalton Heat Shock Proteins in <i>Saccharomyces cerevisiae</i> . <i>Molecular and Cellular Biology</i> , 1990, 10, 1622-1632. | 1.1 | 61 |
| 6637 | Conditional Expression of <i>RPA190</i> , the Gene Encoding the Largest Subunit of Yeast RNA Polymerase I: Effects of Decreased rRNA Synthesis on Ribosomal Protein Synthesis. <i>Molecular and Cellular Biology</i> , 1990, 10, 2049-2059. | 1.1 | 36 |
| 6638 | Courtship in <i>Saccharomyces cerevisiae</i> : an Early Cell-Cell Interaction during Mating. <i>Molecular and Cellular Biology</i> , 1990, 10, 2202-2213. | 1.1 | 63 |
| 6639 | <i>MER1</i> , a Yeast Gene Required for Chromosome Pairing and Genetic Recombination, Is Induced in Meiosis. <i>Molecular and Cellular Biology</i> , 1990, 10, 2379-2389. | 1.1 | 108 |

| # | ARTICLE | IF | CITATIONS |
|------|--|-----|-----------|
| 6640 | Isolation of the Gene Encoding the <i>Saccharomyces cerevisiae</i> Centromere-Binding Protein CP1. <i>Molecular and Cellular Biology</i> , 1990, 10, 2458-2467. | 1.1 | 134 |
| 6641 | <i>RAD10</i> , an Excision Repair Gene of <i>Saccharomyces cerevisiae</i> , Is Involved in the <i>RAD1</i> Pathway of Mitotic Recombination. <i>Molecular and Cellular Biology</i> , 1990, 10, 2485-2491. | 1.1 | 89 |
| 6642 | Mutational Mapping of RAS-Responsive Domains of the <i>Saccharomyces cerevisiae</i> Adenylyl Cyclase. <i>Molecular and Cellular Biology</i> , 1990, 10, 2539-2543. | 1.1 | 35 |
| 6643 | <i>SNF6</i> Encodes a Nuclear Protein That Is Required for Expression of Many Genes in <i>Saccharomyces cerevisiae</i> . <i>Molecular and Cellular Biology</i> , 1990, 10, 2544-2553. | 1.1 | 49 |
| 6644 | Coding and Noncoding Sequences at the 3' End of Yeast Histone H2B mRNA Confer Cell Cycle Regulation. <i>Molecular and Cellular Biology</i> , 1990, 10, 2687-2694. | 1.1 | 32 |
| 6645 | Identification of Positive-Acting Domains in GCN2 Protein Kinase Required for Translational Activation of <i>GCN4</i> Expression. <i>Molecular and Cellular Biology</i> , 1990, 10, 2820-2831. | 1.1 | 85 |
| 6646 | Oncogenic Transformation by vRel Requires an Amino-Terminal Activation Domain. <i>Molecular and Cellular Biology</i> , 1990, 10, 2840-2847. | 1.1 | 102 |
| 6647 | GAL4 Protein: Purification, Association with GAL80 Protein, and Conserved Domain Structure. <i>Molecular and Cellular Biology</i> , 1990, 10, 2916-2923. | 1.1 | 58 |
| 6648 | Mutations in Cell Division Cycle Genes <i>CDC36</i> and <i>CDC39</i> Activate the <i>Saccharomyces cerevisiae</i> Mating Pheromone Response Pathway. <i>Molecular and Cellular Biology</i> , 1990, 10, 2966-2972. | 1.1 | 26 |
| 6649 | TEC1, a Gene Involved in the Activation of Ty1 and Ty1-Mediated Gene Expression in <i>Saccharomyces cerevisiae</i> : Cloning and Molecular Analysis. <i>Molecular and Cellular Biology</i> , 1990, 10, 3541-3550. | 1.1 | 54 |
| 6650 | Molecular Cloning of the Yeast Mitochondrial Aconitase Gene (<i>ACO1</i>) and Evidence of a Synergistic Regulation of Expression by Glucose plus Glutamate. <i>Molecular and Cellular Biology</i> , 1990, 10, 3551-3561. | 1.1 | 57 |
| 6651 | Role of Vacuolar Acidification in Protein Sorting and Zymogen Activation: a Genetic Analysis of the Yeast Vacuolar Proton-Translocating ATPase. <i>Molecular and Cellular Biology</i> , 1990, 10, 3737-3749. | 1.1 | 96 |
| 6652 | Adenylate Cyclase in <i>Saccharomyces cerevisiae</i> Is a Peripheral Membrane Protein. <i>Molecular and Cellular Biology</i> , 1990, 10, 3873-3883. | 1.1 | 26 |
| 6653 | Mutational Analysis of the Consensus Sequence of a Replication Origin from Yeast Chromosome III. <i>Molecular and Cellular Biology</i> , 1990, 10, 3917-3925. | 1.1 | 119 |
| 6654 | Structure and Molecular Analysis of <i>RGR1</i> , a Gene Required for Glucose Repression of <i>Saccharomyces cerevisiae</i> . <i>Molecular and Cellular Biology</i> , 1990, 10, 4130-4138. | 1.1 | 55 |
| 6655 | <i>IRA2</i> , a Second Gene of <i>Saccharomyces cerevisiae</i> That Encodes a Protein with a Domain Homologous to Mammalian <i>ras</i> GTPase-Activating Protein. <i>Molecular and Cellular Biology</i> , 1990, 10, 4303-4313. | 1.1 | 139 |
| 6656 | Expression and Function of a Human Initiator tRNA Gene in the Yeast <i>Saccharomyces cerevisiae</i> . <i>Molecular and Cellular Biology</i> , 1990, 10, 4486-4494. | 1.1 | 17 |
| 6657 | Phosphorylated Forms of GAL4 Are Correlated with Ability To Activate Transcription. <i>Molecular and Cellular Biology</i> , 1990, 10, 4623-4629. | 1.1 | 33 |

| # | ARTICLE | IF | CITATIONS |
|------|--|-----|-----------|
| 6658 | Characterization of Yeast Vps33p, a Protein Required for Vacuolar Protein Sorting and Vacuole Biogenesis. <i>Molecular and Cellular Biology</i> , 1990, 10, 4638-4649. | 1.1 | 69 |
| 6659 | The N-Terminal TPR Region Is the Functional Domain of SSN6, a Nuclear Phosphoprotein of <i>Saccharomyces cerevisiae</i> . <i>Molecular and Cellular Biology</i> , 1990, 10, 4744-4756. | 1.1 | 94 |
| 6660 | Two Systems of Glucose Repression of the <i>GAL1</i> Promoter in <i>Saccharomyces cerevisiae</i> . <i>Molecular and Cellular Biology</i> , 1990, 10, 4757-4769. | 1.1 | 127 |
| 6661 | A Conserved Sequence in Histone H2A Which Is a Ubiquitination Site in Higher Eucaryotes Is Not Required for Growth in <i>Saccharomyces cerevisiae</i> . <i>Molecular and Cellular Biology</i> , 1990, 10, 4905-4911. | 1.1 | 20 |
| 6662 | Determinants for Glycophospholipid Anchoring of the <i>Saccharomyces cerevisiae</i> GAS1 Protein to the Plasma Membrane. <i>Molecular and Cellular Biology</i> , 1991, 11, 27-37. | 1.1 | 108 |
| 6663 | The Strong <i>ADH1</i> Promoter Stimulates Mitotic and Meiotic Recombination at the <i>ADE6</i> Gene of <i>Schizosaccharomyces pombe</i> . <i>Molecular and Cellular Biology</i> , 1991, 11, 289-298. | 1.1 | 65 |
| 6664 | A Poly(dA-dT) Tract Is a Component of the Recombination Initiation Site at the <i>ARG4</i> Locus in <i>Saccharomyces cerevisiae</i> . <i>Molecular and Cellular Biology</i> , 1991, 11, 322-329. | 1.1 | 82 |
| 6665 | Isolation, Nucleotide Sequence Analysis, and Disruption of the <i>MDH2</i> Gene from <i>Saccharomyces cerevisiae</i> : Evidence for Three Isozymes of Yeast Malate Dehydrogenase. <i>Molecular and Cellular Biology</i> , 1991, 11, 370-380. | 1.1 | 38 |
| 6666 | Intramitochondrial Functions Regulate Nonmitochondrial Citrate Synthase (<i>CIT2</i>) Expression in <i>Saccharomyces cerevisiae</i> . <i>Molecular and Cellular Biology</i> , 1991, 11, 38-46. | 1.1 | 99 |
| 6667 | Suppression of Ribosomal Reinitiation at Upstream Open Reading Frames in Amino Acid-Starved Cells Forms the Basis for <i>GCN4</i> Translational Control. <i>Molecular and Cellular Biology</i> , 1991, 11, 486-496. | 1.1 | 150 |
| 6668 | <i>GRR1</i> of <i>Saccharomyces cerevisiae</i> Is Required for Glucose Repression and Encodes a Protein with Leucine-Rich Repeats. <i>Molecular and Cellular Biology</i> , 1991, 11, 5101-5112. | 1.1 | 106 |
| 6669 | In Vivo Analysis of the <i>Saccharomyces cerevisiae</i> Centromere CDEIII Sequence: Requirements for Mitotic Chromosome Segregation. <i>Molecular and Cellular Biology</i> , 1991, 11, 5212-5221. | 1.1 | 36 |
| 6670 | Quantitation of \hat{I} -Factor Internalization and Response during the <i>Saccharomyces cerevisiae</i> Cell Cycle. <i>Molecular and Cellular Biology</i> , 1991, 11, 5251-5258. | 1.1 | 18 |
| 6671 | Heteroduplex Formation and Mismatch Repair of the α -Stuck Mutation during Mating-Type Switching in <i>Saccharomyces cerevisiae</i> . <i>Molecular and Cellular Biology</i> , 1991, 11, 5372-5380. | 1.1 | 57 |
| 6672 | The <i>CDC20</i> Gene Product of <i>Saccharomyces cerevisiae</i> , a \hat{I}^2 -Transducin Homolog, Is Required for a Subset of Microtubule-Dependent Cellular Processes. <i>Molecular and Cellular Biology</i> , 1991, 11, 5592-5602. | 1.1 | 76 |
| 6673 | Mutations in <i>SPT16/CDC68</i> Suppress <i>cis</i> - and <i>trans</i> -Acting Mutations That Affect Promoter Function in <i>Saccharomyces cerevisiae</i> . <i>Molecular and Cellular Biology</i> , 1991, 11, 5710-5717. | 1.1 | 122 |
| 6674 | <i>CDC55</i> , a <i>Saccharomyces cerevisiae</i> Gene Involved in Cellular Morphogenesis: Identification, Characterization, and Homology to the B Subunit of Mammalian Type 2A Protein Phosphatase. <i>Molecular and Cellular Biology</i> , 1991, 11, 5767-5780. | 1.1 | 190 |
| 6675 | Mutations in a Conserved Region of RNA Polymerase II Influence the Accuracy of mRNA Start Site Selection. <i>Molecular and Cellular Biology</i> , 1991, 11, 5781-5791. | 1.1 | 46 |

| # | ARTICLE | IF | CITATIONS |
|------|--|-----|-----------|
| 6676 | A Putative Zinc Finger Protein, <i>Saccharomyces cerevisiae</i> Vps18p, Affects Late Golgi Functions Required for Vacuolar Protein Sorting and Efficient I \pm -Factor Prohormone Maturation. <i>Molecular and Cellular Biology</i> , 1991, 11, 5813-5824. | 1.1 | 61 |
| 6677 | The Fission Yeast <i>dis3⁺</i> Gene Encodes a 110-kDa Essential Protein Implicated in Mitotic Control. <i>Molecular and Cellular Biology</i> , 1991, 11, 5839-5847. | 1.1 | 57 |
| 6678 | A Functional 125-kDa Core Polypeptide of Fission Yeast DNA Topoisomerase II. <i>Molecular and Cellular Biology</i> , 1991, 11, 6093-6102. | 1.1 | 36 |
| 6679 | Expression of the DAL80 Gene, Whose Product Is Homologous to the GATA Factors and Is a Negative Regulator of Multiple Nitrogen Catabolic Genes in <i>Saccharomyces cerevisiae</i> , Is Sensitive to Nitrogen Catabolite Repression. <i>Molecular and Cellular Biology</i> , 1991, 11, 6205-6215. | 1.1 | 95 |
| 6680 | Sequence and Expression of GLN3, a Positive Nitrogen Regulatory Gene of <i>Saccharomyces cerevisiae</i> Encoding a Protein with a Putative Zinc Finger DNA-Binding Domain. <i>Molecular and Cellular Biology</i> , 1991, 11, 6216-6228. | 1.1 | 119 |
| 6681 | Role of the Complex Upstream Region of the GDH2 Gene in Nitrogen Regulation of the NAD-Linked Glutamate Dehydrogenase in <i>Saccharomyces cerevisiae</i> . <i>Molecular and Cellular Biology</i> , 1991, 11, 6229-6247. | 1.1 | 35 |
| 6682 | RPD1 (SIN3/UME4) Is Required for Maximal Activation and Repression of Diverse Yeast Genes. <i>Molecular and Cellular Biology</i> , 1991, 11, 6306-6316. | 1.1 | 117 |
| 6683 | RPD3 Encodes a Second Factor Required To Achieve Maximum Positive and Negative Transcriptional States in <i>Saccharomyces cerevisiae</i> . <i>Molecular and Cellular Biology</i> , 1991, 11, 6317-6327. | 1.1 | 152 |
| 6684 | The Role of RAP1 in the Regulation of the <i>MAT1\pm</i> Locus. <i>Molecular and Cellular Biology</i> , 1991, 11, 1069-1079. | 1.1 | 58 |
| 6685 | Membrane Insertion of Uracil Permease, a Poly topic Yeast Plasma Membrane Protein. <i>Molecular and Cellular Biology</i> , 1991, 11, 1114-1124. | 1.1 | 56 |
| 6686 | The <i>Saccharomyces cerevisiae</i> ADR1 Gene Is a Positive Regulator of Transcription of Genes Encoding Peroxisomal Proteins. <i>Molecular and Cellular Biology</i> , 1991, 11, 699-704. | 1.1 | 77 |
| 6687 | Characterization of <i>RPR1</i> an Essential Gene Encoding the RNA Component of <i>Saccharomyces cerevisiae</i> Nuclear RNase P. <i>Molecular and Cellular Biology</i> , 1991, 11, 721-730. | 1.1 | 61 |
| 6688 | Specific <i>cis</i> -Acting Sequence for <i>PH08</i> Expression Interacts with PH04 Protein, a Positive Regulatory Factor, in <i>Saccharomyces cerevisiae</i> . <i>Molecular and Cellular Biology</i> , 1991, 11, 785-794. | 1.1 | 22 |
| 6689 | Yeast <i>CBP1</i> mRNA 3' End Formation Is Regulated during the Induction of Mitochondrial Function. <i>Molecular and Cellular Biology</i> , 1991, 11, 813-821. | 1.1 | 21 |
| 6690 | The <i>URE2</i> Gene Product of <i>Saccharomyces cerevisiae</i> Plays an Important Role in the Cellular Response to the Nitrogen Source and Has Homology to Glutathione <i>S</i> -Transferases. <i>Molecular and Cellular Biology</i> , 1991, 11, 822-832. | 1.1 | 128 |
| 6691 | Identification and Structure of Four Yeast Genes (<i>SLY</i>) That Are Able To Suppress the Functional Loss of <i>YPT1</i> , a Member of the <i>RAS</i> Superfamily. <i>Molecular and Cellular Biology</i> , 1991, 11, 872-885. | 1.1 | 142 |
| 6692 | p68 RNA Helicase: Identification of a Nucleolar Form and Cloning of Related Genes Containing a Conserved Intron in Yeasts. <i>Molecular and Cellular Biology</i> , 1991, 11, 1326-1333. | 1.1 | 60 |
| 6693 | Identification of DNA Regions Required for Mitotic and Meiotic Functions within the Centromere of <i>Schizosaccharomyces pombe</i> Chromosome I. <i>Molecular and Cellular Biology</i> , 1991, 11, 2206-2215. | 1.1 | 46 |

| # | ARTICLE | IF | CITATIONS |
|------|--|-----|-----------|
| 6694 | <i>MOD5</i> Translation Initiation Sites Determine <i>N⁶-Isopentenyladenosine</i> Modification of Mitochondrial and Cytoplasmic tRNA. <i>Molecular and Cellular Biology</i> , 1991, 11, 2382-2390. | 1.1 | 49 |
| 6695 | Molecular and Genetic Analysis of the Gene Encoding the <i>Saccharomyces cerevisiae</i> Strand Exchange Protein Sep1. <i>Molecular and Cellular Biology</i> , 1991, 11, 2593-2608. | 1.1 | 62 |
| 6696 | The Yeast Homolog to Mouse <i>Tcp-1</i> Affects Microtubule-Mediated Processes. <i>Molecular and Cellular Biology</i> , 1991, 11, 2629-2640. | 1.1 | 70 |
| 6697 | A Mutation in the Yeast Heat-Shock Factor Gene Causes Temperature-Sensitive Defects in Both Mitochondrial Protein Import and the Cell Cycle. <i>Molecular and Cellular Biology</i> , 1991, 11, 2647-2655. | 1.1 | 58 |
| 6698 | Association of RAP1 Binding Sites with Stringent Control of Ribosomal Protein Gene Transcription in <i>Saccharomyces cerevisiae</i> . <i>Molecular and Cellular Biology</i> , 1991, 11, 2723-2735. | 1.1 | 127 |
| 6699 | The Yeast <i>SLY</i> Gene Products, Suppressors of Defects in the Essential GTP-Binding Ypt1 Protein, May Act in Endoplasmic Reticulum-to-Golgi Transport. <i>Molecular and Cellular Biology</i> , 1991, 11, 2980-2993. | 1.1 | 92 |
| 6700 | <i>SPT5</i> , an Essential Gene Important for Normal Transcription in <i>Saccharomyces cerevisiae</i> , Encodes an Acidic Nuclear Protein with a Carboxy-Terminal Repeat. <i>Molecular and Cellular Biology</i> , 1991, 11, 3009-3019. | 1.1 | 120 |
| 6701 | Different Classes of Polyadenylation Sites in the Yeast <i>Saccharomyces cerevisiae</i> . <i>Molecular and Cellular Biology</i> , 1991, 11, 3060-3069. | 1.1 | 60 |
| 6702 | Translation Initiation Factor 5A and Its Hypusine Modification Are Essential for Cell Viability in the Yeast <i>Saccharomyces cerevisiae</i> . <i>Molecular and Cellular Biology</i> , 1991, 11, 3105-3114. | 1.1 | 173 |
| 6703 | GCD2, a Translational Repressor of the <i>GCN4</i> Gene, Has a General Function in the Initiation of Protein Synthesis in <i>Saccharomyces cerevisiae</i> . <i>Molecular and Cellular Biology</i> , 1991, 11, 3203-3216. | 1.1 | 132 |
| 6704 | Complex Formation by Positive and Negative Translational Regulators of <i>GCN4</i> . <i>Molecular and Cellular Biology</i> , 1991, 11, 3217-3228. | 1.1 | 90 |
| 6705 | The PHO84 gene of <i>Saccharomyces cerevisiae</i> encodes an inorganic phosphate transporter. <i>Molecular and Cellular Biology</i> , 1991, 11, 3229-3238. | 1.1 | 189 |
| 6706 | The CYC8 and TUP1 Proteins Involved in Glucose Repression in <i>Saccharomyces cerevisiae</i> Are Associated in a Protein Complex. <i>Molecular and Cellular Biology</i> , 1991, 11, 3307-3316. | 1.1 | 140 |
| 6707 | The <i>Saccharomyces cerevisiae</i> SRK1 Gene, a Suppressor of <i>bcy1</i> and <i>ins1</i> May Be Involved in Protein Phosphatase Function. <i>Molecular and Cellular Biology</i> , 1991, 11, 3369-3373. | 1.1 | 54 |
| 6708 | In vivo characterization of the <i>Saccharomyces cerevisiae</i> centromere DNA element I, a binding site for the helix-loop-helix protein CPF1. <i>Molecular and Cellular Biology</i> , 1991, 11, 3545-3553. | 1.1 | 35 |
| 6709 | <i>byr2</i> , a <i>Schizosaccharomyces pombe</i> gene encoding a protein kinase capable of partial suppression of the <i>ras1</i> mutant phenotype. <i>Molecular and Cellular Biology</i> , 1991, 11, 3554-3563. | 1.1 | 107 |
| 6710 | Complex Transcriptional Regulation of the <i>Saccharomyces cerevisiae</i> CYB2 Gene Encoding Cytochrome <i>b₂</i> : CYP1 (HAP1) Activator Binds to the <i>CYB2</i> Upstream Activation Site UAS1-B2. <i>Molecular and Cellular Biology</i> , 1991, 11, 3762-3772. | 1.1 | 34 |
| 6711 | AAR1/TUP1 Protein, with a Structure Similar to That of the \hat{I}^2 Subunit of G Proteins, Is Required for $\alpha 1$ - \hat{I}^2 and \hat{I}^2 Repression in Cell Type Control of <i>Saccharomyces cerevisiae</i> . <i>Molecular and Cellular Biology</i> , 1991, 11, 3773-3779. | 1.1 | 61 |

| # | ARTICLE | IF | CITATIONS |
|------|--|-----|-----------|
| 6712 | The <i>HXT1</i> Gene Product of <i>Saccharomyces cerevisiae</i> Is a New Member of the Family of Hexose Transporters. <i>Molecular and Cellular Biology</i> , 1991, 11, 3804-3813. | 1.1 | 55 |
| 6713 | The Highly Conserved N-Terminal Domains of Histones H3 and H4 Are Required for Normal Cell Cycle Progression. <i>Molecular and Cellular Biology</i> , 1991, 11, 4111-4120. | 1.1 | 67 |
| 6714 | <i>TRK1</i> and <i>TRK2</i> Encode Structurally Related K ⁺ Transporters in <i>Saccharomyces cerevisiae</i> . <i>Molecular and Cellular Biology</i> , 1991, 11, 4266-4273. | 1.1 | 113 |
| 6715 | Physiological and genetic analysis of the carbon regulation of the NAD-dependent glutamate dehydrogenase of <i>Saccharomyces cerevisiae</i> . <i>Molecular and Cellular Biology</i> , 1991, 11, 4455-4465. | 1.1 | 25 |
| 6716 | Dominant Mutations in a Gene Encoding a Putative Protein Kinase (<i>BCKT</i>) Bypass the Requirement for a <i>Saccharomyces cerevisiae</i> Protein Kinase C Homolog. <i>Molecular and Cellular Biology</i> , 1992, 12, 172-182. | 1.1 | 235 |
| 6717 | <i>MAS5</i> , a Yeast Homolog of DnaJ Involved in Mitochondrial Protein Import. <i>Molecular and Cellular Biology</i> , 1992, 12, 283-291. | 1.1 | 72 |
| 6718 | <i>NAM9</i> Nuclear Suppressor of Mitochondrial Ochre Mutations in <i>Saccharomyces cerevisiae</i> Codes for a Protein Homologous to S4 Ribosomal Proteins from Chloroplasts, Bacteria, and Eucaryotes. <i>Molecular and Cellular Biology</i> , 1992, 12, 402-412. | 1.1 | 19 |
| 6719 | Characterization of the DNA target site for the yeast ARGR regulatory complex, a sequence able to mediate repression or induction by arginine. <i>Molecular and Cellular Biology</i> , 1992, 12, 68-81. | 1.1 | 42 |
| 6720 | The <i>ARS</i> Consensus Sequence Is Required for Chromosomal Origin Function in <i>Saccharomyces cerevisiae</i> . <i>Molecular and Cellular Biology</i> , 1992, 12, 4305-4313. | 1.1 | 79 |
| 6721 | <i>RPC53</i> Encodes a Subunit of <i>Saccharomyces cerevisiae</i> RNA Polymerase C (III) Whose Inactivation Leads to a Predominantly G ¹ Arrest. <i>Molecular and Cellular Biology</i> , 1992, 12, 4314-4326. | 1.1 | 35 |
| 6722 | <i>RPC82</i> Encodes the Highly Conserved, Third-Largest Subunit of RNA Polymerase Q (III) from <i>Saccharomyces cerevisiae</i> . <i>Molecular and Cellular Biology</i> , 1992, 12, 4433-4440. | 1.1 | 25 |
| 6723 | Ectopic Recombination between Ty Elements in <i>Saccharomyces cerevisiae</i> Is Not Induced by DNA Damage. <i>Molecular and Cellular Biology</i> , 1992, 12, 4441-4448. | 1.1 | 27 |
| 6724 | Involvement of the <i>SIN4</i> Global Transcriptional Regulator in the Chromatin Structure of <i>Saccharomyces cerevisiae</i> . <i>Molecular and Cellular Biology</i> , 1992, 12, 4503-4514. | 1.1 | 102 |
| 6725 | The Yeast <i>EUG1</i> Gene Encodes an Endoplasmic Reticulum Protein That Is Functionally Related to Protein Disulfide Isomerase. <i>Molecular and Cellular Biology</i> , 1992, 12, 4601-4611. | 1.1 | 54 |
| 6726 | The osmotic integrity of the yeast cell requires a functional PKC1 gene product. <i>Molecular and Cellular Biology</i> , 1992, 12, 4896-4905. | 1.1 | 138 |
| 6727 | Identification of a Human cDNA Encoding a Protein That Is Structurally and Functionally Related to the Yeast Adenylyl Cyclase-Associated CAP Proteins. <i>Molecular and Cellular Biology</i> , 1992, 12, 5033-5040. | 1.1 | 26 |
| 6728 | Fission Yeast <i>pap1</i> -Dependent Transcription Is Negatively Regulated by an Essential Nuclear Protein, <i>crml</i> . <i>Molecular and Cellular Biology</i> , 1992, 12, 5474-5484. | 1.1 | 42 |
| 6729 | A REB1-Binding Site Is Required for GCN4-Independent <i>ILV1</i> Basal Level Transcription and Can Be Functionally Replaced by an ABFI-Binding Site. <i>Molecular and Cellular Biology</i> , 1992, 12, 5516-5525. | 1.1 | 20 |

| # | ARTICLE | IF | CITATIONS |
|------|--|-----|-----------|
| 6730 | The EGD1 Product, a Yeast Homolog of Human BTF3, May Be Involved in GAL4 DNA Binding. <i>Molecular and Cellular Biology</i> , 1992, 12, 5683-5689. | 1.1 | 24 |
| 6731 | Yeast RHO3 and RHO4 ras Superfamily Genes Are Necessary for Bud Growth, and Their Defect Is Suppressed by a High Dose of Bud Formation Genes CDC42 and BEM1. <i>Molecular and Cellular Biology</i> , 1992, 12, 5690-5699. | 1.1 | 97 |
| 6732 | Truncated Protein Phosphatase GLC7 Restores Translational Activation of GCN4 Expression in Yeast Mutants Defective for the eIF-2 Kinase GCN2. <i>Molecular and Cellular Biology</i> , 1992, 12, 5700-5710. | 1.1 | 58 |
| 6733 | CTF4 (CHL15) Mutants Exhibit Defective DNA Metabolism in the Yeast <i>Saccharomyces cerevisiae</i> . <i>Molecular and Cellular Biology</i> , 1992, 12, 5736-5747. | 1.1 | 70 |
| 6734 | Mutations Activating the Yeast eIF-2a Kinase GCN2: Isolation of Alleles Altering the Domain Related to Histidyl-tRNA Synthetases. <i>Molecular and Cellular Biology</i> , 1992, 12, 5801-5815. | 1.1 | 81 |
| 6735 | Characterization of Double-Strand Break-Induced Recombination: Homology Requirements and Single-Stranded DNA Formation. <i>Molecular and Cellular Biology</i> , 1992, 12, 563-575. | 1.1 | 291 |
| 6736 | A Dominant Activating Mutation in the Effector Region of RAS Abolishes IRA2 Sensitivity. <i>Molecular and Cellular Biology</i> , 1992, 12, 631-637. | 1.1 | 11 |
| 6737 | RelB, a New Rel Family Transcription Activator That Can Interact with p50-NF- κ B. <i>Molecular and Cellular Biology</i> , 1992, 12, 674-684. | 1.1 | 210 |
| 6738 | <i>RSR1</i> , a <i>ras</i> -Like Gene Homologous to Krev-1 (<i>smg21A/rap1A</i>): Role in the Development of Cell Polarity and Interactions with the Ras Pathway in <i>Saccharomyces cerevisiae</i> . <i>Molecular and Cellular Biology</i> , 1992, 12, 758-766. | 1.1 | 61 |
| 6739 | Role of Multifunctional Autonomously Replicating Sequence Binding Factor 1 in the Initiation of DNA Replication and Transcriptional Control in <i>Saccharomyces cerevisiae</i> . <i>Molecular and Cellular Biology</i> , 1992, 12, 1064-1077. | 1.1 | 48 |
| 6740 | <i>IME4</i> , a Gene That Mediates <i>MAT</i> and Nutritional Control of Meiosis in <i>Saccharomyces cerevisiae</i> . <i>Molecular and Cellular Biology</i> , 1992, 12, 1078-1086. | 1.1 | 92 |
| 6741 | Dissection of a Carboxy-Terminal Region of the Yeast Regulatory Protein RAP1 with Effects on Both Transcriptional Activation and Silencing. <i>Molecular and Cellular Biology</i> , 1992, 12, 1209-1217. | 1.1 | 46 |
| 6742 | Molecular and Genetic Analysis of the Yeast Early Meiotic Recombination Genes <i>REC102</i> and <i>REC107/MER2</i> . <i>Molecular and Cellular Biology</i> , 1992, 12, 1248-1256. | 1.1 | 45 |
| 6743 | Two Alternative Pathways of Double-Strand Break Repair That Are Kinetically Separable and Independently Modulated. <i>Molecular and Cellular Biology</i> , 1992, 12, 1292-1303. | 1.1 | 228 |
| 6744 | <i>MEI4</i> , a Meiosis-Specific Yeast Gene Required for Chromosome Synapsis. <i>Molecular and Cellular Biology</i> , 1992, 12, 1340-1351. | 1.1 | 44 |
| 6745 | Human p53 and <i>CDC2Hs</i> Genes Combine To Inhibit the Proliferation of <i>Saccharomyces cerevisiae</i> . <i>Molecular and Cellular Biology</i> , 1992, 12, 1357-1365. | 1.1 | 50 |
| 6746 | Mutational Analysis of Conserved Positions Potentially Important for Initiator tRNA Function in <i>Saccharomyces cerevisiae</i> . <i>Molecular and Cellular Biology</i> , 1992, 12, 1432-1442. | 1.1 | 25 |
| 6747 | <i>ADR1</i> Mutations Enhance the Ability of ADRI To Activate Transcription by a Mechanism That Is Independent of Effects on Cyclic AMP-Dependent Protein Kinase Phosphorylation of Ser-230. <i>Molecular and Cellular Biology</i> , 1992, 12, 1507-1514. | 1.1 | 30 |

| # | ARTICLE | IF | CITATIONS |
|------|--|-----|-----------|
| 6748 | MET4, a Leucine Zipper Protein, and Centromere-Binding Factor 1 Are Both Required for Transcriptional Activation of Sulfur Metabolism in <i>Saccharomyces cerevisiae</i> . <i>Molecular and Cellular Biology</i> , 1992, 12, 1719-1727. | 1.1 | 108 |
| 6749 | A Presumptive Helicase (<i>MOT1</i> Gene Product) Affects Gene Expression and Is Required for Viability in the Yeast <i>Saccharomyces cerevisiae</i> . <i>Molecular and Cellular Biology</i> , 1992, 12, 1879-1892. | 1.1 | 108 |
| 6750 | Yeast telomere repeat sequence (TRS) improves circular plasmid segregation, and TRS plasmid segregation involves the <i>RAP1</i> gene product.. <i>Molecular and Cellular Biology</i> , 1992, 12, 1997-2009. | 1.1 | 61 |
| 6751 | Yeast Telomere Repeat Sequence (TRS) Improves Circular Plasmid Segregation, and TRS Plasmid Segregation Involves the <i>RAP1</i> Gene Product. <i>Molecular and Cellular Biology</i> , 1992, 12, 1997-2009. | 1.1 | 41 |
| 6752 | Hydrolysis of GTP by Sec4 Protein Plays an Important Role in Vesicular Transport and Is Stimulated by a GTPase-Activating Protein in <i>Saccharomyces cerevisiae</i> . <i>Molecular and Cellular Biology</i> , 1992, 12, 2017-2028. | 1.1 | 62 |
| 6753 | <i>SEN1</i> , a Positive Effector of tRNA-Splicing Endonuclease in <i>Saccharomyces cerevisiae</i> . <i>Molecular and Cellular Biology</i> , 1992, 12, 2154-2164. | 1.1 | 58 |
| 6754 | Gene Products That Promote mRNA Turnover in <i>Saccharomyces cerevisiae</i> . <i>Molecular and Cellular Biology</i> , 1992, 12, 2165-2177. | 1.1 | 198 |
| 6755 | Different Sequence Elements Are Required for Function of the Cauliflower Mosaic Virus Polyadenylation Site in <i>Saccharomyces cerevisiae</i> Compared with in Plants. <i>Molecular and Cellular Biology</i> , 1992, 12, 2322-2330. | 1.1 | 28 |
| 6756 | Biochemical and Genetic Characterization of a Yeast TFIID Mutant That Alters Transcription In Vivo and DNA Binding In Vitro. <i>Molecular and Cellular Biology</i> , 1992, 12, 2372-2382. | 1.1 | 28 |
| 6757 | <i>STP1</i> , a Gene Involved in Pre-tRNA Processing, Encodes a Nuclear Protein Containing Zinc Finger Motifs. <i>Molecular and Cellular Biology</i> , 1992, 12, 2633-2643. | 1.1 | 12 |
| 6758 | <i>SRN1</i> , a Yeast Gene Involved in RNA Processing, Is Identical to <i>HEX2/REG1</i> , a Negative Regulator in Glucose Repression. <i>Molecular and Cellular Biology</i> , 1992, 12, 2673-2680. | 1.1 | 41 |
| 6759 | Overproduction of the <i>GAL1</i> or <i>GAL3</i> Protein Causes Galactose-Independent Activation of the <i>GAL4</i> Protein: Evidence for a New Model of Induction for the Yeast <i>GAL/MEL</i> Regulon. <i>Molecular and Cellular Biology</i> , 1992, 12, 2701-2707. | 1.1 | 60 |
| 6760 | Posttranslational Control of Ty1 Retrotransposition Occurs at the Level of Protein Processing. <i>Molecular and Cellular Biology</i> , 1992, 12, 2813-2825. | 1.1 | 65 |
| 6761 | <i>PMI40</i> , an Intron-Containing Gene Required for Early Steps in Yeast Mannosylation. <i>Molecular and Cellular Biology</i> , 1992, 12, 2924-2930. | 1.1 | 31 |
| 6762 | Control of mRNA Turnover as a Mechanism of Glucose Repression in <i>Saccharomyces cerevisiae</i> . <i>Molecular and Cellular Biology</i> , 1992, 12, 2941-2948. | 1.1 | 33 |
| 6763 | Fused Protein Domains Inhibit DNA Binding by LexA. <i>Molecular and Cellular Biology</i> , 1992, 12, 3006-3014. | 1.1 | 108 |
| 6764 | Semidominant Suppressors of <i>Srs2</i> Helicase Mutations of <i>Saccharomyces cerevisiae</i> Map in the <i>RAD51</i> Gene, Whose Sequence Predicts a Protein with Similarities to Prokaryotic RecA Proteins. <i>Molecular and Cellular Biology</i> , 1992, 12, 3224-3234. | 1.1 | 226 |
| 6765 | <i>COT1</i> , a Gene Involved in Cobalt Accumulation in <i>Saccharomyces cerevisiae</i> . <i>Molecular and Cellular Biology</i> , 1992, 12, 3678-3688. | 1.1 | 100 |

| # | ARTICLE | IF | CITATIONS |
|------|---|-----|-----------|
| 6766 | <i>Saccharomyces cerevisiae</i> <i>RAD5</i> -Encoded DNA Repair Protein Contains DNA Helicase and Zinc-Binding Sequence Motifs and Affects the Stability of Simple Repetitive Sequences in the Genome. <i>Molecular and Cellular Biology</i> , 1992, 12, 3807-3818. | 1.1 | 122 |
| 6767 | <i>PRP38</i> Encodes a Yeast Protein Required for Pre-mRNA Splicing and Maintenance of Stable U6 Small Nuclear RNA Levels. <i>Molecular and Cellular Biology</i> , 1992, 12, 3939-3947. | 1.1 | 59 |
| 6768 | A Transcriptionally Active tRNA Gene Interferes with Nucleosome Positioning In Vivo. <i>Molecular and Cellular Biology</i> , 1992, 12, 4015-4025. | 1.1 | 56 |
| 6769 | <i>DPH5</i> , a Methyltransferase Gene Required for Diphthamide Biosynthesis in <i>Saccharomyces cerevisiae</i> . <i>Molecular and Cellular Biology</i> , 1992, 12, 4026-4037. | 1.1 | 47 |
| 6770 | Replication Forks Pause at Yeast Centromeres. <i>Molecular and Cellular Biology</i> , 1992, 12, 4056-4066. | 1.1 | 99 |
| 6771 | Genetic Interaction between Transcription Elongation Factor TFIIS and RNA Polymerase II. <i>Molecular and Cellular Biology</i> , 1992, 12, 4142-4152. | 1.1 | 94 |
| 6772 | Identification of Pre-mRNA Polyadenylation Sites in <i>Saccharomyces cerevisiae</i> . <i>Molecular and Cellular Biology</i> , 1992, 12, 4215-4229. | 1.1 | 44 |
| 6773 | Interactions among the Subunits of the G Protein Involved in <i>Saccharomyces cerevisiae</i> Mating. <i>Molecular and Cellular Biology</i> , 1993, 13, 1-8. | 1.1 | 56 |
| 6774 | Gene <i>RRN4</i> in <i>Saccharomyces cerevisiae</i> encodes the A12.2 subunit of RNA polymerase I and is essential only at high temperatures. <i>Molecular and Cellular Biology</i> , 1993, 13, 114-122. | 1.1 | 78 |
| 6775 | Characterization of <i>HIR1</i> and <i>HIR2</i> , two genes required for regulation of histone gene transcription in <i>Saccharomyces cerevisiae</i> . <i>Molecular and Cellular Biology</i> , 1993, 13, 28-38. | 1.1 | 82 |
| 6776 | An Essential Yeast Gene with Homology to the Exonuclease-Encoding <i>XRN1/KEM1</i> Gene Also Encodes a Protein with Exoribonuclease Activity. <i>Molecular and Cellular Biology</i> , 1993, 13, 341-350. | 1.1 | 70 |
| 6777 | The Effect on Chromosome Stability of Deleting Replication Origins. <i>Molecular and Cellular Biology</i> , 1993, 13, 391-398. | 1.1 | 53 |
| 6778 | A Conserved Alternative Splice in the von Recklinghausen Neurofibromatosis (<i>NF1</i>) Gene Produces Two Neurofibromin Isoforms, Both of Which Have GTPase-Activating Protein Activity. <i>Molecular and Cellular Biology</i> , 1993, 13, 487-495. | 1.1 | 86 |
| 6779 | <i>GCD11</i> , a Negative Regulator of <i>GCN4</i> Expression, Encodes the β^3 Subunit of eIF-2 in <i>Saccharomyces cerevisiae</i> . <i>Molecular and Cellular Biology</i> , 1993, 13, 506-520. | 1.1 | 52 |
| 6780 | Concerted Action of the Transcriptional Activators <i>REB1</i> , <i>RAP1</i> , and <i>GCR1</i> in the High-Level Expression of the Glycolytic Gene <i>TPI</i> . <i>Molecular and Cellular Biology</i> , 1993, 13, 543-550. | 1.1 | 53 |
| 6781 | <i>SPT13 (GAL11)</i> of <i>Saccharomyces cerevisiae</i> Negatively Regulates Activity of the <i>MCM1</i> Transcription Factor in Ty1 Elements. <i>Molecular and Cellular Biology</i> , 1993, 13, 63-71. | 1.1 | 30 |
| 6782 | Roles of multiple glucose transporters in <i>Saccharomyces cerevisiae</i> . <i>Molecular and Cellular Biology</i> , 1993, 13, 638-648. | 1.1 | 59 |
| 6783 | Dominant Missense Mutations in a Novel Yeast Protein related to Mammalian Phosphatidylinositol 3-Kinase and <i>VPS34</i> Abrogate Rapamycin Cytotoxicity. <i>Molecular and Cellular Biology</i> , 1993, 13, 6012-6023. | 1.1 | 155 |

| # | ARTICLE | IF | CITATIONS |
|------|--|-----|-----------|
| 6784 | PUB1: a Major Yeast Poly(A) ⁺ RNA-Binding Protein. <i>Molecular and Cellular Biology</i> , 1993, 13, 6114-6123. | 1.1 | 30 |
| 6785 | <i>HSP78</i> Encodes a Yeast Mitochondrial Heat Shock Protein in the Clp Family of ATP-Dependent Proteases. <i>Molecular and Cellular Biology</i> , 1993, 13, 6304-6313. | 1.1 | 50 |
| 6786 | The Yeast <i>KRE9</i> Gene Encodes an O Glycoprotein involved in cell surface β -Glucan Assembly. <i>Molecular and Cellular Biology</i> , 1993, 13, 6346-6356. | 1.1 | 49 |
| 6787 | DNA Topoisomerase I Controls the Kinetics of Promoter Activation and DNA Topology in <i>Saccharomyces cerevisiae</i> . <i>Molecular and Cellular Biology</i> , 1993, 13, 6702-6710. | 1.1 | 9 |
| 6788 | Short Artificial Hairpins Sequester Splicing Signals and Inhibit Yeast Pre-mRNA Splicing. <i>Molecular and Cellular Biology</i> , 1993, 13, 6841-6848. | 1.1 | 51 |
| 6789 | AFR1 Acts in Conjunction with the β -Factor Receptor To Promote Morphogenesis and Adaptation. <i>Molecular and Cellular Biology</i> , 1993, 13, 6876-6888. | 1.1 | 28 |
| 6790 | The SH2 Domain is Required for Stable Phosphorylation of p56 ^{lck} at Tyrosine 505, the Negative Regulatory Site. <i>Molecular and Cellular Biology</i> , 1993, 13, 7112-7121. | 1.1 | 22 |
| 6791 | A Common Element Involved in Transcriptional Regulation of Two DNA Alkylation Repair Genes (<i>MAG</i> and <i>MGT1</i>) of <i>Saccharomyces cerevisiae</i> . <i>Molecular and Cellular Biology</i> , 1993, 13, 7213-7221. | 1.1 | 20 |
| 6792 | The <i>Saccharomyces cerevisiae</i> Cdc68 Transcription Activator Is Antagonized by San1, a Protein Implicated in Transcriptional Silencing. <i>Molecular and Cellular Biology</i> , 1993, 13, 7553-7565. | 1.1 | 27 |
| 6793 | <i>DRS1</i> to <i>DRS7</i> , Novel Genes Required for Ribosome Assembly and Function in <i>Saccharomyces cerevisiae</i> . <i>Molecular and Cellular Biology</i> , 1993, 13, 7901-7912. | 1.1 | 55 |
| 6794 | p62 ^{cdc23} of <i>Saccharomyces cerevisiae</i> : A Nuclear Tetratricopeptide Repeat Protein With Two Mutable Domains. <i>Molecular and Cellular Biology</i> , 1993, 13, 1212-1221. | 1.1 | 36 |
| 6795 | The rRNA enhancer regulates rRNA transcription in <i>Saccharomyces cerevisiae</i> . <i>Molecular and Cellular Biology</i> , 1993, 13, 1283-1289. | 1.1 | 19 |
| 6796 | A New Transcriptional-Activation Motif Restricted to a Class of Helix-Loop-Helix Proteins Is Functionally Conserved in Both Yeast and Mammalian Cells. <i>Molecular and Cellular Biology</i> , 1993, 13, 792-800. | 1.1 | 84 |
| 6797 | The Promoter Region of the Yeast <i>KAR2</i> (BiP) Gene Contains a Regulatory Domain that Responds to the Presence of Unfolded Proteins in the Endoplasmic Reticulum. <i>Molecular and Cellular Biology</i> , 1993, 13, 877-890. | 1.1 | 193 |
| 6798 | Conservation of Transcriptional Activation Functions of the NF- κ B p50 and p65 Subunits in Mammalian Cells and <i>Saccharomyces cerevisiae</i> . <i>Molecular and Cellular Biology</i> , 1993, 13, 1666-1674. | 1.1 | 48 |
| 6799 | Transcriptional Repression in <i>Saccharomyces cerevisiae</i> by a SIN3-LexA Fusion Protein. <i>Molecular and Cellular Biology</i> , 1993, 13, 1805-1814. | 1.1 | 70 |
| 6800 | PRP19: a Novel Spliceosomal Component. <i>Molecular and Cellular Biology</i> , 1993, 13, 1876-1882. | 1.1 | 48 |
| 6801 | The <i>COT2</i> Gene Is Required for Glucose-Dependent Divalent Cation Transport in <i>Saccharomyces cerevisiae</i> . <i>Molecular and Cellular Biology</i> , 1993, 13, 2041-2049. | 1.1 | 29 |

| # | ARTICLE | IF | CITATIONS |
|------|--|-----|-----------|
| 6802 | Pheromone-Induced Signal Transduction in <i>Saccharomyces cerevisiae</i> Requires the Sequential Function of Three Protein Kinases. <i>Molecular and Cellular Biology</i> , 1993, 13, 2069-2080. | 1.1 | 92 |
| 6803 | Three Downstream Sites Repress Transcription of a Ty2 Retrotransposon in <i>Saccharomyces cerevisiae</i> . <i>Molecular and Cellular Biology</i> , 1993, 13, 2081-2090. | 1.1 | 20 |
| 6804 | Interspersion of an Unusual GCN4 Activation Site with a Complex Transcriptional Repression Site in Ty2 Elements of <i>Saccharomyces cerevisiae</i> . <i>Molecular and Cellular Biology</i> , 1993, 13, 2091-2103. | 1.1 | 12 |
| 6805 | Differential Function and Expression of <i>Saccharomyces cerevisiae</i> B-type Cyclins in Mitosis and Meiosis. <i>Molecular and Cellular Biology</i> , 1993, 13, 2113-2125. | 1.1 | 100 |
| 6806 | <i>GSP1</i> and <i>GSP2</i> , Genetic Suppressors of the <i>prp20-1</i> Mutant in <i>Saccharomyces cerevisiae</i> : GTP-Binding Proteins Involved in the Maintenance of Nuclear Organization. <i>Molecular and Cellular Biology</i> , 1993, 13, 2152-2161. | 1.1 | 72 |
| 6807 | A Complex Regulatory Element from the Yeast Gene <i>ENO2</i> Modulates <i>GCR1</i> -Dependent Transcriptional Activation. <i>Molecular and Cellular Biology</i> , 1993, 13, 2623-2633. | 1.1 | 20 |
| 6808 | Yeast Ribosomal Protein L1 is Required for the Stability of Newly Synthesized 5S rRNA and the Assembly of 60S Ribosomal Subunits. <i>Molecular and Cellular Biology</i> , 1993, 13, 2835-2845. | 1.1 | 96 |
| 6809 | A U5 Small Nuclear Ribonucleoprotein Particle Protein Involved Only in the Second Step of Pre-mRNA Splicing in <i>Saccharomyces cerevisiae</i> . <i>Molecular and Cellular Biology</i> , 1993, 13, 2959-2970. | 1.1 | 49 |
| 6810 | Architecture of a Yeast U6 RNA Gene Promoter. <i>Molecular and Cellular Biology</i> , 1993, 13, 3015-3026. | 1.1 | 64 |
| 6811 | A Yeast Mitogen-Activated Protein Kinase Homolog (Mpk1p) Mediates Signalling by Protein Kinase C. <i>Molecular and Cellular Biology</i> , 1993, 13, 3067-3075. | 1.1 | 207 |
| 6812 | <i>MKK1</i> and <i>MKK2</i> , Which Encode <i>Saccharomyces cerevisiae</i> Mitogen-Activated Protein Kinase-Kinase Homologs, Function in the Pathway Mediated by Protein Kinase C. <i>Molecular and Cellular Biology</i> , 1993, 13, 3076-3083. | 1.1 | 208 |
| 6813 | A 5'-3' Exonuclease From <i>Saccharomyces cerevisiae</i> is Required for in Vitro Recombination Between Linear DNA Molecules with Overlapping Homology. <i>Molecular and Cellular Biology</i> , 1993, 13, 3125-3134. | 1.1 | 27 |
| 6814 | GCN1, a Translational Activator of <i>GCN4</i> in <i>Saccharomyces cerevisiae</i> , is Required for Phosphorylation of Eukaryotic Translation Initiation Factor 2 by Protein Kinase GCN2. <i>Molecular and Cellular Biology</i> , 1993, 13, 3541-3556. | 1.1 | 73 |
| 6815 | Functional Domains of the Yeast STE12 Protein, a Pheromone-Responsive Transcriptional Activator. <i>Molecular and Cellular Biology</i> , 1993, 13, 3765-3772. | 1.1 | 28 |
| 6816 | The Hexokinase Gene Is Required for Transcriptional Regulation of the Glucose Transporter Gene <i>RAG1</i> in <i>Kluyveromyces lactis</i> . <i>Molecular and Cellular Biology</i> , 1993, 13, 3882-3889. | 1.1 | 39 |
| 6817 | Origin Activation and Formation of Single-Strand TG ₁₋₃ Tails Occur Sequentially in Late S Phase on a Yeast Linear Plasmid. <i>Molecular and Cellular Biology</i> , 1993, 13, 4057-4065. | 1.1 | 61 |
| 6818 | Analysis of the Function of the 70-Kilodalton Cyclase-Associated Protein (CAP) by Using Mutants of Yeast Adenylyl Cyclase Defective in CAP Binding. <i>Molecular and Cellular Biology</i> , 1993, 13, 4087-4097. | 1.1 | 15 |
| 6819 | Genetic Evidence for In Vivo Cross-Specificity of the CaaX-Box Protein Prenyltransferases Farnesyltransferase and Geranylgeranyltransferase-I in <i>Saccharomyces cerevisiae</i> . <i>Molecular and Cellular Biology</i> , 1993, 13, 4260-4275. | 1.1 | 65 |

| # | ARTICLE | IF | CITATIONS |
|------|---|-----|-----------|
| 6820 | Evidence that the <i>SKI</i> Antiviral System of <i>Saccharomyces cerevisiae</i> Acts by Blocking Expression of Viral mRNA. <i>Molecular and Cellular Biology</i> , 1993, 13, 4331-4341. | 1.1 | 81 |
| 6821 | A Multicopy Suppressor Gene of the <i>Saccharomyces cerevisiae</i> G ₁ Cell Cycle Mutant Gene <i>dbf4</i> Encodes a Protein Kinase and Is Identified as <i>CDC5</i> . <i>Molecular and Cellular Biology</i> , 1993, 13, 4445-4457. | 1.1 | 130 |
| 6822 | Centromeres of the Fission Yeast <i>Schizosaccharomyces pombe</i> Are Highly Variable Genetic Loci. <i>Molecular and Cellular Biology</i> , 1993, 13, 4578-4587. | 1.1 | 49 |
| 6823 | Guanine Nucleotide Exchange Factor for Eukaryotic Translation Initiation Factor 2 in <i>Saccharomyces cerevisiae</i> : Interactions between the Essential Subunits GCD2, GCD6, and GCD7 and the Regulatory Subunit GCN3. <i>Molecular and Cellular Biology</i> , 1993, 13, 4618-4631. | 1.1 | 35 |
| 6824 | An Essential Yeast Protein, CBF5p, Binds In Vitro to Centromeres and Microtubules. <i>Molecular and Cellular Biology</i> , 1993, 13, 4884-4893. | 1.1 | 91 |
| 6825 | Promoter Elements Determining Weak Expression of the <i>GAL4</i> Regulatory Gene of <i>Saccharomyces cerevisiae</i> . <i>Molecular and Cellular Biology</i> , 1993, 13, 4999-5009. | 1.1 | 24 |
| 6826 | Location and Characterization of Autonomously Replicating Sequences from Chromosome VI of <i>Saccharomyces cerevisiae</i> . <i>Molecular and Cellular Biology</i> , 1993, 13, 5043-5056. | 1.1 | 70 |
| 6827 | Alteration of a Yeast SH3 Protein Leads to Conditional Viability with Defects in Cytoskeletal and Budding Patterns. <i>Molecular and Cellular Biology</i> , 1993, 13, 5070-5084. | 1.1 | 90 |
| 6828 | Translation of the Yeast Transcriptional Activator <i>GCN4</i> Is Stimulated by Purine Limitation: Implications for Activation of the Protein Kinase GCN2. <i>Molecular and Cellular Biology</i> , 1993, 13, 5099-5111. | 1.1 | 63 |
| 6829 | Mutations in a Protein Tyrosine Phosphatase Gene (<i>PTP2</i>) and a Protein Serine/Threonine Phosphatase Gene (<i>PTC1</i>) Cause a Synthetic Growth Defect in <i>Saccharomyces cerevisiae</i> . <i>Molecular and Cellular Biology</i> , 1993, 13, 5408-5417. | 1.1 | 76 |
| 6830 | Inactivation of <i>YME1</i> , a Member of the <i>ftsH-SEC18-PAS1-CDC48</i> Family of Putative ATPase-Encoding Genes, Causes Increased Escape of DNA from Mitochondria in <i>Saccharomyces cerevisiae</i> . <i>Molecular and Cellular Biology</i> , 1993, 13, 5418-5426. | 1.1 | 81 |
| 6831 | Identification and Purification of a Protein That Binds DNA Cooperatively with the Yeast SWI5 Protein. <i>Molecular and Cellular Biology</i> , 1993, 13, 5524-5537. | 1.1 | 29 |
| 6832 | Far1 and Fus3 Link the Mating Pheromone Signal Transduction Pathway to Three G ₁ -Phase Cdc28 Kinase Complexes. <i>Molecular and Cellular Biology</i> , 1993, 13, 5659-5669. | 1.1 | 108 |
| 6833 | Transcriptional Regulator Leu3 of <i>Saccharomyces cerevisiae</i> : Separation of Activator and Repressor Functions. <i>Molecular and Cellular Biology</i> , 1993, 13, 5702-5709. | 1.1 | 14 |
| 6834 | Novel Members of the Mitogen-Activated Protein Kinase Activator Family in <i>Xenopus laevis</i> . <i>Molecular and Cellular Biology</i> , 1993, 13, 5738-5748. | 1.1 | 20 |
| 6835 | <i>SPK1</i> Is an Essential S-Phase-Specific Gene of <i>Saccharomyces cerevisiae</i> That Encodes a Nuclear Serine/Threonine/Tyrosine Kinase. <i>Molecular and Cellular Biology</i> , 1993, 13, 5829-5842. | 1.1 | 55 |
| 6836 | A Pair of Functionally Redundant Yeast Genes (<i>PPZ1</i> and <i>PPZ2</i>) Encoding Type 1-Related Protein Phosphatases Function within the <i>PKC1</i> -Mediated Pathway. <i>Molecular and Cellular Biology</i> , 1993, 13, 5843-5853. | 1.1 | 89 |
| 6837 | Isolation and characterization of a <i>Saccharomyces cerevisiae</i> peptide transport gene. <i>Molecular and Cellular Biology</i> , 1994, 14, 104-115. | 1.1 | 55 |

| # | ARTICLE | IF | CITATIONS |
|------|---|-----|-----------|
| 6838 | The <i>sua8</i> suppressors of <i>Saccharomyces cerevisiae</i> encode replacements of conserved residues within the largest subunit of RNA polymerase II and affect transcription start site selection similarly to <i>sua7</i> (TFIIB) mutations. <i>Molecular and Cellular Biology</i> , 1994, 14, 226-237. | 1.1 | 57 |
| 6839 | CDC44: a putative nucleotide-binding protein required for cell cycle progression that has homology to subunits of replication factor C. <i>Molecular and Cellular Biology</i> , 1994, 14, 255-267. | 1.1 | 35 |
| 6840 | Functional expression of P-glycoprotein in <i>Saccharomyces cerevisiae</i> confers cellular resistance to the immunosuppressive and antifungal agent FK520. <i>Molecular and Cellular Biology</i> , 1994, 14, 277-286. | 1.1 | 20 |
| 6841 | Dual requirement in yeast DNA mismatch repair for <i>MLH1</i> and <i>PMS1</i> , two homologs of the bacterial <i>mutL</i> gene. <i>Molecular and Cellular Biology</i> , 1994, 14, 407-415. | 1.1 | 141 |
| 6842 | Concerted action of RAS and G proteins in the sexual response pathways of <i>Schizosaccharomyces pombe</i> . <i>Molecular and Cellular Biology</i> , 1994, 14, 50-58. | 1.1 | 38 |
| 6843 | Effect of sequence context at stop codons on efficiency of reinitiation in <i>GCN4</i> translational control. <i>Molecular and Cellular Biology</i> , 1994, 14, 606-618. | 1.1 | 76 |
| 6844 | Dissection of the ADR1 protein reveals multiple, functionally redundant activation domains interspersed with inhibitory regions: evidence for a repressor binding to the ADR1 ^c region. <i>Molecular and Cellular Biology</i> , 1994, 14, 629-640. | 1.1 | 32 |
| 6845 | <i>MDS1</i> , a dosage suppressor of an <i>mck1</i> mutant, encodes a putative yeast homolog of glycogen synthase kinase 3. <i>Molecular and Cellular Biology</i> , 1994, 14, 831-839. | 1.1 | 43 |
| 6846 | Two FK506 Resistance-Confering Genes in <i>Saccharomyces cerevisiae</i> , <i>TAT1</i> and <i>TAT2</i> , Encode Amino Acid Permeases Mediating Tyrosine and Tryptophan Uptake. <i>Molecular and Cellular Biology</i> , 1994, 14, 6597-6606. | 1.1 | 61 |
| 6847 | The GLC7 Type 1 Protein Phosphatase Is Required for Glucose Repression in <i>Saccharomyces cerevisiae</i> . <i>Molecular and Cellular Biology</i> , 1994, 14, 6789-6796. | 1.1 | 70 |
| 6848 | Two Alternative Pathways of Transcription Initiation in the Yeast Negative Regulatory Gene <i>GAL80</i> . <i>Molecular and Cellular Biology</i> , 1994, 14, 6819-6828. | 1.1 | 13 |
| 6849 | Differential Effects of Cdc68 on Cell Cycle-Regulated Promoters in <i>Saccharomyces cerevisiae</i> . <i>Molecular and Cellular Biology</i> , 1994, 14, 7455-7465. | 1.1 | 28 |
| 6850 | The Yeast <i>TEM1</i> Gene, Which Encodes a GTP-Binding Protein, Is Involved in Termination of $\hat{\alpha}$ Phase. <i>Molecular and Cellular Biology</i> , 1994, 14, 7476-7482. | 1.1 | 114 |
| 6851 | Domain B of ARS307 contains two functional elements and contributes to chromosomal replication origin function. <i>Molecular and Cellular Biology</i> , 1994, 14, 7652-7659. | 1.1 | 62 |
| 6852 | Are yeast chitin synthases regulated at the transcriptional or the posttranslational level?. <i>Molecular and Cellular Biology</i> , 1994, 14, 7685-7694. | 1.1 | 49 |
| 6853 | Identification and Analysis of a <i>Saccharomyces cerevisiae</i> Copper Homeostasis Gene Encoding a Homeodomain Protein. <i>Molecular and Cellular Biology</i> , 1994, 14, 7792-7804. | 1.1 | 11 |
| 6854 | G ₁ Cyclin Degradation: the PEST Motif of Yeast Cln2 Is Necessary, but Not Sufficient, for Rapid Protein Turnover. <i>Molecular and Cellular Biology</i> , 1994, 14, 7953-7966. | 1.1 | 62 |
| 6855 | Interactions among Three Proteins That Specifically Activate Translation of the Mitochondrial <i>COX3</i> mRNA in <i>Saccharomyces cerevisiae</i> . <i>Molecular and Cellular Biology</i> , 1994, 14, 1045-1053. | 1.1 | 49 |

| # | ARTICLE | IF | CITATIONS |
|------|--|-----|-----------|
| 6856 | Mutational Activation of the <i>STE5</i> Gene Product Bypasses the Requirement for G Protein $\hat{\beta}^2$ and $\hat{\beta}^3$ Subunits in the Yeast Pheromone Response Pathway. <i>Molecular and Cellular Biology</i> , 1994, 14, 1054-1065. | 1.1 | 49 |
| 6857 | tRNA Genes as Transcriptional Repressor Elements. <i>Molecular and Cellular Biology</i> , 1994, 14, 1266-1277. | 1.1 | 74 |
| 6858 | The Mutant Type 1 Protein Phosphatase Encoded by <i>glc7-1</i> from <i>Saccharomyces cerevisiae</i> Fails To Interact Productively with the <i>GAC1</i> -Encoded Regulatory Subunit. <i>Molecular and Cellular Biology</i> , 1994, 14, 896-905. | 1.1 | 65 |
| 6859 | Induction of Pseudohyphal Growth by Overexpression of <i>PHD1</i> , a <i>Saccharomyces cerevisiae</i> Gene Related to Transcriptional Regulators of Fungal Development. <i>Molecular and Cellular Biology</i> , 1994, 14, 2100-2112. | 1.1 | 174 |
| 6860 | Subcellular Locations of <i>MOD5</i> Proteins: Mapping of Sequences Sufficient for Targeting to Mitochondria and Demonstration that Mitochondrial and Nuclear Isoforms Commingle in the Cytosol. <i>Molecular and Cellular Biology</i> , 1994, 14, 2298-2306. | 1.1 | 38 |
| 6861 | The <i>Aspergillus nidulans abaA</i> Gene Encodes a Transcriptional Activator That Acts as a Genetic Switch To Control Development. <i>Molecular and Cellular Biology</i> , 1994, 14, 2503-2515. | 1.1 | 113 |
| 6862 | Requirements for Intercistronic Distance and Level of Eukaryotic Initiation Factor 2 Activity in Reinitiation on <i>GCN4</i> mRNA Vary with the Downstream Cistron. <i>Molecular and Cellular Biology</i> , 1994, 14, 2616-2628. | 1.1 | 63 |
| 6863 | His-154 Is Involved in the Linkage of the <i>Saccharomyces cerevisiae</i> L-A Double-Stranded RNA Virus Gag Protein to the Cap Structure of mRNAs and Is Essential for M ₁ Satellite Virus Expression. <i>Molecular and Cellular Biology</i> , 1994, 14, 2664-2674. | 1.1 | 35 |
| 6864 | Identification and Characterization of a Novel Yeast Gene: the <i>YGP1</i> Gene Product Is a Highly Glycosylated Secreted Protein That Is Synthesized in Response to Nutrient Limitation. <i>Molecular and Cellular Biology</i> , 1994, 14, 2740-2754. | 1.1 | 33 |
| 6865 | <i>MOT2</i> Encodes a Negative Regulator of Gene Expression That Affects Basal Expression of Pheromone-Responsive Genes in <i>Saccharomyces cerevisiae</i> . <i>Molecular and Cellular Biology</i> , 1994, 14, 3139-3149. | 1.1 | 23 |
| 6866 | The Yeast <i>MOT2</i> Gene Encodes a Putative Zinc Finger Protein That Serves as a Global Negative Regulator Affecting Expression of Several Categories of Genes, Including Mating-Pheromone-Responsive Genes. <i>Molecular and Cellular Biology</i> , 1994, 14, 3150-3157. | 1.1 | 23 |
| 6867 | Mutations in the GCD7 Subunit of Yeast Guanine Nucleotide Exchange Factor eIF-2B Overcome the Inhibitory Effects of Phosphorylated eIF-2 on Translation Initiation. <i>Molecular and Cellular Biology</i> , 1994, 14, 3208-3222. | 1.1 | 39 |
| 6868 | Genetic Identification of Residues Involved in Association of $\hat{\alpha}$ and $\hat{\beta}^2$ G-Protein Subunits. <i>Molecular and Cellular Biology</i> , 1994, 14, 3223-3229. | 1.1 | 23 |
| 6869 | The Third Cytoplasmic Loop of a Yeast G-Protein-Coupled Receptor Controls Pathway Activation, Ligand Discrimination, and Receptor Internalization. <i>Molecular and Cellular Biology</i> , 1994, 14, 3339-3349. | 1.1 | 29 |
| 6870 | Mutations in <i>XRS2</i> and <i>RAD50</i> Delay but Do Not Prevent Mating-Type Switching in <i>Saccharomyces cerevisiae</i> . <i>Molecular and Cellular Biology</i> , 1994, 14, 3414-3425. | 1.1 | 149 |
| 6871 | <i>TPD1</i> of <i>Saccharomyces cerevisiae</i> Encodes a Protein Phosphatase 2C-Like Activity Implicated in tRNA Splicing and Cell Separation. <i>Molecular and Cellular Biology</i> , 1994, 14, 3634-3645. | 1.1 | 38 |
| 6872 | Two Types of RAS Mutants That Dominantly Interfere with Activators of RAS. <i>Molecular and Cellular Biology</i> , 1994, 14, 3707-3718. | 1.1 | 26 |
| 6873 | Protein Phosphatase 2C, Encoded by <i>ptc1⁺</i> , Is Important in the Heat Shock Response of <i>Schizosaccharomyces pombe</i> . <i>Molecular and Cellular Biology</i> , 1994, 14, 3742-3751. | 1.1 | 32 |

| # | ARTICLE | IF | CITATIONS |
|------|--|-----|-----------|
| 6874 | Fine-Resolution Mapping of Spontaneous and Double-Strand Break-Induced Gene Conversion Tracts in <i>Saccharomyces cerevisiae</i> Reveals Reversible Mitotic Conversion Polarity. <i>Molecular and Cellular Biology</i> , 1994, 14, 3863-3875. | 1.1 | 72 |
| 6875 | <i>GPD1</i> , Which Encodes Glycerol-3-Phosphate Dehydrogenase, Is Essential for Growth under Osmotic Stress in <i>Saccharomyces cerevisiae</i> , and Its Expression Is Regulated by the High-Osmolarity Glycerol Response Pathway. <i>Molecular and Cellular Biology</i> , 1994, 14, 4135-4144. | 1.1 | 317 |
| 6876 | Biochemical and Genetic Analysis of Dominant-Negative Mutations Affecting a Yeast G-Protein β^3 Subunit. <i>Molecular and Cellular Biology</i> , 1994, 14, 4571-4578. | 1.1 | 17 |
| 6877 | Myocyte Nuclear Factor, a Novel Winged-Helix Transcription Factor under both Developmental and Neural Regulation in Striated Myocytes. <i>Molecular and Cellular Biology</i> , 1994, 14, 4596-4605. | 1.1 | 38 |
| 6878 | Flexibility and Interchangeability of Polyadenylation Signals in <i>Saccharomyces cerevisiae</i> . <i>Molecular and Cellular Biology</i> , 1994, 14, 4633-4642. | 1.1 | 15 |
| 6879 | T7 RNA Polymerase-Dependent Expression of COXII in Yeast Mitochondria. <i>Molecular and Cellular Biology</i> , 1994, 14, 4643-4652. | 1.1 | 30 |
| 6880 | Transcriptional Control of the Yeast <i>PDR5</i> Gene by the <i>PDR3</i> Gene Product. <i>Molecular and Cellular Biology</i> , 1994, 14, 4653-4661. | 1.1 | 83 |
| 6881 | Type 1 Protein Phosphatase Acts in Opposition to Ipl1 Protein Kinase in Regulating Yeast Chromosome Segregation. <i>Molecular and Cellular Biology</i> , 1994, 14, 4731-4740. | 1.1 | 115 |
| 6882 | Induction of recombination between homologous and diverged DNAs by double-strand gaps and breaks and role of mismatch repair. <i>Molecular and Cellular Biology</i> , 1994, 14, 4802-4814. | 1.1 | 50 |
| 6883 | The Yeast and Mammalian Ras Pathways Control Transcription of Heat Shock Genes Independently of Heat Shock Transcription Factor. <i>Molecular and Cellular Biology</i> , 1994, 14, 4929-4937. | 1.1 | 28 |
| 6884 | <i>SPT10</i> and <i>SPT21</i> Are Required for Transcription of Particular Histone Genes in <i>Saccharomyces cerevisiae</i> . <i>Molecular and Cellular Biology</i> , 1994, 14, 5223-5228. | 1.1 | 63 |
| 6885 | <i>GSH1</i> , Which Encodes β^3 -Glutamylcysteine Synthetase, Is a Target Gene for yAP-1 Transcriptional Regulation. <i>Molecular and Cellular Biology</i> , 1994, 14, 5832-5839. | 1.1 | 97 |
| 6886 | Mutation of the Gene Encoding Protein Kinase C 1 Stimulates Mitotic Recombination in <i>Saccharomyces cerevisiae</i> . <i>Molecular and Cellular Biology</i> , 1994, 14, 6039-6045. | 1.1 | 43 |
| 6887 | Two <i>Saccharomyces cerevisiae</i> Genes Which Control Sensitivity to G ₁ Arrest Induced by <i>Kluyveromyces lactis</i> Toxin. <i>Molecular and Cellular Biology</i> , 1994, 14, 6306-6316. | 1.1 | 33 |
| 6888 | Identification of <i>cut8⁺</i> and <i>cek1⁺</i> , a Novel Protein Kinase Gene, Which Complement a Fission Yeast Mutation That Blocks Anaphase. <i>Molecular and Cellular Biology</i> , 1994, 14, 6361-6371. | 1.1 | 20 |
| 6889 | Cloning of <i>Saccharomyces cerevisiae</i> DNA replication genes: isolation of the CDC8 gene and two genes that compensate for the cdc8-1 mutation. <i>Molecular and Cellular Biology</i> , 1983, 3, 1730-1737. | 1.1 | 157 |
| 6890 | Structure and expression of the SNF1 gene of <i>Saccharomyces cerevisiae</i> . <i>Molecular and Cellular Biology</i> , 1984, 4, 54-60. | 1.1 | 53 |
| 6891 | Codon Recognition During Frameshift Suppression in <i>Saccharomyces cerevisiae</i> . <i>Molecular and Cellular Biology</i> , 1984, 4, 2052-2061. | 1.1 | 24 |

| # | ARTICLE | IF | CITATIONS |
|------|--|-----|-----------|
| 6892 | Deletion Mutations Affecting Autonomously Replicating Sequence <i>ARS1</i> of <i>Saccharomyces cerevisiae</i> . <i>Molecular and Cellular Biology</i> , 1984, 4, 2455-2466. | 1.1 | 147 |
| 6893 | Expression of plasmid R388-encoded type II dihydrofolate reductase as a dominant selective marker in <i>Saccharomyces cerevisiae</i> . <i>Molecular and Cellular Biology</i> , 1984, 4, 407-414. | 1.1 | 25 |
| 6894 | Effects of altered 5'-flanking sequences on the in vivo expression of a <i>Saccharomyces cerevisiae</i> tRNATyr gene. <i>Molecular and Cellular Biology</i> , 1984, 4, 657-665. | 1.1 | 98 |
| 6895 | Intrachromosomal movement of genetically marked <i>Saccharomyces cerevisiae</i> transposons by gene conversion. <i>Molecular and Cellular Biology</i> , 1984, 4, 703-711. | 1.1 | 39 |
| 6896 | Regulation of Basal and Induced Levels of the <i>MEL1</i> Transcript in <i>Saccharomyces cerevisiae</i> . <i>Molecular and Cellular Biology</i> , 1984, 4, 1238-1245. | 1.1 | 64 |
| 6897 | Two genes for ribosomal protein 51 of <i>Saccharomyces cerevisiae</i> complement and contribute to the ribosomes. <i>Molecular and Cellular Biology</i> , 1984, 4, 1871-1879. | 1.1 | 80 |
| 6898 | Isolation and DNA Sequence of <i>ADH3</i> , a Nuclear Gene Encoding the Mitochondrial Isozyme of Alcohol Dehydrogenase in <i>Saccharomyces cerevisiae</i> . <i>Molecular and Cellular Biology</i> , 1985, 5, 3024-3034. | 1.1 | 82 |
| 6899 | Cloning and Molecular Analysis of the <i>HAP2</i> Locus: a Global Regulator of Respiratory Genes in <i>Saccharomyces cerevisiae</i> . <i>Molecular and Cellular Biology</i> , 1985, 5, 3410-3416. | 1.1 | 94 |
| 6900 | Effect of <i>RP51</i> Gene Dosage Alterations on Ribosome Synthesis in <i>Saccharomyces cerevisiae</i> . <i>Molecular and Cellular Biology</i> , 1985, 5, 3429-3435. | 1.1 | 79 |
| 6901 | Suppressible and Nonsuppressible +1 G-C Base Pair Insertions Induced by ICR-170 at the <i>his4</i> Locus in <i>Saccharomyces cerevisiae</i> . <i>Molecular and Cellular Biology</i> , 1985, 5, 2247-2256. | 1.1 | 21 |
| 6902 | Identification of Autonomously Replicating Circular Subtelomeric Y' Elements in <i>Saccharomyces cerevisiae</i> . <i>Molecular and Cellular Biology</i> , 1985, 5, 2369-2380. | 1.1 | 36 |
| 6903 | Replicating Plasmids in <i>Schizosaccharomyces pombe</i> : Improvement of Symmetric Segregation by a New Genetic Element. <i>Molecular and Cellular Biology</i> , 1986, 6, 80-89. | 1.1 | 95 |
| 6904 | Expression of the <i>Saccharomyces cerevisiae</i> Inositol-1-Phosphate Synthase (<i>INO1</i>) Gene is Regulated by Factors That Affect Phospholipid Synthesis. <i>Molecular and Cellular Biology</i> , 1986, 6, 3320-3328. | 1.1 | 122 |
| 6905 | Identification of the Crossover Site During <i>FLP</i> -Mediated Recombination in the <i>Saccharomyces cerevisiae</i> Plasmid 2 μ m Circle. <i>Molecular and Cellular Biology</i> , 1986, 6, 3357-3367. | 1.1 | 61 |
| 6906 | Molecular Analysis of <i>SNF2</i> and <i>SNF5</i> , Genes Required for Expression of Glucose-Repressible Genes in <i>Saccharomyces cerevisiae</i> . <i>Molecular and Cellular Biology</i> , 1986, 6, 3643-3651. | 1.1 | 85 |
| 6907 | Genetically Essential and Nonessential $\hat{\pm}$ -Tubulin Genes Specify Functionally Interchangeable Proteins. <i>Molecular and Cellular Biology</i> , 1986, 6, 3722-3733. | 1.1 | 166 |
| 6908 | Glycolytic Gene Expression in <i>Saccharomyces cerevisiae</i> : Nucleotide Sequence of <i>GCR1</i> , Null Mutants, and Evidence for Expression. <i>Molecular and Cellular Biology</i> , 1986, 6, 3774-3784. | 1.1 | 80 |
| 6909 | Structural and Functional Analysis of the <i>MAL1</i> Locus of <i>Saccharomyces cerevisiae</i> . <i>Molecular and Cellular Biology</i> , 1986, 6, 3891-3899. | 1.1 | 61 |

| # | ARTICLE | IF | CITATIONS |
|------|--|-----|-----------|
| 6910 | Developmental Regulation of <i>Dictyostelium discoideum</i> Actin Gene Fusions Carried on Low-Copy and High-Copy Transformation Vectors. <i>Molecular and Cellular Biology</i> , 1986, 6, 3973-3983. | 1.1 | 129 |
| 6911 | Multiple <i>GCD</i> Genes Required for Repression of <i>GCN4</i> , a Transcriptional Activator of Amino Acid Biosynthetic Genes in <i>Saccharomyces cerevisiae</i> . <i>Molecular and Cellular Biology</i> , 1986, 6, 3990-3998. | 1.1 | 75 |
| 6912 | Effects of null mutations in the hexokinase genes of <i>Saccharomyces cerevisiae</i> on catabolite repression. <i>Molecular and Cellular Biology</i> , 1986, 6, 4046-4052. | 1.1 | 77 |
| 6913 | Sequence analysis of temperature-sensitive mutations in the <i>Saccharomyces cerevisiae</i> gene <i>CDC28</i> . <i>Molecular and Cellular Biology</i> , 1986, 6, 4099-4103. | 1.1 | 42 |
| 6914 | Single base-pair mutations in centromere element III cause aberrant chromosome segregation in <i>Saccharomyces cerevisiae</i> . <i>Molecular and Cellular Biology</i> , 1986, 6, 530-538. | 1.1 | 99 |
| 6915 | Sequences Required for Delivery and Localization of the ADP/ATP Translocator to the Mitochondrial Inner Membrane. <i>Molecular and Cellular Biology</i> , 1986, 6, 626-634. | 1.1 | 73 |
| 6916 | Tripartite upstream promoter element essential for expression of <i>Saccharomyces cerevisiae</i> ribosomal protein genes. <i>Molecular and Cellular Biology</i> , 1986, 6, 674-687. | 1.1 | 114 |
| 6917 | <i>RAD7</i> gene of <i>Saccharomyces cerevisiae</i> : transcripts, nucleotide sequence analysis, and functional relationship between the <i>RAD7</i> and <i>RAD23</i> gene products. <i>Molecular and Cellular Biology</i> , 1986, 6, 1497-1507. | 1.1 | 49 |
| 6918 | Sequence and expression of the dCMP deaminase gene (<i>DCD1</i>) of <i>Saccharomyces cerevisiae</i> . <i>Molecular and Cellular Biology</i> , 1986, 6, 1711-1721. | 1.1 | 29 |
| 6919 | <i>Saccharomyces cerevisiae</i> Contains Two Functional Citrate Synthase Genes. <i>Molecular and Cellular Biology</i> , 1986, 6, 1936-1942. | 1.1 | 77 |
| 6920 | An RNA polymerase I enhancer in <i>Saccharomyces cerevisiae</i> . <i>Molecular and Cellular Biology</i> , 1986, 6, 2089-2097. | 1.1 | 121 |
| 6921 | Toxic Effects of Excess Cloned Centromeres. <i>Molecular and Cellular Biology</i> , 1986, 6, 2213-2222. | 1.1 | 86 |
| 6922 | Transcriptional Regulation of Ribosomal Proteins During a Nutritional Upshift in <i>Saccharomyces cerevisiae</i> . <i>Molecular and Cellular Biology</i> , 1986, 6, 2429-2435. | 1.1 | 34 |
| 6923 | The <i>PEP4</i> Gene Encodes an Aspartyl Protease Implicated in the Posttranslational Regulation of <i>Saccharomyces cerevisiae</i> Vacuolar Hydrolases. <i>Molecular and Cellular Biology</i> , 1986, 6, 2500-2510. | 1.1 | 165 |
| 6924 | Transcriptional Control of Glucoamylase Synthesis in Vegetatively Growing and Sporulating <i>Saccharomyces</i> Species. <i>Molecular and Cellular Biology</i> , 1986, 6, 3034-3041. | 1.1 | 28 |
| 6925 | The Membrane-Associated Enzyme Phosphatidylserine Synthase Is Regulated at the Level of mRNA Abundance. <i>Molecular and Cellular Biology</i> , 1987, 7, 167-176. | 1.1 | 63 |
| 6926 | Isolation and Characterization of <i>MODS</i> , a Gene Required for Isopentenylation of Cytoplasmic and Mitochondrial tRNAs of <i>Saccharomyces cerevisiae</i> . <i>Molecular and Cellular Biology</i> , 1987, 7, 177-184. | 1.1 | 86 |
| 6927 | <i>Ty1</i> Sequence with Enhancer and Mating-Type-Dependent Regulatory Activities. <i>Molecular and Cellular Biology</i> , 1987, 7, 258-265. | 1.1 | 64 |

| # | ARTICLE | IF | CITATIONS |
|------|---|-----|-----------|
| 6928 | Primary Structure Requirements for Correct Sorting of the Yeast Mitochondrial Protein ADH III to the Yeast Mitochondrial Matrix Space. <i>Molecular and Cellular Biology</i> , 1987, 7, 294-304. | 1.1 | 23 |
| 6929 | Alterations in the adenine-plus-thymine-rich region of CEN3 affect centromere function in <i>Saccharomyces cerevisiae</i> . <i>Molecular and Cellular Biology</i> , 1987, 7, 68-75. | 1.1 | 84 |
| 6930 | Cloning and characterization of the low-affinity cyclic AMP phosphodiesterase gene of <i>Saccharomyces cerevisiae</i> . <i>Molecular and Cellular Biology</i> , 1987, 7, 3629-3636. | 1.1 | 100 |
| 6931 | The yeast ARD1 gene product is required for repression of cryptic mating-type information at the HML locus. <i>Molecular and Cellular Biology</i> , 1987, 7, 3713-3722. | 1.1 | 91 |
| 6932 | Sequences Required for Transcriptional Initiation of the <i>Saccharomyces cerevisiae</i> CYC7 Gene. <i>Molecular and Cellular Biology</i> , 1987, 7, 3785-3791. | 1.1 | 35 |
| 6933 | Genetic and Biochemical Characterization of Clathrin-Deficient <i>Saccharomyces cerevisiae</i> . <i>Molecular and Cellular Biology</i> , 1987, 7, 3888-3898. | 1.1 | 74 |
| 6934 | Yeast Pre-mRNA Splicing Requires a Minimum Distance between the 5' Splice Site and the Internal Branch Acceptor Site. <i>Molecular and Cellular Biology</i> , 1987, 7, 4010-4016. | 1.1 | 14 |
| 6935 | The Yeast F ₁ -ATPase \hat{I}^2 Subunit Precursor Contains Functionally Redundant Mitochondrial Protein Import Information. <i>Molecular and Cellular Biology</i> , 1987, 7, 4038-4047. | 1.1 | 32 |
| 6936 | Amino Acid Sequences That Determine the Nuclear Localization of Yeast Histone 2B. <i>Molecular and Cellular Biology</i> , 1987, 7, 4048-4057. | 1.1 | 118 |
| 6937 | Trans-acting regulatory mutations that alter transcription of <i>Saccharomyces cerevisiae</i> histone genes. <i>Molecular and Cellular Biology</i> , 1987, 7, 4204-4210. | 1.1 | 85 |
| 6938 | Identification of Healed Terminal DNA Fragments in Linear Minichromosomes of <i>Schizosaccharomyces pombe</i> . <i>Molecular and Cellular Biology</i> , 1987, 7, 4424-4430. | 1.1 | 42 |
| 6939 | Mutational and In Vitro Protein-Binding Studies on Centromere DNA from <i>Saccharomyces cerevisiae</i> . <i>Molecular and Cellular Biology</i> , 1987, 7, 4522-4534. | 1.1 | 83 |
| 6940 | Sequence and Nuclear Localization of the <i>Saccharomyces cerevisiae</i> HAP2 Protein, a Transcriptional Activator. <i>Molecular and Cellular Biology</i> , 1987, 7, 578-585. | 1.1 | 96 |
| 6941 | Role of Transcriptional and Posttranscriptional Regulation in Expression of Histone Genes in <i>Saccharomyces cerevisiae</i> . <i>Molecular and Cellular Biology</i> , 1987, 7, 614-621. | 1.1 | 58 |
| 6942 | The <i>SPT6</i> Gene is Essential for Growth and is Required for \hat{I} -Mediated Transcription in <i>Saccharomyces cerevisiae</i> . <i>Molecular and Cellular Biology</i> , 1987, 7, 679-686. | 1.1 | 134 |
| 6943 | The GCR1 gene encodes a positive transcriptional regulator of the enolase and glyceraldehyde-3-phosphate dehydrogenase gene families in <i>Saccharomyces cerevisiae</i> . <i>Molecular and Cellular Biology</i> , 1987, 7, 813-820. | 1.1 | 83 |
| 6944 | Regulation of <i>RAD54</i> - and <i>RAD52-lacZ</i> Gene Fusions in <i>Saccharomyces cerevisiae</i> in Response to DNA Damage. <i>Molecular and Cellular Biology</i> , 1987, 7, 1078-1084. | 1.1 | 70 |
| 6945 | Concerted Deletions and Inversions Are Caused by Mitotic Recombination between Delta Sequences in <i>Saccharomyces cerevisiae</i> . <i>Molecular and Cellular Biology</i> , 1987, 7, 1198-1207. | 1.1 | 95 |

| # | ARTICLE | IF | CITATIONS |
|------|--|-----|-----------|
| 6946 | Cloning and Characterization of <i>LOS1</i> , a <i>Saccharomyces cerevisiae</i> Gene That Affects tRNA Splicing. <i>Molecular and Cellular Biology</i> , 1987, 7, 1208-1216. | 1.1 | 87 |
| 6947 | mRNA Cap-Binding Protein: Cloning of the Gene Encoding Protein Synthesis Initiation Factor eIF-4E from <i>Saccharomyces cerevisiae</i> . <i>Molecular and Cellular Biology</i> , 1987, 7, 998-1003. | 1.1 | 69 |
| 6948 | Phosphorylation of the <i>Saccharomyces cerevisiae</i> Equivalent of Ribosomal Protein S6 Has No Detectable Effect on Growth. <i>Molecular and Cellular Biology</i> , 1987, 7, 1338-1345. | 1.1 | 21 |
| 6949 | Cloning and Characterization of <i>BCY1</i> , a Locus Encoding a Regulatory Subunit of the Cyclic AMP-Dependent Protein Kinase in <i>Saccharomyces cerevisiae</i> . <i>Molecular and Cellular Biology</i> , 1987, 7, 1371-1377. | 1.1 | 284 |
| 6950 | Structure and Expression of the <i>Saccharomyces Cerevisiae</i> <i>CRY1</i> Gene: a Highly Conserved Ribosomal Protein Gene. <i>Molecular and Cellular Biology</i> , 1987, 7, 1764-1775. | 1.1 | 50 |
| 6951 | Transcriptional regulation of an <i>hsp70</i> heat shock gene in the yeast <i>Saccharomyces cerevisiae</i> . <i>Molecular and Cellular Biology</i> , 1987, 7, 1906-1916. | 1.1 | 96 |
| 6952 | Isolation and Characterization of Temperature-Sensitive Rna Polymerase II Mutants of <i>Saccharomyces cerevisiae</i> . <i>Molecular and Cellular Biology</i> , 1987, 7, 2155-2164. | 1.1 | 41 |
| 6953 | Elements Involved in Oxygen Regulation of the <i>Saccharomyces Cerevisiae</i> <i>CYC7</i> Gene. <i>Molecular and Cellular Biology</i> , 1987, 7, 2212-2220. | 1.1 | 64 |
| 6954 | Fatty acylation is important but not essential for <i>Saccharomyces cerevisiae</i> RAS function. <i>Molecular and Cellular Biology</i> , 1987, 7, 2344-2351. | 1.1 | 63 |
| 6955 | Characterization of <i>Saccharomyces cerevisiae</i> genes encoding subunits of cyclic AMP-dependent protein kinase. <i>Molecular and Cellular Biology</i> , 1987, 7, 2653-2663. | 1.1 | 148 |
| 6956 | <i>Saccharomyces cerevisiae</i> positive regulatory gene <i>PET111</i> encodes a mitochondrial protein that is translated from an mRNA with a long 5' leader. <i>Molecular and Cellular Biology</i> , 1987, 7, 2728-2734. | 1.1 | 47 |
| 6957 | Identification and isolation of the gene encoding the small subunit of ribonucleotide reductase from <i>Saccharomyces cerevisiae</i> : DNA damage-inducible gene required for mitotic viability. <i>Molecular and Cellular Biology</i> , 1987, 7, 2783-2793. | 1.1 | 144 |
| 6958 | Codon Replacement in the <i>PGK1</i> Gene of <i>Saccharomyces Cerevisiae</i> : Experimental Approach to study the role of Biased Codon Usage in Gene Expression. <i>Molecular and Cellular Biology</i> , 1987, 7, 2914-2924. | 1.1 | 109 |
| 6959 | Cell-type-dependent gene activation by yeast transposon <i>Ty1</i> involves multiple regulatory determinants. <i>Molecular and Cellular Biology</i> , 1987, 7, 3205-3211. | 1.1 | 37 |
| 6960 | Isolation and Characterization of Temperature-Sensitive Mutations in <i>RPA190</i> , the Gene Encoding the Largest Subunit of RNA Polymerase I from <i>Saccharomyces cerevisiae</i> . <i>Molecular and Cellular Biology</i> , 1988, 8, 3997-4008. | 1.1 | 43 |
| 6961 | Direction of Chromosome Rearrangements in <i>Saccharomyces cerevisiae</i> by Use of <i>his3</i> Recombinational Substrates. <i>Molecular and Cellular Biology</i> , 1988, 8, 4370-4380. | 1.1 | 29 |
| 6962 | A Regulatory Region Responsible for Proline-Specific Induction of the Yeast <i>PUT2</i> Gene Is Adjacent to Its TATA Box. <i>Molecular and Cellular Biology</i> , 1988, 8, 4634-4641. | 1.1 | 35 |
| 6963 | Characterization of Two Telomeric DNA Processing Reactions in <i>Saccharomyces cerevisiae</i> . <i>Molecular and Cellular Biology</i> , 1988, 8, 4642-4650. | 1.1 | 50 |

| # | ARTICLE | IF | CITATIONS |
|------|---|-----|-----------|
| 6964 | Molecular Analysis of <i>GCN3</i> , a Translational Activator of <i>GCN4</i> : Evidence for Posttranslational Control of <i>GCN3</i> Regulatory Function. <i>Molecular and Cellular Biology</i> , 1988, 8, 4808-4820. | 1.1 | 58 |
| 6965 | Copy Number and Partition of the <i>Saccharomyces cerevisiae</i> 2 μ m Plasmid Controlled by Transcription Regulators. <i>Molecular and Cellular Biology</i> , 1988, 8, 4949-4957. | 1.1 | 21 |
| 6966 | <i>GAL11</i> Protein, an Auxiliary Transcription Activator for Genes Encoding Galactose-Metabolizing Enzymes in <i>Saccharomyces cerevisiae</i> . <i>Molecular and Cellular Biology</i> , 1988, 8, 4991-4999. | 1.1 | 106 |
| 6967 | Connections Between Transcriptional Activators, Silencers, and Telomeres as Revealed by Functional Analysis of a Yeast DNA-Binding Protein. <i>Molecular and Cellular Biology</i> , 1988, 8, 5086-5099. | 1.1 | 238 |
| 6968 | <i>SAM2</i> Encodes the Second Methionine <i>S</i> -Adenosyl Transferase in <i>Saccharomyces cerevisiae</i> : Physiology and Regulation of Both Enzymes. <i>Molecular and Cellular Biology</i> , 1988, 8, 5132-5139. | 1.1 | 67 |
| 6969 | Isolation of a Yeast Gene Involved in Species-Specific Pre-tRNA Processing. <i>Molecular and Cellular Biology</i> , 1988, 8, 5140-5149. | 1.1 | 34 |
| 6970 | Ty3, a Yeast Retrotransposon Associated with tRNA Genes, Has Homology to Animal Retroviruses. <i>Molecular and Cellular Biology</i> , 1988, 8, 5245-5256. | 1.1 | 160 |
| 6971 | Gene Conversion Adjacent to Regions of Double-Strand Break Repair. <i>Molecular and Cellular Biology</i> , 1988, 8, 5292-5298. | 1.1 | 42 |
| 6972 | Cloning of the <i>PYR3</i> Gene of <i>Ustilago maydis</i> and Its Use in DNA Transformation. <i>Molecular and Cellular Biology</i> , 1988, 8, 5417-5424. | 1.1 | 16 |
| 6973 | Structural organization and functional analysis of centromeric DNA in the fission yeast <i>Schizosaccharomyces pombe</i> . <i>Molecular and Cellular Biology</i> , 1988, 8, 754-763. | 1.1 | 76 |
| 6974 | Domain structure and functional analysis of the carboxyl-terminal polyacidic sequence of the RAD6 protein of <i>Saccharomyces cerevisiae</i> . <i>Molecular and Cellular Biology</i> , 1988, 8, 1179-1185. | 1.1 | 52 |
| 6975 | The a-factor pheromone of <i>Saccharomyces cerevisiae</i> is essential for mating. <i>Molecular and Cellular Biology</i> , 1988, 8, 1309-1318. | 1.1 | 174 |
| 6976 | Functional organization of the retrotransposon Ty from <i>Saccharomyces cerevisiae</i> : Ty protease is required for transposition. <i>Molecular and Cellular Biology</i> , 1988, 8, 1421-1431. | 1.1 | 73 |
| 6977 | Regulation of expression and activity of the yeast transcription factor ADR1. <i>Molecular and Cellular Biology</i> , 1988, 8, 1868-1876. | 1.1 | 42 |
| 6978 | Identification of Functional Regions in the Yeast Transcriptional Activator ADR1. <i>Molecular and Cellular Biology</i> , 1988, 8, 2125-2131. | 1.1 | 47 |
| 6979 | Transcription Interferes with Elements Important for Chromosome Maintenance in <i>Saccharomyces cerevisiae</i> . <i>Molecular and Cellular Biology</i> , 1988, 8, 2184-2194. | 1.1 | 115 |
| 6980 | Physical Monitoring of Mating Type Switching in <i>Saccharomyces cerevisiae</i> . <i>Molecular and Cellular Biology</i> , 1988, 8, 2342-2349. | 1.1 | 84 |
| 6981 | Mutations in a Gene Encoding the β Subunit of a <i>Saccharomyces cerevisiae</i> G Protein Indicate a Role in Mating Pheromone Signaling. <i>Molecular and Cellular Biology</i> , 1988, 8, 2484-2493. | 1.1 | 87 |

| # | ARTICLE | IF | CITATIONS |
|------|--|-----|-----------|
| 6982 | Mutational Analysis of Centromere DNA from Chromosome VI of <i>Saccharomyces cerevisiae</i> . Molecular and Cellular Biology, 1988, 8, 2523-2535. | 1.1 | 131 |
| 6983 | Identification of a Ty1 regulatory sequence responsive to STE7 and STE12. Molecular and Cellular Biology, 1988, 8, 2545-2554. | 1.1 | 55 |
| 6984 | <i>ACE1</i> Regulates Expression of the <i>Saccharomyces cerevisiae</i> Metallothionein Gene. Molecular and Cellular Biology, 1988, 8, 2745-2752. | 1.1 | 130 |
| 6985 | <i>TRK1</i> Encodes a Plasma Membrane Protein Required for High-Affinity Potassium Transport in <i>Saccharomyces cerevisiae</i> . Molecular and Cellular Biology, 1988, 8, 2848-2859. | 1.1 | 142 |
| 6986 | Genetic Selection for Mutations That Reduce or Abolish Ribosomal Recognition of the <i>HIS4</i> Translational Initiator Region. Molecular and Cellular Biology, 1988, 8, 2955-2963. | 1.1 | 97 |
| 6987 | Mutational Analysis of the <i>HIS4</i> Translational Initiator Region in <i>Saccharomyces cerevisiae</i> . Molecular and Cellular Biology, 1988, 8, 2964-2975. | 1.1 | 89 |
| 6988 | A Dominant Trifluoperazine Resistance Gene from <i>Saccharomyces cerevisiae</i> Has Homology with F ₀ F ₁ ATP Synthase and Confers Calcium-Sensitive Growth. Molecular and Cellular Biology, 1988, 8, 3094-3103. | 1.1 | 44 |
| 6989 | Genetic Analysis of Small Nuclear RNAs in <i>Saccharomyces cerevisiae</i> : Viable Sextuple Mutant. Molecular and Cellular Biology, 1988, 8, 3150-3159. | 1.1 | 50 |
| 6990 | Characterization of an <i>SNR</i> Gene Locus in <i>Saccharomyces cerevisiae</i> That Specifies Both Dispensible and Essential Small Nuclear RNAs. Molecular and Cellular Biology, 1988, 8, 3282-3290. | 1.1 | 63 |
| 6991 | Transpositional Competence and Transcription of Endogenous Ty Elements in <i>Saccharomyces cerevisiae</i> : Implications for Regulation of Transposition. Molecular and Cellular Biology, 1988, 8, 3571-3581. | 1.1 | 30 |
| 6992 | <i>RAD1</i> , an Excision Repair Gene of <i>Saccharomyces cerevisiae</i> , Is Also Involved in Recombination. Molecular and Cellular Biology, 1988, 8, 3619-3626. | 1.1 | 151 |
| 6993 | Structure and Regulation of a Nuclear Gene in <i>Saccharomyces cerevisiae</i> That Specifies MRP7, a Protein of the Large Subunit of the Mitochondrial Ribosome. Molecular and Cellular Biology, 1988, 8, 3636-3646. | 1.1 | 51 |
| 6994 | Role of STE Genes in the Mating Factor Signaling Pathway Mediated by <i>GPA1</i> in <i>Saccharomyces cerevisiae</i> . Molecular and Cellular Biology, 1988, 8, 3777-3783. | 1.1 | 65 |
| 6995 | Efficient Repair of HO-Induced Chromosomal Breaks in <i>Saccharomyces cerevisiae</i> by Recombination Between Flanking Homologous Sequences. Molecular and Cellular Biology, 1988, 8, 3918-3928. | 1.1 | 133 |
| 6996 | Adjacent upstream activation sequence elements synergistically regulate transcription of ADH2 in <i>Saccharomyces cerevisiae</i> . Molecular and Cellular Biology, 1989, 9, 34-42. | 1.1 | 48 |
| 6997 | Mutations in the Anticodon Stem Affect Removal of Introns from Pre-tRNA in <i>Saccharomyces cerevisiae</i> . Molecular and Cellular Biology, 1989, 9, 4220-4228. | 1.1 | 12 |
| 6998 | Heteroduplex DNA Correction in <i>Saccharomyces cerevisiae</i> Is Mismatch Specific and Requires Functional <i>PMS</i> Genes. Molecular and Cellular Biology, 1989, 9, 4432-4440. | 1.1 | 119 |
| 6999 | Regulation of $\hat{1}\pm$ -Factor Production in <i>Saccharomyces cerevisiae</i> : $\hat{1}\pm$ -Factor Pheromone-Induced Expression of the <i>MF$\hat{1}\pm$</i> and <i>STE13</i> Genes. Molecular and Cellular Biology, 1989, 9, 4507-4514. | 1.1 | 25 |

| # | ARTICLE | IF | CITATIONS |
|------|---|-----|-----------|
| 7000 | The <i>HML</i> Mating-Type Cassette of <i>Saccharomyces cerevisiae</i> Is Regulated by Two Separate but Functionally Equivalent Silencers. <i>Molecular and Cellular Biology</i> , 1989, 9, 4621-4630. | 1.1 | 93 |
| 7001 | Isolation of Constitutive Mutations Affecting the Proline Utilization Pathway in <i>Saccharomyces cerevisiae</i> and Molecular Analysis of the <i>PUT3</i> Transcriptional Activator. <i>Molecular and Cellular Biology</i> , 1989, 9, 4696-4705. | 1.1 | 25 |
| 7002 | The <i>Saccharomyces cerevisiae</i> <i>PUT3</i> Activator Protein Associates with Proline-Specific Upstream Activation Sequences. <i>Molecular and Cellular Biology</i> , 1989, 9, 4706-4712. | 1.1 | 51 |
| 7003 | Basal-Level Expression of the Yeast <i>HSP82</i> Gene Requires a Heat Shock Regulatory Element. <i>Molecular and Cellular Biology</i> , 1989, 9, 4789-4798. | 1.1 | 41 |
| 7004 | Enhancer and Silencerlike Sites within the Transcribed Portion of a Ty2 Transposable Element of <i>Saccharomyces cerevisiae</i> . <i>Molecular and Cellular Biology</i> , 1989, 9, 4824-4834. | 1.1 | 37 |
| 7005 | DNA Damage Induction of Ribonucleotide Reductase. <i>Molecular and Cellular Biology</i> , 1989, 9, 4932-4940. | 1.1 | 71 |
| 7006 | Unusual Enhancer Function in Yeast rRNA Transcription. <i>Molecular and Cellular Biology</i> , 1989, 9, 4986-4993. | 1.1 | 33 |
| 7007 | Mutational Analysis of the <i>Saccharomyces cerevisiae</i> <i>SNF1</i> Protein Kinase and Evidence for Functional Interaction with the <i>SNF4</i> Protein. <i>Molecular and Cellular Biology</i> , 1989, 9, 5034-5044. | 1.1 | 137 |
| 7008 | Molecular Analysis of the <i>SNF4</i> Gene of <i>Saccharomyces cerevisiae</i> : Evidence for Physical Association of the <i>SNF4</i> Protein with the <i>SNF1</i> Protein Kinase. <i>Molecular and Cellular Biology</i> , 1989, 9, 5045-5054. | 1.1 | 120 |
| 7009 | Identification of the DNA Damage-Responsive Element of <i>RNR2</i> and Evidence that Four Distinct Cellular Factors Bind It. <i>Molecular and Cellular Biology</i> , 1989, 9, 5373-5386. | 1.1 | 43 |
| 7010 | Interaction of the H4 Autonomously Replicating Sequence Core Consensus Sequence and Its 3'-Flanking Domain. <i>Molecular and Cellular Biology</i> , 1989, 9, 5464-5472. | 1.1 | 28 |
| 7011 | The <i>Saccharomyces cerevisiae</i> <i>SPT13/GAL11</i> Gene Has Both Positive and Negative Regulatory Roles in Transcription. <i>Molecular and Cellular Biology</i> , 1989, 9, 5602-5609. | 1.1 | 97 |
| 7012 | The Residual Enzymatic Phosphorylation Activity of Hexokinase II Mutants Is Correlated with Glucose Repression in <i>Saccharomyces cerevisiae</i> . <i>Molecular and Cellular Biology</i> , 1989, 9, 5643-5649. | 1.1 | 58 |
| 7013 | Context Affects Nuclear Protein Localization in <i>Saccharomyces cerevisiae</i> . <i>Molecular and Cellular Biology</i> , 1989, 9, 384-389. | 1.1 | 75 |
| 7014 | Selection and Characterization of Ricin Toxin A-Chain Mutations in <i>Saccharomyces cerevisiae</i> . <i>Molecular and Cellular Biology</i> , 1989, 9, 415-420. | 1.1 | 29 |
| 7015 | A Cysteine-Rich Nuclear Protein Activates Yeast Metallothionein Gene Transcription. <i>Molecular and Cellular Biology</i> , 1989, 9, 421-429. | 1.1 | 69 |
| 7016 | Identification of an Upstream Activating Sequence and an Upstream Repressible Sequence of the Pyruvate Kinase Gene of the Yeast <i>Saccharomyces cerevisiae</i> . <i>Molecular and Cellular Biology</i> , 1989, 9, 442-451. | 1.1 | 69 |
| 7017 | Control of the <i>Saccharomyces cerevisiae</i> Regulatory Gene <i>PET494</i> : Transcriptional Repression by Glucose and Translational Induction by Oxygen. <i>Molecular and Cellular Biology</i> , 1989, 9, 484-491. | 1.1 | 42 |

| # | ARTICLE | IF | CITATIONS |
|------|---|-----|-----------|
| 7018 | <i>IRA1</i> , an Inhibitory Regulator of the <i>RAS</i> -Cyclic AMP Pathway in <i>Saccharomyces cerevisiae</i> . <i>Molecular and Cellular Biology</i> , 1989, 9, 757-768. | 1.1 | 145 |
| 7019 | A yeast mutation that stabilizes a plasmid bearing a mutated <i>ARS1</i> element. <i>Molecular and Cellular Biology</i> , 1989, 9, 809-816. | 1.1 | 33 |
| 7020 | Sequence and Structural Requirements of a Mitochondrial Protein Import Signal Defined by Saturation Cassette Mutagenesis. <i>Molecular and Cellular Biology</i> , 1989, 9, 1014-1025. | 1.1 | 27 |
| 7021 | Dominant Effects of Tubulin Overexpression in <i>Saccharomyces cerevisiae</i> . <i>Molecular and Cellular Biology</i> , 1989, 9, 1049-1059. | 1.1 | 115 |
| 7022 | Introduction of Extra Telomeric DNA Sequences into <i>Saccharomyces cerevisiae</i> Results in Telomere Elongation. <i>Molecular and Cellular Biology</i> , 1989, 9, 1488-1497. | 1.1 | 74 |
| 7023 | Introduction of Functional Artificial Introns into the Naturally Intronless <i>ura4</i> Gene of <i>Schizosaccharomyces pombe</i> . <i>Molecular and Cellular Biology</i> , 1989, 9, 1526-1535. | 1.1 | 24 |
| 7024 | Amino-Terminal Extension Generated from an Upstream AUG Codon Increases the Efficiency of Mitochondrial Import of Yeast <i>N²,N²</i> -Dimethylguanosine-Specific tRNA Methyltransferases. <i>Molecular and Cellular Biology</i> , 1989, 9, 1611-1620. | 1.1 | 30 |
| 7025 | Molecular Analysis of <i>GPH1</i> , the Gene Encoding Glycogen Phosphorylase in <i>Saccharomyces cerevisiae</i> . <i>Molecular and Cellular Biology</i> , 1989, 9, 1659-1666. | 1.1 | 43 |
| 7026 | Upstream Activation Sequence-Dependent Alteration of Chromatin Structure and Transcription Activation of the Yeast <i>GAL1-GAL10</i> Genes. <i>Molecular and Cellular Biology</i> , 1989, 9, 1721-1732. | 1.1 | 57 |
| 7027 | Cloning and Sequence Analysis of the <i>Saccharomyces cerevisiae RAD9</i> Gene and Further Evidence that Its Product Is Required for Cell Cycle Arrest Induced by DNA Damage. <i>Molecular and Cellular Biology</i> , 1989, 9, 1882-1896. | 1.1 | 77 |
| 7028 | Positive and Negative Regulation of Basal Expression of a Yeast HSP70 Gene. <i>Molecular and Cellular Biology</i> , 1989, 9, 2025-2033. | 1.1 | 52 |
| 7029 | The <i>Saccharomyces cerevisiae CKS1</i> Gene, a Homolog of the <i>Schizosaccharomyces pombe suc1⁺</i> Gene, Encodes a Subunit of the Cdc28 Protein Kinase Complex. <i>Molecular and Cellular Biology</i> , 1989, 9, 2034-2041. | 1.1 | 107 |
| 7030 | A Transcriptional Cascade Governs Entry into Meiosis in <i>Saccharomyces cerevisiae</i> . <i>Molecular and Cellular Biology</i> , 1989, 9, 2142-2152. | 1.1 | 130 |
| 7031 | Notes: <i>Kluyveromyces lactis</i> Maintains <i>Saccharomyces cerevisiae</i> Intron-Encoded Splicing Signals. <i>Molecular and Cellular Biology</i> , 1989, 9, 2208-2213. | 1.1 | 27 |
| 7032 | GPA1Val-50 mutation in the mating-factor signaling pathway in <i>Saccharomyces cerevisiae</i> . <i>Molecular and Cellular Biology</i> , 1989, 9, 2289-2297. | 1.1 | 40 |
| 7033 | Localization of a Minimal Binding Domain and Activation Regions in Yeast Regulatory Protein ADR1. <i>Molecular and Cellular Biology</i> , 1989, 9, 2360-2369. | 1.1 | 34 |
| 7034 | Cell Size Modulation by <i>CDC25</i> and <i>RAS2</i> Genes in <i>Saccharomyces cerevisiae</i> . <i>Molecular and Cellular Biology</i> , 1989, 9, 2715-2723. | 1.1 | 29 |
| 7035 | Interaction between Transcriptional Activator Protein LAC9 and Negative Regulatory Protein GAL80. <i>Molecular and Cellular Biology</i> , 1989, 9, 2950-2956. | 1.1 | 25 |

| # | ARTICLE | IF | CITATIONS |
|------|--|------|-----------|
| 7036 | Structural and Functional Analyses of <i>Saccharomyces cerevisiae</i> Wild-Type and Mutant <i>RNA1</i> Genes. <i>Molecular and Cellular Biology</i> , 1989, 9, 2989-2999. | 1.1 | 45 |
| 7037 | Functional Domains of a Negative Regulatory Protein, GAL80, of <i>Saccharomyces cerevisiae</i> . <i>Molecular and Cellular Biology</i> , 1989, 9, 3009-3017. | 1.1 | 32 |
| 7038 | Genomic Footprinting of a Yeast tRNA Gene Reveals Stable Complexes over the 5' Flanking Region. <i>Molecular and Cellular Biology</i> , 1989, 9, 3244-3252. | 1.1 | 37 |
| 7039 | Elements Involved in S-Adenosylmethionine-Mediated Regulation of the <i>Saccharomyces cerevisiae</i> MET25 Gene. <i>Molecular and Cellular Biology</i> , 1989, 9, 3292-3298. | 1.1 | 90 |
| 7040 | Failure To Induce a DNA Repair Gene, <i>RAD54</i> , in <i>Saccharomyces cerevisiae</i> Does Not Affect DNA Repair or Recombination Phenotypes. <i>Molecular and Cellular Biology</i> , 1989, 9, 3314-3322. | 1.1 | 43 |
| 7041 | The Yeast Gene <i>ERG6</i> Is Required for Normal Membrane Function but Is Not Essential for Biosynthesis of the Cell-Cycle-Sparking Sterol. <i>Molecular and Cellular Biology</i> , 1989, 9, 3447-3456. | 1.1 | 125 |
| 7042 | Isolation and Characterization of <i>STI1</i> , a Stress-Inducible Gene from <i>Saccharomyces cerevisiae</i> . <i>Molecular and Cellular Biology</i> , 1989, 9, 3638-3646. | 1.1 | 100 |
| 7043 | <i>PRP4</i> (<i>RNA4</i>) from <i>Saccharomyces cerevisiae</i> : Its Gene Product Is Associated with the U4/U6 Small Nuclear Ribonucleoprotein Particle. <i>Molecular and Cellular Biology</i> , 1989, 9, 3698-3709. | 1.1 | 78 |
| 7044 | <i>hsp82</i> Is an Essential Protein That Is Required in Higher Concentrations for Growth of Cells at Higher Temperatures. <i>Molecular and Cellular Biology</i> , 1989, 9, 3919-3930. | 1.1 | 335 |
| 7045 | Regulation of <i>STA1</i> Gene Expression by <i>MAT</i> during the Life Cycle of <i>Saccharomyces cerevisiae</i> . <i>Molecular and Cellular Biology</i> , 1989, 9, 3992-3998. | 1.1 | 23 |
| 7046 | Transformation in fungi. <i>Microbiological Reviews</i> , 1989, 53, 148-170. | 10.1 | 433 |
| 7047 | Genetics of <i>Candida albicans</i> . <i>Microbiological Reviews</i> , 1990, 54, 226-241. | 10.1 | 213 |
| 7048 | Construction of engineered yeast producing ammonia from glutamine and soybean residues (okara). <i>AMB Express</i> , 2020, 10, 70. | 1.4 | 10 |
| 7049 | A <i>Drosophila</i> G1-specific cyclin E homolog exhibits different modes of expression during embryogenesis. <i>Development (Cambridge)</i> , 1993, 119, 673-690. | 1.2 | 128 |
| 7050 | The Dorsal-related immunity factor (Dif) can define the dorsal-ventral axis of polarity in the <i>Drosophila</i> embryo. <i>Development (Cambridge)</i> , 1998, 125, 2159-2169. | 1.2 | 33 |
| 7051 | Modular long-range regulation of <i>Myf5</i> reveals unexpected heterogeneity between skeletal muscles in the mouse embryo. <i>Development (Cambridge)</i> , 2000, 127, 4455-4467. | 1.2 | 102 |
| 7052 | The activity of the <i>Drosophila</i> morphogenetic protein Bicoid is inhibited by a domain located outside its homeodomain. <i>Development (Cambridge)</i> , 2002, 129, 1669-1680. | 1.2 | 38 |
| 7053 | Screening and identification of a gene, <i>PSE-1</i> , that affects protein secretion in <i>Saccharomyces cerevisiae</i> . <i>Journal of Cell Science</i> , 1992, 101, 709-719. | 1.2 | 38 |

| # | ARTICLE | IF | CITATIONS |
|------|--|-----|-----------|
| 7054 | Effects of the expression of mammalian annexins in yeast secretory mutants. <i>Journal of Cell Science</i> , 1992, 103, 1177-1193. | 1.2 | 28 |
| 7055 | The lethality of p60v-src in <i>Saccharomyces cerevisiae</i> and the activation of p34CDC28 kinase are dependent on the integrity of the SH2 domain. <i>Journal of Cell Science</i> , 1993, 105, 519-528. | 1.2 | 19 |
| 7056 | Prp2o, the <i>Saccharomyces cerevisiae</i> homolog of the regulator of chromosome condensation, rcc1, interacts with double-stranded dna through a multi-component complex containing gtp-binding proteins. <i>Journal of Cell Science</i> , 1993, 106, 287-298. | 1.2 | 29 |
| 7057 | Functional consequences in yeast of single-residue alterations in a consensus calmodulin. <i>Journal of Cell Science</i> , 1994, 107, 3235-3249. | 1.2 | 8 |
| 7058 | Mutations in the <i>VPS45</i> gene, a <i>SEC1</i> homologue, result in vacuolar protein sorting defects and accumulation of membrane vesicles. <i>Journal of Cell Science</i> , 1994, 107, 3449-3459. | 1.2 | 134 |
| 7059 | A chromosome breakage assay to monitor mitotic forces in budding yeast. <i>Journal of Cell Science</i> , 1994, 107, 891-902. | 1.2 | 40 |
| 7060 | Comparison of human CAP and CAP2, homologs of the yeast adenylyl cyclase-associated proteins. <i>Journal of Cell Science</i> , 1994, 107, 1671-1678. | 1.2 | 63 |
| 7061 | A calcineurin-like gene <i>ppb1+</i> in fission yeast: mutant defects in cytokinesis, cell polarity, mating and spindle pole body positioning. <i>Journal of Cell Science</i> , 1994, 107, 1725-1735. | 1.2 | 152 |
| 7062 | Yeast homolog of mammalian mitogen-activated protein kinase, FUS3/DAC2 kinase, is required both for cell fusion and for G1 arrest of the cell cycle and morphological changes by the <i>CDC37</i> mutation. <i>Journal of Cell Science</i> , 1994, 107, 2617-2622. | 1.2 | 15 |
| 7063 | A novel essential fission yeast gene <i>Pad1+</i> positively regulates <i>Pap1+</i> -dependent transcription and is implicated in the maintenance of chromosome structure. <i>Journal of Cell Science</i> , 1995, 108, 569-579. | 1.2 | 73 |
| 7064 | Suppression of a conditional mutation in alpha-tubulin by overexpression of two checkpoint genes. <i>Journal of Cell Science</i> , 1995, 108, 1195-1204. | 1.2 | 12 |
| 7065 | Fission yeast TPR-family protein nuc2 is required for G1-arrest upon nitrogen starvation and is an inhibitor of septum formation. <i>Journal of Cell Science</i> , 1995, 108, 895-905. | 1.2 | 54 |
| 7066 | <i>Saccharomyces cerevisiae</i> Apl2p, a homologue of the mammalian clathrin AP $\hat{2}$ subunit, plays a role in clathrin-dependent Golgi functions. <i>Journal of Cell Science</i> , 1995, 108, 1605-1615. | 1.2 | 64 |
| 7067 | Isolation and characterization of <i>SYS</i> genes from yeast, multicopy suppressors of the functional loss of the transport GTPase Ypt6p. <i>Journal of Cell Science</i> , 1996, 109, 2471-2481. | 1.2 | 80 |
| 7068 | Putative GTPase Gtr1p genetically interacts with the RanGTPase cycle in <i>Saccharomyces cerevisiae</i> . <i>Journal of Cell Science</i> , 1996, 109, 2311-2318. | 1.2 | 36 |
| 7069 | The fission yeast <i>STS5+</i> gene is required for maintenance of growth polarity and functionally interacts with protein kinase C and an osmosensing MAP-kinase pathway. <i>Journal of Cell Science</i> , 1996, 109, 2331-2342. | 1.2 | 40 |
| 7070 | Vam3p, a new member of syntaxin related protein, is required for vacuolar assembly in the yeast <i>Saccharomyces cerevisiae</i> . <i>Journal of Cell Science</i> , 1997, 110, 1299-1306. | 1.2 | 97 |
| 7071 | <i>NNF1</i> is an essential yeast gene required for proper spindle orientation, nucleolar and nuclear envelope structure and mRNA export. <i>Journal of Cell Science</i> , 1997, 110, 1615-1624. | 1.2 | 12 |

| # | ARTICLE | IF | CITATIONS |
|------|--|-----|-----------|
| 7072 | Localisation of the <i>Schizosaccharomyces pombe</i> rho1p GTPase and its involvement in the organisation of the actin cytoskeleton. <i>Journal of Cell Science</i> , 1997, 110, 2547-2555. | 1.2 | 84 |
| 7073 | Cloning of <i>SEC61</i> homologues from <i>Schizosaccharomyces pombe</i> and <i>Yarrowia lipolytica</i> reveals the extent of functional conservation within this core component of the ER translocation machinery. <i>Journal of Cell Science</i> , 1997, 110, 2715-2727. | 1.2 | 30 |
| 7074 | Ye1013p (Vac8p), an armadillo repeat protein related to plakoglobin and importin β , is associated with the yeast vacuole membrane. <i>Journal of Cell Science</i> , 1998, 111, 3109-3118. | 1.2 | 44 |
| 7075 | Synaptojanin family members are implicated in endocytic membrane traffic in yeast. <i>Journal of Cell Science</i> , 1998, 111, 3347-3356. | 1.2 | 101 |
| 7076 | Transcription, biochemistry and localization of nematode annexins. <i>Journal of Cell Science</i> , 1999, 112, 1901-1913. | 1.2 | 18 |
| 7077 | <i>Rvs167p</i> , the budding yeast homolog of amphiphysin, colocalizes with actin patches. <i>Journal of Cell Science</i> , 1999, 112, 2529-2537. | 1.2 | 60 |
| 7078 | Caffeine can override the S-M checkpoint in fission yeast. <i>Journal of Cell Science</i> , 1999, 112, 927-937. | 1.2 | 47 |
| 7079 | NH ₄ ⁺ -induced down-regulation of the <i>Saccharomyces cerevisiae</i> Cap1p permease involves its ubiquitination with lysine-63-linked chains. <i>Journal of Cell Science</i> , 1999, 112, 1375-1383. | 1.2 | 121 |
| 7080 | The <i>SHR3</i> homologue from <i>S. pombe</i> demonstrates a conserved function of ER packaging chaperones. <i>Journal of Cell Science</i> , 2000, 113, 4351-4362. | 1.2 | 18 |
| 7081 | Yeast myosin light chain, <i>Mlc1p</i> , interacts with both IQGAP and Class II myosin to effect cytokinesis. <i>Journal of Cell Science</i> , 2000, 113, 4533-4543. | 1.2 | 78 |
| 7082 | Identification of <i>Rgp1p</i> , a novel Golgi recycling factor, as a protein required for efficient localization of yeast casein kinase 1 to the plasma membrane. <i>Journal of Cell Science</i> , 2000, 113, 4545-4555. | 1.2 | 17 |
| 7083 | Interaction of the endoplasmic reticulum α 1,2-mannosidase <i>Mns1p</i> with <i>Rer1p</i> using the split-ubiquitin system. <i>Journal of Cell Science</i> , 2001, 114, 4629-4635. | 1.2 | 26 |
| 7084 | The yeast ADP-ribosylation factor GAP, <i>Gcs1p</i> , is involved in maintenance of mitochondrial morphology. <i>Journal of Cell Science</i> , 2002, 115, 275-282. | 1.2 | 17 |
| 7085 | <i>Saccharomyces cerevisiae</i> mutants lacking a functional vacuole are defective for aspects of the pheromone response. <i>Journal of Cell Science</i> , 1990, 97, 517-525. | 1.2 | 18 |
| 7086 | Identification of a Ubiquitous Family of Membrane Proteins and Their Expression in Mouse Brain. <i>Journal of Experimental Biology</i> , 2000, 203, 447-457. | 0.8 | 29 |
| 7087 | Optimization of Bovine Cathepsin C Production by Cultivation of Recombinant Methylophilic Yeast <i>Candida boidinii</i> . <i>Journal of Chemical Engineering of Japan</i> , 2001, 34, 848-851. | 0.3 | 4 |
| 7088 | A New Method for Production of Combinatorial Libraries by Mating of <i>Saccharomyces cerevisiae</i> Cells. <i>Journal of Chemical Engineering of Japan</i> , 2002, 35, 474-478. | 0.3 | 1 |
| 7089 | Extremely Simple, Rapid, and Highly Efficient Transformation Method for the Yeast <i>Saccharomyces cerevisiae</i> Using Glutathione and Early Log Phase Cells. <i>Journal of Bioscience and Bioengineering</i> , 2002, 94, 166-171. | 1.1 | 24 |

| # | ARTICLE | IF | CITATIONS |
|------|---|-----|-----------|
| 7090 | Construction of uracil and tryptophan auxotrophic mutants from sake yeasts by disruption of URA3 and TRP1 genes.. <i>Agricultural and Biological Chemistry</i> , 1990, 54, 2979-2987. | 0.3 | 51 |
| 7091 | Protein Interaction Mapping of Translational Regulators Affecting Expression of the Critical Stem Cell Factor Nos. <i>Development & Reproduction</i> , 2017, 21, 449-456. | 0.1 | 5 |
| 7092 | A Tradeoff Drives the Evolution of Reduced Metal Resistance in Natural Populations of Yeast. <i>PLoS Genetics</i> , 2011, 7, e1002034. | 1.5 | 27 |
| 7093 | Optimization of Ribosome Structure and Function by rRNA Base Modification. <i>PLoS ONE</i> , 2007, 2, e174. | 1.1 | 123 |
| 7094 | Differentiating between Near- and Non-Cognate Codons in <i>Saccharomyces cerevisiae</i> . <i>PLoS ONE</i> , 2007, 2, e517. | 1.1 | 45 |
| 7095 | Ybp2 Associates with the Central Kinetochore of <i>Saccharomyces cerevisiae</i> and Mediates Proper Mitotic Progression. <i>PLoS ONE</i> , 2008, 3, e1617. | 1.1 | 13 |
| 7096 | Analysis of SEC9 Suppression Reveals a Relationship of SNARE Function to Cell Physiology. <i>PLoS ONE</i> , 2009, 4, e5449. | 1.1 | 7 |
| 7097 | Dopamine-Induced Conformational Changes in Alpha-Synuclein. <i>PLoS ONE</i> , 2009, 4, e6906. | 1.1 | 59 |
| 7098 | Connecting Mutations of the RNA Polymerase II C-Terminal Domain to Complex Phenotypic Changes Using Combined Gene Expression and Network Analyses. <i>PLoS ONE</i> , 2010, 5, e11386. | 1.1 | 2 |
| 7099 | <i>Saccharomyces cerevisiae</i> Bat1 and Bat2 Aminotransferases Have Functionally Diverged from the Ancestral-Like <i>Kluyveromyces lactis</i> Orthologous Enzyme. <i>PLoS ONE</i> , 2011, 6, e16099. | 1.1 | 62 |
| 7100 | Mis17 Is a Regulatory Module of the Mis6-Mal2-Sim4 Centromere Complex That Is Required for the Recruitment of CenH3/CENP-A in Fission Yeast. <i>PLoS ONE</i> , 2011, 6, e17761. | 1.1 | 18 |
| 7101 | Characterization of <i>Leishmania donovani</i> Aquaporins Shows Presence of Subcellular Aquaporins Similar to Tonoplast Intrinsic Proteins of Plants. <i>PLoS ONE</i> , 2011, 6, e24820. | 1.1 | 16 |
| 7102 | Neutral Lipid Metabolism Influences Phospholipid Synthesis and Deacylation in <i>Saccharomyces cerevisiae</i> . <i>PLoS ONE</i> , 2012, 7, e49269. | 1.1 | 29 |
| 7103 | Role of Ess1 in Growth, Morphogenetic Switching, and RNA Polymerase II Transcription in <i>Candida albicans</i> . <i>PLoS ONE</i> , 2013, 8, e59094. | 1.1 | 4 |
| 7104 | Characterization of Chromosome Stability in Diploid, Polyploid and Hybrid Yeast Cells. <i>PLoS ONE</i> , 2013, 8, e68094. | 1.1 | 71 |
| 7105 | SNPs Altering Ammonium Transport Activity of Human Rhesus Factors Characterized by a Yeast-Based Functional Assay. <i>PLoS ONE</i> , 2013, 8, e71092. | 1.1 | 5 |
| 7106 | Identification and Characterization of the Regions Involved in the Nuclear Translocation of the Heterodimeric Leishmanial DNA Topoisomerase IB. <i>PLoS ONE</i> , 2013, 8, e73565. | 1.1 | 10 |
| 7107 | Trans-Kingdom Horizontal DNA Transfer from Bacteria to Yeast Is Highly Plastic Due to Natural Polymorphisms in Auxiliary Nonessential Recipient Genes. <i>PLoS ONE</i> , 2013, 8, e74590. | 1.1 | 19 |

| # | ARTICLE | IF | CITATIONS |
|------|---|-----|-----------|
| 7108 | The N-Terminus of Vps74p Is Essential for the Retention of Glycosyltransferases in the Golgi but Not for the Modulation of Apical Polarized Growth in <i>Saccharomyces cerevisiae</i> . <i>PLoS ONE</i> , 2013, 8, e74715. | 1.1 | 5 |
| 7109 | VelC Positively Controls Sexual Development in <i>Aspergillus nidulans</i> . <i>PLoS ONE</i> , 2014, 9, e89883. | 1.1 | 69 |
| 7110 | Phs1 and the Synthesis of Very Long Chain Fatty Acids Are Required for Ballistospore Formation. <i>PLoS ONE</i> , 2014, 9, e105147. | 1.1 | 9 |
| 7111 | The General Amino Acid Permease FfGap1 of <i>Fusarium fujikuroi</i> Is Sorted to the Vacuole in a Nitrogen-Dependent, but Npr1 Kinase-Independent Manner. <i>PLoS ONE</i> , 2015, 10, e0125487. | 1.1 | 9 |
| 7112 | A Fast and Practical Yeast Transformation Method Mediated by <i>Escherichia coli</i> Based on a Trans-Kingdom Conjugal Transfer System: Just Mix Two Cultures and Wait One Hour. <i>PLoS ONE</i> , 2016, 11, e0148989. | 1.1 | 11 |
| 7113 | A Trypanosomatid Iron Transporter that Regulates Mitochondrial Function Is Required for <i>Leishmania amazonensis</i> Virulence. <i>PLoS Pathogens</i> , 2016, 12, e1005340. | 2.1 | 38 |
| 7114 | The morphogenesis-related NDR kinase pathway of <i>Colletotrichum orbiculare</i> is required for translating plant surface signals into infection-related morphogenesis and pathogenesis. <i>PLoS Pathogens</i> , 2017, 13, e1006189. | 2.1 | 19 |
| 7115 | Amino acid substitution reveals the role of V-shape helix on construction of yeast carboxypeptidase Y. <i>Journal of Biological Macromolecules</i> , 2014, 14, 27-41. | 0.2 | 1 |
| 7116 | VDAC regulates AAC-mediated apoptosis and cytochrome c release in yeast. <i>Microbial Cell</i> , 2016, 3, 500-510. | 1.4 | 20 |
| 7117 | Probing iso-1-cytochrome c structure by site-directed spin labeling and electron paramagnetic resonance techniques.. <i>Acta Biochimica Polonica</i> , 1999, 46, 889-899. | 0.3 | 2 |
| 7118 | Prohibitins and Ras2 protein cooperate in the maintenance of mitochondrial function during yeast aging.. <i>Acta Biochimica Polonica</i> , 2003, 50, 1039-1056. | 0.3 | 26 |
| 7119 | Improving transformation method for industrial yeasts: Construction of ADH1-APT2 gene and using electroporation.. <i>Journal of General and Applied Microbiology</i> , 1993, 39, 285-294. | 0.4 | 13 |
| 7120 | A promoter activity in <i>Saccharomyces cerevisiae</i> of the 3'-noncoding region of the basidiomycetous mushroom gene.. <i>Journal of General and Applied Microbiology</i> , 2002, 48, 223-231. | 0.4 | 6 |
| 7121 | Targeting of insect epicuticular lipids by the entomopathogenic fungus <i>Beauveria bassiana</i> : hydrocarbon oxidation within the context of a host-pathogen interaction. <i>Frontiers in Microbiology</i> , 2013, 4, 24. | 1.5 | 137 |
| 7122 | Enhanced Production of 1,2-Propanediol by <i>tpi1</i> Deletion in <i>Saccharomyces cerevisiae</i> . <i>Journal of Microbiology and Biotechnology</i> , 2008, 18, 1797-1802. | 0.9 | 25 |
| 7123 | Expression of Functional Pentameric Heat-Labile Enterotoxin B Subunit of <i>Escherichia coli</i> in <i>Saccharomyces cerevisiae</i> . <i>Journal of Microbiology and Biotechnology</i> , 2009, 19, 502-510. | 0.9 | 17 |
| 7124 | Production of 1,2-Propanediol from Glycerol in <i>Saccharomyces cerevisiae</i> . <i>Journal of Microbiology and Biotechnology</i> , 2011, 21, 846-853. | 0.9 | 55 |
| 7125 | A versatile vector system for generating recombinant EGFP-tagged proteins in yeast. <i>Advances in Bioscience and Biotechnology (Print)</i> , 2011, 02, 13-19. | 0.3 | 2 |

| # | ARTICLE | IF | CITATIONS |
|------|--|-----|-----------|
| 7126 | Identification and analysis of dihydrofolate reductase alleles from <i>Plasmodium falciparum</i> present at low frequency in polyclonal patient samples.. <i>American Journal of Tropical Medicine and Hygiene</i> , 1999, 61, 131-140. | 0.6 | 16 |
| 7127 | POINT MUTATIONS IN THE DIHYDROFOLATE REDUCTASE AND DIHYDROPTEROATE SYNTHASE GENES OF <i>PLASMODIUM FALCIPARUM</i> AND RESISTANCE TO SULFADOXINE-PYRIMETHAMINE IN SRI LANKA. <i>American Journal of Tropical Medicine and Hygiene</i> , 2006, 74, 198-204. | 0.6 | 12 |
| 7128 | The Shaker Type Potassium Channel, GORK, Regulates Abscisic Acid Signaling in <i>Arabidopsis</i> . <i>Plant Pathology Journal</i> , 2019, 35, 684-691. | 0.7 | 6 |
| 7129 | The Ty1 Retrotransposition Short-Term Test for Selective Detection of Carcinogenic Genotoxins. , 0, , . | | 3 |
| 7130 | <i>Morus</i> species through centuries in pharmacy and as food. <i>Savremene Tehnologije</i> , 2014, 3, 111-115. | 0.0 | 6 |
| 7131 | Introduction and Expression of cDNA Encoding <i>Aspergillus oryzae</i> α -Amylase to Sake Yeast. <i>Journal of the Brewing Society of Japan</i> , 1993, 88, 233-237. | 0.1 | 1 |
| 7132 | Analysis of 3-^{H} -Phosphoadenosine 5-^{H} -Phosphosulfate Transporters: Transporter Activity Assay, Real-Time Reverse Transcription Polymerase Chain Reaction, and. <i>Methods in Molecular Biology</i> , 2022, 2303, 675-685. | 0.4 | 0 |
| 7133 | Derlin rhomboid pseudoproteases employ substrate engagement and lipid distortion to enable the retrotranslocation of ERAD membrane substrates. <i>Cell Reports</i> , 2021, 37, 109840. | 2.9 | 22 |
| 7134 | The stress induced caleosin, RD20/CLO3, acts as a negative regulator of GPA1 in <i>Arabidopsis</i> . <i>Plant Molecular Biology</i> , 2021, 107, 159-175. | 2.0 | 7 |
| 7135 | An NH ₂ -terminal deleted plasma membrane H ⁺ -ATPase is a dominant negative mutant and is sequestered in endoplasmic reticulum derived structures. <i>Biochemistry and Cell Biology</i> , 2000, 78, 51-58. | 0.9 | 6 |
| 7136 | Nitrogen Regulation of <i>Saccharomyces cerevisiae</i> Invertase. , 2000, , 247-254. | | 2 |
| 7137 | Evidence That Single Missense Mutations in the <i>Atp7a</i> Gene of Two Mouse Models for Menkes' Disease Are Responsible for Impaired Copper Transport.. <i>Journal of Clinical Biochemistry and Nutrition</i> , 2000, 29, 37-44. | 0.6 | 0 |
| 7139 | High-Level Production of Prourokinase-Annexin V Chimeras in the Methylophilic Yeast <i>Pichia pastoris</i> .. <i>Journal of Bioscience and Bioengineering</i> , 2002, 94, 467-473. | 1.1 | 1 |
| 7140 | Transformation of <i>Kluyveromyces lactis</i> . , 2003, , 169-174. | | 1 |
| 7143 | Expression and Secretion of <i>Zymomonas mobilis</i> Levansucrase in <i>Saccharomyces cerevisiae</i> .. <i>Journal of Life Science</i> , 2004, 14, 429-434. | 0.2 | 2 |
| 7144 | PKR1 Encodes an Assembly Factor for the Yeast V-Type ATPase. <i>Journal of Biological Chemistry</i> , 2006, 281, 32025-32035. | 1.6 | 4 |
| 7145 | Yeast and Mammalian Two-Hybrid Systems for Studying Protein-Protein Interactions. , 2007, 383, 215-225. | | 2 |
| 7146 | Production of Recombinant Vaccine Cb Peritrophin-42 of Screwworm Fly in <i>Escherichia coli</i> and <i>Saccharomyces cerevisiae</i> . <i>Microbiology Indonesia</i> , 2007, 1, 105-108. | 0.2 | 0 |

| # | ARTICLE | IF | CITATIONS |
|------|---|-----|-----------|
| 7148 | The b $\hat{\epsilon}$ TM x Region of Yeast Protein Disulfide Isomerase is Not Essential for <i>Saccharomyces cerevisiae</i> Viability at 30 $\hat{\text{A}}$ ^o C. <i>Microbiology Indonesia</i> , 2009, 3, 27-32. | 0.2 | 0 |
| 7149 | Regulation of meiotic recombination by a multifunctional ATF/CREB protein. <i>FASEB Journal</i> , 2009, 23, 656.1. | 0.2 | 2 |
| 7150 | Expression of A Gene Encoding A Functional Glycosyl Hydrolase, Trehalase, from <i>Nicotiana Tabacum</i> in <i>Saccharomyces Cerevisiae</i> . <i>Journal of the Faculty of Agriculture, Kyushu University</i> , 2009, 54, 297-303. | 0.1 | 1 |
| 7151 | Multisite Phosphorylation of the Guanine Nucleotide Exchange Factor Cdc24 during Yeast Cell Polarization. , 2014, , 79-106. | | 0 |
| 7152 | Mitochondrial import of cytochrome c oxidase subunit VIIa in <i>Saccharomyces cerevisiae</i> . Identification of sequences required for mitochondrial localization in vivo.. <i>Journal of Biological Chemistry</i> , 1990, 265, 7273-7277. | 1.6 | 6 |
| 7153 | <i>Saccharomyces cerevisiae</i> as a Paradigm for Modern Molecular Genetics of Fungi. , 1985, , 125-160. | | 0 |
| 7154 | Expression of the Gene for Ornithine Decarboxylase of <i>Saccharomyces cerevisiae</i> in <i>Escherichia coli</i> . <i>Molecular and Cellular Biology</i> , 1985, 5, 161-166. | 1.1 | 19 |
| 7155 | Molecular Cloning and Physiological Analysis of the Start Gene <i>cdc25</i> in Budding Yeast. , 1986, , 29-41. | | 2 |
| 7156 | Structure and sequence of the centromeric DNA of chromosome 4 in <i>Saccharomyces cerevisiae</i> . <i>Molecular and Cellular Biology</i> , 1986, 6, 241-245. | 1.1 | 54 |
| 7157 | Duplicate Upstream Activating Sequences in the Promoter Region of the <i>Saccharomyces cerevisiae</i> <i>GAL7</i> Gene. <i>Molecular and Cellular Biology</i> , 1986, 6, 246-256. | 1.1 | 54 |
| 7158 | Tightly Centromere-Linked Gene (<i>SPO15</i>) Essential for Meiosis in the Yeast <i>Saccharomyces cerevisiae</i> . <i>Molecular and Cellular Biology</i> , 1986, 6, 158-167. | 1.1 | 25 |
| 7159 | In Vivo Genetic Manipulation. , 1987, , 3-53. | | 2 |
| 7162 | The Stability of Chromosomes in Yeast. , 1987, , 1-1. | | 6 |
| 7164 | Molecular Cloning of Chromosome I DNA from <i>Saccharomyces cerevisiae</i> : Isolation and Analysis OF THE <i>CEN1-ADE1-CDC15</i> Region. <i>Molecular and Cellular Biology</i> , 1987, 7, 410-419. | 1.1 | 28 |
| 7165 | An Adenylate Cyclase from <i>Saccharomyces cerevisiae</i> That Is Stimulated by <i>RAS</i> Proteins with Effector Mutations. <i>Molecular and Cellular Biology</i> , 1988, 8, 52-61. | 1.1 | 38 |
| 7166 | Recombinant DNA Research in Phytopathogenic Fungi. <i>Advances in Plant Pathology</i> , 1988, 6, 1-26. | 0.3 | 5 |
| 7167 | Genetic Analysis and Transformation of Distiller's Yeast. <i>Modern Methods of Plant Analysis</i> , 1988, , 145-163. | 0.1 | 1 |
| 7169 | Plasmid Vectors for the Analysis of Regulatory Sequences in Yeast. , 1988, 10, 405-417. | | 0 |

| # | ARTICLE | IF | CITATIONS |
|------|---|-----|-----------|
| 7170 | The General Control Activator Protein GCN4 Is Essential for a Basal Level of <i>ARO3</i> Gene Expression in <i>Saccharomyces cerevisiae</i> . <i>Molecular and Cellular Biology</i> , 1989, 9, 144-151. | 1.1 | 20 |
| 7171 | Transformation of plant pathogenic fungi. , 1989, , 195-207. | | 0 |
| 7172 | A Synthetic Intron in a Naturally Intronless Yeast Pre-tRNA Is Spliced Efficiently In Vivo. <i>Molecular and Cellular Biology</i> , 1989, 9, 329-331. | 1.1 | 5 |
| 7173 | Mutations at the RAS2 Locus that, in a <i>ras1</i> - Background, Impair the Growth of Yeast on Nonfermentable Carbon Sources. , 1989, , 251-256. | | 0 |
| 7174 | RNA Isolation. , 1989, , 94-97. | | 0 |
| 7175 | DNA-Mediated Transformation of Phytopathogenic Fungi. , 1989, , 127-143. | | 0 |
| 7176 | The Yeast G-Protein Homolog Is Involved in the Mating Pheromone Signal Transduction System. <i>Molecular and Cellular Biology</i> , 1989, 9, 152-158. | 1.1 | 16 |
| 7177 | Transformation of <i>Saccharomyces cerevisiae</i> . , 1990, , 103-107. | | 0 |
| 7178 | Mouse Zfx Protein Is Similar to Zfy-2: Each Contains an Acidic Activating Domain and 13 Zinc Fingers. <i>Molecular and Cellular Biology</i> , 1990, 10, 681-688. | 1.1 | 26 |
| 7179 | Genetic and Biochemical Evaluation of Eucaryotic Membrane Protein Topology: Multiple Transmembrane Domains of <i>Saccharomyces cerevisiae</i> 3-Hydroxy-3-Methylglutaryl Coenzyme A Reductase. <i>Molecular and Cellular Biology</i> , 1990, 10, 672-680. | 1.1 | 38 |
| 7180 | Enhancer and Promoter Elements from Simian Virus 40 and Polyomavirus Can Substitute for an Upstream Activation Sequence in <i>Saccharomyces cerevisiae</i> . <i>Molecular and Cellular Biology</i> , 1990, 10, 947-957. | 1.1 | 4 |
| 7181 | Association of Catalytic and Regulatory Subunits of Cyclic AMP-Dependent Protein Kinase Requires a Negatively Charged Side Group at a Conserved Threonine. <i>Molecular and Cellular Biology</i> , 1990, 10, 1066-1075. | 1.1 | 28 |
| 7182 | Functional analysis of a centromere from fission yeast: a role for centromere-specific repeated DNA sequences. <i>Molecular and Cellular Biology</i> , 1990, 10, 1863-1872. | 1.1 | 66 |
| 7183 | A Gene Family for Acidic Ribosomal Proteins in <i>Schizosaccharomyces pombe</i> : Two Essential and Two Nonessential Genes. <i>Molecular and Cellular Biology</i> , 1990, 10, 2341-2348. | 1.1 | 20 |
| 7184 | Null Alleles of <i>SAC7</i> Suppress Temperature-Sensitive Actin Mutations in <i>Saccharomyces cerevisiae</i> . <i>Molecular and Cellular Biology</i> , 1990, 10, 2308-2314. | 1.1 | 32 |
| 7185 | The <i>SLP1</i> Gene of <i>Saccharomyces cerevisiae</i> Is Essential for Vacuolar Morphogenesis and Function. <i>Molecular and Cellular Biology</i> , 1990, 10, 2214-2223. | 1.1 | 55 |
| 7186 | BET1, BOS1, and SEC22 Are Members of a Group of Interacting Yeast Genes Required for Transport from the Endoplasmic Reticulum to the Golgi Complex. <i>Molecular and Cellular Biology</i> , 1990, 10, 3405-3414. | 1.1 | 116 |
| 7187 | Modification and Transfer into an Embryonal Carcinoma Cell Line of a 360-Kilobase Human-Derived Yeast Artificial Chromosome. <i>Molecular and Cellular Biology</i> , 1990, 10, 4163-4169. | 1.1 | 36 |

| # | ARTICLE | IF | CITATIONS |
|------|---|-----|-----------|
| 7188 | Sequences within an Upstream Activation Site in the Yeast Enolase Gene <i>ENO2</i> Modulate Repression of <i>ENO2</i> Expression in Strains Carrying a Null Mutation in the Positive Regulatory Gene <i>GCR1</i> . <i>Molecular and Cellular Biology</i> , 1990, 10, 4863-4871. | 1.1 | 19 |
| 7189 | The Mouse <i>c-rel</i> Protein Has an N-Terminal Regulatory Domain and a C-Terminal Transcriptional Transactivation Domain. <i>Molecular and Cellular Biology</i> , 1990, 10, 5473-5485. | 1.1 | 105 |
| 7190 | Functional analysis of a duplicated linked pair of ribosomal protein genes in <i>Saccharomyces cerevisiae</i> . <i>Molecular and Cellular Biology</i> , 1990, 10, 6097-6100. | 1.1 | 7 |
| 7191 | In Vivo Expression and Mitochondrial Targeting of Yeast Apoiso-1-Cytochrome <i>c</i> Fusion Proteins. <i>Molecular and Cellular Biology</i> , 1990, 10, 5753-5762. | 1.1 | 5 |
| 7192 | Structural and functional characters of plasmid-like mitochondrial maize DNA. <i>Biopolymers and Cell</i> , 1990, 6, 100-105. | 0.1 | 0 |
| 7193 | Molecular and Expression Analysis of the Negative Regulators Involved in the Transcriptional Regulation of Acid Phosphatase Production in <i>Saccharomyces cerevisiae</i> . <i>Molecular and Cellular Biology</i> , 1990, 10, 5950-5957. | 1.1 | 22 |
| 7194 | Molecular Genetic Analysis of <i>Saccharomyces cerevisiae</i> C ₁ -Tetrahydrofolate Synthase Mutants Reveals a Noncatalytic Function of the <i>ADE3</i> Gene Product and an Additional Folate-Dependent Enzyme. <i>Molecular and Cellular Biology</i> , 1990, 10, 5679-5687. | 1.1 | 17 |
| 7195 | Sequence and Functional Similarity between a Yeast Ribosomal Protein and the <i>Escherichia coli</i> S5 <i>ram</i> Protein. <i>Molecular and Cellular Biology</i> , 1990, 10, 6544-6553. | 1.1 | 32 |
| 7196 | Mutational Analysis of the Functional Domains of Yeast K1 Killer Toxin. <i>Molecular and Cellular Biology</i> , 1991, 11, 175-181. | 1.1 | 17 |
| 7198 | The Yeast <i>rad18</i> Mutator Specifically Increases G \hat{A} -C \hat{A} T \hat{T} A Transversions without Reducing Correction of G-A or C-T Mismatches to G \hat{A} -C Pairs. <i>Molecular and Cellular Biology</i> , 1991, 11, 218-225. | 1.1 | 21 |
| 7199 | The <i>CDC25</i> Protein of <i>Saccharomyces cerevisiae</i> Promotes Exchange of Guanine Nucleotides Bound to Ras. <i>Molecular and Cellular Biology</i> , 1991, 11, 2641-2646. | 1.1 | 82 |
| 7200 | Genetic Evidence that Different Functional Domains of the <i>PET54</i> Gene Product Facilitate Expression of the Mitochondrial Genes <i>COX1</i> and <i>COX3</i> in <i>Saccharomyces cerevisiae</i> . <i>Molecular and Cellular Biology</i> , 1991, 11, 2399-2405. | 1.1 | 17 |
| 7201 | Viability of Clathrin Heavy-Chain-Deficient <i>Saccharomyces cerevisiae</i> Is Compromised by Mutations at Numerous Loci: Implications for the Suppression Hypothesis. <i>Molecular and Cellular Biology</i> , 1991, 11, 3868-3878. | 1.1 | 33 |
| 7202 | Transformation of the yeast <i>Saccharomyces kluyveri</i> by <i>Saccharomyces cerevisiae</i> -based plasmids. <i>FEMS Microbiology Letters</i> , 1991, 82, 149-152. | 0.7 | 7 |
| 7203 | Cloning and Characterization of a Gene Which Determines Osmotic Stability in <i>Saccharomyces cerevisiae</i> . <i>Molecular and Cellular Biology</i> , 1991, 11, 4235-4243. | 1.1 | 16 |
| 7204 | Assessment of the Transcriptional Activation Potential of the HMG Chromosomal Proteins. <i>Molecular and Cellular Biology</i> , 1991, 11, 4483-4489. | 1.1 | 32 |
| 7205 | <i>AAR2</i> , a Gene for Splicing Pre-mRNA of the <i>MATa1</i> Cistron in Cell Type Control of <i>Saccharomyces cerevisiae</i> . <i>Molecular and Cellular Biology</i> , 1991, 11, 5693-5700. | 1.1 | 17 |
| 7206 | Protocol for High-Efficiency Yeast Transformation. , 1992, , 501-505. | | 1 |

| # | ARTICLE | IF | CITATIONS |
|------|--|-----|-----------|
| 7207 | Connections between RNA Splicing and DNA Intron Mobility in Yeast Mitochondria: RNA Maturase and DNA Endonuclease Switching Experiments. <i>Molecular and Cellular Biology</i> , 1992, 12, 696-705. | 1.1 | 21 |
| 7208 | Regulation of Yeast COX6 by the General Transcription Factor ABF1 and Separate HAP2- and Heme-Responsive Elements. <i>Molecular and Cellular Biology</i> , 1992, 12, 2302-2314. | 1.1 | 16 |
| 7209 | Role of GCR2 in Transcriptional Activation of Yeast Glycolytic Genes. <i>Molecular and Cellular Biology</i> , 1992, 12, 3834-3842. | 1.1 | 42 |
| 7210 | A Delay in the <i>Saccharomyces cerevisiae</i> Cell Cycle That Is Induced by a Dicentric Chromosome and Dependent upon Mitotic Checkpoints. <i>Molecular and Cellular Biology</i> , 1992, 12, 3857-3864. | 1.1 | 28 |
| 7211 | The 70-Kilodalton Adenylyl Cyclase-Associated Protein Is Not Essential for Interaction of <i>Saccharomyces cerevisiae</i> Adenylyl Cyclase with RAS Proteins. <i>Molecular and Cellular Biology</i> , 1992, 12, 4937-4945. | 1.1 | 11 |
| 7212 | Suppressors of Clathrin Deficiency: Overexpression of Ubiquitin Rescues Lethal Strains of Clathrin-Deficient <i>Saccharomyces cerevisiae</i> . <i>Molecular and Cellular Biology</i> , 1993, 13, 521-532. | 1.1 | 34 |
| 7213 | Yeast as a Vehicle for the Expression of Heterologous Genes. , 1993, , 325-356. | | 1 |
| 7214 | Antibody Mimicking the Action of RAS Proteins on Yeast Adenylyl Cyclase: Implication for RAS-Effector Interaction. <i>Molecular and Cellular Biology</i> , 1993, 13, 769-774. | 1.1 | 5 |
| 7215 | Differential Transcriptional Activation by v-Myb and c-Myb in Animal Cells and <i>Saccharomyces cerevisiae</i> . <i>Molecular and Cellular Biology</i> , 1993, 13, 4423-4431. | 1.1 | 22 |
| 7216 | Expression of p60 ^{v-src} in <i>Saccharomyces cerevisiae</i> Results in Elevation of p34 ^{CDC28} Kinase Activity and Release of the Dependence of DNA Replication on Mitosis. <i>Molecular and Cellular Biology</i> , 1993, 13, 5112-5121. | 1.1 | 9 |
| 7217 | Yeast Phenotype Classifies Mammalian Protein Kinase C cDNA Mutants. <i>Molecular and Cellular Biology</i> , 1993, 13, 4728-4735. | 1.1 | 5 |
| 7218 | NPK1, a tobacco gene that encodes a protein with a domain homologous to yeast BCK1, STE11, and Byr2 protein kinases. <i>Molecular and Cellular Biology</i> , 1993, 13, 4745-4752. | 1.1 | 44 |
| 7221 | Histone H1 Expressed in <i>Saccharomyces cerevisiae</i> Binds to Chromatin and Affects Survival, Growth, Transcription, and Plasmid Stability but Does Not Change Nucleosomal Spacing. <i>Molecular and Cellular Biology</i> , 1994, 14, 2822-2835. | 1.1 | 13 |
| 7222 | Special Peptidyl-tRNA Molecules Can Promote Translational Frameshifting without Slippage. <i>Molecular and Cellular Biology</i> , 1994, 14, 8107-8116. | 1.1 | 35 |
| 7223 | A practical method for fission yeast transformation by electroporation. <i>Genes and Genetic Systems</i> , 1995, 70, 1-6. | 0.2 | 0 |
| 7224 | Expression and Screening in Yeast of Genes Mutagenized in Vitro. , 1995, , 373-384. | | 0 |
| 7225 | Structural and functional analysis of methylotrophic yeast <i>Hansenula polymorpha</i> alcohol oxydase gene promoter. <i>Biopolymers and Cell</i> , 1995, 11, 39-45. | 0.1 | 0 |
| 7226 | Methods for Expressing Recombinant Proteins in Yeast. , 1996, , 105-153. | | 0 |

| # | ARTICLE | IF | CITATIONS |
|------|--|-----|-----------|
| 7227 | Recombinant Protein Techniques. , 1996, , 325-363. | | 0 |
| 7229 | Expression of small synthetic genes coding for hEGF, human epidermal growth factor, and CPTI II, serine proteinase inhibitor from Cucurbitacea, cloned in a novel expression/secretion vector in <i>Saccharomyces cerevisiae</i> .. <i>Acta Biochimica Polonica</i> , 1996, 43, 255-264. | 0.3 | 1 |
| 7230 | Dosage Suppressors of a Benomyl-Dependent Tubulin Mutant: Evidence for a Link Between Microtubule Stability and Cellular Metabolism. <i>Genetics</i> , 1996, 144, 1363-1373. | 1.2 | 1 |
| 7231 | Asparaginase II of <i>Saccharomyces cerevisiae</i> . , 1997, 63-65, 203-212. | | 17 |
| 7232 | Identification of proteins that interact with a protein of interest: Applications of the yeast two-hybrid system. , 1997, , 67-79. | | 28 |
| 7234 | Yeast Meiotic Mutants Proficient for the Induction of Ectopic Recombination. <i>Genetics</i> , 1998, 148, 581-598. | 1.2 | 27 |
| 7235 | Functional Dissection of Yeast Hir1p, a WD Repeatâ€“Containing Transcriptional Corepressor. <i>Genetics</i> , 1998, 148, 657-667. | 1.2 | 35 |
| 7236 | A Novel Asn344 Deletion in the Core Domain of Coagulation Factor XIII A Subunit: Its Effects on Protein Structure and Function. <i>Blood</i> , 1998, 92, 481-487. | 0.6 | 0 |
| 7237 | Yeast genetics. , 1999, , 43-82. | | 0 |
| 7238 | BTF3 is a potential new substrate of protein kinase CK2. , 1999, , 121-128. | | 5 |
| 7239 | Frequent Meiotic Recombination Between the Ends of Truncated Chromosome Fragments of <i>Saccharomyces cerevisiae</i> . <i>Genetics</i> , 1999, 153, 1583-1590. | 1.2 | 5 |
| 7240 | Evaluation of Competence Phenomenon of Yeast <i>Saccharomyces cerevisiae</i> by Lipophilic Cations Accumulation and FT-IR Spectroscopy. Relation of Competence to Cell Cycle. <i>Fungal Biology</i> , 2015, , 239-252. | 0.3 | 0 |
| 7242 | Transformation of Intact Cells of <i>Saccharomyces cerevisiae</i> : Lithium Methods and Possible Underlying Mechanism. <i>Fungal Biology</i> , 2015, , 187-192. | 0.3 | 1 |
| 7243 | Unique C-terminal region of Hap3 is required for methanol-regulated gene expression in the methylotrophic yeast <i>Candida boidinii</i> . <i>Microbiology (United Kingdom)</i> , 2016, 162, 898-907. | 0.7 | 1 |
| 7248 | Hefen. , 2019, , 63-91. | | 0 |
| 7250 | Electrotransformation of <i>Saccharomyces cerevisiae</i> . <i>Methods in Molecular Biology</i> , 2020, 2050, 187-193. | 0.4 | 2 |
| 7253 | Genomic Promoter Shuffling by Using Recyclable Cassettes. <i>Methods in Molecular Biology</i> , 2021, 2196, 39-51. | 0.4 | 0 |
| 7255 | Physical, genetic and functional interactions between the eisosome protein Pil1 and the MBOAT O-acyltransferase Gup1. <i>FEMS Yeast Research</i> , 2021, 21, . | 1.1 | 0 |

| # | ARTICLE | IF | CITATIONS |
|------|--|-----|-----------|
| 7256 | <i>rad3</i> gene of <i>Saccharomyces cerevisiae</i> : Nucleotide Sequence of Wild-Type and Mutant Alleles, Transcript Mapping, and Aspects of Gene Regulation. <i>Molecular and Cellular Biology</i> , 1985, 5, 17-26. | 1.1 | 57 |
| 7257 | Identification of the <i>ergC</i> gene involved in polyene drug sensitivity in the Mucorales species <i>Phycomyces blakesleeanus</i> . <i>Molecular Biology Reports</i> , 2022, 49, 981-987. | 1.0 | 1 |
| 7258 | Plasmid DNA transformation of yeast cells treated with chelating agents. The role of lipid peroxidation in inducing competence in yeast. <i>Biopolymers and Cell</i> , 1985, 1, 147-153. | 0.1 | 1 |
| 7260 | Study Essential Gene Functions by Plasmid Shuffling. <i>Methods in Molecular Biology</i> , 2021, 2196, 53-62. | 0.4 | 3 |
| 7261 | PHO5-LACZ hybrid proteins block translocation of native acid phosphatase in <i>Saccharomyces cerevisiae</i> . <i>Journal of Biological Chemistry</i> , 1988, 263, 6908-6915. | 1.6 | 4 |
| 7262 | Characterization of an .ALPHA.-Ketoglutarate-Resistant Sake Yeast Mutant with High Organic Acid Productivity. <i>Journal of Bioscience and Bioengineering</i> , 2003, 96, 332-336. | 1.1 | 1 |
| 7263 | Repeated Chromosome Splitting Targeted to .DELTA. Sequences in <i>Saccharomyces cerevisiae</i> . <i>Journal of Bioscience and Bioengineering</i> , 2003, 96, 397-400. | 1.1 | 1 |
| 7264 | Invertase β -Galactosidase Hybrid Proteins Fail To Be Transported from the Endoplasmic Reticulum in <i>Saccharomyces cerevisiae</i> . <i>Molecular and Cellular Biology</i> , 1984, 4, 2347-2355. | 1.1 | 41 |
| 7265 | Repetitive <i>Dictyostelium</i> Heat-Shock Promotor Functions in <i>Saccharomyces cerevisiae</i> . <i>Molecular and Cellular Biology</i> , 1984, 4, 591-598. | 1.1 | 20 |
| 7266 | Transformation of Protoplasted Yeast Cells Is Directly Associated with Cell Fusion. <i>Molecular and Cellular Biology</i> , 1984, 4, 771-778. | 1.1 | 45 |
| 7267 | Micronuclear DNA of <i>Oxytricha nova</i> Contains Sequences with Autonomously Replicating Activity in <i>Saccharomyces cerevisiae</i> . <i>Molecular and Cellular Biology</i> , 1984, 4, 1725-1729. | 1.1 | 2 |
| 7268 | Point Mutations Implicate Repeated Sequences as Essential Elements of the <i>CYC7</i> Negative Upstream Site in <i>Saccharomyces cerevisiae</i> . <i>Molecular and Cellular Biology</i> , 1985, 5, 2951-2958. | 1.1 | 24 |
| 7269 | Structure of the Promoter of the <i>Dictyostelium discoideum</i> Prespore EB4 Gene. <i>Molecular and Cellular Biology</i> , 1985, 5, 1465-1472. | 1.1 | 22 |
| 7270 | Effect of <i>ARS1</i> Mutations on Chromosome Stability in <i>Saccharomyces cerevisiae</i> . <i>Molecular and Cellular Biology</i> , 1985, 5, 1676-1684. | 1.1 | 59 |
| 7271 | Recombination Hot Spot in the Human β -Globin Gene Cluster: Meiotic Recombination of Human DNA Fragments in <i>Saccharomyces cerevisiae</i> . <i>Molecular and Cellular Biology</i> , 1985, 5, 2029-2038. | 1.1 | 40 |
| 7272 | <i>Saccharomyces cerevisiae</i> nuclear fusion requires prior activation by alpha factor. <i>Molecular and Cellular Biology</i> , 1986, 6, 3490-3497. | 1.1 | 49 |
| 7273 | Site-Specific Mutagenesis of <i>cdc2⁺</i> , a Cell Cycle Control Gene of the Fission Yeast <i>Schizosaccharomyces pombe</i> . <i>Molecular and Cellular Biology</i> , 1986, 6, 3523-3530. | 1.1 | 104 |
| 7274 | Expression of the <i>denV</i> Gene of Coliphage T4 in UV-Sensitive <i>rad</i> Mutants of <i>Saccharomyces cerevisiae</i> . <i>Molecular and Cellular Biology</i> , 1986, 6, 3559-3562. | 1.1 | 6 |

| # | ARTICLE | IF | CITATIONS |
|------|---|-----|-----------|
| 7275 | Mating type-like conversion promoted by the 2 micrograms circle site-specific recombinase: implications for the double-strand-gap repair model. <i>Molecular and Cellular Biology</i> , 1986, 6, 3831-3837. | 1.1 | 6 |
| 7276 | A nuclear gene of <i>Saccharomyces cerevisiae</i> needed for stable maintenance of plasmids. <i>Molecular and Cellular Biology</i> , 1986, 6, 4053-4059. | 1.1 | 7 |
| 7277 | In vivo evidence for posttranslational translocation and signal cleavage of the killer preprotoxin of <i>Saccharomyces cerevisiae</i> . <i>Molecular and Cellular Biology</i> , 1986, 6, 4274-4280. | 1.1 | 20 |
| 7278 | Isolation and characterization of PRT1, a gene required for the initiation of protein biosynthesis in <i>Saccharomyces cerevisiae</i> . <i>Molecular and Cellular Biology</i> , 1986, 6, 4419-4424. | 1.1 | 9 |
| 7279 | Activation of a cryptic TACTAAC box in the <i>Saccharomyces cerevisiae</i> actin intron. <i>Molecular and Cellular Biology</i> , 1986, 6, 1571-1578. | 1.1 | 20 |
| 7280 | Differential expressions of essential and nonessential alpha-tubulin genes in <i>Schizosaccharomyces pombe</i> . <i>Molecular and Cellular Biology</i> , 1986, 6, 2168-2178. | 1.1 | 52 |
| 7281 | Isolation and functional analysis of sporulation-induced transcribed sequences from <i>Saccharomyces cerevisiae</i> . <i>Molecular and Cellular Biology</i> , 1986, 6, 2185-2197. | 1.1 | 42 |
| 7282 | Identification of a regulatory region that mediates glucose-dependent induction of the <i>Saccharomyces cerevisiae</i> enolase gene ENO2. <i>Molecular and Cellular Biology</i> , 1986, 6, 2287-2297. | 1.1 | 54 |
| 7283 | Effect of Intron Mutations on Processing and Function of <i>Saccharomyces cerevisiae</i> SUP53 tRNA In Vitro and In Vivo. <i>Molecular and Cellular Biology</i> , 1986, 6, 2663-2673. | 1.1 | 61 |
| 7284 | Identification of a Second <i>trans</i> -Acting Gene Controlling Maltose Fermentation in <i>Saccharomyces carlsbergensis</i> . <i>Molecular and Cellular Biology</i> , 1986, 6, 2757-2765. | 1.1 | 26 |
| 7285 | Differential effectiveness of yeast cytochrome c oxidase subunit genes results from differences in expression not function. <i>Molecular and Cellular Biology</i> , 1987, 7, 3520-3526. | 1.1 | 37 |
| 7286 | Insertions of up to 17 Amino Acids into a Region of α -Tubulin Do Not Disrupt Function In Vivo. <i>Molecular and Cellular Biology</i> , 1987, 7, 3799-3805. | 1.1 | 37 |
| 7287 | <i>RAD4</i> Gene of <i>Saccharomyces cerevisiae</i> : Molecular Cloning and Partial Characterization of a Gene That Is Inactivated in <i>Escherichia coli</i> . <i>Molecular and Cellular Biology</i> , 1987, 7, 1180-1192. | 1.1 | 24 |
| 7288 | Effect of Deletion and Insertion on Double-Strand-Break Repair in <i>Saccharomyces cerevisiae</i> . <i>Molecular and Cellular Biology</i> , 1987, 7, 1300-1303. | 1.1 | 13 |
| 7289 | Guanine Nucleotide Activation of, and Competition between, Ras Proteins from <i>Saccharomyces Cerevisiae</i> . <i>Molecular and Cellular Biology</i> , 1987, 7, 2128-2133. | 1.1 | 72 |
| 7290 | Regulatory function of the <i>Saccharomyces cerevisiae</i> RAS C-terminus. <i>Molecular and Cellular Biology</i> , 1987, 7, 2309-2315. | 1.1 | 48 |
| 7291 | Genetic manipulation of centromere function. <i>Molecular and Cellular Biology</i> , 1987, 7, 2397-2405. | 1.1 | 147 |
| 7292 | Upstream regulatory regions controlling the expression of the yeast maltase gene. <i>Molecular and Cellular Biology</i> , 1987, 7, 2477-2483. | 1.1 | 18 |

| # | ARTICLE | IF | CITATIONS |
|------|---|-----|-----------|
| 7293 | Phycomyces blakesleeanus TRP1 gene: organization and functional complementation in Escherichia coli and Saccharomyces cerevisiae. Molecular and Cellular Biology, 1987, 7, 2664-2670. | 1.1 | 11 |
| 7294 | Transcription of the constitutively expressed yeast enolase gene ENO1 is mediated by positive and negative cis-acting regulatory sequences. Molecular and Cellular Biology, 1987, 7, 2753-2761. | 1.1 | 65 |
| 7295 | Identification of sequence elements that confer cell-type-specific control of MF alpha 1 expression in Saccharomyces cerevisiae. Molecular and Cellular Biology, 1987, 7, 3185-3193. | 1.1 | 34 |
| 7296 | Comparison of tRNA Gene Transcription Complexes Formed In Vitro and in Nuclei. Molecular and Cellular Biology, 1987, 7, 3212-3220. | 1.1 | 28 |
| 7297 | The Yeast Acid Phosphatase Can Enter the Secretory Pathway without Its N-Terminal Signal Sequence. Molecular and Cellular Biology, 1987, 7, 3306-3314. | 1.1 | 16 |
| 7298 | Properties of Promoters Cloned Randomly from the <i>Saccharomyces cerevisiae</i> Genome. Molecular and Cellular Biology, 1988, 8, 4217-4224. | 1.1 | 27 |
| 7299 | Genetic Evidence for Promoter Competition in <i>Saccharomyces cerevisiae</i> . Molecular and Cellular Biology, 1988, 8, 4608-4615. | 1.1 | 65 |
| 7300 | Transcriptional Regulation of the Cell Cycle-Dependent Thymidylate Synthase Gene of <i>Saccharomyces cerevisiae</i> . Molecular and Cellular Biology, 1988, 8, 4616-4624. | 1.1 | 29 |
| 7301 | A Ty1 Cell-Type-Specific Regulatory Sequence Is a Recognition Element for a Constitutive Binding Factor. Molecular and Cellular Biology, 1988, 8, 5299-5309. | 1.1 | 31 |
| 7302 | Role of <i>SGP2</i> , a Suppressor of a <i>gpa1</i> Mutation, in the Mating-Factor Signaling Pathway of <i>Saccharomyces cerevisiae</i> . Molecular and Cellular Biology, 1988, 8, 5410-5416. | 1.1 | 19 |
| 7303 | The First and Fourth Upstream Open Reading Frames in <i>GCN4</i> mRNA Have Similar Initiation Efficiencies but Respond Differently in Translational Control to Change in Length and Sequence. Molecular and Cellular Biology, 1988, 8, 5439-5447. | 1.1 | 22 |
| 7304 | SPT3 is required for normal levels of a-factor and alpha-factor expression in <i>Saccharomyces cerevisiae</i> . Molecular and Cellular Biology, 1988, 8, 822-827. | 1.1 | 25 |
| 7305 | Nematode Repetitive DNA with <i>ARS</i> and Segregation Function in <i>Saccharomyces cerevisiae</i> . Molecular and Cellular Biology, 1988, 8, 875-883. | 1.1 | 7 |
| 7306 | Gene conversion associated with site-specific recombination in yeast plasmid pSR1. Molecular and Cellular Biology, 1988, 8, 955-962. | 1.1 | 19 |
| 7307 | Constitutive expression of the maltose fermentative enzymes in <i>Saccharomyces carlsbergensis</i> is dependent upon the mutational activation of a nonessential homolog of MAL63. Molecular and Cellular Biology, 1988, 8, 1027-1035. | 1.1 | 21 |
| 7308 | The <i>Saccharomyces cerevisiae</i> ACP2 gene encodes an essential HMG1-like protein. Molecular and Cellular Biology, 1988, 8, 1282-1289. | 1.1 | 26 |
| 7309 | Single-Amino-Acid Substitutions within the Signal Sequence of Yeast Prepro- α -Factor Affect Membrane Translocation. Molecular and Cellular Biology, 1988, 8, 1915-1922. | 1.1 | 16 |
| 7310 | A New Means of Inducibly Inactivating a Cellular Protein. Molecular and Cellular Biology, 1988, 8, 2638-2646. | 1.1 | 43 |

| # | ARTICLE | IF | CITATIONS |
|------|--|-----|-----------|
| 7311 | Bent DNA Functions as a Replication Enhancer in <i>Saccharomyces cerevisiae</i> . <i>Molecular and Cellular Biology</i> , 1988, 8, 2763-2769. | 1.1 | 50 |
| 7312 | The Positive Regulatory Function of the 5' Proximal Open Reading Frames in <i>GCN4</i> mRNA Can Be Mimicked by Heterologous, Short Coding Sequences. <i>Molecular and Cellular Biology</i> , 1988, 8, 3827-3836. | 1.1 | 31 |
| 7313 | Differential Repression of GAL4 and Adjacent Transcription Activators by Operators in the Yeast <i>GAL</i> Upstream Activating Sequence. <i>Molecular and Cellular Biology</i> , 1989, 9, 4282-4290. | 1.1 | 13 |
| 7314 | Mating-Type Control in <i>Saccharomyces cerevisiae</i> : Isolation and Characterization of Mutants Defective in Repression by <i>a1-2</i> . <i>Molecular and Cellular Biology</i> , 1989, 9, 4523-4530. | 1.1 | 20 |
| 7315 | Identification of a Yeast Protein with Properties Similar to Those of the Immunoglobulin Heavy-Chain Enhancer-Binding Protein NF- κ B. <i>Molecular and Cellular Biology</i> , 1989, 9, 4535-4540. | 1.1 | 10 |
| 7316 | Elongation Factor EF-1 Gene Dosage Alters Translational Fidelity in <i>Saccharomyces cerevisiae</i> . <i>Molecular and Cellular Biology</i> , 1989, 9, 4571-4575. | 1.1 | 41 |
| 7317 | Mutations in the Signal Sequence of Prepro- α -Factor Inhibit Both Translocation into the Endoplasmic Reticulum and Processing by Signal Peptidase in Yeast Cells. <i>Molecular and Cellular Biology</i> , 1989, 9, 4977-4985. | 1.1 | 6 |
| 7318 | Role of glutamic acid 177 of the ricin toxin A chain in enzymatic inactivation of ribosomes. <i>Molecular and Cellular Biology</i> , 1989, 9, 5012-5021. | 1.1 | 31 |
| 7319 | Identification of an Upstream Activation Sequence and Other cis-Acting Elements Required for Transcription of <i>COX6</i> from <i>Saccharomyces cerevisiae</i> . <i>Molecular and Cellular Biology</i> , 1989, 9, 5350-5358. | 1.1 | 21 |
| 7320 | Upstream regulatory sequences of the yeast <i>RNR2</i> gene include a repression sequence and an activation site that binds the RAP1 protein. <i>Molecular and Cellular Biology</i> , 1989, 9, 5359-5372. | 1.1 | 37 |
| 7321 | Isolation and Characterization of Mutations in the <i>HXK2</i> Gene of <i>Saccharomyces cerevisiae</i> . <i>Molecular and Cellular Biology</i> , 1989, 9, 5630-5642. | 1.1 | 20 |
| 7322 | <i>Sec59</i> Encodes a Membrane Protein Required for Core Glycosylation in <i>Saccharomyces cerevisiae</i> . <i>Molecular and Cellular Biology</i> , 1989, 9, 1191-1199. | 1.1 | 25 |
| 7323 | Sequences within the Spacer Region of Yeast rRNA Cistrons That Stimulate 35S rRNA Synthesis In Vivo Mediate RNA Polymerase I-Dependent Promoter and Terminator Activities. <i>Molecular and Cellular Biology</i> , 1989, 9, 1243-1254. | 1.1 | 31 |
| 7324 | Intragenic Revertants of Yeast Invertase Variants with Secretion-Defective Leader Sequences. <i>Molecular and Cellular Biology</i> , 1989, 9, 1452-1464. | 1.1 | 5 |
| 7325 | The OBF1 Protein and Its DNA-Binding Site Are Important for the Function of an Autonomously Replicating Sequence in <i>Saccharomyces cerevisiae</i> . <i>Molecular and Cellular Biology</i> , 1989, 9, 2914-2921. | 1.1 | 36 |
| 7326 | Use of Yeast Nuclear DNA Sequences To Define the Mitochondrial RNA Polymerase Promoter In Vitro. <i>Molecular and Cellular Biology</i> , 1989, 9, 3193-3202. | 1.1 | 20 |
| 7327 | A Gene Product Needed for Induction of Allantoin System Genes in <i>Saccharomyces cerevisiae</i> but Not for Their Transcriptional Activation. <i>Molecular and Cellular Biology</i> , 1989, 9, 3869-3877. | 1.1 | 14 |
| 7329 | <i>Saccharomyces cerevisiae</i> cells lacking the homologous pairing protein p175 SEP1 arrest at pachytene during meiotic prophase. <i>Chromosoma</i> , 1994, 103, 129-141. | 1.0 | 6 |

| # | ARTICLE | IF | CITATIONS |
|------|--|-----|-----------|
| 7330 | Mutational analysis of the \hat{I}^2 -subunit of yeast geranylgeranyl transferase I. <i>Molecular Genetics and Genomics</i> , 1996, 252, 1-10. | 2.4 | 0 |
| 7401 | Mcm10 and the MCM2-7 complex interact to initiate DNA synthesis and to release replication factors from origins. <i>Genes and Development</i> , 2000, 14, 913-26. | 2.7 | 85 |
| 7402 | Extranuclear gene expression in yeast: evidence for a plasmid-encoded RNA polymerase of unique structure. <i>Nucleic Acids Research</i> , 1988, 16, 8097-112. | 6.5 | 89 |
| 7405 | Nin1p, a regulatory subunit of the 26S proteasome, is necessary for activation of Cdc28p kinase of <i>Saccharomyces cerevisiae</i> . <i>EMBO Journal</i> , 1995, 14, 3105-15. | 3.5 | 32 |
| 7406 | GCN20, a novel ATP binding cassette protein, and GCN1 reside in a complex that mediates activation of the eIF-2 alpha kinase GCN2 in amino acid-starved cells. <i>EMBO Journal</i> , 1995, 14, 3184-99. | 3.5 | 62 |
| 7407 | A novel protein kinase gene <i>ssp1+</i> is required for alteration of growth polarity and actin localization in fission yeast. <i>EMBO Journal</i> , 1995, 14, 3325-38. | 3.5 | 33 |
| 7408 | Functionality and specific membrane localization of transport GTPases carrying C-terminal membrane anchors of synaptobrevin-like proteins. <i>EMBO Journal</i> , 1995, 14, 3645-53. | 3.5 | 16 |
| 7409 | The <i>Saccharomyces cerevisiae</i> translation initiation factor Tif3 and its mammalian homologue, eIF-4B, have RNA annealing activity. <i>EMBO Journal</i> , 1995, 14, 3820-7. | 3.5 | 40 |
| 7410 | Identification of the cis-elements mediating the autogenous control of ribosomal protein L2 mRNA stability in yeast. <i>EMBO Journal</i> , 1995, 14, 4022-30. | 3.5 | 18 |
| 7411 | Termination of translation in eukaryotes is governed by two interacting polypeptide chain release factors, eRF1 and eRF3. <i>EMBO Journal</i> , 1995, 14, 4065-72. | 3.5 | 295 |
| 7412 | A downstream target of RHO1 small GTP-binding protein is PKC1, a homolog of protein kinase C, which leads to activation of the MAP kinase cascade in <i>Saccharomyces cerevisiae</i> . <i>EMBO Journal</i> , 1995, 14, 5931-8. | 3.5 | 160 |
| 7413 | REG1 binds to protein phosphatase type 1 and regulates glucose repression in <i>Saccharomyces cerevisiae</i> . <i>EMBO Journal</i> , 1995, 14, 5939-46. | 3.5 | 131 |
| 7414 | <i>Schizosaccharomyces pombe atf1+</i> encodes a transcription factor required for sexual development and entry into stationary phase. <i>EMBO Journal</i> , 1995, 14, 6193-208. | 3.5 | 131 |
| 7415 | MSG5, a novel protein phosphatase promotes adaptation to pheromone response in <i>S. cerevisiae</i> . <i>EMBO Journal</i> , 1994, 13, 61-70. | 3.5 | 117 |
| 7416 | Functional differences between mammalian transcription activation domains at the yeast GAL1 promoter. <i>EMBO Journal</i> , 1994, 13, 641-5. | 3.5 | 31 |
| 7417 | A dominant negative mutation in the conserved RNA helicase motif 'SAT' causes splicing factor PRP2 to stall in spliceosomes. <i>EMBO Journal</i> , 1994, 13, 879-87. | 3.5 | 66 |
| 7418 | The Dbf2 and Dbf20 protein kinases of budding yeast are activated after the metaphase to anaphase cell cycle transition. <i>EMBO Journal</i> , 1994, 13, 1103-13. | 3.5 | 60 |
| 7419 | VPS21 encodes a rab5-like GTP binding protein that is required for the sorting of yeast vacuolar proteins. <i>EMBO Journal</i> , 1994, 13, 1297-309. | 3.5 | 96 |

| # | ARTICLE | IF | CITATIONS |
|------|---|-----|-----------|
| 7420 | The <i>Saccharomyces cerevisiae</i> APS1 gene encodes a homolog of the small subunit of the mammalian clathrin AP-1 complex: evidence for functional interaction with clathrin at the Golgi complex. <i>EMBO Journal</i> , 1994, 13, 1706-17. | 3.5 | 43 |
| 7421 | GDI1 encodes a GDP dissociation inhibitor that plays an essential role in the yeast secretory pathway. <i>EMBO Journal</i> , 1994, 13, 1718-28. | 3.5 | 71 |
| 7422 | A mitochondrial homolog of bacterial GrpE interacts with mitochondrial hsp70 and is essential for viability. <i>EMBO Journal</i> , 1994, 13, 1998-2006. | 3.5 | 52 |
| 7423 | Characterization of two protein kinases from <i>Schizosaccharomyces pombe</i> involved in the regulation of DNA repair. <i>EMBO Journal</i> , 1994, 13, 2777-88. | 3.5 | 47 |
| 7424 | Molecular characterization of SIG1, a <i>Saccharomyces cerevisiae</i> gene involved in negative regulation of G-protein-mediated signal transduction. <i>EMBO Journal</i> , 1994, 13, 3050-64. | 3.5 | 11 |
| 7425 | The yeast NOP4 gene product is an essential nucleolar protein required for pre-rRNA processing and accumulation of 60S ribosomal subunits. <i>EMBO Journal</i> , 1994, 13, 3127-35. | 3.5 | 61 |
| 7426 | Cloning and expression of the MEP1 gene encoding an ammonium transporter in <i>Saccharomyces cerevisiae</i> . <i>EMBO Journal</i> , 1994, 13, 3456-63. | 3.5 | 101 |
| 7427 | A functional interaction between Rev and yeast pre-mRNA is related to splicing complex formation. <i>EMBO Journal</i> , 1994, 13, 4096-104. | 3.5 | 37 |
| 7428 | The HOG pathway controls osmotic regulation of transcription via the stress response element (STRE) of the <i>Saccharomyces cerevisiae</i> CTT1 gene. <i>EMBO Journal</i> , 1994, 13, 4382-9. | 3.5 | 154 |
| 7429 | A nucleosome precludes binding of the transcription factor Pho4 in vivo to a critical target site in the PHO5 promoter. <i>EMBO Journal</i> , 1994, 13, 4848-55. | 3.5 | 62 |
| 7430 | Fission yeast cut3 and cut14, members of a ubiquitous protein family, are required for chromosome condensation and segregation in mitosis. <i>EMBO Journal</i> , 1994, 13, 4938-52. | 3.5 | 168 |
| 7431 | A zinc finger protein, essential for chromosome segregation, constitutes a putative DNA binding subunit of the <i>Saccharomyces cerevisiae</i> kinetochore complex, Cbf3. <i>EMBO Journal</i> , 1994, 13, 5203-11. | 3.5 | 53 |
| 7432 | Phosphorylation of dis2 protein phosphatase at the C-terminal cdc2 consensus and its potential role in cell cycle regulation. <i>EMBO Journal</i> , 1994, 13, 5310-8. | 3.5 | 45 |
| 7433 | Fission yeast cut5 links nuclear chromatin and M phase regulator in the replication checkpoint control. <i>EMBO Journal</i> , 1994, 13, 5319-29. | 3.5 | 79 |
| 7434 | Complete DNA sequence of yeast chromosome II. <i>EMBO Journal</i> , 1994, 13, 5795-809. | 3.5 | 61 |
| 7435 | The bovine papillomavirus 1 E2 protein contains two activation domains: one that interacts with TBP and another that functions after TBP binding. <i>EMBO Journal</i> , 1995, 14, 329-40. | 3.5 | 57 |
| 7436 | Sm and Sm-like proteins belong to a large family: identification of proteins of the U6 as well as the U1, U2, U4 and U5 snRNPs. <i>EMBO Journal</i> , 1995, 14, 2089-98. | 3.5 | 138 |
| 7437 | Mutations in GCD11, the structural gene for eIF-2 gamma in yeast, alter translational regulation of GCN4 and the selection of the start site for protein synthesis. <i>EMBO Journal</i> , 1995, 14, 2239-49. | 3.5 | 36 |

| # | ARTICLE | IF | CITATIONS |
|------|---|-----|-----------|
| 7438 | Interpathway regulation of the TRP4 gene of yeast. EMBO Journal, 1989, 8, 939-45. | 3.5 | 34 |
| 7439 | Characterization of the END1 gene required for vacuole biogenesis and gluconeogenic growth of budding yeast. EMBO Journal, 1989, 8, 1349-59. | 3.5 | 34 |
| 7440 | The ras-related mouse ypt1 protein can functionally replace the YPT1 gene product in yeast. EMBO Journal, 1989, 8, 1427-32. | 3.5 | 60 |
| 7441 | Mutational analysis of SEC4 suggests a cyclical mechanism for the regulation of vesicular traffic. EMBO Journal, 1989, 8, 1685-93. | 3.5 | 131 |
| 7442 | A point mutation in the CYC1 UAS1 creates a new combination of regulatory elements that activate transcription synergistically. EMBO Journal, 1989, 8, 1801-8. | 3.5 | 9 |
| 7443 | The cloned human oestrogen receptor contains a mutation which alters its hormone binding properties. EMBO Journal, 1989, 8, 1981-6. | 3.5 | 119 |
| 7444 | Isolation and DNA-binding characteristics of a protein involved in transcription activation of two divergently transcribed, essential yeast genes. EMBO Journal, 1989, 8, 3029-37. | 3.5 | 48 |
| 7445 | Mutants of H-ras that interfere with RAS effector function in <i>Saccharomyces cerevisiae</i> . EMBO Journal, 1989, 8, 3039-44. | 3.5 | 23 |
| 7446 | <i>Saccharomyces cerevisiae</i> STE6 gene product: a novel pathway for protein export in eukaryotic cells. EMBO Journal, 1989, 8, 3973-84. | 3.5 | 186 |
| 7447 | A yeast nucleolar protein related to mammalian fibrillarin is associated with small nucleolar RNA and is essential for viability. EMBO Journal, 1989, 8, 4015-24. | 3.5 | 184 |
| 7448 | Sequence, expression and mutational analysis of BAF1, a transcriptional activator and ARS1-binding protein of the yeast <i>Saccharomyces cerevisiae</i> . EMBO Journal, 1989, 8, 4265-72. | 3.5 | 53 |
| 7449 | A human homologue of the yeast GST1 gene codes for a GTP-binding protein and is expressed in a proliferation-dependent manner in mammalian cells. EMBO Journal, 1989, 8, 3807-14. | 3.5 | 60 |
| 7450 | Isolation and characterization of <i>S. cerevisiae</i> mutants defective in somatostatin expression: cloning and functional role of a yeast gene encoding an aspartyl protease in precursor processing at monobasic cleavage sites. EMBO Journal, 1993, 12, 285-94. | 3.5 | 12 |
| 7451 | Evidence for an interaction between the CYP1(HAP1) activator and a cellular factor during heme-dependent transcriptional regulation in the yeast <i>Saccharomyces cerevisiae</i> . EMBO Journal, 1993, 12, 1209-18. | 3.5 | 26 |
| 7452 | The control in cis of the position and the amount of the ARG4 meiotic double-strand break of <i>Saccharomyces cerevisiae</i> . EMBO Journal, 1993, 12, 1459-66. | 3.5 | 51 |
| 7453 | Two novel protein kinase C-related genes of fission yeast are essential for cell viability and implicated in cell shape control. EMBO Journal, 1993, 12, 1987-95. | 3.5 | 47 |
| 7454 | A <i>Saccharomyces cerevisiae</i> UAS element controlled by protein kinase A activates transcription in response to a variety of stress conditions. EMBO Journal, 1993, 12, 1997-2003. | 3.5 | 167 |
| 7455 | A membrane-associated complex containing the Vps15 protein kinase and the Vps34 PI 3-kinase is essential for protein sorting to the yeast lysosome-like vacuole. EMBO Journal, 1993, 12, 2195-204. | 3.5 | 144 |

| # | ARTICLE | IF | CITATIONS |
|------|--|-----|-----------|
| 7456 | GCR1, a transcriptional activator in <i>Saccharomyces cerevisiae</i> , complexes with RAP1 and can function without its DNA binding domain. <i>EMBO Journal</i> , 1993, 12, 2431-7. | 3.5 | 48 |
| 7457 | A U3 snoRNP protein with homology to splicing factor PRP4 and G beta domains is required for ribosomal RNA processing. <i>EMBO Journal</i> , 1993, 12, 2549-58. | 3.5 | 78 |
| 7458 | Vps1p, a member of the dynamin GTPase family, is necessary for Golgi membrane protein retention in <i>Saccharomyces cerevisiae</i> . <i>EMBO Journal</i> , 1993, 12, 3049-59. | 3.5 | 69 |
| 7459 | Salt tolerance and methionine biosynthesis in <i>Saccharomyces cerevisiae</i> involve a putative phosphatase gene. <i>EMBO Journal</i> , 1993, 12, 3105-10. | 3.5 | 49 |
| 7460 | Properties of <i>Saccharomyces cerevisiae</i> wee1 and its differential regulation of p34CDC28 in response to G1 and G2 cyclins. <i>EMBO Journal</i> , 1993, 12, 3417-26. | 3.5 | 198 |
| 7461 | A new pair of B-type cyclins from <i>Saccharomyces cerevisiae</i> that function early in the cell cycle. <i>EMBO Journal</i> , 1993, 12, 3437-47. | 3.5 | 52 |
| 7462 | A <i>Saccharomyces cerevisiae</i> homologue of mammalian translation initiation factor 4B contributes to RNA helicase activity. <i>EMBO Journal</i> , 1993, 12, 3997-4003. | 3.5 | 66 |
| 7463 | A new yeast translation initiation factor suppresses a mutation in the eIF-4A RNA helicase. <i>EMBO Journal</i> , 1993, 12, 4005-11. | 3.5 | 45 |
| 7464 | The yeast SSS1 gene is essential for secretory protein translocation and encodes a conserved protein of the endoplasmic reticulum. <i>EMBO Journal</i> , 1993, 12, 4083-93. | 3.5 | 34 |
| 7465 | Functional cooperation of mitochondrial protein import receptors in yeast. <i>EMBO Journal</i> , 1993, 12, 4115-23. | 3.5 | 68 |
| 7466 | The <i>Schizosaccharomyces pombe</i> cwg2+ gene codes for the beta subunit of a geranylgeranyltransferase type I required for beta-glucan synthesis. <i>EMBO Journal</i> , 1993, 12, 5245-54. | 3.5 | 30 |
| 7467 | Characterization of NGG1, a novel yeast gene required for glucose repression of GAL4p-regulated transcription. <i>EMBO Journal</i> , 1993, 12, 5255-65. | 3.5 | 29 |
| 7468 | A DNA unwinding element and an ARS consensus comprise a replication origin within a yeast chromosome. <i>EMBO Journal</i> , 1993, 12, 4521-31. | 3.5 | 55 |
| 7469 | Chromatin transitions during activation and repression of galactose-regulated genes in yeast. <i>EMBO Journal</i> , 1993, 12, 4603-13. | 3.5 | 53 |
| 7470 | Phospholipid transfer activity is relevant to but not sufficient for the essential function of the yeast SEC14 gene product. <i>EMBO Journal</i> , 1993, 12, 4775-84. | 3.5 | 48 |
| 7471 | <i>Drosophila</i> protein phosphatase V functionally complements a SIT4 mutant in <i>Saccharomyces cerevisiae</i> and its amino-terminal region can confer this complementation to a heterologous phosphatase catalytic domain. <i>EMBO Journal</i> , 1993, 12, 4833-42. | 3.5 | 20 |
| 7472 | Genetic analysis of glucose regulation in <i>saccharomyces cerevisiae</i> : control of transcription versus mRNA turnover. <i>EMBO Journal</i> , 1996, 15, 363-74. | 3.5 | 32 |
| 7473 | Reading two bases twice: mammalian antizyme frameshifting in yeast. <i>EMBO Journal</i> , 1996, 15, 1360-70. | 3.5 | 37 |

| # | ARTICLE | IF | CITATIONS |
|------|---|-----|-----------|
| 7474 | A suppressor of mutations in the class III transcription system encodes a component of yeast TFIIIB. EMBO Journal, 1996, 15, 1941-9. | 3.5 | 39 |
| 7475 | A novel Hsp70 of the yeast ER lumen is required for the efficient translocation of a number of protein precursors. EMBO Journal, 1996, 15, 2640-50. | 3.5 | 60 |
| 7476 | Utilizing the GCN4 leader region to investigate the role of the sequence determinants in nonsense-mediated mRNA decay. EMBO Journal, 1996, 15, 2810-9. | 3.5 | 28 |
| 7477 | Pip2p: a transcriptional regulator of peroxisome proliferation in the yeast <i>Saccharomyces cerevisiae</i> . EMBO Journal, 1996, 15, 2924-34. | 3.5 | 47 |
| 7478 | A pathway in the yeast cell division cycle linking protein kinase C (Pkc1) to activation of Cdc28 at START. EMBO Journal, 1996, 15, 3040-52. | 3.5 | 30 |
| 7479 | Fission yeast <i>cdc21</i> , a member of the MCM protein family, is required for onset of S phase and is located in the nucleus throughout the cell cycle. EMBO Journal, 1996, 15, 861-72. | 3.5 | 52 |
| 7480 | Copper-dependent degradation of the <i>Saccharomyces cerevisiae</i> plasma membrane copper transporter Ctr1p in the apparent absence of endocytosis. EMBO Journal, 1996, 15, 3515-23. | 3.5 | 88 |
| 7481 | Competition between folding and glycosylation in the endoplasmic reticulum. EMBO Journal, 1996, 15, 3538-46. | 3.5 | 19 |
| 7482 | The HeLa 200 kDa U5 snRNP-specific protein and its homologue in <i>Saccharomyces cerevisiae</i> are members of the DEXH-box protein family of putative RNA helicases. EMBO Journal, 1996, 15, 4001-15. | 3.5 | 68 |
| 7483 | Single point mutations in various domains of a plant plasma membrane H(+)-ATPase expressed in <i>Saccharomyces cerevisiae</i> increase H(+)-pumping and permit yeast growth at low pH. EMBO Journal, 1996, 15, 5513-26. | 3.5 | 51 |
| 7484 | Kes1p shares homology with human oxysterol binding protein and participates in a novel regulatory pathway for yeast Golgi-derived transport vesicle biogenesis. EMBO Journal, 1996, 15, 6447-59. | 3.5 | 93 |
| 7485 | Two GTPase isoforms, Ypt31p and Ypt32p, are essential for Golgi function in yeast. EMBO Journal, 1996, 15, 6460-75. | 3.5 | 81 |
| 7486 | The Exocyst is a multiprotein complex required for exocytosis in <i>Saccharomyces cerevisiae</i> . EMBO Journal, 1996, 15, 6483-94. | 3.5 | 404 |
| 7487 | Requirement for PP1 phosphatase and 20S cyclosome/APC for the onset of anaphase is lessened by the dosage increase of a novel gene <i>sds23+</i> . EMBO Journal, 1996, 15, 6629-40. | 3.5 | 33 |
| 7488 | Functional analysis of the interaction between the small GTP binding protein Cdc42 and the Ste20 protein kinase in yeast. EMBO Journal, 1996, 15, 7046-59. | 3.5 | 126 |
| 7489 | UBA 1: an essential yeast gene encoding ubiquitin-activating enzyme. EMBO Journal, 1991, 10, 227-36. | 3.5 | 100 |
| 7490 | Identification of regulatory residues of the yeast adenylyl cyclase. EMBO Journal, 1991, 10, 349-59. | 3.5 | 4 |
| 7491 | Identification of the structural gene for glucose-6-phosphate dehydrogenase in yeast. Inactivation leads to a nutritional requirement for organic sulfur. EMBO Journal, 1991, 10, 547-53. | 3.5 | 52 |

| # | ARTICLE | IF | CITATIONS |
|------|--|-----|-----------|
| 7492 | Proteinase yscE, the yeast proteasome/multicatalytic-multifunctional proteinase: mutants unravel its function in stress induced proteolysis and uncover its necessity for cell survival. EMBO Journal, 1991, 10, 555-62. | 3.5 | 143 |
| 7493 | Distinct cis-acting signals enhance 3' endpoint formation of CYC1 mRNA in the yeast <i>Saccharomyces cerevisiae</i> . EMBO Journal, 1991, 10, 563-71. | 3.5 | 81 |
| 7494 | The small nucleolar RNP protein NOP1 (fibrillarin) is required for pre-rRNA processing in yeast. EMBO Journal, 1991, 10, 573-83. | 3.5 | 202 |
| 7495 | Negative regulation of transcription of the <i>Saccharomyces cerevisiae</i> catalase T (CTT1) gene by cAMP is mediated by a positive control element. EMBO Journal, 1991, 10, 585-92. | 3.5 | 42 |
| 7496 | Transcription-induced nucleosome 'splitting': an underlying structure for DNase I sensitive chromatin. EMBO Journal, 1991, 10, 607-15. | 3.5 | 66 |
| 7497 | Mutational analysis of the putative effector domain of the GTP-binding Ypt1 protein in yeast suggests specific regulation by a novel GAP activity. EMBO Journal, 1991, 10, 785-92. | 3.5 | 32 |
| 7498 | Double-strand-break repair and recombination catalyzed by a nuclear extract of <i>Saccharomyces cerevisiae</i> . EMBO Journal, 1991, 10, 987-96. | 3.5 | 21 |
| 7499 | Exceptional codon recognition by the glutamine tRNAs in <i>Saccharomyces cerevisiae</i> . EMBO Journal, 1991, 10, 1481-91. | 3.5 | 15 |
| 7500 | Multiple Ca ²⁺ /calmodulin-dependent protein kinase genes in a unicellular eukaryote. EMBO Journal, 1991, 10, 1511-22. | 3.5 | 48 |
| 7501 | ABC1, a novel yeast nuclear gene has a dual function in mitochondria: it suppresses a cytochrome b mRNA translation defect and is essential for the electron transfer in the bc ₁ complex. EMBO Journal, 1991, 10, 2023-31. | 3.5 | 41 |
| 7502 | A specific DNA sequence is required for high frequency of recombination in the <i>ade6</i> gene of fission yeast. EMBO Journal, 1991, 10, 2157-63. | 3.5 | 83 |
| 7503 | The 3' to 5' exonuclease activity located in the DNA polymerase delta subunit of <i>Saccharomyces cerevisiae</i> is required for accurate replication. EMBO Journal, 1991, 10, 2165-70. | 3.5 | 113 |
| 7504 | Cloning of a yeast U1 snRNP 70K protein homologue: functional conservation of an RNA-binding domain between humans and yeast. EMBO Journal, 1991, 10, 2627-34. | 3.5 | 39 |
| 7505 | A novel switch-activating site (SAS1) and its cognate binding factor (SAP1) required for efficient mat1 switching in <i>Schizosaccharomyces pombe</i> . EMBO Journal, 1991, 10, 3025-32. | 3.5 | 73 |
| 7506 | Site-directed mutagenesis of the yeast actin gene: a test for actin function in vivo. EMBO Journal, 1991, 10, 3951-8. | 3.5 | 18 |
| 7507 | A genetic and structural analysis of the yeast Vps15 protein kinase: evidence for a direct role of Vps15p in vacuolar protein delivery. EMBO Journal, 1991, 10, 4049-60. | 3.5 | 45 |
| 7508 | 7SL RNA from <i>Schizosaccharomyces pombe</i> is encoded by a single copy essential gene. EMBO Journal, 1988, 7, 231-7. | 3.5 | 40 |
| 7509 | The NAM2 proteins from <i>S. cerevisiae</i> and <i>S. douglasii</i> are mitochondrial leucyl-tRNA synthetases, and are involved in mRNA splicing. EMBO Journal, 1988, 7, 473-83. | 3.5 | 86 |

| # | ARTICLE | IF | CITATIONS |
|------|---|-----|-----------|
| 7510 | Dual regulation of the expression of the polyubiquitin gene by cyclic AMP and heat shock in yeast. EMBO Journal, 1988, 7, 495-502. | 3.5 | 66 |
| 7511 | A yeast gene required for the G1-to-S transition encodes a protein containing an A-kinase target site and GTPase domain. EMBO Journal, 1988, 7, 1175-82. | 3.5 | 57 |
| 7512 | Mutation of lysine-48 to arginine in the yeast RAD3 protein abolishes its ATPase and DNA helicase activities but not the ability to bind ATP. EMBO Journal, 1988, 7, 3263-9. | 3.5 | 119 |
| 7513 | Yeast mutants temperature-sensitive for growth after random mutagenesis of the chromosomal RAS2 gene and deletion of the RAS1 gene. EMBO Journal, 1988, 7, 3375-83. | 3.5 | 15 |
| 7514 | The processing peptidase of yeast mitochondria: the two co-operating components MPP and PEP are structurally related. EMBO Journal, 1988, 7, 3493-500. | 3.5 | 61 |
| 7515 | Import of proteins into yeast mitochondria: the nuclear MAS2 gene encodes a component of the processing protease that is homologous to the MAS1-encoded subunit. EMBO Journal, 1988, 7, 3863-71. | 3.5 | 62 |
| 7516 | A novel nucleoskeletal-like protein located at the nuclear periphery is required for the life cycle of <i>Saccharomyces cerevisiae</i> . EMBO Journal, 1988, 7, 4323-34. | 3.5 | 84 |
| 7517 | The WHI1+ gene of <i>Saccharomyces cerevisiae</i> tethers cell division to cell size and is a cyclin homolog. EMBO Journal, 1988, 7, 4335-46. | 3.5 | 295 |
| 7518 | A U1 snRNA:pre-mRNA base pairing interaction is required early in yeast spliceosome assembly but does not uniquely define the 5' cleavage site. EMBO Journal, 1988, 7, 2533-8. | 3.5 | 202 |
| 7519 | Biogenesis of the yeast lysosome (vacuole): biosynthesis and maturation of proteinase yscB. EMBO Journal, 1988, 7, 1705-10. | 3.5 | 27 |
| 7520 | Dissection of functional domains of the yeast proton-pumping ATPase by directed mutagenesis. EMBO Journal, 1988, 7, 1793-8. | 3.5 | 31 |
| 7521 | Cold-sensitive and caffeine-supersensitive mutants of the <i>Schizosaccharomyces pombe</i> dis genes implicated in sister chromatid separation during mitosis. EMBO Journal, 1988, 7, 1465-73. | 3.5 | 113 |
| 7522 | Peroxisomal protein import is conserved between yeast, plants, insects and mammals. EMBO Journal, 1990, 9, 85-90. | 3.5 | 113 |
| 7523 | In vitro reconstitution of <i>cdc25</i> regulated <i>S. cerevisiae</i> adenylyl cyclase and its kinetic properties. EMBO Journal, 1990, 9, 641-51. | 3.5 | 18 |
| 7524 | Intermediates of recombination during mating type switching in <i>Saccharomyces cerevisiae</i> . EMBO Journal, 1990, 9, 663-73. | 3.5 | 252 |
| 7525 | A yeast tRNA precursor containing a pre-mRNA intron is spliced via the pre-mRNA splicing mechanism. EMBO Journal, 1990, 9, 705-9. | 3.5 | 7 |
| 7526 | Accumulation of pre-tRNA splicing '2/3' intermediates in a <i>Saccharomyces cerevisiae</i> mutant. EMBO Journal, 1990, 9, 1245-52. | 3.5 | 27 |
| 7527 | The <i>rhp6+</i> gene of <i>Schizosaccharomyces pombe</i> : a structural and functional homolog of the RAD6 gene from the distantly related yeast <i>Saccharomyces cerevisiae</i> . EMBO Journal, 1990, 9, 1423-30. | 3.5 | 37 |

| # | ARTICLE | IF | CITATIONS |
|------|--|-----|-----------|
| 7528 | The plant G box promoter sequence activates transcription in <i>Saccharomyces cerevisiae</i> and is bound in vitro by a yeast activity similar to GBF, the plant G box binding factor. <i>EMBO Journal</i> , 1990, 9, 1727-35. | 3.5 | 33 |
| 7529 | Effect of transcription of yeast chromatin on DNA topology in vivo. <i>EMBO Journal</i> , 1990, 9, 1873-81. | 3.5 | 19 |
| 7530 | Structural and functional analysis of <i>ypt2</i> , an essential ras-related gene in the fission yeast <i>Schizosaccharomyces pombe</i> encoding a Sec4 protein homologue. <i>EMBO Journal</i> , 1990, 9, 1957-62. | 3.5 | 36 |
| 7531 | An essential <i>Saccharomyces cerevisiae</i> single-stranded DNA binding protein is homologous to the large subunit of human RP-A. <i>EMBO Journal</i> , 1990, 9, 2321-9. | 3.5 | 93 |
| 7532 | Three GCN4 responsive elements act synergistically as upstream and as TATA-like elements in the yeast TRP4 promoter. <i>EMBO Journal</i> , 1990, 9, 2951-7. | 3.5 | 12 |
| 7533 | Primary structure, genomic organization and heterologous expression of a glucose transporter from <i>Arabidopsis thaliana</i> . <i>EMBO Journal</i> , 1990, 9, 3045-50. | 3.5 | 86 |
| 7534 | The recognition component of the N-end rule pathway. <i>EMBO Journal</i> , 1990, 9, 3179-89. | 3.5 | 166 |
| 7535 | The adenylate cyclase/protein kinase cascade regulates entry into meiosis in <i>Saccharomyces cerevisiae</i> through the gene <i>IME1</i> . <i>EMBO Journal</i> , 1990, 9, 3225-32. | 3.5 | 47 |
| 7536 | <i>Drosophila cdc2</i> homologs: a functional homolog is coexpressed with a cognate variant. <i>EMBO Journal</i> , 1990, 9, 3573-81. | 3.5 | 82 |
| 7537 | Signal sequence for generation of mRNA 3' end in the <i>Saccharomyces cerevisiae</i> <i>GAL7</i> gene. <i>EMBO Journal</i> , 1990, 9, 3691-7. | 3.5 | 43 |
| 7538 | <i>Saccharomyces cerevisiae</i> protein phosphatase 2A performs an essential cellular function and is encoded by two genes. <i>EMBO Journal</i> , 1990, 9, 4339-46. | 3.5 | 92 |
| 7539 | The <i>CHL 1</i> (CTF 1) gene product of <i>Saccharomyces cerevisiae</i> is important for chromosome transmission and normal cell cycle progression in G2/M. <i>EMBO Journal</i> , 1990, 9, 4347-58. | 3.5 | 92 |
| 7540 | Role of trans-activating proteins in the generation of active chromatin at the <i>PHO5</i> promoter in <i>S. cerevisiae</i> . <i>EMBO Journal</i> , 1990, 9, 2523-8. | 3.5 | 90 |
| 7541 | Regulation of a yeast <i>HSP70</i> gene by a cAMP responsive transcriptional control element. <i>EMBO Journal</i> , 1990, 9, 2543-53. | 3.5 | 47 |
| 7542 | Common signal transduction system shared by <i>STE2</i> and <i>STE3</i> in haploid cells of <i>Saccharomyces cerevisiae</i> : autocrine cell-cycle arrest results from forced expression of <i>STE2</i> . <i>EMBO Journal</i> , 1987, 6, 249-54. | 3.5 | 47 |
| 7543 | Independent mutations at the amino terminus of a protein act as surrogate signals for mitochondrial import. <i>EMBO Journal</i> , 1987, 6, 705-11. | 3.5 | 21 |
| 7544 | Three suppressor mutations which cure a mitochondrial RNA maturase deficiency occur at the same codon in the open reading frame of the nuclear <i>NAM2</i> gene. <i>EMBO Journal</i> , 1987, 6, 713-21. | 3.5 | 40 |
| 7545 | Efficient splicing of two yeast mitochondrial introns controlled by a nuclear-encoded maturase. <i>EMBO Journal</i> , 1987, 6, 1085-91. | 3.5 | 24 |

| # | ARTICLE | IF | CITATIONS |
|------|---|-----|-----------|
| 7546 | Nuclear pre-mRNA splicing in the fission yeast <i>Schizosaccharomyces pombe</i> strictly requires an intron-contained, conserved sequence element. <i>EMBO Journal</i> , 1987, 6, 1757-63. | 3.5 | 42 |
| 7547 | Transport of proteins to the mitochondrial intermembrane space: the 'matrix-targeting' and the 'sorting' domains in the cytochrome c1 presequence. <i>EMBO Journal</i> , 1987, 6, 2433-9. | 3.5 | 32 |
| 7548 | Specific accessory sequences in <i>Saccharomyces cerevisiae</i> introns control assembly of pre-mRNAs into spliceosomes. <i>EMBO Journal</i> , 1987, 6, 3833-9. | 3.5 | 29 |
| 7549 | Molecular cloning and characterization of the yeast RAD10 gene and expression of RAD10 protein in <i>E. coli</i> . <i>EMBO Journal</i> , 1985, 4, 1575-82. | 3.5 | 28 |
| 7550 | Role of conserved sequence elements in yeast centromere DNA. <i>EMBO Journal</i> , 1985, 4, 1867-74. | 3.5 | 87 |
| 7551 | The first twelve amino acids (less than half of the pre-sequence) of an imported mitochondrial protein can direct mouse cytosolic dihydrofolate reductase into the yeast mitochondrial matrix. <i>EMBO Journal</i> , 1985, 4, 2061-8. | 3.5 | 114 |
| 7552 | Nucleotide sequences of STE2 and STE3, cell type-specific sterile genes from <i>Saccharomyces cerevisiae</i> . <i>EMBO Journal</i> , 1985, 4, 2643-8. | 3.5 | 182 |
| 7553 | In vivo modulation of yeast tRNA gene expression by 5'-flanking sequences. <i>EMBO Journal</i> , 1985, 4, 2649-56. | 3.5 | 35 |
| 7554 | Structure, expression and regulation of a nuclear gene encoding a mitochondrial protein: the yeast L(+)-lactate cytochrome c oxidoreductase (cytochrome b2). <i>EMBO Journal</i> , 1985, 4, 3265-72. | 3.5 | 69 |
| 7555 | Yeast mRNA initiation sites are determined primarily by specific sequences, not by the distance from the TATA element. <i>EMBO Journal</i> , 1985, 4, 3273-80. | 3.5 | 103 |
| 7556 | The first twelve amino acids of a yeast mitochondrial outer membrane protein can direct a nuclear-coded cytochrome oxidase subunit to the mitochondrial inner membrane. <i>EMBO Journal</i> , 1985, 4, 3509-18. | 3.5 | 60 |
| 7557 | Import of proteins into mitochondria: a 70 kilodalton outer membrane protein with a large carboxy-terminal deletion is still transported to the outer membrane. <i>EMBO Journal</i> , 1983, 2, 2161-8. | 3.5 | 67 |
| 7558 | GAC1 may encode a regulatory subunit for protein phosphatase type 1 in <i>Saccharomyces cerevisiae</i> . <i>EMBO Journal</i> , 1992, 11, 87-96. | 3.5 | 73 |
| 7559 | Coordinate genetic control of yeast fatty acid synthase genes FAS1 and FAS2 by an upstream activation site common to genes involved in membrane lipid biosynthesis. <i>EMBO Journal</i> , 1992, 11, 107-14. | 3.5 | 56 |
| 7560 | Ubiquitin as a degradation signal. <i>EMBO Journal</i> , 1992, 11, 497-505. | 3.5 | 102 |
| 7561 | U14 function in <i>Saccharomyces cerevisiae</i> can be provided by large deletion variants of yeast U14 and hybrid mouse-yeast U14 RNAs. <i>EMBO Journal</i> , 1992, 11, 683-9. | 3.5 | 20 |
| 7562 | Species specific protein-DNA interactions may determine the chromatin units of genes in <i>S.cerevisiae</i> and in <i>S.pombe</i> . <i>EMBO Journal</i> , 1992, 11, 1177-85. | 3.5 | 14 |
| 7563 | Artificial nucleosome positioning sequences tested in yeast minichromosomes: a strong rotational setting is not sufficient to position nucleosomes in vivo. <i>EMBO Journal</i> , 1992, 11, 1187-93. | 3.5 | 11 |

| # | ARTICLE | IF | CITATIONS |
|------|--|-----|-----------|
| 7564 | Identification and functional analysis of two U3 binding sites on yeast pre-ribosomal RNA. EMBO Journal, 1992, 11, 1531-42. | 3.5 | 158 |
| 7565 | Cloning by functional complementation of a mouse cDNA encoding a homologue of CDC25, a <i>Saccharomyces cerevisiae</i> RAS activator. EMBO Journal, 1992, 11, 2151-7. | 3.5 | 86 |
| 7566 | OCH1 encodes a novel membrane bound mannosyltransferase: outer chain elongation of asparagine-linked oligosaccharides. EMBO Journal, 1992, 11, 2511-9. | 3.5 | 116 |
| 7567 | A 22 bp cis-acting element is necessary and sufficient for the induction of the yeast KAR2 (BiP) gene by unfolded proteins. EMBO Journal, 1992, 11, 2583-93. | 3.5 | 150 |
| 7568 | A role for clathrin in the sorting of vacuolar proteins in the Golgi complex of yeast. EMBO Journal, 1992, 11, 2811-8. | 3.5 | 87 |
| 7569 | A novel and conserved salt-induced protein is an important determinant of salt tolerance in yeast. EMBO Journal, 1992, 11, 3157-64. | 3.5 | 67 |
| 7570 | The <i>Saccharomyces cerevisiae</i> NPS1 gene, a novel CDC gene which encodes a 160 kDa nuclear protein involved in G2 phase control. EMBO Journal, 1992, 11, 4017-26. | 3.5 | 35 |
| 7571 | Two distinct yeast transcriptional activators require the function of the GCN5 protein to promote normal levels of transcription. EMBO Journal, 1992, 11, 4145-52. | 3.5 | 130 |
| 7572 | Dominant-negative mutants of a yeast G-protein beta subunit identify two functional regions involved in pheromone signalling. EMBO Journal, 1992, 11, 4805-13. | 3.5 | 40 |
| 7573 | The protein kinase homologue Ste20p is required to link the yeast pheromone response G-protein beta gamma subunits to downstream signalling components. EMBO Journal, 1992, 11, 4815-24. | 3.5 | 198 |
| 7574 | A new subclass of nucleoporins that functionally interact with nuclear pore protein NSP1. EMBO Journal, 1992, 11, 5051-61. | 3.5 | 98 |
| 7575 | Sequence-specific initiator elements focus initiation of transcription to distinct sites in the yeast TRP4 promoter. EMBO Journal, 1992, 11, 4583-90. | 3.5 | 10 |
| 7576 | Localization and sequence analysis of chloroplast DNA sequences of <i>Chlamydomonas reinhardtii</i> that promote autonomous replication in yeast. EMBO Journal, 1984, 3, 415-21. | 3.5 | 35 |
| 7577 | The expression of cDNA clones of yeast M1 double-stranded RNA in yeast confers both killer and immunity phenotypes. EMBO Journal, 1984, 3, 1383-7. | 3.5 | 22 |
| 7578 | Intracellular sorting of alcohol dehydrogenase isoenzymes in yeast: a cytosolic location reflects absence of an amino-terminal targeting sequence for the mitochondrion. EMBO Journal, 1986, 5, 161-5. | 3.5 | 24 |
| 7579 | Heme regulates the expression in <i>Saccharomyces cerevisiae</i> of chimaeric genes containing 5'-flanking soybean leghemoglobin sequences. EMBO Journal, 1986, 5, 843-7. | 3.5 | 8 |
| 7580 | Mitotic spindle pulls but fails to separate chromosomes in type II DNA topoisomerase mutants: uncoordinated mitosis. EMBO Journal, 1986, 5, 1003-10. | 3.5 | 112 |
| 7581 | Chromosome walking shows a highly homologous repetitive sequence present in all the centromere regions of fission yeast. EMBO Journal, 1986, 5, 1011-21. | 3.5 | 127 |

| # | ARTICLE | IF | CITATIONS |
|------|--|-----|-----------|
| 7582 | Yeast pre-messenger RNA splicing efficiency depends on critical spacing requirements between the branch point and 3' splice site. <i>EMBO Journal</i> , 1986, 5, 1023-30. | 3.5 | 27 |
| 7583 | Expression of murine and human granulocyte-macrophage colony-stimulating factors in <i>S. cerevisiae</i> : mutagenesis of the potential glycosylation sites. <i>EMBO Journal</i> , 1986, 5, 1193-7. | 3.5 | 26 |
| 7584 | The cleavable pre-sequence of an imported chloroplast protein directs attached polypeptides into yeast mitochondria. <i>EMBO Journal</i> , 1986, 5, 1343-50. | 3.5 | 56 |
| 7585 | Mutations in conserved intron sequences affect multiple steps in the yeast splicing pathway, particularly assembly of the spliceosome. <i>EMBO Journal</i> , 1986, 5, 1683-95. | 3.5 | 117 |
| 7586 | The nucleotide sequence of the fission yeast DNA topoisomerase II gene: structural and functional relationships to other DNA topoisomerases. <i>EMBO Journal</i> , 1986, 5, 2355-61. | 3.5 | 49 |
| 7587 | Molecular cloning and transcriptional analysis of the start gene <i>CDC25</i> of <i>Saccharomyces cerevisiae</i> . <i>EMBO Journal</i> , 1986, 5, 2363-2369. | 3.5 | 16 |
| 7588 | Isolation and characterization of <i>Schizosaccharomyces pombe</i> cutmutants that block nuclear division but not cytokinesis. <i>EMBO Journal</i> , 1986, 5, 2973-9. | 3.5 | 148 |
| 7589 | At least two nuclear gene products are specifically required for translation of a single yeast mitochondrial mRNA. <i>EMBO Journal</i> , 1986, 5, 3637-41. | 3.5 | 73 |
| 7590 | Suppression of defective <i>RAS1</i> and <i>RAS2</i> functions in yeast by an adenylate cyclase activated by a single amino acid change. <i>EMBO Journal</i> , 1986, 5, 3657-63. | 3.5 | 45 |
| 7671 | Extrachromosomal recombination is deranged in the <i>rec2</i> mutant of <i>Ustilago maydis</i> . <i>Genetics</i> , 1991, 129, 1052-60. | 1.2 | 12 |
| 7771 | Functional production and reconstitution of the human equilibrative nucleoside transporter (hENT1) in <i>Saccharomyces cerevisiae</i> . Interaction of inhibitors of nucleoside transport with recombinant hENT1 and a glycosylation-defective derivative (hENT1/N48Q). <i>Biochemical Journal</i> , 1999, 339 (Pt 1), 21-32. | 1.7 | 27 |
| 7772 | Molecular and enzymic properties of recombinant 1, 2-alpha-mannosidase from <i>Aspergillus saitoi</i> overexpressed in <i>Aspergillus oryzae</i> cells. <i>Biochemical Journal</i> , 1999, 339 (Pt 3), 589-97. | 1.7 | 8 |
| 7773 | <i>CLN3</i> expression is sufficient to restore G1-to-S-phase progression in <i>Saccharomyces cerevisiae</i> mutants defective in translation initiation factor eIF4E. <i>Biochemical Journal</i> , 1999, 340 (Pt 1), 135-41. | 1.7 | 26 |
| 7774 | Identification of copper ligands in <i>Aspergillus oryzae</i> tyrosinase by site-directed mutagenesis. <i>Biochemical Journal</i> , 2000, 350 Pt 2, 537-45. | 1.7 | 10 |
| 7775 | Definition of a minimal <i>munc18c</i> domain that interacts with syntaxin 4. <i>Biochemical Journal</i> , 2000, 350 Pt 3, 741-6. | 1.7 | 2 |
| 7776 | A carbon-source-responsive element is required for regulation of the hypoxic ADP/ATP carrier (AAC3) isoform in <i>Saccharomyces cerevisiae</i> . <i>Biochemical Journal</i> , 2000, 352 Pt 3, 893-8. | 1.7 | 2 |
| 7777 | A translational fidelity mutation in the universally conserved sarcin/ricin domain of 25S yeast ribosomal RNA. <i>Rna</i> , 1996, 2, 254-63. | 1.6 | 35 |
| 7778 | Effect of frameshift-inducing mutants of elongation factor 1alpha on programmed +1 frameshifting in yeast. <i>Rna</i> , 1998, 4, 38-46. | 1.6 | 16 |

| # | ARTICLE | IF | CITATIONS |
|------|--|-----|-----------|
| 7779 | Genetic interaction between U6 snRNA and the first intron nucleotide in <i>Saccharomyces cerevisiae</i> . <i>Rna</i> , 1998, 4, 167-80. | 1.6 | 24 |
| 7780 | Construction of an in vivo-regulated U6 snRNA transcription unit as a tool to study U6 function. <i>Rna</i> , 1998, 4, 231-8. | 1.6 | 9 |
| 7781 | U3 snoRNA genes with and without intron in the <i>Kluyveromyces</i> genus: yeasts can accommodate great variations of the U3 snoRNA 3'-terminal domain. <i>Rna</i> , 1998, 4, 285-302. | 1.6 | 6 |
| 7782 | Transcription termination downstream of the <i>Saccharomyces cerevisiae</i> FBP1 [changed from FPB1] poly(A) site does not depend on efficient 3'end processing. <i>Rna</i> , 1998, 4, 303-18. | 1.6 | 19 |
| 7783 | A comprehensive biochemical and genetic analysis of the yeast U1 snRNP reveals five novel proteins. <i>Rna</i> , 1998, 4, 374-93. | 1.6 | 130 |
| 7832 | The final stages of spliceosome maturation require Spp2p that can interact with the DEAH box protein Prp2p and promote step 1 of splicing. <i>Rna</i> , 1995, 1, 375-90. | 1.6 | 78 |
| 7833 | Studies in the development of Japanese encephalitis vaccine: expression of virus envelope glycoprotein V3 (E) gene in yeast. <i>Bulletin of the World Health Organization</i> , 1987, 65, 303-8. | 1.5 | 7 |
| 7836 | CTD kinase large subunit is encoded by CTK1, a gene required for normal growth of <i>Saccharomyces cerevisiae</i> . <i>Gene Expression</i> , 1991, 1, 149-67. | 0.5 | 153 |
| 7837 | Identification of novel steroid-response elements. <i>Gene Expression</i> , 1992, 2, 39-47. | 0.5 | 12 |
| 7838 | Interferon regulatory factor-1 (IRF-1) activates the synthetic IRF-1-responsive sequence (GAAAGT) ₄ in <i>Saccharomyces cerevisiae</i> . <i>Gene Expression</i> , 1992, 2, 329-37. | 0.5 | 7 |
| 7839 | The MRS1 gene of <i>S. douglasii</i> : co-evolution of mitochondrial introns and specific splicing proteins encoded by nuclear genes. <i>Gene Expression</i> , 1992, 2, 203-14. | 0.5 | 23 |
| 7840 | Differential modes of activation define orphan subclasses within the steroid/thyroid receptor superfamily. <i>Gene Expression</i> , 1992, 2, 273-83. | 0.5 | 27 |
| 7841 | Role of <i>Saccharomyces cerevisiae</i> Rap1 protein in Ty1 and Ty1-mediated transcription. <i>Gene Expression</i> , 1993, 3, 237-51. | 0.5 | 16 |
| 7842 | Mcm10 and the MCM2-7 complex interact to initiate DNA synthesis and to release replication factors from origins. <i>Genes and Development</i> , 2000, 14, 913-926. | 2.7 | 163 |
| 7843 | Microbial Efflux Pump Inhibitors: A Journey around Quinoline and Indole Derivatives. <i>Molecules</i> , 2021, 26, 6996. | 1.7 | 14 |
| 7844 | Artemisinin Binds and Inhibits the Activity of <i>Plasmodium falciparum</i> Ddi1, a Retroviral Aspartyl Protease. <i>Pathogens</i> , 2021, 10, 1465. | 1.2 | 5 |
| 7846 | Possible Role for Allelic Variation in Yeast MED15 in Ecological Adaptation. <i>Frontiers in Microbiology</i> , 2021, 12, 741572. | 1.5 | 1 |
| 7847 | Slt2 Is Required to Activate ER-Stress-Protective Mechanisms through TORC1 Inhibition and Hexosamine Pathway Activation. <i>Journal of Fungi (Basel, Switzerland)</i> , 2022, 8, 92. | 1.5 | 8 |

| # | ARTICLE | IF | CITATIONS |
|------|--|-----|-----------|
| 7848 | Fermentation Efficiency of Genetically Modified Yeasts in Grapes Must. <i>Foods</i> , 2022, 11, 413. | 1.9 | 1 |
| 7849 | <i>Drosophila melanogaster</i> Uncoupling Protein-4A (UCP4A) Catalyzes a Unidirectional Transport of Aspartate. <i>International Journal of Molecular Sciences</i> , 2022, 23, 1020. | 1.8 | 12 |
| 7850 | Mutant phosphatidate phosphatase Pah1-W637A exhibits altered phosphorylation, membrane association, and enzyme function in yeast. <i>Journal of Biological Chemistry</i> , 2022, 298, 101578. | 1.6 | 6 |
| 7851 | Characterization of Arabidopsis Post-Glycosylphosphatidylinositol Attachment to Proteins Phospholipase 3 Like Genes. <i>Frontiers in Plant Science</i> , 2022, 13, 817915. | 1.7 | 1 |
| 7852 | Proteasome activity modulates amyloid toxicity. <i>FEMS Yeast Research</i> , 2022, 22, . | 1.1 | 1 |
| 7853 | Complete coding sequence of wheat phosphoribulokinase: developmental and light-dependent expression of the mRNA. <i>Molecular Genetics and Genomics</i> , 1989, 220, 43-8. | 2.4 | 18 |
| 7854 | A novel promoter, derived from the isocitrate lyase gene of <i>Candida tropicalis</i> , inducible with acetate in <i>Saccharomyces cerevisiae</i> . <i>Applied Microbiology and Biotechnology</i> , 1995, 43, 489-92. | 1.7 | 7 |
| 7855 | FOG1 and FOG2 genes, required for the transcriptional activation of glucose-repressible genes of <i>Kluyveromyces lactis</i> , are homologous to GAL83 and SNF1 of <i>saccharomyces cerevisiae</i> . <i>Current Genetics</i> , 1996, 29, 316-26. | 0.8 | 35 |
| 7856 | Mutational analysis of the beta-subunit of yeast geranylgeranyl transferase I. <i>Molecular Genetics and Genomics</i> , 1996, 252, 1-10. | 2.4 | 5 |
| 7857 | Identification of proteins that interact with a protein of interest: applications of the yeast two-hybrid system. <i>Molecular and Cellular Biochemistry</i> , 1997, 172, 67-79. | 1.4 | 48 |
| 7858 | BTF3 is a potential new substrate of protein kinase CK2. <i>Molecular and Cellular Biochemistry</i> , 1999, 191, 121-8. | 1.4 | 7 |
| 7859 | Two alpha isopropylmalate synthase isozymes with similar kinetic properties are extant in the yeast <i>Lachancea kluyveri</i> . <i>FEMS Yeast Research</i> , 2022, 22, . | 1.1 | 2 |
| 7860 | BrpNAC895 and BrpABI449 coregulate the transcription of the efflux-type cadmium transporter <i>BrpHMA2</i> in <i>Brassica parachinensis</i> . <i>Horticulture Research</i> , 2022, 9, . | 2.9 | 11 |
| 7861 | Cleavage-defective Topoisomerase I mutants sharply increase G-quadruplex-associated genomic instability. <i>Microbial Cell</i> , 2022, 9, 52-68. | 1.4 | 3 |
| 7862 | The toxic effects of yeast Ppz1 phosphatase are counteracted by subcellular relocalization mediated by its regulatory subunit Hal3. <i>FEBS Letters</i> , 2022, 596, 1556-1566. | 1.3 | 5 |
| 7863 | Golgin Imh1 and GARP complex cooperate to restore the impaired SNARE recycling transport induced by ER stress. <i>Cell Reports</i> , 2022, 38, 110488. | 2.9 | 5 |
| 7864 | Neurons undergo pathogenic metabolic reprogramming in models of familial ALS. <i>Molecular Metabolism</i> , 2022, 60, 101468. | 3.0 | 6 |
| 7865 | Disruption of the Snf1 Gene Enhances Cell Growth and Reduces the Metabolic Burden in Cellulase-Expressing and Lipid-Accumulating <i>Yarrowia lipolytica</i> . <i>Frontiers in Microbiology</i> , 2021, 12, 757741. | 1.5 | 6 |

| # | ARTICLE | IF | CITATIONS |
|------|---|-----|-----------|
| 7866 | Effects of abolishing Whi2 on the proteome and nitrogen catabolite repression-sensitive protein production. <i>G3: Genes, Genomes, Genetics</i> , 2022, 12, . | 0.8 | 0 |
| 7867 | Weakening the IF2-fMet-tRNA Interaction Suppresses the Lethal Phenotype Caused by GTPase Inactivation. <i>International Journal of Molecular Sciences</i> , 2021, 22, 13238. | 1.8 | 1 |
| 7868 | PUFA synthase-independent DHA synthesis pathway in <i>Parietichytrium</i> sp. and its modification to produce EPA and n-3DPA. <i>Communications Biology</i> , 2021, 4, 1378. | 2.0 | 12 |
| 7869 | Utilizing chemically induced dimerization of FKBP to analyze endocytosis by live-cell imaging in budding yeast. <i>STAR Protocols</i> , 2022, 3, 101323. | 0.5 | 1 |
| 7887 | Improvement of S-adenosylmethionine production by integration of the ethionine-resistance gene into chromosomes of the yeast <i>Saccharomyces cerevisiae</i> . <i>Applied Microbiology and Biotechnology</i> , 1995, 42, 730-733. | 1.7 | 2 |
| 7888 | Generation of a stable non-reverting Leu ⁻ mutant of <i>Kluyveromyces lactis</i> by gene disruption. <i>Applied Microbiology and Biotechnology</i> , 1995, 42, 734-737. | 1.7 | 0 |
| 7889 | Secretion by <i>Saccharomyces cerevisiae</i> of rat apolipoprotein E as a fusion to <i>Mucor</i> rennin. <i>Applied Microbiology and Biotechnology</i> , 1995, 42, 865-870. | 1.7 | 0 |
| 7890 | CEN14 sequences cause slower proliferation, reduced cell size and asporogeny in <i>Saccharomyces cerevisiae</i> . <i>Applied Microbiology and Biotechnology</i> , 1995, 43, 877-879. | 1.7 | 0 |
| 7891 | Cloning of <i>Corticium rolfii</i> glucoamylase cDNA and its expression in <i>Saccharomyces cerevisiae</i> . <i>Applied Microbiology and Biotechnology</i> , 1995, 44, 451-458. | 1.7 | 1 |
| 7892 | Application of Ty1 for cloned gene insertion: amplification of a large regulated expression cassette in <i>Saccharomyces cerevisiae</i> . <i>Applied Microbiology and Biotechnology</i> , 1996, 44, 620-623. | 1.7 | 0 |
| 7893 | Detection of antagonistic cellular regulatory functions by the gene-gene interference method in yeast. <i>Current Genetics</i> , 1996, 29, 114-121. | 0.8 | 0 |
| 7894 | Construction of the complete rat fatty acid synthase cDNA and its expression in <i>Saccharomyces cerevisiae</i> . <i>Current Genetics</i> , 1996, 29, 219-226. | 0.8 | 0 |
| 7895 | Immunogold labeling of yeast cells: an efficient tool for the study of protein targeting and morphological alterations due to overexpression and inactivation of genes. <i>Histochemistry and Cell Biology</i> , 1996, 106, 115-130. | 0.8 | 3 |
| 7896 | HAM1, the gene controlling 6-N-hydroxylaminopurine sensitivity and mutagenesis in the yeast <i>Saccharomyces cerevisiae</i> . <i>Yeast</i> , 1996, 12, 17-29. | 0.8 | 32 |
| 7897 | Separate roles for N- and C-termini of the STE4 (beta) subunit of the <i>Saccharomyces cerevisiae</i> G protein in the mediation of the growth arrest. Lack of growth-arresting activity of mammalian beta gamma complexes. <i>Yeast</i> , 1996, 12, 41-51. | 0.8 | 0 |
| 7898 | Cloning of <i>Candida albicans</i> SEC14 gene homologue coding for a putative essential function. <i>Yeast</i> , 1996, 12, 1097-105. | 0.8 | 22 |
| 7899 | Genetic aspects of carbon catabolite repression of the STA2 glucoamylase gene in <i>Saccharomyces cerevisiae</i> . <i>Yeast</i> , 1996, 12, 1297-300. | 0.8 | 14 |
| 7900 | Phosphorylation of glucosamine-6-phosphate synthase is important but not essential for germination and mycelial growth of <i>Candida albicans</i> . <i>FEMS Microbiology Letters</i> , 2004, 235, 73-80. | 0.7 | 8 |

| # | ARTICLE | IF | CITATIONS |
|------|---|-----|-----------|
| 7901 | Simultaneous Display of Multiple Kinds of Enzymes on the Yeast Cell Surface for Multistep Reactions. <i>Methods in Molecular Biology</i> , 2022, 2491, 627-641. | 0.4 | 4 |
| 7902 | Lipofection mediated transfection fails for sea urchin coelomocytes. <i>PLoS ONE</i> , 2022, 17, e0267911. | 1.1 | 2 |
| 7903 | Differential translation elongation directs protein synthesis in response to acute glucose deprivation in yeast. <i>RNA Biology</i> , 2022, 19, 636-649. | 1.5 | 5 |
| 7904 | Heterologous fusion gene expression and characterization of a novel carbohydrate binding module (Cbm36) to laccase (Lcc2). <i>Biocatalysis and Agricultural Biotechnology</i> , 2022, 42, 102377. | 1.5 | 5 |
| 7905 | Host-specific activation of a pathogen effector Aave_4606 from <i>Acidovorax citrulli</i> , the causal agent for bacterial fruit blotch. <i>Biochemical and Biophysical Research Communications</i> , 2022, 616, 41-48. | 1.0 | 2 |
| 7912 | Genetic evidence for functional specificity of the yeast GCN2 kinase. <i>Molecular Genetics and Genomics</i> , 1996, 251, 613. | 2.4 | 0 |
| 7916 | Generation of Arming Yeasts with Active Proteins and Peptides via Cell Surface Display System: Cell Surface Engineering, <i>Bio-Arming Technology. Methods in Molecular Biology</i> , 2022, , 59-77. | 0.4 | 1 |
| 7918 | Cloning and screening of the putative hexokinase genes from <i>Rhizopus oryzae</i> and their heterologous expression in <i>Saccharomyces cerevisiae</i> . <i>Molecular Biology Reports</i> , 0, , . | 1.0 | 1 |
| 7919 | Construction of HGF-Displaying Yeast by Cell Surface Engineering. <i>Microorganisms</i> , 2022, 10, 1373. | 1.6 | 0 |
| 7920 | Glycogen synthase kinase homolog Rim11 regulates lipid synthesis through the phosphorylation of Pah1 phosphatidate phosphatase in yeast. <i>Journal of Biological Chemistry</i> , 2022, 298, 102221. | 1.6 | 9 |
| 7921 | Transcriptional activation of budding yeast DDI2/3 through chemical modifications of Fzf1. <i>Cell Biology and Toxicology</i> , 0, , . | 2.4 | 0 |
| 7922 | Mnt1, an α -(1 \rightarrow 2)-mannosyltransferase responsible for the elongation of N-glycans and O-glycans in <i>Aspergillus fumigatus</i> . <i>Glycobiology</i> , 2022, 32, 1137-1152. | 1.3 | 5 |
| 7924 | Metabolic engineering of the l-serine biosynthetic pathway improves glutathione production in <i>Saccharomyces cerevisiae</i> . <i>Microbial Cell Factories</i> , 2022, 21, . | 1.9 | 2 |
| 7925 | Glutathione degradation activity of γ -glutamyl peptidase 1 manifests its dual roles in primary and secondary sulfur metabolism in <i>Arabidopsis</i> . <i>Plant Journal</i> , 2022, 111, 1626-1642. | 2.8 | 10 |
| 7926 | Recent advances in genetic technology development of oleaginous yeasts. <i>Applied Microbiology and Biotechnology</i> , 2022, 106, 5385-5397. | 1.7 | 4 |
| 7928 | Heterologous (Over) Expression of Human Solute Carrier (SLC) in Yeast: A Well-Recognized Tool for Human Transporter Function/Structure Studies. <i>Life</i> , 2022, 12, 1206. | 1.1 | 1 |
| 7929 | Microbial electrosynthesis from carbon dioxide feedstock linked to yeast growth for the production of high-value isoprenoids. <i>Bioresource Technology</i> , 2022, 363, 127906. | 4.8 | 5 |
| 7930 | Rpd3 regulates single-copy origins independently of the rDNA array by opposing Fkh1-mediated origin stimulation. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 2022, 119, . | 3.3 | 9 |

| # | ARTICLE | IF | CITATIONS |
|------|--|-----|-----------|
| 7931 | Neomycin Interferes with Phosphatidylinositol-4,5-Bisphosphate at the Yeast Plasma Membrane and Activates the Cell Wall Integrity Pathway. <i>International Journal of Molecular Sciences</i> , 2022, 23, 11034. | 1.8 | 1 |
| 7932 | Sir2 and Reb1 antagonistically regulate nucleosome occupancy in subtelomeric X-elements and repress TERRAs by distinct mechanisms. <i>PLoS Genetics</i> , 2022, 18, e1010419. | 1.5 | 2 |
| 7933 | Galactaric acid production by engineering substrate specificity in glucose oxidase from <i>Aspergillus niger</i> . <i>Biochemical Engineering Journal</i> , 2022, 187, 108646. | 1.8 | 1 |
| 7934 | Trehalose 6-Phosphate/SnRK1 Signaling Participates in Harvesting-Stimulated Rubber Production in the Hevea Tree. <i>Plants</i> , 2022, 11, 2879. | 1.6 | 1 |
| 7935 | Role of pepper MYB transcription factor CaDIM1 in regulation of the drought response. <i>Frontiers in Plant Science</i> , 0, 13, . | 1.7 | 3 |
| 7936 | Genetic analysis of DNA-damage tolerance pathways in <i>Arabidopsis</i> . <i>Plant Cell Reports</i> , 0, , . | 2.8 | 0 |
| 7937 | The caleosin CLO7 and its role in the heterotrimeric G-protein signalling network. <i>Journal of Plant Physiology</i> , 2022, 279, 153841. | 1.6 | 2 |
| 7938 | An ERAD-independent role for rhomboid pseudoprotease Dfm1 in mediating sphingolipid homeostasis. <i>EMBO Journal</i> , 2023, 42, . | 3.5 | 5 |
| 7939 | Dbf4 Zn-Finger Motif Is Specifically Required for Stimulation of Ctf19-Activated Origins in <i>Saccharomyces cerevisiae</i> . <i>Genes</i> , 2022, 13, 2202. | 1.0 | 0 |
| 7940 | The human pathobiont <i>Malassezia furfur</i> secreted protease Mfsap1 regulates cell dispersal and exacerbates skin inflammation. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 2022, 119, . | 3.3 | 12 |
| 7942 | The ChaC1 active site: Defining the residues and determining the role of ChaC1-exclusive residues in the structural and functional stability. <i>Proteins: Structure, Function and Bioinformatics</i> , 2023, 91, 567-580. | 1.5 | 1 |
| 7943 | Evidence That the Igl ^h Gene MAR Regulates the Probability of Premature V-J Joining and Somatic Hypermutation. <i>Journal of Immunology</i> , 1999, 162, 6029-6039. | 0.4 | 25 |
| 7945 | Biocatalyst collection and heterologous expression of sesquiterpene synthases from basidiomycetous fungi: Discovery of a novel sesquiterpene hydrocarbon. <i>Microbial Biotechnology</i> , 2023, 16, 632-644. | 2.0 | 5 |
| 7946 | Construction of Constitutive <i>Saccharomyces Cerevisiae</i> Engineered Strain of Î ² -glucosidase. <i>Lecture Notes in Computer Science</i> , 2023, , 190-208. | 1.0 | 0 |
| 7947 | Haem transporter HRG-1 is essential in the barber's pole worm and an intervention target candidate. <i>PLoS Pathogens</i> , 2023, 19, e1011129. | 2.1 | 5 |
| 7948 | Conserved regions of the regulatory subunit Spo7 are required for Nem1-Spo7/Pah1 phosphatase cascade function in yeast lipid synthesis. <i>Journal of Biological Chemistry</i> , 2023, 299, 104683. | 1.6 | 3 |
| 7949 | Pepper clade A PP2C, CaSIP1, negatively modulates drought resistance by suppressing CaSnRK2.6 kinase activity. <i>Environmental and Experimental Botany</i> , 2023, 209, 105275. | 2.0 | 1 |
| 7950 | Changed life course upon defective replication of ribosomal RNA genes. <i>Genes and Genetic Systems</i> , 2022, 97, 285-295. | 0.2 | 3 |

| # | ARTICLE | IF | CITATIONS |
|------|---|-----|-----------|
| 7951 | The ENA1 Na ⁺ -ATPase Gene Is Regulated by the SPS Sensing Pathway and the Stp1/Stp2 Transcription Factors. <i>International Journal of Molecular Sciences</i> , 2023, 24, 5548. | 1.8 | 1 |
| 7953 | Multiple intermolecular interactions facilitate rapid evolution of essential genes. <i>Nature Ecology and Evolution</i> , 2023, 7, 745-755. | 3.4 | 2 |
| 7960 | Mitogen-activated protein kinase (MAPK) cascadesâ€”A yeast perspective. <i>The Enzymes</i> , 2023, , 137-170. | 0.7 | 0 |
| 7970 | Proteinâ€™Protein Interactions: Yeast Two Hybrid. <i>Methods in Molecular Biology</i> , 2024, , 235-246. | 0.4 | 0 |